

Potential for analytic breeding in allopolyploids: an illustration from Longiflorum × Asiatic hybrid lilies (*Lilium*)

Nadeem Khan · Shujun Zhou · M. S. Ramanna ·
Paul Arens · Jeronimo Herrera ·
Richard G. F. Visser · Jaap M. Van Tuyl

Received: 2 June 2008 / Accepted: 18 September 2008 / Published online: 10 October 2008
© The Author(s) 2008. This article is published with open access at Springerlink.com

Abstract Ploidy level and intergenomic recombination was studied in interspecific hybrids between Longiflorum × Asiatic lilies (LA hybrid) backcross to Asiatic parents in order to assess the possibility for analytic breeding in lily. By backcrossing the diploid ($2n = 2x = 24$) F1 interspecific hybrid between Longiflorum × Asiatic lilies to Asiatic parents, 104 BC1 progeny plants were produced. Among these, there were 27 diploids, 73 triploids ($2n = 2x = 36$) and 4 aneuploids ($2x - 1$, $2x + 2$ or $2x + 3$). In addition, by backcrossing triploid BC1 (LAA) plants to diploid Asiatic parents in $2x - 3x$ and reciprocal combinations, 14 diploid BC2 progenies were produced. Genomic in situ hybridization (GISH) was performed to study the intergenomic recombination and karyotype composition. GISH indicated extensive intergenomic recombination among the chromosomes in LA hybrids. A large number of Longiflorum chromosomes were transmitted to the BC1 progenies from LA hybrids. However, very few Longiflorum chromosomes were transmitted from the BC1 triploid

(LAA) plants to the BC2 progenies. The occurrence of diploid plants in the BC progenies of LA hybrids has opened the prospects of analytic breeding in lilies. In this approach, the selection of superior genotypes can be carried out at the diploid level and polyploid forms are synthesized from superior diploid parents. The advantages of analytic breeding are evident: (a) a maximum level of heterozygosity can be attained in the synthetic polyploids and (b) introgression can be achieved with a minimum of linkage drag. Based on GISH results the potential application of analytic breeding in lily allopolyploids has been discussed.

Keywords Back cross progenies · Haploid gametes · GISH · Intergenomic recombination · Lily · Polyploids

Introduction

Generally, it is difficult to breed polyploid crops as compared to diploid forms. This is true for both auto- and allopolyploids for different reasons. Autopolyploids display polysomic inheritance, the segregation of characters is not clear cut and this complicates the process of selection for desirable combinations of traits (Peloquin 1981). On the other hand, allopolyploids which are also called functional diploids, behave as “permanent hybrids” because of the preferential chromosome pairing between the parental genomes during meiosis (Barba-Gonzalez et al. 2005; Feldman

N. Khan (✉) · M. S. Ramanna · P. Arens · J. Herrera ·
R. G. F. Visser · J. M. Van Tuyl
Plant Breeding, Wageningen University and Research,
P.O. Box 16, 6700AA Wageningen, The Netherlands
e-mail: nadeem.khan@wur.nl; khan_m_nadeem@yahoo.com

S. Zhou
Department of Horticulture, Agricultural and
Technological College, Zhejiang University,
No. 258 KaiXuan Road, Hangzhou City,
Zhejiang Province 310029, China

and Levy 2005). As a result, hardly any genetic variation occurs in the progenies of allopolyploids due to fixed heterozygosity and this is not favourable for the process of selection. In view of these differences, auto- and allopolyploids require different breeding approaches.

For autopolyploids, Chase (1963) proposed a novel approach, the so called “analytic breeding”, which was applied in the case of cultivated potato, *Solanum tuberosum* L. (Mendiburu et al. 1974). This approach involves the reduction of tetraploids to diploid level followed by intensive selection of superior genotypes through breeding at this level and in the final step tetraploid cultivars are produced (through sexual polyploidization) from the selected diploid parents (Jacobsen et al. 1991). Many practical advantages of breeding at the diploid level, such as relatively simple Mendelian segregation of characters and the requirement of smaller size of population for selection, have been highlighted along with other theoretical considerations (Bingham et al. 1994). Analytic breeding has also been relevant in other crops like alfalfa (Mariani and Tavoletti 1992; Barcaccia et al. 2000) and cranberry (Lyrene et al. 2003; Zeldin and McCown 2004). In view of the attractiveness of analytic breeding in autopolyploids, the question arises whether such an approach can also be relevant for breeding allopolyploids.

Breeding at the diploid level implies that the genotypes used in this approach necessarily produce haploid (n) gametes so that they give rise to diploids in the subsequent generations (Zeldin and McCown 2004). In the hybrids of species that possess very closely related or similar genomes (the case for autopolyploids) it is not a problem to have diploid forms that can produce functional n gametes due to normal segregation of chromosomes during meiosis (Weiss and Maluszynska 2000; Lopez-Lavalle and Orjeda 2002; Gu et al. 2005; Hoya et al. 2007; Zhang et al. 2007). On the contrary, in interspecific hybrids of distantly related species (allopolyploid types) it is very difficult to have genotypes that can produce normal n gametes. This is because such diploid hybrids are mostly sterile due to a disturbed segregation of chromosomes during meiosis (Estilai 1977; Van Tuyl and De Jeu 1997; Lim et al. 2000; Barba-Gonzalez et al. 2006b). However, if diploid (distant) interspecific hybrids that can produce both n as well as $2n$ gametes become available, there might be a potential

for analytic breeding even in the case of allopolyploids. Once the selection has been made at the diploid level, then the unilateral sexual polyploids can be achieved by using $2n$ gametes to reach the optimum ploidy level, e.g., triploid in lilies (Van Tuyl and Lim 2003; Zhou et al. 2008). Hence, the analytic breeding scheme might be possible when genotypes producing both n and $2n$ gametes are available. The cultivated lilies appear to possess such a potential. There are three important groups of cultivated lilies (*Lilium*), viz., Longiflorum, Asiatic and Oriental, which were predominantly diploid ($2n = 2x = 24$) until recently (Lim et al. 2000). During the last few decades, polyploids have been replacing the diploid forms due to their superior characteristics of the former. These polyploid cultivars have originated from hybridization of cultivars that belong to different taxonomic sections of lilies such as Longiflorum \times Asiatic (LA) [Lim et al. 2001; Zhou et al. 2008] or Oriental \times Asiatic (OA) [Barba-Gonzalez et al. 2004; 2005]. Because the three groups possess clearly differentiated genomes, their hybrids have given rise to typical allotriploid (ALA and AOA) or allotetraploid (LLAA) cultivars (Barba-Gonzalez et al. 2005; Zhou 2007; Zhou et al. 2008). These polyploids are difficult to use as parents in breeding because the triploids are mostly sterile and the allotetraploids are permanent hybrids as was mentioned earlier. A salient feature of some of the diploid LA hybrids was that they produced not only $2n$ gametes but also n gametes (Zhou 2007). This provides unique opportunities for generating allotriploid ($2n = 3x = 36$) as well as diploid ($2n = 2x = 24$) BC1 progenies from backcrossing LA hybrids to Asiatic parents. In addition, diploid BC2 progenies were also generated from backcrossing allotriploid (ALA or LAA) to diploid Asiatic parents, i.e., $3x - 2x$ or reciprocal crosses. These initial attempts indicated that it might be possible to generate fairly large numbers of diploid BC1 and BC2 progenies from LA hybrids and their backcross progenies. With the aim of evaluating the prospects of using diploid backcross progenies in analytic breeding, quite a number of diploid BC progenies from LA hybrids were produced and the genome composition of the diploid BC1 and BC2 progenies were analysed by GISH. Based on these results the potential application and significance of these hybrids is discussed with particular reference to analytic breeding in allopolyploids lilies.

Materials and methods

Plant material

Most of the diploid F1 Longiflorum × Asiatic hybrids ($2n = 2x = 24$), and allotriploids i.e., LAA and ALA ($2n = 3x = 36$) cultivars were supplied by Dutch lily breeding companies: De Jong Lilies BV, Royal Van Zanten BV, Testcentrum BV, Vletter and Den Haan BV and World Breeding BV. One F1 LA hybrids i.e., 024004-5 was developed at Plant Breeding, Wageningen UR. As the two groups which belong to two different taxonomic sections, the cultivars of different sections could be hybridized, or backcrossed, only through special techniques (Van Tuyl et al. 1991; Lim et al. 2001; Barba-Gonzalez et al. 2004). For backcrossing, the F1 LA hybrids were used both as female and male parent and crossed with eight different Asiatic parents to get LAA or ALA progenies (Table 1). Furthermore, the triploid genotypes from the breeders (which were BC1s) were backcrossed to diploid Asiatic parents by using both $2x - 3x$ or reciprocal combinations (Table 2). All the plant material is being maintained vegetatively at Wageningen UR, Plant Breeding, Wageningen, The Netherlands.

Pollen germination

Pollen was collected on day of anthesis and cultured for 5–20 h at 25°C in an artificial agar medium containing 100 g/l sucrose, 5 g/l of bacteriological agar, 20 mg/l of boric acid and 200 mg/l of calcium nitrate. The percentage of germinating pollen was determined based on the growth of pollen tube under an anatomical microscope.

Flow cytometry

Flow cytometry was done to evaluate the ploidy level of the BC1 and BC2 progenies. The germinating embryos were transferred into the propagation medium and allowed to grow until leaves developed. One leaf or scale was collected from each seedling for testing ploidy level as described by Van Tuyl and Boon (1997).

Mitotic chromosome preparation

For mitotic metaphase chromosome analysis, root tips were collected early in the morning, incubated in

0.7 mM cycloheximide solution for 4–6 h and then fixed in ethanol–acetic acid (3:1) solution for 12–24 h and stored at 4°C until use. The root tips were washed in distilled water and incubated in a pectolytic enzyme mixture containing 0.2% (w/v) pectinase Y23, 0.2% (w/v) cytohellicase and 0.2% (w/v) cellulase RS in 10 mM citrate buffer (pH 4.5) at 37°C for about 1 h. Squash preparations were made in a drop of 45% acetic acid and frozen in liquid nitrogen. The cover slips were removed by using a razor blade. The slides were then dehydrated in absolute ethanol and air dried.

Genomic in situ hybridization (GISH)

Genomic DNA of Longiflorum cultivar ‘White Fox’ was used as probe in GISH and labelled with Digoxigenin-11-dUTP by nick translation (Roche Diagnostics GmbH, Mannheim, Germany). The GISH procedure was performed as described by Lim et al. (2003) and Zhou et al. (2008). Briefly, the hybridization mixture contained 50% (v/v) deionized formamide, 10% (w/v) sodium dextran sulphate, $2\times$ SSC, 0.25% (w/v) sodium dodecyl sulphate, 1–1.5 ng/μl Digoxigenin labelled DNA from Longiflorum cultivars and 25–100 ng/μl block DNA of the Asiatic cultivar. DNA was denatured by heating the hybridization mixture at 70°C for 10 min followed by incubation in ice for 5 min. The hybridization mixture (40 μl) was then applied on each slide. The chromosome preparations were denatured at 80°C for 5 min immediately prior to incubation. After overnight hybridization in a humid chamber at 37°C the slides were washed at room temperature in $2\times$ SSC for 15 min followed by stringent washing with $0.1\times$ SSC for 30 min at 42°C. Digoxigenin labelled DNA was detected with antidigoxigenin–fluorescein raised in sheep (Boehringer, Mannheim, Germany) and amplified with fluorescein anti-sheep immunoglobulin raised in rabbit (Vector Laboratories). Preparations were analysed using a Zeiss Axiophot epifluorescence microscope and photographed by Canon digital camera. For each plant, the total number of chromosomes and the number of recombinant points were determined.

Chromosome identification, measurement and determination of genome contribution

For karyotyping in both genomes (L and A), chromosomes are arranged in sequence of decreasing short-arm

Table 1 The frequencies of diploid ($2n = 2x = 24$), or aneuploid, and triploid ($2n = 3x = 36$) BC1 progenies resulting from crosses between different genotypes of LA hybrids and the Asiatic parents

Cross	Genotype code	Parents		No. of plants analysed	Ploidy level ^b		
		Female	Male		2x	3x	Others
LAA	044511	041543	Mont Blanc	2	2		
LAA ^a	044512	041558	Mont Blanc	1		1	
LAA	044518	041543	Pollyanna	1		1	
LAA	044525	041556	Mont Blanc	1		1	
LAA	044535	041562	Pollyanna	1		1	
LAA	044536	041563	Pollyanna	1	1		
LAA	044538	041560	Pollyanna	4	4		
LAA	044539	041558	Pollyanna	5		5	
LAA	044541	041549	Pollyanna	1		1	
LAA	044567	041548	Pollyanna	1		1	
LAA	044571	041557	Mont Blanc	4		4	
LAA	044611	041559	Pollyanna	1		1	
LAA	044624	041565	Pollyanna	1		1	
LAA	044630	041502	Mont Blanc	1		1	
LAA	062035	041560	Sarina	2		2	
LAA	062071	041560	Montreux	4	1	3	
LAA	062074	041560	Montreux	5	2	3	
LAA	065051	024004-5	061095	4	2	1	1 (2x - 1)
LAA	066828	041543	051072	5	1	4	
LAA	066960	041543	Lanzarote	18	3	14	1 (2x + 2)
LAA	066963	041543	Gironde	7	2	5	
LAA	066994	041560	Alaska	13	6	5	2 (2x + 3)
LAA	066995	041560	031040	1		1	
ALA ^a	044595	Pollyanna	041519	3	1	2	
ALA	044601	Mont Blanc	041502	9		9	
ALA	044602	Mont Blanc	041557	2	2		
ALA	044608	Vivaldi	041502	6		6	
Total				104	27	73	4

^a Denotes only the direction of the crosses but not the ploidy level. LAA indicates that the hybrid LA was the female parent and ALA indicates that LA hybrid was the male parent

^b Estimated cytologically

length according to Stewart (1947). Some of the chromosomes in the somatic karyotype could be identified on the basis of total length and arm ratios (e.g., 1, 2, 3, 10, 11 and 12). In other cases, however, the differences in the lengths of short-arms were used for identification. By identifying recombination sites analysed by GISH, the same chromosome could be accurately identified in 5–10 cells and compared with the previous and the succeeding chromosomes in the karyotype. Furthermore, the centromeric index (short-arm length/short-arm length + long-arm length) and the relative chromosome

length index (individual chromosome length/total length of all chromosomes) were determined for all genotypes (Barthes and Ricroch 2001). For the measurements and determination of genome contribution, images of mitotic metaphase chromosomes from each genotype were collected and were measured in micrometres (μm). The contribution of the amount of both L and A genomes (in terms of percentage) in BC progenies was determined using the computer programme MicroMeasure (Reeves and Tear 2000; <http://www.colostate.edu/Depts/Biology/MicroMeasure>).

Table 2 The ploidy level of BC2 progenies resulting from crossing triploid BC1 plants with diploid Asiatic parents

Interploid crosses	Parents		No. of plants analysed	No. of diploid plants ^a
	Female	Male		
2x – 3x (BC2)				
044529	Pollyanna	041513	3	3
044530	Mont Blanc	041513	5	5
3x – 2x (BC2)				
044519	041513	Pollyanna	3	3
044524	041513	Mont Blanc	1	1
044634	041552	Pollyanna	3	2
Total			15	14

^a Estimated cytologically

Results

Production of BC1 and BC2 progenies from LA hybrids

In order to assess the possibility to produce relatively large number of diploid backcross progenies from the LA hybrids or from its BC1 progenies a large number of crosses were made. For this purpose, several LA hybrids and allotriploid BC1 (LAA) genotypes were backcrossed to diploid Asiatic parents and the ploidy levels of these progenies were determined through flow cytometry (data not presented) as well as by cytological chromosome counting (Tables 1 and 2).

Not all LA hybrids were successful as male and female parents. Viable pollen grains (5–30%) were produced in some cases (e.g., F1 hybrids 041502, 041519, 041557) and these could be used successfully as male parents (Table 1). Other genotypes could only be used as female parent. By crossing LA hybrids both as female and male parent, a total of 104 BC1 progenies were obtained of which 27 (26%) were diploid, 73 (70.2%) were triploid and four were aneuploid (3.8%) with $2x - 1$, $2x + 2$ and $2x + 3$, respectively. This indicated the occurrence of haploid (n), diploid ($2n$) and aneuploid gametes in the LA hybrids (Table 1). Analysis of the different BC1 types produced by each genotype indicated that three genotypes produced only n eggs. All other produced both n and $2n$ eggs although in the majority of cases $2n$ eggs are produced. Similarly, there was one genotype that produced only n pollen (041557), one that produced both n and $2n$ pollen (041519) and one genotype produced only $2n$ pollen (041502). The observations

indicated that it was possible to produce reasonable numbers of diploid BC1 progenies along with a large number of triploid progenies from LA hybrids despite possessing highly differentiated genomes and abnormal meiosis in these hybrids.

Apart from producing BC1 progenies from diploid LA hybrids, an attempt was also made to obtain diploid progenies by backcrossing allotriploid (LAA and ALA) BC1 plants with different diploid Asiatic parents. From the flow cytometric determination of the ploidy levels of the progenies of $2x - 3x$ and reciprocal crosses, it was established that 90.33% of the BC2 plants showed diploid chromosome numbers and this was confirmed cytologically (Table 2; Fig. 1c). Obviously, despite being allotriploids, they produced balanced haploid gametes both male and female, giving rise to diploid BC2 progenies on back crossing.

Genome composition and the extent of crossing-over in BC progenies

Through GISH analysis, the genome composition was determined in 24 diploid BC1 (Table 3) and five BC2 plants (Table 4). Among BC1 progenies, 20 had the expected 24 chromosomes and four were aneuploid ($2x - 1$, $2x + 2$ or $2x + 3$). Because the Asiatic parent was used for backcrossing, the number of A genome chromosomes (chromosomes of which the centromere was of Asiatic genome) predominated in the BC1 progenies and varied from 15 to 23 (Table 3). On the other hand, the chromosomes with the centromere of the L genome varied from only one (044602-2) to nine (066960-2 and 066963-2) (Table 3). There were obviously four type of chromosomes; two




Fig. 1 Chromosome identification and identification of recombination sites with GISH. **a** and **b** BC1 LA diploids (044538-2; 066963-7). **c** BC2 LA diploid (044529-2). Recombinant chromo-

somes are marked with arrow heads indicating both types of recombinant chromosomes either L/A (centromere of Longiflorum genome with Asiatic recombinant segment) or vice versa i.e., A/L

non-recombinant chromosomes, L and A and two recombinant chromosomes L/A and A/L. Those with a centromere of Asiatic chromosome with a recombinant segment of Longiflorum, are indicated as A/L, and vice versa, i.e., L/A (Table 3; Fig. 1a–c). In all, there were 71 L/A and 86 A/L types of recombinant chromosomes found in the total of progenies analysed. Taking the total length of both A and L chromosomes in the BC1 progenies, the percentages of each genome present in the BC1 progenies was estimated (Table 3). The percentage of L genome deviated from the expected (25%) and it varied from 3.3% (044602-2) to as high as 32.5% (066960-15), nearly a ten fold variation. In order to estimate the number and types of recombinant chromosomes and the lengths of recombinant segments, the karyotypes of some of the BC1 progenies were determined as shown in Fig. 2. It was found that a considerable amount of Longiflorum genome was transmitted from LA hybrids to BC1 progenies, however, the BC2

progenies possessed hardly any of the alien genome or segments of L genome (Fig. 1c).

Discussion

Nearly 30% of the 104 BC1 progenies produced from LA hybrids in this investigation turned out to be diploid or near diploid (i.e., aneuploid with almost diploid chromosome count). This shows that there is a prospect for producing sufficient number of diploid BC1 progenies suitable for analytic breeding in LA hybrids. An attractive feature of these progenies is that it might be possible to monitor the segregation of distinctive genetic traits that distinguish the two parental species. The inheritance patterns of some of the desirable horticultural traits in these progenies might provide an insight into how such traits can be utilized in breeding lilies. Because the generation time of lilies is long (two or more years) the segregation of

Table 3 Genome composition of BC1 (diploid and aneuploid) LA hybrids and the number of recombinant chromosomes among different genotypes

Cross	Genotype code	Ploidy level	Genome composition		Genome percentage		No. of recombinant chromosomes
			L ^(L/A)	A ^(A/L)	L (%)	A (%)	
LAA	044511-1	2x	3(1)	21(2)	12.55	87.45	3
LAA	044538-1	2x	7(7)	17(4)	24.37	75.62	11
LAA	044538-2	2x	7(6)	17(3)	28.6	71.4	9
LAA	044538-3	2x	4(2)	20(4)	17.84	82.15	6
LAA	044538-4	2x	5(2)	19(5)	24.32	75.67	7
LAA	062704-2	2x	4(2)	20(4)	25.6	74.63	6
LAA	062704-5	2x	6(4)	18(4)	16	84	8
LAA	065051-3	2x	3(3)	21(2)	8.81	91.18	5
LAA	065051-5	2x – 1	3(2)	20(3)	17.80	82.20	5
LAA	065051-6	2x	7(6)	17(3)	27.5	72.5	9
LAA	066828-5	2x	5(4)	19(2)	15.2	84.8	6
LAA	066960-2	2x	9(5)	15	26.88	73.11	5
LAA	066960-15	2x + 2	7(2)	19(1)	32.52	67.47	3
LAA	066960-17	2x	8(4)	16(3)	31.88	68.11	7
LAA	066960-22	2x	5(5)	19(4)	13.9	86	9
LAA	066963-2	2x	9(5)	15	28.9	71.09	5
LAA	066963-7	2x	7(3)	17	25.2	74.8	3
LAA	066994-1	2x	6(3)	18(4)	27.81	72.18	7
LAA	066994-2	2x	4(4)	20(5)	29.28	70.72	9
LAA	066994-5	2x	7(5)	17(4)	21	79	9
LAA	066994-6	2x + 3	7(4)	20(4)	35.9	64.1	8
LAA	066994-7	2x + 3	7(3)	20(4)	24.37	75.63	7
LAA	066994-10	2x	6(3)	18(4)	26.2	73.8	7
ALA	044602-2	2x	1(1)	23(2)	3.34	96.66	3


Table 4 Genome composition of BC2 (diploid and aneuploid) LA hybrids and the number of recombinant chromosomes among different genotypes analysed by GISH

Cross	Genotype code	Ploidy level	Genome composition		No. of recombinant chromosomes
			L ^(L/A)	A ^(A/L)	
ALAA	044529-2	2x	1(1)	23	1
ALAA	044529-3	2x	–	24	0
ALAA	044529-4	2x	–	24	0
ALAA	044530-1	2x	–	24(1)	1
LAAA	044634-1	2x + 1	1	24(2)	2

traits has not yet been analysed in the present study. Nevertheless, a cursory look at the karyotypes of BC progenies (Fig. 2) does indicate that there is a potential for genetic variation and consequently a possibility for selection at the diploid level. From the point of view of breeding allopolyploid crops, the results of

the present investigation are relevant to other ornamental. Just as in *Lilium*, an important group of tulips, the so called “Darwin Hybrids”, have originated from interspecific hybrids between *Tulipa gesneriana* × *T. fosteriana* (Marasek et al. 2006; Marasek and Okazaki 2008). Despite having genetically differentiated

Fig. 2 Diagrammatic representation of recombinant chromosomes in BC diploid LA hybrid. 044538-2 and 065051-6 (BC1 LAA type hybrids); 044602-2 (BC1 ALA hybrid) and 044529-2 (A LAA BC2 hybrid). In this figure the black colour represents the Longiflorum genome while grey colour represents the Asiatic one



genomes, these F1 hybrids have been shown to produce both n and $2n$ gametes so that both diploid ($2n = 2x = 24$) and triploid ($2n = 3x = 36$) BC1 progenies can be obtained by backcrossing with one of the parent i.e., *T. gesneriana* (Marasek and Okazaki 2008).

Similarly, occurrence of both n and $2n$ gametes has also been reported in other ornamentals like *Alstroemeria aurea* \times *A. inodora* (Kamstra et al. 1999). This indicates that there are prospects of using analytic breeding in more ornamental crops other than lily. The potential for genetic variation and consequently the possibility for selection at the diploid level has been practiced in some other crops of economic importance like *Lycopersicon esculentum* \times *Solanum lycopersicoides* (Chetelat et al. 1997) and

Festuca \times *Lolium* (Zwierzykowski et al. 1998; Thomas et al. 2003).

Three important aspects of analytic breeding are, (a) conventional breeding at diploid level for improved quality and high yield traits (b) the maximization of heterozygosity in the polyploids that might be produced from the selected diploid genotypes and (c) introgression of desirable traits from one species to the other. For good reasons, it has been argued that a higher degree of heterozygosity can be achieved at the polyploid rather than at the diploid level (Bingham 1980; Sanford 1983). The reason being that polyploids have a greater probability of possessing three or more different alleles at the same locus whereas it is restricted to only two in diploids. The importance of

poly allelic loci for inducing vigour and fertility in autopolyploids has been demonstrated in the case of *Medicago sativa* (Busbice and Wilsie 1966). This aspect can be verified by using diploid BC progenies for breeding and selection followed by polyploidization in lilies. A hypothetical analytic breeding scheme for lily LA hybrids has been represented in Fig. 3.

The traditional method of introgressing traits into allopolyploids was through the re-synthesis of the crop by using the putative parental species as has been done in *Ipomea* spp. (Orjeda et al. 1991), *Aegilops-Triticum* (Feldman et al. 1997; Ozkan et al. 2001), *Spartina* spp. (Baumel et al. 2002), *Musa* Spp. (Ortiz 1997), and *Brassicacae* (Lukens et al. 2006; Nicolas et al. 2007). Hybridization of such re-synthesized allopolyploid with elite cultivars adds the entire genomes of the alien species rather than only the chromosome segments that possess the desirable traits. The addition of entire genomes, obviously, adds many undesirable traits along with the desirable ones into the cultivars enhancing so called, linkage drag (Hospital 2001). One effective way of preventing linkage drag is to add specific recombinant segments that possess desirable traits (Servin et al. 2004). An important requirement for this is that, (a) homoeologous recombination must be accomplished in the hybrids of the putative parental species and (b) the recombinant chromosomes

must be recovered in the progenies of the F1 hybrids. These two requirements are most ideally achieved in the case of LA hybrids studied here and a wide range of recombinant chromosomes with varying numbers and lengths of recombinant segments have been recovered in the BC1 progenies (Table 2; Figs. 1a, b and 2). Besides, it has also been possible to produce BC2 progenies from the triploid BC1 genotypes (Table 2). In a previous study it was found that in the progenies of $3x - 2x$ crosses of Longiflorum \times Asiatic and $3x - 2x$ and reciprocal crosses of Oriental \times Asiatic hybrids lilies, predominantly near diploid (i.e., aneuploid) or diploid progenies occurred indicating that triploid BC1 plants produced both balanced n as well as near haploid (e.g., $n + 1$ or $n + 2$ etc.) gametes (Lim et al. 2003; Barba-Gonzalez et al. 2006a). Similarly, present investigation shows that most of the BC2 progenies obtained after $2x - 3x$ and in reciprocal crosses produced mostly diploid progenies. However, there is no convincing explanation for the occurrence of balanced n gametes in triploids used in the present study. This appears to be genotype dependent that needs to be confirmed by further analysis. Therefore, more progenies from this cross have to be analysed. A notable feature of the BC2 progenies obtained from $2x - 3x$ and reciprocal crosses is that they transmit very short recombinant segments or very few alien chromosomes to the progenies. These might be especially favourable in avoiding linkage drag. The scarcity of L genome chromosomes or segments in these BC2 progenies might have resulted from a process of selection of gametes during fertilization or embryo development or due to non-inclusion of L genome chromosomes during gamete formation. This implies that introgression of some desirable Longiflorum traits into an Asiatic cultivar could be realized at BC2 level. Similar results have been found in *Festuca-Lolium* intergeneric hybrids where freezing tolerance genes were successfully introgressed into *Lolium multiflorum* from *Festuca pratensis* in a back cross breeding programme (Kosmala et al. 2006).

Besides their relevance to analytic breeding, the diploid BC progenies can be useful for molecular mapping of useful traits as exemplified in other crops like banana (Fauré et al. 1993; Raboin et al. 2005), wheat (Dubcovsky et al. 1996), alfalfa (Barcaccia et al. 1999) and cotton (Desai et al. 2008). Especially when quantitative traits are to be transferred, diploids




Fig. 3 Hypothetical analytic breeding scheme for LA hybrids, L—Longiflorum genome, A—Asiatic genome

might more conducive for tagging the quantitative loci that might be used in marker assisted selection (Dirlewanger et al. 2004). Diploid BC1 ALA and LAA plants can be backcrossed with the recurrent Asiatic parents to get the genes of interest in backcross progenies. Finally, the concept of hybrid vigour resulting from the degree of heterozygosity of the polyploid progenies can be assessed through the use of molecular methods and the advantages of analytical breeding can be determined.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Barba-Gonzalez R, Lokker BH, Lim K-B, Ramanna MS, Van Tuyl JM (2004) Use of $2n$ gametes for the production of sexual polyploids from sterile Oriental \times Asiatic hybrids of lilies (*Lilium*). *Theor Appl Genet* 109:1125–1132. doi:10.1007/s00122-004-1739-0
- Barba-Gonzalez R, Lim K-B, Ramanna MS, Visser RGF, Van Tuyl JM (2005) The occurrence of intergenomic recombination in the F1 hybrid of Oriental \times Asiatic lily hybrids (*Lilium*) and its significance for genetic variation in the BC1 progenies as revealed by GISH and FISH analysis. *Genome* 48:884–894. doi:10.1139/g05-064
- Barba-Gonzalez R, Alex AVS, Visser RGF, Ramanna MS, Van Tuyl JM (2006a) Progenies of allotripleids of Oriental \times Asiatic lilies (*Lilium*) examined by GISH analysis. *Euphytica* 151:243–250. doi:10.1007/s10681-006-9148-x
- Barba-Gonzalez R, Miller CT, Ramanna MS, Van Tuyl JM (2006b) Induction of $2n$ gametes for overcoming F1-sterility in lily and tulip. *Acta Hort* 714:99–106
- Barcaccia G, Albertini E, Tavoletti S, Falcinelli M, Veronesi F (1999) AFLP fingerprinting in *Medicago* spp.: its development and application in linkage mapping. *Plant Breed* 118:335–340. doi:10.1046/j.1439-0523.1999.00376.x
- Barcaccia G, Albertini E, Rosellini D, Tavoletti S, Veronesi F (2000) Inheritance and mapping of $2n$ egg production in diploid alfalfa. *Genome* 43:528–537. doi:10.1139/gen-43-3-528
- Barthes L, Ricroch A (2001) Interspecific chromosomal rearrangements in monosomic addition lines of *Allium*. *Genome* 44:929–935. doi:10.1139/gen-44-5-929
- Baumel A, Ainouche ML, Kalendar R, Schulman AH (2002) Retrotransposons and genomic stability in populations of the young allopolyploid species *Spartina anglica* C.E. Hubbard (Poaceae). *Mol Biol Evol* 19:1218–1227
- Bingham ET (1980) Maximising heterozygosity in autopolyploids. In: Lewis WH (ed) Polyploidy-biological relevance. Plenum Press, New York, pp 471–490
- Bingham ET, Groose RW, Woodfield DR, Kidwell KK (1994) Complementary gene interactions in alfalfa are greater in autotetraploids than diploids. *Crop Sci* 34:823–829
- Busbice TH, Wilsie CP (1966) Genetics of *Medicago sativa* L. I. Inheritance of dwarf character, Dw_1^1 . *Crop Sci* 6:327–330
- Chase SA (1963) Androgenesis—its use for transfer of maize cytoplasm. *J Hered* 54:152–158
- Chetelat RT, Cisneros P, Stamova L, Rick CM (1997) A male-fertile *Lycopersicon esculentum \times *Solanum lycopersicon* hybrid enables direct backcrossing to tomato at the diploid level. *Euphytica* 95:99–108. doi:10.1023/A:1002958030799*
- Desai A, Chee PW, May OL, Paterson AH (2008) Correspondence of trichome mutations in diploid and tetraploid cottons. *J Hered* 99:182–186. doi:10.1093/jhered/esm112
- Dirlewanger E, Graziano E, Joobeur T, Garriga-Calderé F, Cosson P, Howad W et al (2004) Comparative mapping and marker-assisted selection in Rosaceae fruit crops. *Proc Natl Acad Sci USA* 101:9891–9896. doi:10.1073/pnas.0307937101
- Dubcovsky J, Luo M-C, Zhong G-Y, Bransteiter R, Desai A, Kilian A et al (1996) Genetic map of diploid wheat, *Triticum monococcum* L., and its comparison with maps of *Hordeum vulgare* L. *Genetics* 143:983–999
- Estilai A (1977) Interspecific hybrids between *Carthamus tinctorius* and *C. alexandrinus*¹. *Crop Sci* 17:800–802
- Fauré S, Noyer J-L, Horry JP, Bakry F, Lanaud C, González-de-León D (1993) A molecular marker-based linkage map of diploid bananas (*Musa acuminata*). *Theor Appl Genet* 87:517–526. doi:10.1007/BF00215098
- Feldman M, Levy AA (2005) Allopolyploidy—a shaping force in the evolution of wheat genomes. *Cytogenet Genome Res* 109:250–258. doi:10.1159/000082407
- Feldman M, Liu B, Segal G, Abbo S, Levy AA, Vega JM (1997) Rapid elimination of low-copy DNA sequences in polyploid wheat: a possible mechanism for differentiation of homoeologous chromosomes. *Genetics* 147:1381–1387
- Gu XF, Yang AF, Meng H, Zhang JR (2005) In vitro induction of tetraploid plants from diploid *Zizyphus jujuba* Mill cv. Zhanhua. *Plant Cell Rep* 24:671–676. doi:10.1007/s00299-005-0017-1
- Hospital F (2001) Size of donor chromosome segments around introgressed loci and reduction of linkage drag in marker-assisted backcross programs. *Genetics* 158:1363–1379
- Hoya A, Shibaie H, Morita T, Ito M (2007) Germination characteristics of native Japanese dandelion autopolyploids and their putative diploid parent species. *J Plant Res* 120:139–147. doi:10.1007/s10265-006-0034-3
- Jacobsen E, Ramanna MS, Huigen DJ, Sawor Z (1991) Introduction of an amylose free (*amf*) mutant into breeding of cultivated potato *Solanum tuberosum* L. *Euphytica* 53:247–253. doi:10.1007/BF00023276
- Kamstra SA, Kuipers AGJ, De Jeu MJ, Ramanna MS, Jacobsen E (1999) The extent and position of homoeologous recombination in a distant hybrid of *Alstroemeria*: a molecular cytogenetic assessment of first generation backcross progenies. *Chromosoma* 108:52–63. doi:10.1007/s0041200 50351
- Kosmala A, Zwierzykowski Z, Gasior D, Rapacz M, Zwierzykowska E, Humphreys MW (2006) GISH/FISH mapping of genes for freezing tolerance transferred from *Festuca pratensis* to *Lolium multiflorum*. *Heredity* 96:243–251. doi:10.1038/sj.hdy.6800787
- Lim K-B, Chung J-D, Van Kronenburg BCE, Ramanna MS, De Jong JH, Van Tuyl JM (2000) Introgression of *Lilium*

- rubellum* Baker chromosomes into *L. longiflorum* Thunb.: a genome painting study of the F1 hybrid, BC1 and BC2 progenies. Chromosome Res 8:119–125. doi:10.1023/A:1009290418889
- Lim K-B, Ramanna MS, De Jong JH, Jacobsen E, Van Tuyl JM (2001) Indeterminate meiotic restitution (IMR): a novel type of meiotic nuclear restitution mechanism detected in interspecific lily hybrids by GISH. Theor Appl Genet 103:219–230. doi:10.1007/s001220100638
- Lim K-B, Ramanna MS, Jacobsen E, Van Tuyl JM (2003) Evaluation of BC2 progenies derived from $3x - 2x$ and $3x - 4x$ crosses of *Lilium* hybrids: a GISH analysis. Theor Appl Genet 106:568–574
- Lopez-Lavalle BLA, Orjeda G (2002) Occurrence and cytological mechanism of $2n$ pollen formation in a tetraploid accession of *Ipomoea batatas* (sweet potato). J Hered 93:185–192. doi:10.1093/jhered/93.3.185
- Lukens LN, Pires JC, Leon E, Vogelzang R, Oslach L, Osborn T (2006) Patterns of sequence loss and cytosine methylation within a population of newly resynthesized *Brassica napus* allopolyploids. Plant Physiol 140:336–348. doi:10.1104/pp.105.066308
- Lyrene PM, Vorsa N, Ballington JR (2003) Polyploidy and sexual polyploidization in the genus *Vaccinium*. Euphytica 133:27–36. doi:10.1023/A:1025608408727
- Marasek A, Okazaki K (2008) Analysis of introgression of *Tulipa fosteriana* genome into *Tulipa gesneriana* using GISH and FISH. Euphytica 160:217–230. doi:10.1007/s10681-007-9539-7
- Marasek A, Mizuochi H, Okazaki K (2006) The origin of Darwin hybrid tulips analyzed by flow cytometry, karyotype analyses and genomic in situ hybridization. Euphytica 151:279–290. doi:10.1007/s10681-006-9147-y
- Mariani A, Tavoletti S (1992) Gametes with somatic chromosome number in the evolution and breeding of polyploid polysomic species. In: Proc Workshop, Perugia, Tipolitothographia Porziuncola-Assisi (PG) Italy, pp 1–103
- Mendiburu AO, Peloquin SJ, Mok DWS (1974) Potato breeding with haploids and $2n$ gametes. In: Kasha KJ (ed) Haploids in higher plants. Guelph University Press, Guelph, pp 249–258
- Nicolas SD, Mignon GL, Eber F, Coriton O, Monod H, Clouet V et al (2007) Homoeologous recombination plays a major role in chromosome rearrangements that occur during meiosis of *Brassica napus* haploids. Genetics 175:487–503. doi:10.1534/genetics.106.062968
- Orjeda G, Freye R, Iwanaga M (1991) Use of *Ipomoea trifida* germ plasm for sweet potato improvement. 3. Development of $4x$ interspecific hybrids between *Ipomoea batatas* (L.) Lam. ($2n = 6x = 90$) and *I. trifida* (H.B.K.) G. Don. ($2x = 2x = 30$) as storage-root initiators for wild species. Theor Appl Genet 83:159–163. doi:10.1007/BF00226245
- Ortiz R (1997) Occurrence and inheritance of $2n$ pollen in *Musa*. Ann Bot (Lond) 79:449–453. doi:10.1006/anbo.1996.0367
- Ozkan H, Levy AA, Feldman M (2001) Allopolyploidy-induced rapid genome evolution in the wheat (*Aegilops-Triticum*) group. Plant Cell 13:1735–1747
- Peloquin SJ (1981) Chromosomal and cytoplasmic manipulations. In: Frey K (ed) Plant breeding symposium II. The Iowa State University Press, Iowa, pp 117–150
- Raboin L-M, Carreel F, Noyer JL, Baurens FC, Horry JP, Bakry F et al (2005) Diploid ancestors of triploid export banana cultivars: molecular identification of $2n$ restitution gamete donors and n gamete donors. Mol Breed 16:333–341. doi:10.1007/s11032-005-2452-7
- Reeves A, Tear J (2000) MicroMeasure for windows, version 3.3. Free program distributed by the authors over the internet from <http://www.colostate.edu/Depts/Biology/MicroMeasure>
- Sanford JC (1983) Ploidy manipulations. In: Janick J (ed) Methods in fruit breeding. Perdue University Press, West Lafayette
- Servin B, Martin OC, Mézard M, Hospital F (2004) Toward a theory of marker-assisted gene pyramiding. Genetics 168:513–523. doi:10.1534/genetics.103.023358
- Stewart RN (1947) The morphology of somatic chromosomes in *Lilium*. Am J Bot 34:9–26. doi:10.2307/2437190
- Thomas HM, Morgan WG, Humphreys MW (2003) Designing grasses with a future-combining the attributes of *Lolium* and *Festuca*. Euphytica 133:19–26. doi:10.1023/A:1025694819031
- Van Tuyl JM, Boon E (1997) Variation in DNA-content in the genus *Lilium*. Acta Hort 430:829–835
- Van Tuyl JM, De Jeu MJ (1997) Methods for overcoming interspecific crossing barriers. In: Shivanna KR, Sawhney VK (eds) Pollen biotechnology for crop production and improvement. Cambridge University Press, Cambridge, pp 273–292
- Van Tuyl JM, Lim K-B (2003) Interspecific hybridization and polyploidization as tools in ornamental plant breeding. Acta Hort 612:13–22
- Van Tuyl JM, Van Diën MP, Van Creij MGM, Van Kleinwee TCM, Franken J, Bino RJ (1991) Application of in vitro pollination, ovary culture, ovule culture and embryo rescue for overcoming incongruity barriers in interspecific *Lilium* crosses. Plant Sci 74:115–126. doi:10.1016/0168-9452(91)90262-7
- Weiss H, Maluszynska J (2000) Chromosomal rearrangement in autotetraploid plants of *Arabidopsis thaliana*. Hereditas 133:255–261. doi:10.1111/j.1601-5223.2000.00255.x
- Zeldin EL, McCown BH (2004) Polyploid breeding and the potential of its use for transgene containment with American cranberry (*Vaccinium macrocarpon*). Acta Hort 663: 838–840
- Zhang Z, Dai H, Xiao M, Liu X (2007) In vitro induction of tetraploids in *Phlox subulata* L. Euphytica 159:59–65. doi:10.1007/s10681-007-9457-8
- Zhou S (2007) Intergenomic recombination and introgression breeding in Longiflorum × Asiatic lilies. PhD Thesis, University Wageningen, 111 pp
- Zhou S, Ramanna MS, Visser RGF, Van Tuyl JM (2008) Genome composition of triploid lily cultivars derived from sexual polyploidization of Longiflorum × Asiatic hybrids (*Lilium*). Euphytica 160:207–215. doi:10.1007/s10681-007-9538-8
- Zwierzykowski Z, Tayyar R, Brunell M, Lukaszewski AJ (1998) Genome recombination in intergeneric hybrids between tetraploid *Festuca pratensis* and *Lolium multiflorum*. J Hered 89:324–328. doi:10.1093/jhered/89.4.324