

This is a "Post-Print" accepted manuscript, which has been published in "Environmental Microbiology".

Please cite this publication as follows:

Mols, M., Van Kranenburg, R., Van Melis, C.C.J., Moezelaar, R., Abee, T. (2010) Analysis of acid-stressed *Bacillus cereus* reveals a major oxidative response and inactivation-associated radical formation. *Environmental Microbiology* 12 (4), 873-885.

You can download the published version at:

<http://dx.doi.org/10.1111/j.1462-2920.2009.02132.x>

1 **Title**

2 Analysis of acid-stressed *Bacillus cereus* reveals a major oxidative response and inactivation-  
3 associated radical formation

4

5 **Running title**

6 Acid-induced radical formation in *B. cereus*

7

8 **Authors**

9 Maarten Mols<sup>1,2\*</sup>, Richard van Kranenburg<sup>1,3</sup>, Clint C.J. van Melis<sup>2</sup>, Roy Moezelaar<sup>1,4</sup>, and  
10 Tjakko Abee<sup>1,2</sup>

11

12 **Affiliations**

13 <sup>1</sup> Top Institute Food and Nutrition, Wageningen, the Netherlands

14 <sup>2</sup> Laboratory of Food Microbiology, Wageningen University, Wageningen, the Netherlands

15 <sup>3</sup> PURAC, Gorinchem, the Netherlands

16 <sup>4</sup> Food Technology Centre, Wageningen University and Research Centre, Wageningen, the  
17 Netherlands

18

19 \* For correspondence. E-mail maarten.mols@wur.nl; Tel. (+31) 317 484983; Fax (+31) 317  
20 484978.

21

22

1 **Summary**

2

3 Acid stress resistance of the food-borne human pathogen *Bacillus cereus* may contribute to its  
4 survival in acidic environments, such as encountered in soil, food, and the human  
5 gastrointestinal tract. The acid stress responses of *B. cereus* strains ATCC 14579 and ATCC  
6 10987 were analysed in aerobically grown cultures acidified to pH values ranging from pH  
7 5.4 to pH 4.4 with HCl. Comparative phenotype and transcriptome analyses revealed three  
8 acid stress-induced responses in this pH range: growth rate reduction, growth arrest and loss  
9 of viability. These physiological responses showed to be associated with metabolic shifts and  
10 the induction of general stress response mechanisms with a major oxidative component,  
11 including up-regulation of catalases and superoxide dismutases. Flow cytometry analysis in  
12 combination with the hydroxyl (OH·) and peroxynitrite (ONOO<sup>-</sup>) -specific fluorescent probe  
13 3'-(p-hydroxyphenyl) fluorescein (HPF), showed excessive radicals to be formed in both *B.*  
14 *cereus* strains in bactericidal conditions only. Our study shows that radicals can indicate acid-  
15 induced malfunctioning of cellular processes that lead to cell death.

16

17

## 1 **Introduction**

2

3 *Bacillus cereus* is a Gram-positive, spore-forming, facultative anaerobic, rod-shaped food-  
4 borne human pathogen that appears to be well-equipped to survive in various adverse  
5 conditions. The spores and vegetative cells of *B. cereus* can be found in a range of  
6 environments, such as soil (Von Stetten *et al.*, 1999; Vilain *et al.*, 2006), plant rhizosphere  
7 (Berg *et al.*, 2005), and various foods (Choma *et al.*, 2000; Rosenquist *et al.*, 2005). Besides  
8 being notorious for causing spoilage of dairy products, *B. cereus* is a food-borne pathogen  
9 that can cause two distinct types of disease, i.e., emesis and diarrhoea (Kotiranta *et al.*, 2000).  
10 The emetic syndrome occurs upon ingestion of the heat-stable toxin cereulide, which is  
11 produced in food by emetic *B. cereus* strains (Agata *et al.*, 2002). The diarrheal syndrome is  
12 associated with the action of enterotoxins, such as non-haemolytic enterotoxin (NHE) and  
13 cytotoxin K (CytK) (Granum and Lund, 1997), that are produced by vegetative cells inside  
14 the human small intestine (Stenfors Arnesen *et al.*, 2008). Before entering the small intestine  
15 and subsequent production of enterotoxins, *B. cereus* cells have to survive the low pH of the  
16 human stomach. Therefore, acid resistance is a key parameter in the pathogenic potential of  
17 enterotoxic *B. cereus* strains. Obviously, the highly resistant dormant spores of *B. cereus* can  
18 pass the stomach unaffected and germination in the acid environment of the small intestine is  
19 an important aspect of their pathogenic potential (Wijnands *et al.*, 2007; Hornstra *et al.*,  
20 2009). Outside the human host, *B. cereus* may also be frequently exposed to acidic conditions  
21 including a range of low pH foods, where in specific cases organic acids have been added as  
22 preservatives (Brul and Coote, 1999). In conclusion, coping with low pH stress is an  
23 important feature in the performance of *B. cereus* in a variety of environments as described  
24 above, but also in other ecological niches such as soil and plant rhizosphere (Neumann and  
25 Martinoia, 2002).

1 Acid stress responses have mainly been studied in Gram-negative organisms, such as  
2 *Escherichia coli* and *Salmonella* Typhimurium (Richard and Foster, 2003), and in a select  
3 number of Gram-positive bacteria, such as lactic acid bacteria and *Listeria monocytogenes*  
4 (van de Guchte *et al.*, 2002; Cotter and Hill, 2003; Ryan *et al.*, 2008). These reviews highlight  
5 the importance of proton pumps, i.e., F<sub>1</sub>F<sub>0</sub>-ATPase, transcriptional regulators, such as RpoS  
6 (Gram-negatives) and  $\sigma^B$  (Gram-positives), proteins involved in protection of  
7 macromolecules, such as DnaK and GroESL, and enzymes that produce alkaline compounds,  
8 such as the ammonium forming enzymes urease and arginine deiminase. In contrast, the acid  
9 stress response of *B. cereus* has not been studied extensively. Available information is limited  
10 to alternative sigma factor  $\sigma^B$  expression upon exposure to a low pH (van Schaik *et al.*,  
11 2004), and the acid tolerance response, which includes modulation of intracellular pH and  
12 protein synthesis (Browne and Dowds, 2002; Jobin *et al.*, 2002; Thomassin *et al.*, 2006).  
13 Additionally, the role of urease in acid resistance of a large number of *B. cereus* strains has  
14 been studied (Mols and Abee, 2008), and revealed that its role in acid resistance of *B. cereus*  
15 was limited.

16 Therefore, we set out to investigate the molecular mechanisms involved in acid stress  
17 response of *B. cereus* and to identify possible acid-induced inactivation mechanisms, by  
18 comparing responses of cells exposed to selected pHs leading to mild, bacteriostatic and  
19 bactericidal acid stress. To determine both general and phenotype-associated transcriptional  
20 responses, two model strains ATCC 14579, isolated from air (Ivanova *et al.*, 2003) and ATCC  
21 10987, a food-isolate (Rasko *et al.*, 2004) were investigated. Recently, Kohanski and  
22 colleagues (2007) reported that the formation of reactive oxygen species (ROS), such as  
23 hydroxyl radicals (OH $\cdot$ ), plays a role in antibiotic-induced inactivation of aerobically grown  
24 *E. coli* and *Staphylococcus aureus* cells. These ROS were suggested to originate from  
25 antibiotic-induced perturbation of the electron transfer chain resulting in the production of

1 superoxide ( $O_2^-$ ).  $O_2^-$  can damage iron-sulphur clusters and subsequently react with the  
2 released iron, resulting in  $OH\cdot$  formation via the Fenton reaction. Therefore, flow cytometry  
3 analysis, in combination with the  $OH\cdot$  and peroxynitrite ( $ONOO^-$ )-specific fluorescent probe  
4 3'-(p-hydroxyphenyl) fluorescein (HPF) (Setsukinai *et al.*, 2003), was included in our study to  
5 detect ROS in (sub)lethally acid-stressed *B. cereus* cells. Our study provides evidence that  
6 radicals can indicate acid-induced malfunctioning of cellular processes and the stress-induced  
7 formation of reactive oxygen species as a common theme in bacterial stress response and  
8 cellular death is discussed.

9

## 10 **Results**

11

### 12 *Physiological response to acid stress*

13 The physiological response to acid stress was studied using *B. cereus* strains ATCC 14579  
14 and ATCC 10987 by acidifying aerobically grown cultures to pH values ranging from pH 5.4  
15 to pH 4.4 by addition of HCl (Fig. 1). Upon exposure to the different acid shocks, the growth  
16 of exponentially growing *B. cereus* cells was instantly affected. The two strains used showed  
17 different phenotypic responses to different levels of acidity. *B. cereus* strain ATCC 14579  
18 showed to continue growth as reflected in an increase of colony forming units upon exposure  
19 to pH shocks as low as pH 5.0. This response is hereafter referred to as growth phenotype.  
20 Strain ATCC 14579 was inactivated at pH 4.6 and lower as shown by the inability to form  
21 colonies on BHI plates incubated at 30°C for 16 hours. This response is hereafter referred to  
22 as inactivation phenotype and the condition as bactericidal. Upon exposure to pHs between  
23 pH 5.0 and 4.7, ATCC 14579 showed a stable number of viable counts within the first hour of  
24 exposure. This response is hereafter referred to as survival phenotype and the condition as  
25 bacteriostatic. However, prolonged exposure (overnight) resulted in a decrease of viable

1 counts (Fig. 2). The growth boundary of the other strain tested, *B. cereus* ATCC 10987, was  
2 determined at pH 5.0. Upon exposure to pHs higher than pH 5.0, ATCC 10987 was able to  
3 grow and at pHs lower than pH 5.0 this strain was inactivated. There was no apparent survival  
4 phenotype in the ATCC 10987 acid shock response using 0.1 pH unit intervals, as observed  
5 for ATCC 14579 between pH 5.0 and pH 4.7. The display of an intermediate physiological  
6 survival response by strain ATCC 14579 over the pH range 4.7 to 5.0, and the different  
7 inactivation boundaries were the main differences between the two strains tested.

8

### 9 *Microarray analysis using hierarchical clustering*

10 To investigate the impact of mild, bacteriostatic and bactericidal acid shocks on the gene  
11 expression of the two *B. cereus* strains, four pHs were selected based on the different  
12 phenotypic responses displayed by the two strains (Fig. 2A and 2B). At 0, 10, 30 and 60  
13 minutes after the exposure to the different pH shocks, i.e., pH 5.4, pH 5.0, pH 4.8 and pH 4.5,  
14 RNA samples were collected and subsequent microarray analyses were performed. To  
15 compare the transcriptomes of both strains, data obtained of orthologous genes that are  
16 present in the genomes of both ATCC 14579 and ATCC 10987 (Ivanova *et al.*, 2003; Rasko  
17 *et al.*, 2004; Mols *et al.*, 2007) were collected and subjected to hierarchical clustering. The  
18 transcriptome profiles clustered in two different groups, with one cluster including samples  
19 obtained of the growth phenotype and the other cluster encompassing samples of the survival  
20 (strain ATCC 14579 only) and inactivation phenotypes (Fig. 3). The transcriptome profiles of  
21 cultures that showed growth after acid shock exposure clustered together independent of the  
22 exposure time and strain. Within these two major groups, the different branches of the  
23 hierarchical clustering were separated mostly depending on strain and exposure pH rather  
24 than exposure time. In conclusion, the exposure of *B. cereus* ATCC 14579 and ATCC 10987

1 to mild, bacteriostatic and bactericidal acid stress led to phenotype specific transcriptome  
2 profiles independent of exposure time.

3 The pH- and phenotype-specific responses were investigated by analyzing the transcriptome  
4 data obtained for genes showing significant differential expression in one or more conditions  
5 per strain. Groups of genes with similar expression profiles were identified using hierarchical  
6 clustering (*Supplementary material*). The results obtained for a selection of genes putatively  
7 involved in low pH or oxidative responses are presented and discussed below.

8

### 9 *Acid shock response of low pH associated genes*

10 A selection of genes, based on their putative role in acid stress response of other Gram-  
11 positive organisms (Cotter and Hill, 2003; Ter Beek *et al.*, 2008), was monitored profoundly.

12 The selection includes transcription regulators, proton pumps, glutamate decarboxylase,  
13 production of alkaline compounds, protection of macromolecules, membrane synthesis, and  
14 multidrug transporters. The ratios of these genes obtained from cells exposed to pH 5.4 and  
15 pH 4.5 for 10, 30 and 60 minutes were averaged and plotted per gene (Fig. 4). In general, the  
16 average ratios, showing the up- or down-regulation, were less pronounced in inactivated cells.

17 The genes encoding sigma factors  $\sigma^B$  and  $\sigma^H$ , involved in the global adaptive response to  
18 stress, were slightly up-regulated in growing cells of both strains. On the other hand, *codY*,  
19 which is a key regulator in the nutrient starvation response of Gram-positive organisms,  
20 showed no significant up-regulation. The major oxidative stress response regulator *perR* was  
21 one of the most up-regulated transcription regulators in both growing and inactivated cells,  
22 indicating an oxidative response upon low pH exposure. Previously, proton pumps, i.e., F<sub>1</sub>F<sub>0</sub>-  
23 ATPase, were shown to contribute to pH homeostasis in fermenting Gram-positives exposed  
24 to mild acid conditions (Cotter and Hill, 2003). In this study, genes encoding subunits of the  
25 F<sub>1</sub>F<sub>0</sub>-ATPase (represented by *atpA* in Fig. 4) were highly down-regulated in aerobically



1 grown and exposed *B. cereus* cells upon exposure to sub-lethal pHs. Upon exposure to lethal  
2 acid shocks, genes encoding sodium-proton antiporters *napA* and *nhaC* were not down-  
3 regulated and *napA* even showed to be up-regulated. In *Listeria monocytogenes* (Cotter *et al.*,  
4 2001) and *Lactococcus lactis* (Sanders *et al.*, 1998) acid-induced glutamate decarboxylase  
5 (*gad*), which catalyzes the decarboxylation of glutamate with concomitant consumption of  
6 protons, was found to play an important role in low pH survival. In *B. cereus* ATCC 10987,  
7 however, the *gad* gene, that is not present in the genome of ATCC 14579, showed not to be  
8 up-regulated upon low pH exposure. This is in line with the notion that *B. cereus* ATCC  
9 10987 lacks a glutamate/GABA exchanger (Mols *et al.*, 2007), that is required to supply  
10 glutamate decarboxylase with its substrate (Cotter and Hill, 2003). Alkaline compound  
11 forming mechanisms, such as the arginine deiminase (ADI) pathway and the urease enzyme,  
12 are involved in acid tolerance of Gram-positive organisms (Cotter and Hill, 2003). Arginine  
13 deiminase (*arcA*), which is involved in acid resistance of streptococci (Curran *et al.*, 1995)  
14 and *L. monocytogenes* (Ryan *et al.*, 2009), showed significant up-regulation in both *B. cereus*  
15 strains upon exposure to sub-lethal acid shocks, whereas exposure to bactericidal acid shocks  
16 revealed no significant induction. Urease encoding genes, specific for ATCC 10987 (Mols *et*  
17 *al.*, 2007), were induced upon exposure to pH 5.4 (represented by *ureA* in Fig. 4A and 4B),  
18 but not in bactericidal conditions. Macromolecules are easily damaged during stress exposure,  
19 and their protection and repair is crucial for bacterial survival. DnaK and GroES are  
20 chaperones, preventing misfolding of proteins, and in *Streptococcus mutans* deletion of these  
21 chaperones resulted in less resistant cells (Lemos *et al.*, 2001). Notably, chaperone encoding  
22 genes *dnaK* and *groES* and protease encoding gene *clpC* were up-regulated upon exposure to  
23 sub-lethal acid shocks, whereas exposure to lethal pHs did not induce these genes in *B.*  
24 *cereus*.

1 Mild sorbic acid stress induces the expression of the fatty acid biosynthesis genes (*fab*), *bkdR*  
2 and a multidrug transport gene in *B. subtilis* (Ter Beek *et al.*, 2008). Two homologous  
3 multidrug systems (*mdr1* and *mdr2*) showed to be also up-regulated in sub-lethal inorganic  
4 acid stress in *B. cereus*. In contrast to sorbic acid stressed *B. subtilis*, fatty acid biosynthesis  
5 (represented by *fabF* in Fig. 4A and 4B) was down-regulated at pH 5.4 and no significant  
6 induction was found for branched-chain fatty acid biosynthesis (*bkdR*) in *B. cereus* upon  
7 exposure to (sub)lethal inorganic acid stress.

8

### 9 *Oxidative response and rearrangements in energy metabolism*

10 The response of several genes involved in oxidative stress and energy production were  
11 investigated in more detail (Fig. 5). Two distinct types of cytochrome oxidases showed  
12 different expression patterns. Cytochrome C oxidase, which acts as complex IV in aerobic  
13 conditions, was repressed in sub-lethal conditions in both strains. Cytochrome D ubiquinol  
14 oxidase, which can act as an alternative complex IV, was also down-regulated upon exposure  
15 to pH 5.4. However, it was highly induced in bacteriostatic and bactericidal conditions. Genes  
16 involved in oxidative stress, such as *sodA*, *katB* (Fig. 5) and *perR* (Fig. 4) were highly up-  
17 regulated in all acid shock conditions tested. The induction of these genes indicates that a low  
18 pH may induce the formation of oxidative compounds, such as H<sub>2</sub>O<sub>2</sub>. Nitric oxide (NO),  
19 formed from arginine by nitric oxide synthase (*nos*), putatively protects cells from H<sub>2</sub>O<sub>2</sub>-  
20 induced DNA damage by inhibition of the Fenton reaction and activation of catalase (Gusarov  
21 and Nudler, 2005; Shatalin *et al.*, 2008). Although *nos* was only slightly up-regulated in  
22 bactericidal conditions, the formation of nitric oxide may be inferred from the induction of  
23 nitric oxide dioxygenase (*hmp*) and a nitric oxide dependant transcriptional regulator (*dnrN*,  
24 *Supplementary material*). Nitric oxide dioxygenase facilitates the reaction of nitric oxide with  
25 oxygen to form nitrate. Nitrate reductase (*nar*) and nitrite reductase (*nas*) are involved in

1 nitrogen metabolism and may serve as an alternative for aerobic respiration. Nitrate reductase  
2 and nitrite reductase genes are unique for strain ATCC 14579 and cluster together with nitrite  
3 extrusion protein *narK* (Mols *et al.*, 2007). The cluster, including *nar*, *nas* and *narK*, was  
4 highly up-regulated upon exposure to all acid shocks tested.

5 Besides genes involved in oxidative responses or energy metabolism, other genes were also  
6 up-regulated upon exposure to all the different acid shocks (see *Supplementary material*).

7 Both strains showed to induce the expression of iron transporting and iron binding proteins,  
8 such as *feoB* (BC0709, BCE0783) and *dps* (BC2011, BC5044, BCE2092, BC5191).  
9 Furthermore, *mntH* (BC1803, BCE1960), encoding for manganese transport protein, also  
10 belonged to the group of up-regulated genes. Manganese and iron ions may play a role in  
11 oxidative stress response, conceivably acting as co-factors for superoxide dismutase proteins,  
12 and via other redox balancing mechanisms.

13

#### 14 *Inactivation associated radical formation*

15 The induction of oxidative stress associated genes and a recent publication that showed that  
16 hydroxyl radicals (OH $\cdot$ ) were formed upon exposure to bactericidal antibiotics in *Escherichia*  
17 *coli* and *Staphylococcus aureus* (Kohanski *et al.*, 2007), prompted us to investigate the  
18 formation of radicals upon low pH exposure of *B. cereus*. The formation of OH $\cdot$  and/or  
19 peroxyxynitrite (ONOO $\cdot$ ) in ATCC 14579 and ATCC 10987 cells was tested upon exposure to  
20 selected pHs (pH 5.4, pH 5.0, pH 4.8, and pH 4.5) at different intervals using the fluorescent  
21 probe 3'-(p-hydroxyphenyl) fluorescein (Fig. 6 and Fig. 7, respectively). Upon exposure to  
22 pH 4.5, ATCC 14579 was inactivated and this pH induced an increase of fluorescence  
23 indicating the formation of OH $\cdot$  and/or ONOO $\cdot$ . The exposure to the other pHs tested, i.e., pH  
24 5.4, 5.0, and 4.8, did not result in inactivation of the cells and also did not induce excess  
25 radical formation. Strain ATCC 10987 showed excess radical formation corresponding to the

1 inactivation observed at pH 5.0, pH 4.8 and pH 4.5. At pH 5.4, where this strain was able to  
2 resume growth, no excess radical formation was measured.

3 The formation of the oxygen derived radicals  $\text{OH}\cdot$  and  $\text{ONOO}^-$  should be prevented when  
4 oxygen is not available during acid exposure. Indeed, no  $\text{OH}\cdot$  and  $\text{ONOO}^-$  formation was  
5 observed in both strains anaerobically exposed to similar low pH values. Correspondingly,  
6 increased acid resistance was observed in both strains when exposed anaerobically (data now  
7 shown). In addition, the generation of superoxide, a key precursor in  $\text{OH}\cdot$  and  $\text{ONOO}^-$   
8 formation, was monitored upon low pH exposure using a superoxide-specific fluorescent  
9 probe. Detectable levels of superoxide were only found in aerobically exposed cells, and not  
10 in anaerobically exposed cells (Mols *et al.*, unpublished results). Taken together, these data  
11 provide evidence that the formation of hydroxyl radicals and/or peroxynitrite is associated  
12 with inactivation of *B. cereus* strains ATCC 14579 and ATCC 10987 exposed to low pH  
13 environments in the presence of oxygen.

14

## 15 **Discussion**

16 In this study, we describe the physiological and transcriptional responses of *Bacillus cereus*  
17 strains ATCC 14579 and ATCC 10987 to sub-lethal and lethal acid shocks. The two model  
18 strains were subjected to a range of pHs demonstrating that ATCC 14579 was more acid-  
19 resistant than ATCC 10987. ATCC 14579 survived acid conditions between pH 5.0 and pH  
20 4.7 without growth or inactivation in the first hour of exposure. However, a prolonged  
21 exposure of ATCC 14579 to pH 4.8 resulted in a decrease of viable cells. In contrast, ATCC  
22 10987 did not display this survival phenotype and was inactivated within the first hour of  
23 exposure to pHs lower than pH 5.0.

24 The exposure of *B. cereus* to sub-lethal and lethal acid stress resulted in distinct transcriptome  
25 profiles related to the physiological response displayed by the cultures. The concurrent

1 analysis of two strains thus enables for distinguishing between phenotype-specific, stress  
2 level-specific and strain-specific transcriptome responses. Furthermore, the approach used  
3 showed not only the well-studied responses to mild pHs, including the induction of several  
4 general stress response genes, but also the response to lethal levels of acidity, an issue that has  
5 up to now mostly been neglected, as exemplified in recent studies on mild acid stress response  
6 of *Bacillus subtilis* (Wilks *et al.*, 2009). Cotter and Hill (2003) have reviewed the response of  
7 Gram-positive organisms to mild levels of acidity and mechanisms of acid resistance were  
8 described for fermentative lactic acid bacteria and *L. monocytogenes*, including roles of  
9 proton pumps, regulators, altered metabolism, protein and DNA repair, cell envelope  
10 alterations and alkali production. Using two model strains of *B. cereus*, we have demonstrated  
11 that protein and DNA repair, stress related transcriptional regulators, altered metabolism and  
12 alkali production were indeed induced at low pH. In contrast to fermentative lactic acid  
13 bacteria, F<sub>1</sub>F<sub>0</sub>-ATPase was not up-regulated in these respiring *B. cereus* strains upon exposure  
14 to acid, indicating that *B. cereus* does not use F<sub>1</sub>F<sub>0</sub>-ATPase to extrude protons under the  
15 conditions tested. Down-regulation of F<sub>1</sub>F<sub>0</sub>-ATPase is best explained by the cells trying to  
16 prevent excessive inward flux of protons via this ATPase upon exposure to acid conditions.  
17 Furthermore, no indications were found in the transcriptome analyses for low pH-induced  
18 membrane damage or rearrangement of membrane composition. For example, our  
19 experiments did not show an induction of fatty acid biosynthesis (*fab* genes), as was shown  
20 for *B. subtilis* exposed to mild sorbic acid stress (Ter Beek *et al.*, 2008). Furthermore, Ter  
21 Beek and colleagues (2008) reported that a putative multidrug resistance (*mdr*) transporter  
22 was induced in *B. subtilis* exposed to mild sorbic acid stress and they proposed this  
23 transporter to export sorbate anions from the cell. Two homologous genes in *B. cereus* were  
24 up-regulated upon exposure to inorganic acid stress at pH 5.4 that were not induced in  
25 response to lethal pH exposures. Since there is no apparent connection with sorbic acid stress

1 and the induction of these putative *mdr* systems in *B. cereus*, their role in acid resistance of *B.*  
2 *cereus*, if any, remains to be elucidated.

3 The transcriptome analyses of the phenotypic responses to various levels of acidity revealed a  
4 major oxidative response. In bactericidal conditions, the oxidative response could be linked to  
5 the formation of  $\text{OH}\cdot$  and/or  $\text{ONOO}^-$  using flow cytometry in combination with the  
6 fluorescent probe HPF that specifically targets these reactive oxygen species. The observed  
7 oxidative burst in *B. cereus* may originate in a similar way as described for the formation of  
8  $\text{OH}\cdot$  radicals in *Escherichia coli* and *Staphylococcus aureus* upon exposure to bactericidal  
9 antibiotics in aerobic conditions (Kohanski *et al.*, 2007). The formation of  $\text{OH}\cdot$  and  $\text{ONOO}^-$   
10 was not observed in anaerobically acid-stressed *B. cereus* cells, and correspondingly,  
11 increased acid resistance was observed under these conditions for both strains (data not  
12 shown). Based on phenotype and transcriptome analyses we propose a model for acid-induced  
13 radical formation, including  $\text{OH}\cdot$  and  $\text{ONOO}^-$ , in *B. cereus* (Fig. 8). Acid stress may cause  
14 perturbation of the aerobic electron transfer chain (ETC) in *B. cereus* indicated by the  
15 differential expression of several genes potentially involved in ETC activity. This disturbance  
16 may cause premature leakage of electrons to oxygen leading to the formation of superoxide  
17 ( $\text{O}_2^-$ ). Indeed, elevated levels of  $\text{O}_2^-$  could be detected in *B. cereus* cells upon exposure to  
18 lethal levels of acidity as indicated by staining of these cells with a superoxide-specific  
19 fluorescent probe (Mols *et al.*, unpublished results). Furthermore, the formation of  $\text{O}_2^-$  can be  
20 inferred from the induction of superoxide dismutase and catalase genes. Iron-sulphur clusters  
21 may subsequently be damaged by  $\text{O}_2^-$  releasing iron in the cytoplasm (Imlay, 2006). Free iron  
22 can react with hydrogen peroxide, originating from the dismutation of  $\text{O}_2^-$ , forming the highly  
23 toxic  $\text{OH}\cdot$  radicals in the Fenton reaction (Imlay *et al.*, 1988). Furthermore,  $\text{O}_2^-$  can rapidly  
24 react with nitric oxide (NO) to form another highly toxic oxidative compound,  $\text{ONOO}^-$   
25 (Beckman and Koppenol, 1996). NO is formed by a reaction catalyzed by nitric oxide

1 synthase (bNOS). Indirect indications for the formation of NO upon low pH exposure can be  
2 inferred from the up-regulation of nitric oxide dioxygenase and nitric oxide dependant  
3 regulator *dnrN*. The induction of bNOS activity, which is possibly regulated at protein level  
4 (Shatalin *et al.*, 2008), may initially have a positive effect on surviving oxidative stress.  
5 bNOS-derived NO may inhibit thiol reduction leading to the inhibition of the OH· forming  
6 Fenton reaction (Gusarov and Nudler, 2005; Sudhamsu and Crane, 2009). Furthermore, NO  
7 induces catalase activity in *B. anthracis* (Shatalin *et al.*, 2008) and inhibits the aerobic ETC  
8 (Husain *et al.*, 2008). On the other hand, NO facilitates the formation of ONOO<sup>-</sup>, which may  
9 have a damaging effect that could lead to cell death. Nitric oxide dioxygenase and nitrite  
10 reductase are described to be possible NO dissipation routes (Payne *et al.*, 1997; Gardner,  
11 2005). The genome of ATCC 14579 encodes both mechanisms and this strain showed to be  
12 more acid resistant than the nitrite/nitrate reductase deficient ATCC 10987 strain.

13 The phenomenon that exposure to stresses such as salt, heat, acid, and bile, results in  
14 secondary oxidative stress, has been described earlier for *B. cereus* and numerous other  
15 bacteria (Aldsworth *et al.*, 1999; Clements *et al.*, 1999; Hecker and Volker, 2001; Airo *et al.*,  
16 2004; Banjerdkij *et al.*, 2005; Latifi *et al.*, 2005; Dodd *et al.*, 2007; Kim *et al.*, 2008), but up  
17 to now, this secondary oxidative stress response has not been linked to radical-associated cell  
18 death. Moreover, our findings are supported by earlier observations in amongst others *S.*  
19 *aureus* and *Vibrio vulnificus*, where acid resistance was found to be superoxide dismutase  
20 (and catalase) dependant (Clements and Foster, 1999; Kim *et al.*, 2005).

21 In conclusion, the results obtained in our study provide evidence for the origin of acid stress-  
22 induced oxidative stress. In extension to the antibiotic study of Kohanski and colleagues  
23 (2007), we now propose that in aerobic conditions, the formation of radicals such as OH· and  
24 ONOO<sup>-</sup> may be a common mechanism of cellular death in bacteria exposed to severe stress  
25 conditions.

1

## 2 **Experimental procedures**

3

### 4 *Bacterial strains and growth conditions*

5 *B. cereus* strains ATCC 14579 and ATCC 10987 were obtained from the American Type  
6 Culture Collection (ATCC). Stock cultures, grown in brain heart infusion (BHI, Becton  
7 Dickinson, France) broth, were stored at -80°C in 33% glycerol. To prepare pre-cultures, 10  
8 ml BHI in a 100 ml Erlenmeyer flask was inoculated with a droplet from the glycerol stock  
9 and incubated overnight at 30°C, with shaking at 200 rpm.

10 To study the effect of pH on *B. cereus* cells and the corresponding transcriptome profiles, 100  
11 ml BHI in a 500 ml Erlenmeyer flask was inoculated with 0.5 ml pre-culture and incubated at  
12 30°C, with shaking at 200 rpm. Upon reaching an optical density of 0.5 measured at 600 nm  
13 (OD, Novaspec II, Pharmacia Biotech, Germany), the culture pH was measured (PHM 240  
14 pH/ION Meter, Radiometer, Denmark) and serial dilutions were made in peptone  
15 physiological salt solution (PPS, 1g/l neutralized bacteriological peptone (Oxoid, England)  
16 and 8.5 g/l NaCl in water) and plated with a spiral-plater (Eddy Jet; IUL Instruments, Spain)  
17 on BHI agar plates (15 g/l bacteriological agar, Oxoid, England). 20 ml of the culture was  
18 used to extract RNA (sample t = 0). The remaining volume of the culture was acidified with  
19 hydrochloric acid (HCl 37%, Merck, Germany) to pH 5.4, 5.0, 4.8 or 4.5 and incubated at  
20 30°C, with shaking at 200 rpm. At designated time points (10, 30 and 60 minutes), samples  
21 were taken to measure the OD, to determine the viable counts, and to extract RNA.

22

23

### 24 *RNA isolation*



1 RNA isolation was performed by transferring 20 ml of the cultures into a 50-ml Falcon tube  
2 (Greiner Bio-one, Germany) at the designated time points. Subsequently, the cultures were  
3 pelleted at maximum speed at 4°C for 30 s (Eppendorf centrifuge 5804 R, Eppendorf,  
4 Germany). After decanting the supernatant, the cell pellets were frozen in liquid nitrogen.  
5 Within 10 min after freezing the cell pellets, 1 ml TRI-reagent (Ambion, United Kingdom)  
6 was added to the pellets. The samples were stored at -80°C until RNA extraction. RNA was  
7 extracted as described previously (van Schaik *et al.*, 2004). Residual chromosomal DNA was  
8 removed by treating the samples with DNA-free (Ambion, United Kingdom). The RNA  
9 concentration was measured in 2 ml cuvettes (UVettes, Eppendorf, Germany) with a  
10 BioPhotometer (Eppendorf, Germany) by determining the OD<sub>260</sub> and OD<sub>280</sub>. The quality of  
11 the RNA was monitored using the RNA 6000 Nano Assay (Agilent, United States) and the  
12 Agilent 2100 Bio-analyzer (Agilent, United States) according to the provided protocol. The  
13 extracted RNA samples were stored in 70% ethanol with 83 mM sodium acetate buffer (pH  
14 5.2) at -20°C.

15

#### 16 *cDNA synthesis, labelling and microarray hybridization and design*

17 Complementary DNA with amino-allyl-labelled dUTP (Ambion, United Kingdom) from the  
18 extracted RNA was prepared in reverse transcription reactions using Superscript III  
19 (Invitrogen, The Netherlands). Cy3 and Cy5 labelling of the cDNAs was performed with the  
20 CyScribe Post-Labeling kit (GE Healthcare, Belgium) as previously described (den Hengst *et*  
21 *al.*, 2005). The labelled cDNAs were purified using the CyScribe GFX purification kit (GE  
22 Healthcare) according to the provided protocol. To conduct the microarray hybridization, the  
23 Cy5-labelled cDNA samples were combined with the corresponding Cy3-labelled t0 reference  
24 samples (1:1 ratio). The microarray experiments for the comparison of the transcriptomes of  
25 the cultures exposed to various pHs were performed in two independent biological replicates,

1 where the replicate was performed with the dyes swapped. *B. cereus* ATCC 14579 and *B.*  
2 *cereus* ATCC 10987 microarrays (details below) were hybridized with 200 to 300 ng labelled  
3 cDNA following the 60-mer oligo microarray processing protocol (Agilent, United States).  
4 The microarrays used in this study were custom-made *B. cereus* ATCC 14579 and custom-  
5 made *B. cereus* ATCC 10987 microarrays developed by Agilent Technologies (United  
6 States). The *B. cereus* ATCC 14579 microarray design was based on the 11K platform of  
7 Agilent Technologies (GEO accession number GPL7680). A total of 10,262 spots represented  
8 5,131 chromosomal open reading frames, meaning that 98.0% of the predicted chromosomal  
9 open reading frames (NCBI accession number NC\_004722) are represented on the  
10 microarray. 99.6 % of the open reading frames for which probes could be designed were  
11 represented by two non-overlapping probes on the array. The remaining 0.4% of the open  
12 reading frames was represented by a single oligonucleotide spotted in duplicate on the array.  
13 The *B. cereus* ATCC 10987 microarray design was based on the 22K platform of Agilent  
14 Technologies (GEO accession number GPL7681). A total of 17,697 spots represented 5,578  
15 chromosomal open reading frames, 240 plasmid open reading frames and 81 putative small-  
16 RNAs, meaning that 99.6% of the predicted chromosomal and the plasmid open reading  
17 frames (NCBI accession numbers NC\_003909 and NC\_005707, respectively) were  
18 represented on the microarray. All features (chromosomal and plasmid open reading frames  
19 and small-RNAs) were represented by three individual spots. For 4,914 features three non-  
20 overlapping probes were designed, for 488 features two probes were designed (one probe  
21 spotted in duplicate) and for 497 only one oligonucleotide could be designed (one probe  
22 spotted in triplicate).  
23 After hybridization at 60°C for 17 hours, the microarrays were washed with 6 × SSC (0.9 M  
24 NaCl and 0.09 M sodium citrate) supplemented with 0.005% Triton X-102 at room

1 temperature for 10 min. Subsequently, the microarray slides were washed at 4°C with  
2 prechilled  $0.1 \times$  SSC with 0.005% Triton X-102 for 5 min and dried with nitrogen gas.

3

#### 4 *Microarray scanning and data analysis*

5 The microarray slides were scanned using an Agilent microarray scanner (G2565BA), and  
6 data were extracted from the scanned microarrays with Agilent's Feature Extraction software  
7 (version 8.1.1.1), which includes a LOWESS (locally weighted scatterplot smoothing)  
8 normalization step for the raw data. After removal of the data for the control spots, the  
9 normalized data for each spot from the microarrays were analyzed for statistical significance  
10 using the web-based VAMPIRE microarray suite (Hsiao *et al.*, 2005). A spot was found to be  
11 differentially expressed between two samples when the false discovery rate was smaller than  
12 0.05. Subsequently, the data for the single spots were integrated to obtain expression ratios for  
13 a corresponding feature (i.e., open reading frame or sRNA). A feature was found to be  
14 differentially expressed when all spots representing the feature were significantly  
15 differentially expressed between samples.

16 Hierarchical clustering (Eisen *et al.*, 1998) was performed per strain to identify groups of  
17 genes showing similar expression patterns. In Genemaths XT (version 1.6.1, Applied Maths,  
18 Belgium) genes that were significantly differentially expressed in one or more conditions  
19 were  $\log_2$  transformed and clustered using the complete linkage method and the Euclidian  
20 distance matrix. The groups identified from the hierarchical clustering were based on an  
21 arbitrary cut-off value. To identify relevant biological processes significantly overrepresented  
22 in a group, the genes corresponding to a group were analyzed using FIVA (Blom *et al.*, 2007).  
23 To visualize the relation between the different acidic conditions independent of the strains, the  
24  $\log_2$  transformed expression ratios from genes present on both microarrays were hierarchically

1 clustered using the average linkage method and the Euclidian distance matrix (Genemaths  
2 XT, 1.6.1).

3

#### 4 *Flow cytometry and radical measurements*

5 To detect radical formation, the fluorescent reporter dye 3'-(p-hydroxyphenyl) fluorescein  
6 (HPF, Invitrogen, The Netherlands) was used (Setsukinai *et al.*, 2003). At designated time  
7 points (0, 10, 30, 60 minutes) after adding HCl to the exponentially growing cultures  
8 ( $OD_{600nm} \sim 0.5$ ), samples were obtained by centrifuging ( $15,000 \times g$ , 30 s) 1 ml of culture and  
9 resuspension of the cell pellet in 1 ml filtered phosphate buffer saline (PBS). The samples  
10 were washed once and diluted with filtered PBS to obtain a concentration of approximately  
11  $10^6$  cells per ml, subsequently HPF was added at a final concentration of 5 mM. Samples were  
12 run on a Becton Dickinson FACSCalibur flow cytometer with the following photomultiplier  
13 tube (PMT) voltage settings: E00 (FSC), 360 (SSC) and 825 (FL1). Data were obtained from  
14 20,000 events (cells) at medium flow rate using Cellquest Pro (version 4.0.2), subsequently  
15 analyzed with WinMDI 2.9 (Joseph Trotter, Salk Institute for Biological Studies, La Jolla,  
16 California, USA; <http://facs.scripps.edu/software.html>) and graphically presented using  
17 Adobe Illustrator CS2 (version 12.0.1).

18

#### 19 **Acknowledgements**

20

21 We would like to thank Marcel H. Zwietering (Laboratory of Food Microbiology,  
22 Wageningen University, the Netherlands) for fruitful discussions and critical reading of the  
23 manuscript. We kindly acknowledge Menno van der Voort (Laboratory of Food  
24 Microbiology, Wageningen University, the Netherlands) for technical assistance regarding the  
25 microarray analyses.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**References**

Agata, N., Ohta, M., and Yokoyama, K. (2002) Production of *Bacillus cereus* emetic toxin (cereulide) in various foods. *Int J Food Microbiol* **73**: 23-27.

Airo, A., Chan, S.L., Martinez, Z., Platt, M.O., and Trent, J.D. (2004) Heat shock and cold shock in *Deinococcus radiodurans*. *Cell Biochem Biophys* **40**: 277-288.

Aldsworth, T.G., Sharman, R.L., and Dodd, C.E. (1999) Bacterial suicide through stress. *Cell Mol Life Sci* **56**: 378-383.

Banjerdkij, P., Vattanaviboon, P., and Mongkolsuk, S. (2005) Exposure to cadmium elevates expression of genes in the OxyR and OhrR regulons and induces cross-resistance to peroxide killing treatment in *Xanthomonas campestris*. *Appl Environ Microbiol* **71**: 1843-1849.

Beckman, J.S., and Koppenol, W.H. (1996) Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol* **271**: C1424-C1437.

Berg, G., Eberl, L., and Hartmann, A. (2005) The rhizosphere as a reservoir for opportunistic human pathogenic bacteria. *Environ Microbiol* **7**: 1673-1685.

Blom, E.J., Bosman, D.W., van Hijum, S.A., Breitling, R., Tijmsma, L., Silvis, R. *et al.* (2007) FIVA: Functional Information Viewer and Analyzer extracting biological knowledge from transcriptome data of prokaryotes. *Bioinformatics* **23**: 1161-1163.

Browne, N., and Dowds, B.C. (2002) Acid stress in the food pathogen *Bacillus cereus*. *J Appl Microbiol* **92**: 404-414.

Brul, S., and Coote, P. (1999) Preservative agents in foods. Mode of action and microbial resistance mechanisms. *Int J Food Microbiol* **50**: 1-17.

1 Choma, C., Guinebretiere, M.H., Carlin, F., Schmitt, P., Velge, P., Granum, P.E., and  
2 Nguyen-The, C. (2000) Prevalence, characterization and growth of *Bacillus cereus* in  
3 commercial cooked chilled foods containing vegetables. *J Appl Microbiol* **88**: 617-  
4 625.

5 Clements, M.O., and Foster, S.J. (1999) Stress resistance in *Staphylococcus aureus*. *Trends*  
6 *Microbiol* **7**: 458-462.

7 Clements, M.O., Watson, S.P., and Foster, S.J. (1999) Characterization of the major  
8 superoxide dismutase of *Staphylococcus aureus* and its role in starvation survival,  
9 stress resistance, and pathogenicity. *J Bacteriol* **181**: 3898-3903.

10 Cotter, P.D., and Hill, C. (2003) Surviving the acid test: responses of Gram-positive bacteria  
11 to low pH. *Microbiol Mol Biol Rev* **67**: 429-453.

12 Cotter, P.D., Gahan, C.G., and Hill, C. (2001) A glutamate decarboxylase system protects  
13 *Listeria monocytogenes* in gastric fluid. *Mol Microbiol* **40**: 465-475.

14 Curran, T.M., Lieou, J., and Marquis, R.E. (1995) Arginine deiminase system and acid  
15 adaptation of oral streptococci. *Appl Environ Microbiol* **61**: 4494-4496.

16 den Hengst, C.D., van Hijum, S.A., Geurts, J.M., Nauta, A., Kok, J., and Kuipers, O.P. (2005)  
17 The *Lactococcus lactis* CodY regulon: identification of a conserved cis-regulatory  
18 element. *J Biol Chem* **280**: 34332-34342.

19 Dodd, C.E., Richards, P.J., and Aldsworth, T.G. (2007) Suicide through stress: a bacterial  
20 response to sub-lethal injury in the food environment. *Int J Food Microbiol* **120**: 46-  
21 50.

22 Eisen, M.B., Spellman, P.T., Brown, P.O., and Botstein, D. (1998) Cluster analysis and  
23 display of genome-wide expression patterns. *Proc Natl Acad Sci U S A* **95**: 14863-  
24 14868.

1 Gardner, P.R. (2005) Nitric oxide dioxygenase function and mechanism of flavohemoglobin,  
2 hemoglobin, myoglobin and their associated reductases. *J Inorg Biochem* **99**: 247-266.

3 Granum, P.E., and Lund, T. (1997) *Bacillus cereus* and its food poisoning toxins. *FEMS*  
4 *Microbiol Lett* **157**: 223-228.

5 Gusarov, I., and Nudler, E. (2005) NO-mediated cytoprotection: instant adaptation to  
6 oxidative stress in bacteria. *Proc Natl Acad Sci U S A* **102**: 13855-13860.

7 Hecker, M., and Volker, U. (2001) General stress response of *Bacillus subtilis* and other  
8 bacteria. *Adv Microb Physiol* **44**: 35-91.

9 Hornstra, L.M., van der Voort, M., Wijnands, L.M., Roubos-van den Hil, P.J., and Abee, T.  
10 (2009) Role of germinant receptors in Caco-2 cell-initiated germination of *Bacillus*  
11 *cereus* ATCC 14579 endospores. *Appl Environ Microbiol* **75**: 1201-1203.

12 Hsiao, A., Ideker, T., Olefsky, J.M., and Subramaniam, S. (2005) VAMPIRE microarray  
13 suite: a web-based platform for the interpretation of gene expression data. *Nucleic*  
14 *Acids Res* **33**: W627-W632.

15 Husain, M., Bourret, T.J., McCollister, B.D., Jones-Carson, J., Laughlin, J., and Vazquez-  
16 Torres, A. (2008) Nitric oxide evokes an adaptive response to oxidative stress by  
17 arresting respiration. *J Biol Chem* **283**: 7682-7689.

18 Imlay, J.A. (2006) Iron-sulphur clusters and the problem with oxygen. *Mol Microbiol* **59**:  
19 1073-1082.

20 Imlay, J.A., Chin, S.M., and Linn, S. (1988) Toxic DNA damage by hydrogen peroxide  
21 through the Fenton reaction in vivo and in vitro. *Science* **240**: 640-642.

22 Ivanova, N., Sorokin, A., Anderson, I., Galleron, N., Candelon, B., Kapatral, V. *et al.* (2003)  
23 Genome sequence of *Bacillus cereus* and comparative analysis with *Bacillus*  
24 *anthracis*. *Nature* **423**: 87-91.

- 1 Jobin, M.P., Clavel, T., Carlin, F., and Schmitt, P. (2002) Acid tolerance response is low-pH  
2 and late-stationary growth phase inducible in *Bacillus cereus* TZ415. *Int J Food*  
3 *Microbiol* **79**: 65-73.
- 4 Kim, J.S., Sung, M.H., Kho, D.H., and Lee, J.K. (2005) Induction of manganese-containing  
5 superoxide dismutase is required for acid tolerance in *Vibrio vulnificus*. *J Bacteriol*  
6 **187**: 5984-5995.
- 7 Kim, Y.J., Moon, M.H., Song, J.Y., Smith, C.P., Hong, S.K., and Chang, Y.K. (2008) Acidic  
8 pH shock induces the expressions of a wide range of stress-response genes. *BMC*  
9 *Genomics* **9**: 604.
- 10 Kohanski, M.A., Dwyer, D.J., Hayete, B., Lawrence, C.A., and Collins, J.J. (2007) A  
11 common mechanism of cellular death induced by bactericidal antibiotics. *Cell* **130**:  
12 797-810.
- 13 Kotiranta, A., Lounatmaa, K., and Haapasalo, M. (2000) Epidemiology and pathogenesis of  
14 *Bacillus cereus* infections. *Microbes Infect* **2**: 189-198.
- 15 Latifi, A., Jeanjean, R., Lemeille, S., Havaux, M., and Zhang, C.C. (2005) Iron starvation  
16 leads to oxidative stress in *Anabaena* sp. strain PCC 7120. *J Bacteriol* **187**: 6596-  
17 6598.
- 18 Lemos, J.A., Chen, Y.Y., and Burne, R.A. (2001) Genetic and physiologic analysis of the  
19 *groE* operon and role of the HrcA repressor in stress gene regulation and acid  
20 tolerance in *Streptococcus mutans*. *J Bacteriol* **183**: 6074-6084.
- 21 Mols, M., and Abee, T. (2008) Role of ureolytic activity in *Bacillus cereus* nitrogen  
22 metabolism and acid survival. *Appl Environ Microbiol* **74**: 2370-2378.
- 23 Mols, M., de Been, M., Zwietering, M.H., Moezelaar, R., and Abee, T. (2007) Metabolic  
24 capacity of *Bacillus cereus* strains ATCC 14579 and ATCC 10987 interlinked with  
25 comparative genomics. *Environ Microbiol* **9**: 2933-2944.



- 1 Neumann, G., and Martinoia, E. (2002) Cluster roots--an underground adaptation for survival  
2 in extreme environments. *Trends Plant Sci* **7**: 162-167.
- 3 Payne, W.J., Liu, M.Y., Bursakov, S.A., and Le Gall, J. (1997) Microbial and plant  
4 metabolism of NO. *Biofactors* **6**: 47-52.
- 5 Rasko, D.A., Ravel, J., Okstad, O.A., Helgason, E., Cer, R.Z., Jiang, L. *et al.* (2004) The  
6 genome sequence of *Bacillus cereus* ATCC 10987 reveals metabolic adaptations and a  
7 large plasmid related to *Bacillus anthracis* pXO1. *Nucleic Acids Res* **32**: 977-988.
- 8 Richard, H.T., and Foster, J.W. (2003) Acid resistance in *Escherichia coli*. *Adv Appl*  
9 *Microbiol* **52**: 167-186.
- 10 Rosenquist, H., Smidt, L., Andersen, S.R., Jensen, G.B., and Wilcks, A. (2005) Occurrence  
11 and significance of *Bacillus cereus* and *Bacillus thuringiensis* in ready-to-eat food.  
12 *FEMS Microbiol Lett* **250**: 129-136.
- 13 Ryan, S., Hill, C., and Gahan, C.G. (2008) Acid stress responses in *Listeria monocytogenes*.  
14 *Adv Appl Microbiol* **65**: 67-91.
- 15 Ryan, S., Begley, M., Gahan, C.G., and Hill, C. (2009) Molecular characterization of the  
16 arginine deiminase system in *Listeria monocytogenes*: regulation and role in acid  
17 tolerance. *Environ Microbiol* **11**: 432-445.
- 18 Sanders, J.W., Leenhouts, K., Burghoorn, J., Brands, J.R., Venema, G., and Kok, J. (1998) A  
19 chloride-inducible acid resistance mechanism in *Lactococcus lactis* and its regulation.  
20 *Mol Microbiol* **27**: 299-310.
- 21 Setsukinai, K., Urano, Y., Kakinuma, K., Majima, H.J., and Nagano, T. (2003) Development  
22 of novel fluorescence probes that can reliably detect reactive oxygen species and  
23 distinguish specific species. *J Biol Chem* **278**: 3170-3175.

1 Shatalin, K., Gusarov, I., Avetissova, E., Shatalina, Y., McQuade, L.E., Lippard, S.J., and  
2 Nudler, E. (2008) *Bacillus anthracis*-derived nitric oxide is essential for pathogen  
3 virulence and survival in macrophages. *Proc Natl Acad Sci U S A* **105**: 1009-1013.

4 Stenfors Arnesen, L.P., Fagerlund, A., and Granum, P.E. (2008) From soil to gut: *Bacillus*  
5 *cereus* and its food poisoning toxins. *FEMS Microbiol Rev* **32**: 579-606.

6 Sudhamsu, J., and Crane, B.R. (2009) Bacterial nitric oxide synthases: what are they good  
7 for? *Trends Microbiol* **17**: 212-218.

8 Ter Beek, A., Keijser, B.J., Boorsma, A., Zakrzewska, A., Orij, R., Smits, G.J., and Brul, S.  
9 (2008) Transcriptome analysis of sorbic acid-stressed *Bacillus subtilis* reveals a  
10 nutrient limitation response and indicates plasma membrane remodeling. *J Bacteriol*  
11 **190**: 1751-1761.

12 Thomassin, S., Jobin, M.P., and Schmitt, P. (2006) The acid tolerance response of *Bacillus*  
13 *cereus* ATCC14579 is dependent on culture pH, growth rate and intracellular pH.  
14 *Arch Microbiol* **186**: 229-239.

15 van de Guchte, M., Serror, P., Chervaux, C., Smokvina, T., Ehrlich, S.D., and Maguin, E.  
16 (2002) Stress responses in lactic acid bacteria. *Antonie Van Leeuwenhoek* **82**: 187-216.

17 van Schaik, W., Tempelaars, M.H., Wouters, J.A., de Vos, W.M., and Abee, T. (2004) The  
18 alternative sigma factor sigma-B of *Bacillus cereus*: response to stress and role in heat  
19 adaptation. *J Bacteriol* **186**: 316-325.

20 Vilain, S., Luo, Y., Hildreth, M.B., and Brozel, V.S. (2006) Analysis of the life cycle of the  
21 soil saprophyte *Bacillus cereus* in liquid soil extract and in soil. *Appl Environ*  
22 *Microbiol* **72**: 4970-4977.

23 Von Stetten, F., Mayr, R., and Scherer, S. (1999) Climatic influence on mesophilic *Bacillus*  
24 *cereus* and psychrotolerant *Bacillus weihenstephanensis* populations in tropical,  
25 temperate and alpine soil. *Environ Microbiol* **1**: 503-515.

1 Wijnands, L.M., Dufrenne, J.B., van Leusden, F.M., and Abee, T. (2007) Germination of  
2 *Bacillus cereus* spores is induced by germinants from differentiated Caco-2 Cells, a  
3 human cell line mimicking the epithelial cells of the small intestine. *Appl Environ*  
4 *Microbiol* **73**: 5052-5054.

5 Wilks, J.C., Kitko, R.D., Cleeton, S.H., Lee, G.E., Ugwu, C.S., Jones, B.D. *et al.* (2009) Acid  
6 and base stress and transcriptomic responses in *Bacillus subtilis*. *Appl Environ*  
7 *Microbiol* **75**: 981-990.

8  
9

## 10 **Figure legends**

11

12 Fig. 1. Physiological response of *B. cereus* ATCC 14579 (filled squares) and ATCC 10987  
13 (open diamonds) upon exposure to a range of acidity levels. The colony forming units were  
14 determined after 0 and 60 minutes of exposure, the difference between the t0 and t60 is  
15 depicted. Data points represent single experiments, indicating a large variability of the  
16 responses of ATCC 10987 exposed to pH 5.0. The grey line at 0 corresponds to no growth  
17 (growth above the line) and no inactivation (inactivation below the line). The filled area  
18 between the dotted lines depicts the pH values where ATCC 14579 displayed no growth and  
19 no inactivation, i.e., survival, and where ATCC 10987 already showed to be inactivated.

20

21 Fig. 2. Physiological response of *B. cereus* ATCC 14579 (A) and ATCC 10987 (B) upon  
22 exposure to low pH. The colony forming units determined at different time points upon  
23 exposure to pH 5.4 (squares), pH 5.0 (diamonds), pH 4.8 (triangles) and pH 4.5 (circles) are  
24 depicted. At 0, 10, 30 and 60 minutes samples were taken for microarray analysis, indicated

1 with arrows and the error bars represent the standard deviation between duplicate  
2 experiments.

3

4 Fig. 3. Hierarchical clustering of the transcriptome profiles of different pH exposures based  
5 on the common genes of *B. cereus* ATCC 14579 and ATCC 10987. Samples with similar  
6 expression patterns were clustered using Euclidean distance and complete linkage. The  
7 corresponding phenotypic responses are shown at the right. Samples obtained from growing  
8 cultures are depicted in black, samples from non-growing (“survival” and “inactivation”  
9 phenotypes) cultures are shown in grey. Relative distance in similarity between the branches  
10 is shown at the top and bootstrap values are indicated at each branch.

11

12 Fig. 4. Average ratios of low pH and stress associated genes from *B. cereus* ATCC 14579  
13 (closed bars) and ATCC 10987 (open bars) upon exposure to pH 5.4 (A) and pH 4.5 (B). The  
14 global adaptive response is represented by *sigB* (BCE1086 and BC1004), *sigH* (BCE0093 and  
15 BC0114) and *codY* (BCE3869 and BC3826), encoding  $\sigma^B$ ,  $\sigma^H$  and CodY respectively.  
16 Additionally, *perR* (BCE0592 and BC0518), a major oxidative stress response regulator is  
17 shown. *atpA* (BCE5432 and BC5308), *napA* (BCE1729 and BC1612), and *nhaC* (BCE1840  
18 and BC1709) represent  $F_1F_0$ -ATPase and two proton antiporters. *gad* (glutamate  
19 decarboxylase, BCE2691), *arcA* (arginine deiminase, BCE0472 and BC0406), and *ureA*  
20 (urease, BCE3664) represent systems that are described to be involved in alkaline production.  
21 Glutamate decarboxylase and urease are specific for ATCC 10987 and are indicated with an  
22 asterisk. The general stress response chaperones and proteases are depicted by *dnaK*  
23 (BCE4395 and BC4312), *groES* (BCE0288 and BC0294), and *clpC* (BCE0079 and BC  
24 BC0100). Additionally, mechanisms involved in sorbic acid stress of *B. subtilis* (Ter Beek *et*  
25 *al.*, 2008), such as, multidrug transporters (*mdr1*, BCE4699 and BC4568 and *mdr2*, BCE1943

1 and BC1786), fatty acid biosynthesis (*fabF*, BCE1294 and BC1174), and branched-chain fatty  
2 acid biosynthesis (*bkdR*, BCE4239 and BC4165) are shown.

3

4 Fig. 5. Average ratios of selected genes associated with respiration and oxidative responses  
5 from *B. cereus* ATCC 14579 (closed bars) and ATCC 10987 (open bars) upon exposure to pH  
6 5.4 (A), pH 5.0 (B), pH 4.8 (C) and pH 4.5 (D). *ETC-4a* and *ETC-4b* depict two distinct types  
7 of complex IV of the aerobic electron transfer chain. Cytochrome C oxidase polypeptide I  
8 gene *ctaD* (BCE3990 and BC3943) represents *ETC-4a* and cytochrome D ubiquinol oxidase  
9 subunit I gene *cydA* (BCE4949 and BC4792) represent *ETC-4b*. The genomes of ATCC  
10 14579 and ATCC 10987 harbour four different superoxide dismutase genes and three  
11 different catalase genes, the data of *sodA* (BCE5579 and BC5445) and *katB* (BCE1261 and  
12 BC1155) are shown here. Nitric oxide synthase and nitric oxide dioxygenase are represented  
13 by the corresponding genes, *nos* (BCE5578 and BC5444) and *hmp* (BCE1571 and BC1448),  
14 respectively. Nitrate and nitrite reductases are encoded by multiple genes on the genome of  
15 ATCC 14579, therefore *narI* (BC2121), *nasD* (BC2136) and *nark* (BC2128) are shown. This  
16 nitrate and nitrite reductase cluster of ATCC 14579 is absent in ATCC 10987 and therefore  
17 indicated with an asterisk.

18

19 Fig. 6. Radical formation in *B. cereus* ATCC 14579 upon exposure to pH 5.4, pH 5.0, pH 4.8  
20 and 4.5. Samples were taken at 0 (green), 10 (light blue), 30 (blue) and 60 (dark blue)  
21 minutes. The pH and corresponding physiological response are indicated at each graph. The  
22 shift in fluorescent signal to the right indicates the formation of hydroxyl and/or peroxy nitrite  
23 radicals.

24

1 Fig. 7. Radical formation in *B. cereus* ATCC 10987 upon exposure to pH 5.4, pH 5.0, pH 4.8  
2 and 4.5. Samples were taken at 0 (green), 10 (light blue), 30 (blue) and 60 (dark blue)  
3 minutes. The pH and corresponding physiological response are indicated at each graph. The  
4 shift in fluorescent signal to the right indicates the formation of hydroxyl and/or peroxynitrite  
5 radicals.

6

7 Fig. 8. Low pH induced oxidative stress response and radical forming mechanisms in *B.*  
8 *cereus* ATCC 14579 and ATCC 10987. Schematic representation of radical formation  
9 conceivably induced upon exposure to lethal acid stress. Acid stress may cause perturbation  
10 of the electron transfer chain and an excess of superoxide radicals ( $O_2^-$ ) may be formed.  
11 Superoxide radicals can be converted to hydrogen peroxide and water by superoxide  
12 dismutase (*sod*) and catalase (*kat*). However, when the capacity to dismutate superoxide is not  
13 sufficient, free superoxide radicals can cause damage to iron-sulphur (Fe-S) cluster containing  
14 enzymes supplying unbound iron ions. These free iron ions and hydrogen peroxide can react  
15 (Fenton reaction) and produce hydroxyl radicals ( $OH\cdot$ ). Another possible route in forming  
16 highly damaging radicals may occur via nitric oxide. Nitric oxide can be formed by nitric  
17 oxide synthase (*nos*) and can react with superoxide radicals to form peroxynitrite ( $ONOO^-$ ).  
18 Nitric oxide can be converted to nitrate by nitric oxide dioxygenase (*hmp*). Subsequently,  
19 nitrate can be converted to nitrite and ammonium by nitrate (*nar*) and nitrite (*nas*) reductase,  
20 respectively. Nitrite can also be transported outside the bacterial cell by a Nitrite extrusion  
21 protein (*nark*). *nar*, *nas* and *narK* are ATCC 14579 specific, the reactions they catalyze are  
22 indicated with dotted lines.

23

24 **Supplementary material**

25

1 The following supplementary material is available for this article online:

2 **14579\_Allgroups.tar.** A compressed folder containing the results of the hierarchical  
3 clustering of the significant differentially expressed genes of strain ATCC 14579. Opening the  
4 `htmlOutputFiva.html` file shows the graphical output of all used annotation modules and can  
5 be used to navigate through the results. By clicking the cluster names, the corresponding  
6 transcriptome profiles of the groups are revealed. The size of each group is displayed in blue  
7 underneath the group name and by clicking the list of genes will appear. Numbers in each  
8 rectangle represent absolute values of occurrences. The significance of occurrences is  
9 visualized in a colour gradient and with symbols which are displayed at the bottom of the plot.  
10 On the left the categories showing over representation in one or more groups are listed with  
11 the corresponding size. Clicking the categories will reveal the genes present in the  
12 corresponding category per group. The description of each category is placed at the right and  
13 more information about the categories can be displayed by clicking the description.


14 **10987\_Allgroups.tar.** A compressed folder containing the results of the hierarchical  
15 clustering of the significant differentially expressed genes of strain ATCC 10987. A detailed  
16 description is given above.

17 **Supplementary results.pdf.** Supplementary results based on the microarray data obtained for  
18 *B. cereus* strains ATCC 14579 and ATCC 10987 exposed to pH 5.4, pH 5.0, pH 4.8 and pH  
19 4.5. The data of each strain was clustered hierarchically and different groups of genes were  
20 identified based on their common transcriptome profile. The FIVA analysis showed that  
21 several function categories were overrepresented in certain groups.

22

23 Raw and processed microarray data are available for reviewers following the link to the GEO  
24 database below.


25 <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?token=hfsfreqageqakzi&acc=GSE13773>



1

2 Fig 1

3




4

5 Fig 2

6






1

2 Fig 3


3



1

2 Fig 4


3



4

5 Fig 5


6



1

2 Fig 6


3



4

1 Fig 7

2



3

4 Fig 8