

Cytochromes c_{550} , c_{552} , and c_1 in the Electron Transport Network of *Paracoccus denitrificans*: Redundant or Subtly Different in Function?

MARIJKE F. OTTEN,¹ JOHN VAN DER OOST,² WILLEM N. M. REIJNDERS,¹ HANS V. WESTERHOFF,^{1,3}
BERND LUDWIG,⁴ AND ROB J. M. VAN SPANNING^{1*}

*Department of Molecular Cell Physiology, Faculty of Biology, BioCentrum Amsterdam, Free University, 1081 HV Amsterdam,¹
Department of Microbiology, Wageningen University, 6703 CT Wageningen,² and Department of Mathematical Biochemistry,
Swammerdam Institute of Life Sciences, BioCentrum Amsterdam, University of Amsterdam, 1018 TV Amsterdam,³
The Netherlands, and Biozentrum der Universitaet, D-60439 Frankfurt, Germany⁴*

Received 24 July 2001/Accepted 19 September 2001

Paracoccus denitrificans strains with mutations in the genes encoding the cytochrome c_{550} , c_{552} , or c_1 and in combinations of these genes were constructed, and their growth characteristics were determined. Each mutant was able to grow heterotrophically with succinate as the carbon and free-energy source, although their specific growth rates and maximum cell numbers fell variably behind those of the wild type. Maximum cell numbers and rates of growth were also reduced when these strains were grown with methylamine as the sole free-energy source, with the triple cytochrome c mutant failing to grow on this substrate. Under anaerobic conditions in the presence of nitrate, none of the mutant strains lacking the cytochrome bc_1 complex reduced nitrite, which is cytotoxic and accumulated in the medium. The cytochrome c_{550} -deficient mutant did denitrify provided copper was present. The cytochrome c_{552} mutation had no apparent effect on the denitrifying potential of the mutant cells. The studies show that the cytochromes c have multiple tasks in electron transfer. The cytochrome bc_1 complex is the electron acceptor of the Q-pool and of amicyanin. It is also the electron donor to cytochromes c_{550} and c_{552} and to the cbb_3 -type oxidase. Cytochrome c_{552} is an electron acceptor both of the cytochrome bc_1 complex and of amicyanin, as well as a dedicated electron donor to the aa_3 -type oxidase. Cytochrome c_{550} can accept electrons from the cytochrome bc_1 complex and from amicyanin, whereas it is also the electron donor to both cytochrome c oxidases and to at least the nitrite reductase during denitrification. Deletion of the c -type cytochromes also affected the concentrations of remaining cytochromes c , suggesting that the organism is plastic in that it adjusts its infrastructure in response to signals derived from changed electron transfer routes.

Many bacteria that live in soil, sewage or sludge possess a genome that encodes a highly branched respiratory network. The encoded network comprises various types of dehydrogenase and of terminal oxidoreductase, which are capable of electron input and output flow, respectively, during oxidative phosphorylation (4). Such versatility allows these bacteria to use a variety of electron donors and terminal electron acceptors once they become available in their natural habitats. The genes or gene clusters that encode the different redox enzymes are usually tightly regulated. They are expressed only under the growth conditions at which they are required. One of the best-studied organisms in that respect is *Paracoccus denitrificans*. This gram-negative bacterium is able to grow heterotrophically with a variety of organic carbon and free-energy sources, as well as autotrophically, by using hydrogen, or so-called C1 substrates such as methylamine or methanol, as free-energy sources (2, 3, 17, 41, 42, 46). The electron transport chain used for aerobic heterotrophic growth possesses a full complement of proteins with counterparts in the mitochondrial respiratory chain (20, 41). Upon depletion of the organic substrates, the bacterium induces hydrogenase and methanol or methylamine dehydrogenases, as well as their dedicated elec-

tron acceptors, provided that their substrates are present (16, 49).

P. denitrificans is able to reduce molecular oxygen at a wide range of oxygen concentrations. This flexibility is the result of its potential to synthesize three types of terminal oxidase (9, 10, 32, 33, 35), which all belong to the superfamily of heme copper oxidases (15, 38, 44). The one that is most expressed at atmospheric oxygen concentrations is the aa_3 -type cytochrome c oxidase, which has a relatively low affinity for oxygen (31, 36). Its expression level decreases with decreasing oxygen concentrations (5). The second type of oxidase is the cbb_3 -type cytochrome c oxidase, which has a relatively high affinity for oxygen and which is increasingly synthesized at decreasing oxygen concentrations (10, 31). The third type of oxidase is a ba_3 -type quinol oxidase, which is the counterpart of the bo_3 -type quinol oxidase found in *Escherichia coli* (8, 35). The ba_3 -type oxidase receives electrons from ubiquinol, is expressed and increasingly active under conditions that give rise to high reduction levels of the Q-pool, and apparently serves to prevent that by rapidly transferring electrons from ubiquinol to oxygen (27).

P. denitrificans is also able to synthesize four types of N-oxide oxidoreductase, which catalyze the sequential reduction of nitrate to dinitrogen gas via the intermediates nitrite, nitric oxide, and nitrous oxide. This denitrification process takes place at nearly anoxic conditions, during which nitrate substitutes for oxygen as a terminal electron acceptor. Nitrate, ni-

* Corresponding author. Mailing address: Department of Molecular Cell Physiology, Faculty of Biology, Free University, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands. Phone: 31-20-4447179. Fax: 31-20-4447229. E-mail: spanning@bio.vu.nl.

TABLE 1. Strains and plasmids

| Strain or plasmid | Relevant genotype ^a | Source or reference |
|-------------------------|---|---------------------|
| Bacteria | | |
| <i>E. coli</i> | | |
| TG1 | <i>supE hsdD5 thi D(lac-proAB) F' (traD36 proAB lacI^q lacZΔM15)</i> | 37 |
| S17-1 | <i>Sm^r pro r⁻ m⁺ RP4-2 integrated (Tc::Mu) (Km^r::Tn7)</i> | 39 |
| HB101 | <i>F⁻ hsdS20 (r_B⁻ m_B⁻) lacY1 proA2 recA13</i> | 6 |
| <i>P. denitrificans</i> | | |
| Pd1222 | DSM413 derivative, Rif ^r , enhanced conjugation frequencies, m ⁻ | 13 |
| Pd21.21 | Pd1222 derivative, <i>cycA</i> ::Km ^r | 11 |
| Pd21.31 | Pd21.21 derivative, Δ <i>cycA</i> | 11 |
| Pd28.22 | Pd1222 derivative, <i>cycM</i> ::Km ^r | This study |
| Pd24.21 | Pd1222 derivative, <i>fbcC</i> ::Km ^r | 11 |
| Pd24.31 | Pd24.21 derivative, Δ <i>fbcC</i> | This study |
| Pd92.06 | Pd21.31 derivative, Δ <i>cycA fbcC</i> ::Km ^r | 11 |
| Pd92.07 | Pd92.06 derivative, Δ <i>cycA ΔfbcC</i> | This study |
| Pd92.26 | Pd21.31 derivative, Δ <i>cycA cycM</i> ::Km ^r | This study |
| Pd92.28 | Pd24.31 derivative, Δ <i>fbcC cycM</i> ::Km ^r | This study |
| Pd93.14 | Pd92.07 derivative, Δ <i>cycA ΔfbcC cycM</i> ::Km ^r | This study |
| Pd93.12 | Pd1222 derivative, Δ <i>ctaDI ΔctaDII ccoN</i> ::Km ^r | 10 |
| Pd93.13 | Pd1222 derivative, Δ <i>cycA ΔctaDII cycM</i> ::Km ^r | This study |
| Plasmids | | |
| pGRPd1 | <i>oriV (colE1) Amp^r oriT Sm^r (Tn1831)</i> | 51 |
| pRVS1 | pGRPd1 derivative, PTn5 <i>lacZ</i> | 52 |
| pMP190 | pRK2 derivative, mcs, promoterless <i>lacZ</i> | 40 |
| pBL250 | pBR322 derivative, <i>fbcC</i> | 22 |
| PRVS2 | pBSII (KS) derivative, <i>oriT</i> (RK2), Sm ^r | 48 |
| pAT3 | pBSII (KS) derivative, <i>cycM</i> | 43 |
| pAT8 | pAT3 derivative, <i>cycM</i> ::Km ^r | 43 |
| pRTd28.22 | pRVS2 derivative, <i>cycM</i> ::Km ^r | This study |
| pRTd24.31 | pRVS1 derivative, Δ <i>fbcC</i> | This study |

^a P, promoter. Sm^r, streptomycin resistance.

trite, nitric oxide, and nitrous oxide reductases are expressed when nitrate is present and oxygen is virtually absent (2, 3, 14, 42).

Much of our knowledge on how electrons are transferred from the dehydrogenases to the terminal oxidoreductases stems from the analyses of mutants with mutations in genes encoding electron carriers that make up the respiratory network (9, 11, 50, 51, 52, 53). However, it has thus far been difficult to understand the exact role of the different types of cytochrome *c* in situ since mutations in their corresponding genes did not give clear phenotypes if at all. Apparently, cytochromes *c* or other small redox carriers can substitute for each other. The present study sought to elucidate the roles of the cytochromes *c*₅₅₀ and *c*₅₅₂ and the cytochrome *bc*₁ complex in electron transport in *P. denitrificans*. The approach was to isolate *P. denitrificans* strains with mutations in the genes encoding these three types of cytochrome *c* and in combinations of these genes in order to study the resultant effects of these mutations on their growth characteristics.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. The strains and plasmids used in this study are listed in Table 1. *E. coli* strains were cultivated in yeast-tryptone medium at 37°C. *P. denitrificans* strains were grown at 34°C in batch cultures either aerobically, i.e., in 2-liter flasks filled with 300 ml of medium and shaken vigorously; semiaerobically, i.e., in 1-liter flasks filled with 500 ml of medium and slowly shaken; or anaerobically, i.e., in flasks almost completely filled with medium and not shaken. The *P. denitrificans* strains were grown on

mineral batch medium containing Lawford trace solution and methylamine (100 mM), succinate (25 mM), or butyrate (25 mM) as carbon and free-energy sources at pH 7.0 (7, 24). Anaerobic cultures were supplemented with 100 mM potassium nitrate as an electron acceptor. Inhibition of the cytochrome *bc*₁ complex was achieved by addition of 5 μM myxothiazol to the cultures (24). Cells were harvested in the mid-exponential phase of growth when the turbidity at 660 nm was ca. 1.0 cm⁻¹. When appropriate, antibiotics were added, i.e., rifampin (40 mg liter⁻¹), streptomycin (50 mg liter⁻¹), kanamycin (50 mg liter⁻¹), or ampicillin (100 mg liter⁻¹). Residual nitrite in denitrifying cultures was determined by using nitrite strips from Boehringer.

Isolation of intact cells and membrane fractions. Cells were washed twice in 0.1 M of Tris (Tris hydroxymethyl aminomethaan)-HCl at pH 7.5 at 4°C and then suspended in the same buffer to a turbidity of 50 at 660 nm. After the addition of MnCl₂ to 10 mM and DNase I to 2 mg ml⁻¹, cell extracts were obtained by using a French pressure cell (American Instrument Company, Silver Spring, Md.) at 10,000 lb/in². Cell debris was spun down for 5 min at 17,000 × *g*, and the resulting supernatant was centrifuged at 438,000 × *g* for 30 min at 4°C in order to separate the soluble and membrane fractions. The pellet containing the membrane fraction was resuspended in 0.1 M Tris-HCl at pH 7.5. Protein concentrations were determined by using the BCA Kit (Pierce) with bovine serum albumin as a standard. Samples (at 10 mg of protein ml⁻¹) were routinely stored at -80°C.

PAGE and heme staining. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was carried out by using the Bio-Rad Mini-Protein II Gel System with 13% slab gels (23). The samples (100 μg of protein each) were diluted in sample buffer (60 mM Tris-HCl, pH 6.8; 25% glycerol [vol/vol]; 2% SDS; 50 mM 2-mercaptoethanol; 50 mM dithiothreitol) and incubated at room temperature for 10 min. The heme-binding proteins were detected by using chemiluminescent substrate "Lumi-Light Plus" (Boehringer Mannheim) on a high-performance chemiluminescence film (Amersham) (29). The program NIH-Image/ppc 1.56b61 was used for densitometric analyses.

Spectral analyses. Spectra of dithionite (5 mg/ml) reduced cell suspensions (turbidity at 660 nm of 50) were recorded in the dual-wavelength mode of an

automated Aminco/SLM DW-2 UV/Vis spectrophotometer with the reference set at 578 nm. During kinetic measurements, the reduction level of cytochromes *c* as a function of time was recorded at 552 nm with the reference set at 578 nm. Cuvettes were thermostated at 20°C. Oxygen was released from 20 μ l of 0.3% hydrogen peroxide by intracellular catalase activity. Dithionite was added to a final concentration of 5 mg/ml.

Oxygen consumption. Oxygen consumption rates of cells were measured polarographically with a Clark-type oxygen electrode (Yellow Springs, Ohio) in 1 ml of Tris-HCl (pH 7.5) at 25°C. Succinate (10 mM) was added as an electron donor. Where indicated, the electron flow via cytochrome *bc*₁ was inhibited by adding antimycin A (6 μ M) and myxothiazol (6 μ M).

DNA manipulations. Routine cloning procedures were performed according to standard protocols (1). The construction of marked and unmarked mutations in chromosomal genes is described in Results and was performed as described earlier (52). *E. coli* TG1 was used routinely for cloning procedures. *E. coli* HB101(pRK2020) was used as the helper strain in the conjugative transfer of plasmids via triparental mating.

Construction of cytochrome *c* single and multiple mutant strains. The following strains were constructed in an earlier study (11): Pd21.21 (*cycA::Km^r*) with an interrupted cytochrome *c*₅₅₀ encoding gene, Pd21.31 (*Δ cycA*) with an unmarked mutation in the cytochrome *c*₅₅₀ encoding gene, Pd24.21 (*fbcC::Km^r*) with an interrupted cytochrome *c*₁ encoding gene, and the double mutant strain Pd92.06 (*Δ cycA fbcC::Km^r*). In this study, an internal part of the chromosomal *fbcC* gene that contained the gene cartridge encoding kanamycin resistance (*Km^r*) in strains Pd24.21 and Pd92.06 was removed in order to make the *fbcC* mutations in these strains unmarked. For this, plasmid pBL250, which is a pBR322 derivative containing the *fbcC* gene and flanking regions (22), was cut with *Xho*I and *Sal*I and then religated. As a result, a 1,060-bp fragment within the *fbcC* gene was removed. The fragment with the mutated gene was then cloned in suicide vector pRVS1, yielding plasmid pRTd24.31. This plasmid was used in the gene exchange technique described earlier (52) in order to isolate strains Pd24.31 (*Δ fbcC*) and Pd92.07 (*Δ cycA Δ fbcC*), which have unmarked mutations in their chromosomal *fbcC* genes. In this gene exchange technique, mutants with unmarked mutations are selected by using the *E. coli lacZ* gene as a suicide gene (52). Plasmid pAT8 harbors the cytochrome *c*₅₅₂ encoding *cycM* gene in which the gene cartridge encoding kanamycin resistance was inserted in the internal *Stu*I site (43). A 4.6-kb *Not*I fragment of this plasmid, which contains the mutated *cycM* gene, was cloned in suicide vector pRVS2, yielding vector pRTd28.22. This plasmid was used in the gene exchange technique described earlier (51) in order to isolate strains Pd28.22 (*cycM::Km^r*), Pd92.26 (*Δ cycA cycM::Km^r*), Pd92.28 (*Δ fbcC cycM::Km^r*), and Pd93.14 (*Δ cycA Δ fbcC cycM::Km^r*), which all have marked mutations in their chromosomal *cycM* genes. The correctnesses of the mutations were confirmed by Southern analyses of their chromosomal DNA. As a result of these constructions, we had mutant strains with mutations in one, two, or three of the genes encoding cytochromes *c*₅₅₀, *c*₅₅₂, and *c*₁.

RESULTS

Spectral analyses of the cytochrome *c* single and multiple mutant strains. *P. denitrificans* wild-type and mutants with single and multiple mutations in the *cycA*, *cycM*, and *fbcC* genes were cultured aerobically in mineral medium containing succinate as the sole carbon and free-energy source. Cell suspensions from these cultures exhibited composite α -bands of ca. 555 nm at which cytochromes *b* (at ca. 560 nm) and *c* (at ca. 550 nm) have their maximum absorbance (Fig. 1). It should be noted that the absorbance at that wavelength reflects not only all three cytochromes *c* that are important here, i.e., cytochromes *c*₅₅₀, *c*₅₅₂, and *c*₁, but also further contributions from other cytochromes *c* as well. Both the shape and the height of that composite band differed between the strains. Only the cytochrome *c*₅₅₂-deficient mutant displayed a spectrum very similar to that of the wild-type strain. Paradoxically, strains Pd24.31 (*c*₁ deficient) and Pd92.28 (*c*₅₅₂/*c*₁ deficient) had a higher cytochrome *c* peak than the wild-type strain, whereas that peak is lower in strains Pd21.31 and Pd92.26, which are cytochrome *c*₅₅₀ and cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutants, respectively. A sharp decrease in the height of that band is

observed in the spectra of strains Pd92.07 and Pd93.14, which are cytochrome *c*₅₅₀/*c*₁- and cytochrome *c*₅₅₀/*c*₅₅₂/*c*₁-deficient mutants. Another remarkable observation is that the peak at ca. 605 nm where hemes *a* of cytochrome *aa*₃ have their maximum absorbance is decreased to ca. 20% of the wild-type signal in the spectra of strains Pd21.31 (cytochrome *c*₅₅₀ deficient), Pd92.26 (cytochrome *c*₅₅₀/*c*₅₅₂ deficient), and Pd93.14 (cytochrome *c*₅₅₀/*c*₅₅₂/*c*₁ deficient).

In order to study the effect of the oxygen concentration on the spectral properties of the mutant cells, the spectra of cells grown semiaerobically were also recorded (results not shown). These spectra show that the cytochrome *c* peak of strain Pd24.31 (cytochrome *c*₁ deficient) is ca. four to five times higher than that of the wild-type strain and that the cytochrome *b* peak is also slightly increased. The cytochrome *c*₅₅₀/*c*₁-deficient double-mutant strain, in contrast, showed a decrease in both the cytochrome *c* and cytochrome *b* absorption bands. Apparently, the cytochrome *c* mutations, as well as the oxygen concentration, affect the concentrations of cytochromes *c* other than the ones that were mutated. The consequences of oxygen deprivation, as well as of the cytochrome *c* mutations, include reduced electron flow through the respiratory system, resulting in reduction of its constituent components. Perhaps *P. denitrificans* possesses a signal transduction mechanism that governs the expression of cytochrome *c* genes in response to that altered redox state. In order to test that hypothesis, *P. denitrificans* wild-type cells were cultured aerobically under different conditions, between which we expected the reduction levels of the redox enzymes to differ. In one approach, *P. denitrificans* wild-type cells were grown aerobically in mineral medium with either succinate or butyrate as the sole carbon and energy source. Since butyrate is a more reduced substrate than succinate, one would expect a rise in the electron-input rates and thus higher reduction states of the redox enzymes in butyrate-grown cells compared to succinate-grown cells. The cytochrome *c* peak of butyrate-grown cells was almost five times higher than that of the succinate-grown cells (results not shown). In another approach, *P. denitrificans* wild-type cells were cultured aerobically in mineral medium with succinate as the sole carbon and free-energy source and in the presence of increasing concentrations of myxothiazol, which is an inhibitor of the cytochrome *bc*₁ complex. The growth rates of these cultures did not differ significantly from one another, indicating that the myxothiazol-treated cells had sufficiently adapted their mode of respiration in response to the presence of the inhibitor. Spectral analyses of these cultures showed an increase in the cytochrome *c* peak with increasing concentrations of myxothiazol ultimately resulting in a more than fivefold increase of the cytochrome *c* peak at a concentration of 5 μ M myxothiazol (results not shown).

PAGE and heme staining of cytochrome *c* single and multiple mutant strains. In order to study the cytochrome *c* composition of the mutants in more detail, cells from cultures grown aerobically on succinate were broken and centrifuged to yield soluble and membrane fractions. For each fraction, the proteins were separated by PAGE, after which the cytochromes *c* were visualized by a heme staining approach. Two important conclusions can be drawn from the results (Fig. 2). First, they confirm that the mutations in the *cycA*, *cycM*, and *fbcC* genes of the mutants had resulted in the inability to




FIG. 1. Dithionite-reduced absorption spectra at room temperature of cell suspensions from the wild-type strain (Pd1222 [C]) and from mutant strains deficient in cytochrome c_1 (Pd24.31 [A]), cytochromes c_{552}/c_1 (Pd92.28 [B]), cytochrome c_{552} (Pd28.22 [D]), cytochromes c_{550}/c_1 (Pd92.07 [E]), cytochrome c_{550} (Pd21.31 [F]), cytochromes c_{550}/c_{552} (Pd92.26 [G]), or cytochromes $c_{550}/c_{552}/c_1$ (Pd93.14 [H]). The cells had been grown aerobically to their mid-exponential phase of growth (optical density at 660 nm [OD_{660}] = $\sim 1.0 \text{ cm}^{-1}$) in batch cultures with succinate as the carbon and free-energy source.

express the corresponding cytochromes c as judged by the absence of heme-stained proteins of 14 kDa (cytochrome c_{550}) in the periplasmic fractions (Fig. 2A) and of 22 kDa (cytochrome c_{552}) and ca. 60 kDa (cytochrome c_1) in the membrane fractions (Fig. 2B) of the corresponding mutants. The faint band at ca. 14 kDa observed in the lane of the cytochrome c_{550}/c_{552} -deficient double mutant in Fig. 2A probably originates from a cytochrome c different from cytochrome c_{550} and which is specifically expressed in this mutant. This band is not present in the lane of its parent strain Pd21.31 (Fig. 2A, lane 2). We monitored the staining of the cytochromes c present in that part of the gel over time (Fig. 3) and noticed that cytochrome c_{550} stained much faster than the unknown one (maximally stained after 1 min and after 15 min, respectively). The second conclusion is that the knockout of cytochromes c_{550} , c_{552} , and bc_1 singly and in various combinations had resulted in the increased expression of other types of cytochrome c . The periplasmic fractions of the wild-type strain and the cytochrome c_{552} -deficient strain contained moderate levels of cytochrome c_{550} , but its concentration was more than 10-fold increased in the cytochrome c_1 - and cytochrome c_{552}/c_1 -deficient mutant strains. In addition, all of the strains that lacked

the cytochrome bc_1 complex induced high levels of cytochrome c peroxidase (45 kDa), as well as two cytochromes c of ca. 25 and 35 kDa. The 35-kDa cytochrome c was abundant in the cytochrome c_{550}/c_{552} -deficient mutant strain. Since this abundance correlated with the appearance of the unknown cytochrome c of 14 kDa, the latter might be a heme containing breakdown product of the former. In contrast to the cytochrome bc_1 mutants, the cytochrome c_{550}/c_{552} -deficient mutant strain did not show a significantly increased induction of cytochrome c peroxidase nor of the 25-kDa cytochrome c . This correlation could indicate that the 25-kDa cytochrome c was a heme-containing breakdown product of cytochrome c peroxidase. Inspection of the cytochromes c in the membrane fraction indicated that the concentrations of the cytochrome bc_1 complex and of cytochrome c_{552} in the mutants still able to express them were not significantly different from those of the wild-type strain. In contrast, the intensities of the bands containing the cytochrome c containing subunits of the cbb_3 -type oxidase, CcoO and, to a lesser extent, CcoP, were up to five times higher in the cytochrome bc_1 mutants compared to the wild-type strain.




FIG. 2. SDS-PAGE and subsequent heme staining of cytochromes *c* in the soluble fractions (A) and membrane fractions (B) of *P. denitrificans* wild-type (lanes 1) and mutant strains deficient in cytochrome c_{550} (lanes 2), cytochrome c_{552} (lanes 3), cytochrome c_1 (lanes 4), cytochromes c_{550}/c_1 (lanes 5), cytochromes c_{550}/c_{552} (lanes 6), cytochromes c_{552}/c_1 (lanes 7), and cytochromes $c_{550}/c_{552}/c_1$ (lanes 8). Growth and harvesting conditions were as shown in the legend of Fig. 1. The fractions, containing 100 μ g of protein each, were obtained from one set of experiments and analyzed in parallel. The relative concentrations of heme *c*-containing proteins within each gel were determined by densitometric analyses. The *c*-type cytochromes indicated are cytochromes c_{550} and c_{552} (CycA and CycM), cytochrome *c* peroxidase (Ccp), subunits of the *cbb*₃-type oxidase (CcoO and CcoP), and cytochrome c_1 (FbcC). Sizes of the molecular mass markers are indicated on the right of the gels in kilodaltons.

Growth characteristics of the cytochrome *c* mutants. *P. denitrificans* wild-type strain and the cytochrome c_{550} , c_{552} , and bc_1 single and multiple mutants were grown in mineral medium under three different growth conditions. The increase in cell density with time was determined, as well as their final turbidity at 660 nm. The growth conditions were (i) aerobic with succinate as the sole electron donor, (ii) aerobic with methylamine as the sole electron donor, and (iii) anaerobic with succinate as the electron donor and nitrate as the electron acceptor. The data were used to estimate the maximum specific growth rates and the maximum cell numbers (as determined from turbidity measurements) of the strains. These values are presented in Table 2.

The maximum specific growth rates of succinate-grown cells did not differ significantly between the strains. The final turbidities of the mutant cultures, however, were all ca. 10% less compared to that of the wild type. The only exception was the

cytochrome c_{552} -deficient mutant, which had wild-type growth characteristics.

More-dramatic differences between the strains were observed when they were cultured in medium with methylamine as the sole free-energy source. Except for the cytochrome c_{552} -deficient mutant, all of the strains had decreased maximum specific growth rates and final turbidities compared to the wild-type strain. The most profound decrease in growth rate was observed for the cytochrome c_1 -, cytochrome c_{550}/c_1 -, and cytochrome c_{550}/c_{552} -deficient mutant strains, which grew up to 50% more slowly than the wild-type cells. The cytochrome $c_{550}/c_{552}/c_1$ -deficient triple mutant was unable to grow on methylamine. Apparently, a possible electron transfer route from amicyanin, the dedicated electron acceptor of methylamine dehydrogenase (53), to any of the cytochrome *c* oxidases cannot support growth of the mutant strain under this condition. Since all of the cytochrome *c* double mutants (cyto-




FIG. 3. Time-dependent staining of soluble cytochromes *c* run into the gel to the position of the 14-kDa marker. Lane 1, the cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutant; lane 2, the cytochrome *c*₁-deficient mutant. The exposure times of the film are indicated at the left in minutes.

chrome *c*₅₅₀/*c*₅₅₂, cytochrome *c*₅₅₀/*c*₁, and cytochrome *c*₅₅₂/*c*₁-deficient mutant strains) were able to grow on methylamine, cytochrome *c*₅₅₀, cytochrome *c*₅₅₂, or the cytochrome *bc*₁ complex all suffice for electron flow from amicyanin to oxygen.

Electrons that are passed on via amicyanin to the cytochrome *bc*₁ complex may either be directed to oxygen via the Q-pool and the *ba*₃-type quinol oxidase in a reversed electron flow mechanism (45) or via cytochrome *c*₁ to any of the cytochrome *c* oxidases. In order to study the fate of electrons in a cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutant, this strain was grown with methylamine as an electron donor in the presence of 10 μ M antimycin A and myxothiazol in order to inactivate the heme *b*-containing subunit of cytochrome *bc*₁. As a control, we also cultured strain Pd93.11, a strain that lacks the *cbb*₃- and *aa*₃-type oxidases. This strain has been shown to be able to grow on methylamine by redirecting the electrons from cytochromes *c* via the cytochrome *bc*₁ complex and the Q-pool to the *ba*₃-type quinol oxidase (45). Both the wild-type strain and the cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutant were still able to grow under this condition, indicating that the inhibition of the

heme *b*-containing subunit of the cytochrome *bc*₁ complex did not prevent electron transport in these strains. The double cytochrome *c* oxidase mutant, however, was unable to grow under this condition, indicating that the reversed electron transfer route from methylamine to oxygen in this strain was completely blocked. The experiments described above imply that, in the absence of cytochromes *c*₅₅₀ and *c*₅₅₂, electrons can flow from the cytochrome *bc*₁ complex directly to one or both of the cytochrome *c* oxidases, most probably to the *cbb*₃-type oxidase since the *aa*₃-type oxidase is hardly present in the cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutant. In order to test this hypothesis, we constructed a triple-mutant strain, which lacks cytochromes *c*₅₅₀ and *c*₅₅₂, as well as the *aa*₃-type oxidase. This strain grew as fast on methylamine as the cytochrome *c*₅₅₀/*c*₅₅₂-deficient strain, indicating that electrons from the cytochrome *bc*₁ complex can indeed be directly transferred to the *cbb*₃-type oxidase.

All mutant strains were also able to grow on plates supplemented with methanol as the sole source of free energy, except for the triple mutant lacking cytochromes *c*₅₅₀ and *c*₅₅₂ and the cytochrome *bc*₁ complex. Quantitative data are not included since the cells grew poorly in batch cultures with methanol as sole source of carbon and free energy.

In the course of our studies on the role of the cytochromes *c* in electron transport under denitrifying conditions, we noticed that cytochrome *c*₅₅₀-deficient mutants that were cultured in brain heart infusion (BHI) broth were unable to reduce nitrite and grew to low turbidities. This may well have been due to the cytotoxicity of nitrite, which accumulated in these cultures. Earlier studies with that mutant, however, had indicated that its growth characteristics during denitrification were similar to those of the wild-type strain when they were raised in mineral medium supplemented with succinate and nitrate (51). Apparently, a component essential for denitrifying growth in the latter medium was absent from the BHI broth. We therefore added the individual ingredients of the mineral medium to BHI broth and studied the resultant effects on denitrifying growth. We found that the cultures of the cytochrome *c*₅₅₀-deficient mutant that had copper added to it were able to denitrify again. In order to study the importance of that metal

TABLE 2. Growth characteristics of *P. denitrificans* wild-type and cytochrome *c* mutant strains

| Strain | Characteristic(s) | Growth condition ^a | | | | | | | | | | |
|---------|-----------------------------|-------------------------------|---------------------------------|-----|-------------------------------|-----|--|-----|---------|---|-----|---------|
| | | Methanol, + O ₂ | Methylamine + O ₂ | | Succinate + O ₂ | | Succinate + NO ₃ , with copper | | | Succinate + NO ₃ , without copper | | |
| | | | μ (h ⁻¹) | OD | μ (h ⁻¹) | OD | μ (h ⁻¹) | OD | Nitrite | μ (h ⁻¹) | OD | Nitrite |
| Pd1222 | Wild type | G | 0.15 | 2.0 | 0.55 | 2.7 | 0.33 | 2.2 | — | 0.33 | 2.2 | — |
| Pd21.31 | CycA ⁻ | G | 0.12 | 1.8 | 0.53 | 2.5 | 0.33 | 2.2 | — | DG | 0.5 | + |
| Pd28.22 | CycM ⁻ | G | 0.14 | 2.0 | 0.53 | 2.7 | 0.33 | 2.2 | — | 0.33 | 2.2 | — |
| Pd24.21 | FbcC ⁻ | G | 0.09 | 1.7 | 0.53 | 2.4 | DG | 0.5 | + | DG | 0.5 | + |
| Pd92.06 | CycA.FbcC ⁻ | G | 0.07 | 1.6 | 0.55 | 2.5 | DG | 0.5 | + | DG | 0.5 | + |
| Pd92.26 | CycA.CycM ⁻ | G | 0.09 | 1.8 | 0.53 | 2.5 | 0.33 | 2.2 | — | DG | 0.5 | + |
| Pd92.28 | FbcC.CycM ⁻ | G | 0.08 | 1.6 | 0.55 | 2.4 | DG | 0.5 | + | DG | 0.5 | + |
| Pd93.14 | CycA.FbcC.CycM ⁻ | NG | NG | NG | 0.53 | 2.4 | DG | 0.5 | + | DG | 0.5 | + |

^a Growth in batch cultures with mineral medium and the electron donors and acceptors indicated. G, growth; NG, no growth; DG, disturbed growth, presumably as a consequence of nitrite accumulation, which is cytotoxic (assayed with nitur strips: —, no nitrite; +, nitrite at >80 mg/liter). Growth on methanol was studied by plating on mineral medium plates with methanol. μ , Maximum specific growth rate (per hour). The means of at least three independent experiments are shown, with standard error of the mean values of <5%. OD, the OD₆₆₀ at the end of growth. The means of at least three independent experiments are shown, with standard error of the mean values of <5%.

TABLE 3. Oxygen consumption rates of whole-cell suspensions of *P. denitrificans* strains

| Strain | Characteristics | O ₂ uptake rate ^a (nmol/min/mg of protein): | |
|---------|--------------------------------------|--|-----------------|
| | | Without AA-MYXO | With AA-MYXO |
| Pd1222 | Wild-type | 132 | 65 |
| Pd21.31 | $\Delta cycA$ | 135 | 86 |
| Pd28.22 | $cycM::Km^r$ | 130 | 60 |
| Pd24.31 | $\Delta fbcC$ | 152 | 148 |
| Pd92.07 | $\Delta cycA \Delta fbcC$ | 150 | 146 |
| Pd92.26 | $\Delta cycA cycM::Km^r$ | 136 | 80 |
| Pd92.28 | $\Delta fbcC cycM::Km^r$ | 148 | 140 |
| Pd93.14 | $\Delta cycA \Delta fbcC cycM::Km^r$ | 155 | 150 |

^a Rates were determined after the addition of succinate as an electron donor and either without or with of 5 μ M antimycin A and 5 μ M myxothiazol (AA-MYXO).

on the denitrifying growth of the mutant strains, we determined their characteristics of growth in media with or without added copper. The results presented in Table 2 show that the cytochrome *bc*₁ mutant strains had disturbed growth, accumulated nitrite, and reached final turbidities of ca. 25% of the wild-type value. All other mutants showed characteristics comparable to those of the wild-type when they were cultured in medium that contained copper. In the absence of copper, however, both the cytochrome *c*₅₅₀ and cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutant strains displayed characteristics similar to those of the cytochrome *bc*₁ mutants. They showed disturbed growth as a consequence of nitrite accumulation and did not produce gas in the culture medium, which would be diagnostic for the operation of a fully competent denitrification pathway. Apparently, electron transfer to at least the nitrite reductase requires a copper-containing component, which compensates for the loss of cytochrome *c*₅₅₀. We also noticed that the supernatants of the cytochrome *c*₅₅₀ and cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutants, which were raised under denitrifying growth conditions in mineral medium without copper, were yellow, whereas the supernatants of the other strains were not. The component responsible for this yellow color could not be precipitated with trichloroacetic acid, indicating that it was not a protein. The spectrum of the yellow component displayed an absorption band with a maximum at ca. 360 nm.

Kinetics of electron transport in cytochrome *c* mutants. *P. denitrificans* wild-type strain and cytochrome *c* mutants were grown under aerobic conditions with succinate as the sole carbon and free-energy source. Cells grown to their mid-exponential phase of growth were harvested, washed, and resuspended. The oxygen consumption rates of these suspensions were determined with succinate as the electron donor (Table 3). The data show that the oxygen consumption rates of the strains that still had the potential to synthesize the cytochrome *bc*₁ complex were comparable. Those that lacked the complex had rates ca. 15% higher than that of the wild-type strain. In the presence of antimycin A and myxothiazol, inhibitors of the cytochrome *bc*₁ complex, the oxygen consumption rates of the wild-type and cytochrome *c*₅₅₂-deficient mutant were almost halved compared to the uninhibited condition, whereas they were ca. 25% decreased in cytochrome *c*₅₅₀- and cytochrome *c*₅₅₀/*c*₅₅₂-deficient mutants. The respiratory inhibitor hardly if

at all affected the oxygen consumption rates of the cytochrome *c*₁-deficient strains.

We also studied the kinetics of oxidation and reduction of cytochromes *c* present in the wild-type and the cytochrome *c*₁-deficient mutant strains. The suspensions were incubated with succinate as reductant to the cytochromes, the reduction level of which was recorded spectrophotometrically in time (results not shown). Oxygen was supplied to these suspensions at a concentration of 1 mM by adding hydrogen peroxide, which is rapidly converted into oxygen and water by intracellular catalase activity (12). The traces show that the cytochromes *c* became more oxidized upon the addition of oxygen and became reduced again after oxygen had been fully consumed. The wild-type strain consumed the supplied oxygen at a rate of ca. 130 nmol of oxygen min⁻¹ mg of protein⁻¹ (results not shown). Cytochrome *bc*₁-deficient strains did so between 10 to 15% faster. The subsequent reduction of the cytochromes *c* in the latter strains occurred at an at least 40-fold-lower rate than that of the wild-type strain (results not shown).

DISCUSSION

The studies presented in this study have shed more light on the roles of cytochromes *c*₅₅₀, *c*₅₅₂, and *c*₁ in controlling the rates of electron transfer in *P. denitrificans* grown at various growth conditions. The data also indicate that cytochrome *c* deficiencies by the mutations are in many cases accompanied by changed levels of expression of other respiratory enzymes.

Much of our knowledge on the makeup and function of the *P. denitrificans* respiratory system at various growth conditions has been described in earlier studies (reviewed in references 2, 3, and 46). The roles of cytochromes *c*₅₅₀, *c*₅₅₂, and *c*₁ in that electron transport have now been defined by careful examination of the growth characteristics of mutant strains lacking one or more of these types of cytochrome. A scheme of the flexible respiratory network of methylamine grown cells as based on these studies is presented in Fig. 4. It shows that cytochrome *c*₅₅₀ is a potential electron donor to any of the cytochrome *c* oxidases. Cytochrome *c*₅₅₂ is a dedicated donor to only the *aa*₃-type oxidase, in agreement with the observation that cytochrome *c*₅₅₂ made part of a purified supercomplex containing the cytochrome *bc*₁-complex and the *aa*₃-type oxidase, which was functional in electron transport (4). Our results further show that cytochrome *c*₅₅₂ is not a donor to the *cbb*₃-type oxidase in agreement with an earlier observation that a mutant strain lacking cytochromes *c*₁ and *c*₅₅₀, as well as the *aa*₃-type oxidase, was unable to grow with methylamine as the sole free-energy source, whereas it could still express cytochrome *c*₅₅₂ and the *cbb*₃-type oxidase (11).

The studies described here present a qualitative view on all possible electron transfer pathways during methylamine oxidation as they operate in the various mutant strains. Our observation do not prove that all of these pathways operate to the same extent in the wild-type cells as they do in these mutants. The deletions of cytochromes *c*₁ and *c*₅₅₀ resulted in decreased maximum specific growth rates of the mutant strains by 40 and 20%, respectively. These cytochromes are apparently essential for optimal performance of the methylamine oxidizing pathways since they have multiple tasks in connecting quinol and amicyanin oxidation to cytochrome *c* reduction. However, it is




FIG. 4. Scheme of the respiratory network of *P. denitrificans* during growth with methylamine as sole source of free energy. The sequential oxidation of methylamine to carbon dioxide (dashed arrows) is mediated by methylamine dehydrogenase (MADH), formaldehyde dehydrogenase (FDH), and formate dehydrogenase (FDH). Physiologically important electron transfer routes (straight arrows) from NADH dehydrogenase (NADHDH) and MADH to the terminal oxidases were deduced from the studies in this study. amic, amicyanin.

not possible to deduce from these numbers the relative electron fluxes through the parallel electron transfer pathways in the wild-type cells. This is because every mutation in these mutants gives rise to changes in the kinetics of the remaining electron transfer reactions as a consequence of (i) changes in the expression levels of the respiratory components (28) and (ii) changes in the reduction levels of them (27).

The respiratory network of methanol-grown cells is quite similar to that of methylamine-grown cells. The difference is that methylamine dehydrogenase and its dedicated electron acceptor amicyanin (19, 53) are replaced by a methanol dehydrogenase and its dedicated electron acceptor cytochrome c_{551i} (18, 50). Just like amicyanin, this cytochrome requires another cytochrome c as electron acceptor and is unable to donate to any of the cytochrome c oxidases.

When the cytochrome c -deficient strains were grown with succinate as the carbon and free-energy source, their maximum specific growth rates were quite comparable to that of the wild-type strain, indicating that these mutant cells adapt completely in that respect to compensate for the loss of any of these cytochromes c . That adaptation involves higher electron transfer fluxes through the ba_3 -type quinol oxidases since these mutants display lower maximum cell numbers (since this pathway is less efficient with respect to free-energy transduction) and higher maximum oxygen uptake activity via the ba_3 -type oxidase. This result is in agreement with earlier studies on cytochrome bc_1 -deficient mutants (27).

The analyses of the mutants grown under denitrifying conditions indicated that cytochrome c_1 is essential for the reduction of nitrite to dinitrogen gas. This result is in agreement with the view that the cytochrome bc_1 complex is a compulsory intermediate in electron transfer from the Q-pool to the nitrite, nitric oxide, and nitrous oxide reductases (reviewed in reference 34). At least up to and including nitrite reductase, the pathway also involves cytochrome c_{550} , which appeared to

be an essential electron donor to this enzyme during copper limitation. In the presence of copper, however, the mutation in the gene encoding cytochrome c_{550} had hardly any effect on the denitrifying potential of that mutant strain, indicating that a copper-dependent electron carrier substituted for cytochrome c_{550} (51). A likely candidate is pseudoazurin, which is a type I blue copper protein and able to mediate electron transfer between the cytochrome bc_1 complex and the nitrite, nitric oxide, and nitrous oxide reductases in vitro (21, 25, 26). Indeed, it has been shown that a mutant strain deficient in cytochrome c_{550} and pseudoazurin failed to reduce nitrite (30; Stuart Ferguson, unpublished data). Cytochrome c_{552} is apparently redundant in the denitrifying pathways since we observed virtually no effect of its absence on the denitrifying potential of the mutant cells and since it was unable to compensate for the loss of cytochrome c_{550} under copper limitation.

In the course of our studies, we noticed that the expression levels of certain types of cytochrome c was increased in aerobically grown mutant strains, especially in those that lacked the cytochrome bc_1 complex. Apparently, the mutations in the cytochrome c -deficient mutants cause metabolic changes that result in changes in transcription activation of the genes encoding these cytochromes c . One of these metabolic changes affects the activity of the FnrP protein, which is a transcription activator regulating the expression of genes in response to oxygen limitation (47). This conclusion is based on the observation that the concentrations of the cytochrome c -containing subunits of the cbb_3 -type oxidase and cytochrome c peroxidase, the genes of which are regulated by FnrP (47), are increased in the mutant strains. Some of the mutants, i.e., the cytochrome c_{550} single mutant, the cytochrome c_{550}/c_{552} double mutant, and the cytochrome $c_{550}/c_{552}/c_1$ triple mutant, also displayed lower levels of the aa_3 -type oxidase as judged by spectroscopic analyses. That phenomenon has been described earlier for the cytochrome c_{550} -deficient mutant (51). The reason for the ob-

served decrease is not clear at present but cannot be merely ascribed to the *cycA* mutation alone since the cytochrome *c*₅₅₀/*c*₁-deficient double mutant has wild-type levels of the *aa*₃-type oxidase.

Taken together, these observations indicate that the performance of the respiratory network is continuously monitored and that its makeup adapted in response to changes in that performance apparently in order to maintain an optimal electron flux at each given growth condition.

ACKNOWLEDGMENTS

This work was supported by the Chemical Sciences Foundation (CW), with additional financial aid from The Netherlands Organization for Scientific Research (NWO).

REFERENCES

- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1993. Current protocols in molecular biology. John Wiley & Sons, New York, N.Y.
- Baker, S. C., S. J. Ferguson, B. Ludwig, M. D. Page, O. M. H. Richter, and R. J. M. Van Spanning. 1998. Molecular genetics of the genus *Paracoccus*: metabolically versatile bacteria with bioenergetic flexibility. *Microbiol. Mol. Biol. Rev.* **62**:1046–1078.
- Berks, B. C., S. J. Ferguson, J. W. B. Moir, and D. J. Richardson. 1995. Enzymes and associated electron transport systems that catalyze the respiratory reduction of nitrogen oxides and oxyanions. *Bba-Bioenergetics* **1232**: 97–173.
- Berry, E. A., and B. L. Trumpower. 1985. Isolation of ubiquinol oxidase from *Paracoccus denitrificans* and resolution into cytochrome *bc*₁ and cytochrome *c-aa*₃ complexes. *J. Biol. Chem.* **260**:2458–2467.
- Bosma, G., M. Braster, A. H. Stouthamer, and H. W. Van Verseveld. 1987. Isolation and characterization of ubiquinol oxidase complexes from *Paracoccus denitrificans* cells cultured under various limiting conditions in the chemostat. *Eur. J. Biochem.* **165**:657–663.
- Boyer, H. W., and D. Roulland-Dussoix. 1969. A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J. Mol. Biol.* **41**:459–472.
- Chang, J. P., and J. G. Morris. 1962. Studies on the utilization of nitrate by *Micrococcus denitrificans*. *J. Gen. Microbiol.* **29**:301–310.
- De Gier, J. W. 1995. The terminal oxidases of *Paracoccus denitrificans*. Ph.D. thesis. Free University, Amsterdam, The Netherlands.
- De Gier, J. W. L., M. Lubben, W. N. M. Reijnders, C. A. Tipker, D. J. Slotboom, R. J. M. Van Spanning, A. H. Stouthamer, and J. Van der Oost. 1994. The terminal oxidases of *Paracoccus denitrificans*. *Mol. Microbiol.* **13**:183–196.
- De Gier, J. W. L., M. Schepper, W. N. M. Reijnders, S. J. Van Dyck, D. J. Slotboom, A. Warne, M. Saraste, K. Krab, M. Finel, A. H. Stouthamer, R. J. M. Van Spanning, and J. Van der Oost. 1996. Structural and functional analysis of *aa*₃-type and *cbb*₃-type cytochrome *c* oxidases of *Paracoccus denitrificans* reveals significant differences in proton-pump design. *Mol. Microbiol.* **20**:1247–1260.
- De Gier, J. W. L., J. Van der Oost, N. Harms, A. H. Stouthamer, and R. J. M. Van Spanning. 1995. The oxidation of methylamine in *Paracoccus denitrificans*. *Eur. J. Biochem.* **229**:148–154.
- De Gier, J. W. L., R. J. M. Van Spanning, L. F. Oltmann, and A. H. Stouthamer. 1992. Oxidation of methylamine by a *Paracoccus denitrificans* mutant impaired in the synthesis of the *bc*₁ complex and the *aa*₃-type oxidase. Evidence for the existence of an alternative cytochrome *c* oxidase in this bacterium. *FEBS Lett.* **306**:23–26.
- De Vries, G. E., N. Harms, J. Hoogendijk, and A. H. Stouthamer. 1989. Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of a n(GATC)_n DNA-modifying property. *Arch. Microbiol.* **152**:52–57.
- Ferguson, S. J. 1994. Denitrification and its control. *Anton Leeuwenhoek Int. J. Gen. M.* **66**:89–110.
- Garciahorsman, J. A., B. Barquera, J. Rumbley, J. X. Ma, and R. B. Gennis. 1994. The superfamily of heme-copper respiratory oxidases. *J. Bacteriol.* **176**:5587–5600.
- Harms, N., J. Ras, S. Koning, W. N. M. Reijnders, A. H. Stouthamer, and R. J. M. Van Spanning. 1996. Genetics of C1 metabolism regulation in *Paracoccus denitrificans*, p. 126–132. In M. E. Lidstrom and F. R. Tabita (ed.), *Microbial growth on C1 compounds*. Kluwer Academic Publishers, Amsterdam, The Netherlands.
- Harms, N., and R. J. M. Van Spanning. 1991. C₁ metabolism in *Paracoccus denitrificans*: genetics of *Paracoccus denitrificans*. *J. Bioenerg. Biomembr.* **23**:187–210.
- Husain, M., and V. L. Davidson. 1986. Characterization of two inducible *c*-type cytochromes from *Paracoccus denitrificans*. *J. Biol. Chem.* **261**:8577–8580.
- Husain, M., and V. L. Davidson. 1985. An inducible periplasmic blue copper protein from *Paracoccus denitrificans*: purification, properties, and physiological role. *J. Biol. Chem.* **260**:14626–14629.
- John, P., and F. R. Whatley. 1975. *Paracoccus denitrificans* and the evolutionary origin of the mitochondrion. *Nature* **254**:495–498.
- Koutny, M., I. Kucera, R. Tesarik, J. Turanek, and R. J. M. Van Spanning. 1999. Pseudoazurin mediates periplasmic electron flow in a mutant strain of *Paracoccus denitrificans* lacking cytochrome *c*₅₅₀. *FEBS Lett.* **448**:157–159.
- Kurowski, B., and B. Ludwig. 1987. The genes of the *Paracoccus denitrificans* *bc*₁ complex. Nucleotide sequence and homologies between bacterial and mitochondrial subunits. *J. Biol. Chem.* **262**:13805–13811.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680–685.
- Lawford, H. G., J. C. Cox, P. B. Garland, and B. A. Hadock. 1976. Electron transport in aerobically grown *Paracoccus denitrificans*: kinetic characterization of the membrane-bound cytochromes and the stoichiometry of respiratory driven-proton translocation. *FEBS Lett.* **64**:369–374.
- Moir, J. W. B., D. Baratta, D. J. Richardson, and S. J. Ferguson. 1993. The purification of a *cd*₁-type nitrite reductase from, and the absence of a copper-type nitrite reductase from, the aerobic denitrifier *Thiosphaera pantotropa*: the role of pseudoazurin as an electron donor. *Eur. J. Biochem.* **212**:377–385.
- Moir, J. W. B., and S. J. Ferguson. 1994. Properties of a *Paracoccus denitrificans* mutant deleted in cytochrome *c*₅₅₀ indicate that a copper protein can substitute for this cytochrome in electron transport to nitrite, nitric oxide and nitrous oxide. *Microbiology* **140**:389–397.
- Otten, M. F., W. N. Reijnders, J. J. Bedaux, H. V. Westerhoff, K. Krab, and R. J. Van Spanning. 1999. The reduction state of the Q-pool regulates the electron flux through the branched respiratory network of *Paracoccus denitrificans*. *Eur. J. Biochem.* **261**:767–774.
- Otten, M. F., D. M. Stork, W. N. Reijnders, H. V. Westerhoff, and R. J. Van Spanning. 2001. Regulation of expression of terminal oxidases in *Paracoccus denitrificans*. *Eur. J. Biochem.* **268**:2486–2497.
- Otto, B. R., S. J. van Dooren, J. H. Nuijens, J. Luirink, and B. Oudega. 1998. Characterization of a hemoglobin protease secreted by the pathogenic *Escherichia coli* strain EB1. *J. Exp. Med.* **188**:1091–1103.
- Pearson, I. 2000. Cloning, mutagenesis and expression studies of the *pazS* gene encoding pseudoazurin from *Paracoccus denitrificans*. Ph.D. thesis. Oxford University, Oxford, England.
- Preisig, O., R. Zufferey, L. Thonymeyer, C. A. Appleby, and H. Hennecke. 1996. A high-affinity *cbb*₃-type cytochrome oxidase terminates the symbiosis-specific respiratory chain of *Bradyrhizobium japonicum*. *J. Bacteriol.* **178**: 1532–1538.
- Raitio, M., T. Jalli, and M. Saraste. 1987. Isolation and analysis of the genes for cytochrome *c* oxidase in *Paracoccus denitrificans*. *EMBO J.* **6**:2825–2833.
- Raitio, M., J. M. Pispas, T. Metso, and M. Saraste. 1990. Are there isoenzymes of cytochrome *c* oxidase in *Paracoccus denitrificans*? *FEBS Lett.* **261**: 431–435.
- Richardson, D. J. 2000. Bacterial respiration: a flexible process for a changing environment. *Microbiology* **146**:551–571.
- Richter, O. M. H., J. S. Tao, A. Turba, and B. Ludwig. 1994. A cytochrome *ba*₃ functions as a quinol oxidase in *Paracoccus denitrificans*—Purification, cloning, and sequence comparison. *J. Biol. Chem.* **269**:23079–23086.
- Riistama, S., A. Puustinen, M. I. Verkhovskiy, J. E. Morgan, and M. Wikstrom. 2000. Binding of O₂ and its reduction are both retarded by replacement of valine 279 by isoleucine in cytochrome *c* oxidase from *Paracoccus denitrificans*. *Biochemistry* **39**:6365–6372.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Saraste, M., and J. Castresana. 1994. Cytochrome oxidase evolved by tinkering with denitrification enzymes. *FEBS Lett.* **341**:1–4.
- Simon, R., U. Priefer, and A. Pühler. 1983. Vector plasmids for *in vivo* and *in vitro* manipulations of gram-negative bacteria, p. 98–106. In A. Pühler (ed.), *Molecular genetics of the bacteria-plant interactions*. Springer-Verlag KG, Berlin, Germany.
- Spaink, H. P., R. J. H. Okker, C. A. Wijffelman, E. Pees, and B. J. J. Lugtenberg. 1987. Promoters in the nodulation region of *Rhizobium leguminosarum* Sym plasmid pRL1J1. *Plant Mol. Biol.* **9**:27–39.
- Stouthamer, A. H. 1992. Metabolic pathways in *Paracoccus denitrificans* and closely related bacteria in relation to the phylogeny of prokaryotes. *Antonie Leeuwenhoek Int. J. Gen. M.* **61**:1–33.
- Stouthamer, A. H. 1991. Metabolic regulation including anaerobic metabolism in *Paracoccus denitrificans*. *J. Bioenerg. Biomembr.* **23**:163–185.
- Turba, A., M. Jetzek, and B. Ludwig. 1995. Purification of *Paracoccus denitrificans* cytochrome *c*₅₅₂ and sequence analysis of the gene. *Eur. J. Biochem.* **231**:259–265.
- Van der Oost, J., A. P. N. Deboer, J. W. L. Degier, W. G. Zumft, A. H. Stouthamer, and R. J. M. Van Spanning. 1994. The heme-copper oxidase family consists of three distinct types of terminal oxidases and is related to nitric oxide reductase. *FEMS Microbiol. Lett.* **121**:1–9.

45. Van der Oost, J., M. Schepper, A. H. Stouthamer, H. V. Westerhoff, R. J. M. Van Spanning, and J. W. L. De Gier. 1995. Reversed electron transfer through the *bc*₁ complex enables a cytochrome *c* oxidase mutant ($\Delta aa_3/cbb_3$) of *Paracoccus denitrificans* to grow on methylamine. *FEBS Lett.* **371**:267–270.
46. Van Spanning, R. J. M., A. P. N. De Boer, W. N. M. Reijnders, J. W. L. De Gier, C. O. Delorme, A. H. Stouthamer, H. V. Westerhoff, N. Harms, and J. Van der Oost. 1995. Regulation of oxidative phosphorylation: The flexible respiratory network of *Paracoccus denitrificans*. *J. Bioenerg Biomembr.* **27**: 499–512.
47. Van Spanning, R. J. M., A. P. N. De Boer, W. N. M. Reijnders, H. V. Westerhoff, A. H. Stouthamer, and J. Van der Oost. 1997. FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR family of transcriptional activators but have distinct roles in respiratory adaptation in response to oxygen limitation. *Mol. Microbiol.* **23**:893–907.
48. Van Spanning, R. J. M., A. P. N. De Boer, D.-J. Slotboom, W. N. M. Reijnders, and A. H. Stouthamer. 1995. Isolation and characterization of a novel insertion sequence element, *ISI248*, in *Paracoccus denitrificans*. *Plasmid* **34**:11–21.
49. Van Spanning, R. J. M., S. De Vries, and N. Harms. 2000. Coping with formaldehyde during C1 metabolism of *Paracoccus denitrificans*. *J. Mol. Catal. B Enzymatic* **8**:37–50.
50. Van Spanning, R. J. M., C. W. Wansell, T. De Boer, M. J. Hazelaar, H. Anazawa, N. Harms, L. F. Oltmann, and A. H. Stouthamer. 1991. Isolation and characterization of the *moxJ*, *moxG*, *moxI*, and *moxR* genes of *Paracoccus denitrificans*: inactivation of *moxJ*, *moxG*, and *moxR* and the resultant effect on methylotrophic growth. *J. Bacteriol.* **173**:6948–6961.
51. Van Spanning, R. J. M., C. W. Wansell, N. Harms, L. F. Oltmann, and A. H. Stouthamer. 1990. Mutagenesis of the gene encoding cytochrome *c*₅₅₀ of *Paracoccus denitrificans* and analysis of the resultant physiological effects. *J. Bacteriol.* **172**:986–996.
52. Van Spanning, R. J. M., C. W. Wansell, W. N. M. Reijnders, N. Harms, J. Ras, L. F. Oltmann, and A. H. Stouthamer. 1991. A method for introduction of unmarked mutations in the genome of *Paracoccus denitrificans*: construction of strains with multiple mutations in the genes encoding periplasmic cytochromes *c*₅₅₀, *c*_{551a}, and *c*_{553a}. *J. Bacteriol.* **173**:6962–6970.
53. Van Spanning, R. J. M., C. W. Wansell, W. N. M. Reijnders, L. F. Oltmann, and A. H. Stouthamer. 1990. Mutagenesis of the gene encoding amicyanin of *Paracoccus denitrificans* and the resultant effect on methylamine oxidation. *FEBS Lett.* **275**:217–220.