

Landscape - Soilscape Evolution Modelling: LAPSUS

M.P.W. Sonneveld^A, A.J.A.M. Temme^A, J.M. Schoorl^A, L. Claessens^{A,B}, W. Viveen^A, J.E.M. Baartman^A, J.P. Lesschen^C and W. van Gorp^A

^A Wageningen University, Land Dynamics Group, P.O. Box 47, 6700 AA Wageningen, The Netherlands. Emails Marthijn.Sonneveld@wur.nl, Arnaud.Temme@wur.nl, Jeroen.Schoorl@wur.nl, Jantiene.Baartman@wur.nl, Wouter.vanGorp@wur.nl

^B International Potato Center (CIP), Nairobi, Kenya. Email l.claessens@CGIAR.ORG

^C Alterra, Wageningen UR, Wageningen, The Netherlands. Email JanPeter.Lesschen@wur.nl

Abstract

Landscape evolution modelling can make landscape evolution hypotheses explicit and theoretically allows for their falsification and improvement. Ideally, landscape evolution models (LEMs) combine the results of all relevant landscape forming processes into an ever-adapting digital landscape (e.g. DEM). These processes may act on different spatial and temporal scales. LAPSUS is an example of such a LEM. In multiple study cases different landscape processes have been included in LAPSUS: water erosion and deposition, landslide activity, creep, solifluction, weathering, tectonics and tillage. Besides properties of soils influencing landscape forming processes, vegetation effects can also be included. Process descriptions are kept as simple and generic as possible, ensuring wide applicability of the modelling approach. Interactions between processes are turn-based: soil redistribution caused by one process are calculated and used to adapt the DEM before another process is simulated. LAPSUS uses multiple flow techniques to model flows of water and sediment over the landscape. Though computationally costly, this gives a more realistic result than steepest descent methods. In addition, the combination of different processes may create sinks during modelling. Since these sinks are not spurious, the model has been adapted to deal with sinks in natural ways. This is crucial for several purposes, for instance when studying damming of valleys by landslides, and subsequent infilling of the resulting lake with sediments from upstream.

Key Words

Landscape Evolution Modelling, LAPSUS, soil redistribution, erosion.

Introduction

This short paper summarizes ongoing and completed work with the LAPSUS model and foreseen developments in the near future. LAPSUS is a landscape evolution model (LEM) that combines the effects of multiple landscape forming processes, including soil formation, into one dynamic landscape modelling framework. Spatial and temporal extent and resolution may vary from slope, catchment to basin, processed grid cells from 1 to 1000 m², timesteps of multiple events, seasons, years, decades and simulation periods from years to millennial time scales.

Interactions between processes are turn-based: soil redistribution caused by one process are calculated and used to adapt the DEM before another process is simulated. In multiple study cases different landscape processes have been included in LAPSUS : water erosion and deposition, landslide activity, creep, solifluction, weathering, tectonics and tillage. (Figure 1).




Figure 1. Overview of processes incorporated in the LAPSUS modelling framework (see also www.lapsusmodel.nl).

Besides properties of soils influencing landscape forming processes, vegetation effects can also be included. Process descriptions are kept as simple and generic as possible, ensuring wide applicability of the modelling approach. LAPSUS uses multiple flow routing techniques to model the flow of water and sediment over the landscape. This is computationally costly, but yields a more realistic result than steepest descent methods, especially when combining multiple processes over multiple timesteps.

The combination of different processes may create sinks during modelling. Since these sinks are not spurious, the model has been adapted to deal with them in a natural way. This is crucial when studying damming of valleys by landslides, and subsequent infilling of the resulting lake with sediments from upstream.

Results and discussion

LAPSUS has been used for erosion and landscape evolution studies in many landscapes in many countries over the last years. The development of LAPSUS started in 2000 with the programming, calibration and validation of the LAPSUS model and applications concerning land use in Spain and Ecuador (Schoorl *et al.* 2000, 2002, 2004, 2006; Schoorl and Veldkamp 2001, 2006). Later, the model has been extended in order to include soil redistribution by landsliding in New Zealand and Taiwan (Claessens *et al.* 2005, 2006a, 2006b, 2007a, 2007b). In addition, issues of DEM resolution and the treatment of sinks and pits in the landscape have been investigated (Temme *et al.* 2006, 2009a) as well as stretching the models time scale to landscape evolution time spans, for example in South Africa (Temme and Veldkamp 2009; Temme *et al.* 2009b). Different applications with individual processes have been developed, for example, the model has been used in regional nutrient balance studies in Africa (Haileslassie *et al.* 2005, 2006, 2007; Roy *et al.* 2004; Lesschen *et al.* 2005). The model has also been applied in desert environments in Israel (Buis and Veldkamp 2008); it has been used in combination with geostatistical tools and tillage in Canada (Heuvelink *et al.* 2006), and to investigate the faith of phosphor in the landscapes of the Netherlands (Sonneveld *et al.* 2006).

Recent developments and directions with the LAPSUS model are:

- Connectivity, agricultural terraces and land abandonment (Lesschen *et al.* 2007, 2009).
- Interactions and feedback mechanisms between land use and soil redistribution (Claessens *et al.* 2009).
- Effects of hydrological engineering on soil redistribution in large fluvial systems (Viveen *et al.* 2009)
- Erosion in a landscape evolution context, comparing event based and long term based models: LISEM and LAPSUS (Baartman *et al.* 2009).
- Refining the LAPSUS temporal resolution. Modelling daily sediment yield from a meso-scale catchment, a case study in SW Poland. (Covert-Keesstra *et al.* 2009)
- Land sliding in mountainous areas. Landscape Dynamics: Calibrating landscape process modelling with Caesium-137 data, separating water driven erosion from landslides? See (Schoorl *et al.* 2009).
- 3D river gradient modelling. Quaternary tectonics, sea level and climate change: the case of the river Miño (Viveen *et al.* 2009).
- Coupling and interaction with TOA modelling. A novel site-specific methodology to assess the supply curve of environmental services (Stoorvogel *et al.* 2009; Claessens *et al.* in prep).

Conclusion

Landscape evolution modelling allows for confirmation, falsification or improvement of landscape evolution hypotheses and can make the consequences temporally and spatially explicit. Ideally, landscape evolution models (LEMs) combine the results of all relevant landscape forming processes into an ever-adapting digital landscape model. These processes may act and interact on different spatial and temporal scales. The LAPSUS modelling framework is an example of a LEM that has embedded multiple landscape forming processes and their interactions in a generic tool that can be used to study many landscapes of the world at multiple temporal and spatial scales.

References

- Baartman JEM, Schoorl JM, Veldkamp A, Ritsema CJ (2009) Erosion in a landscape evolution context: LISEM and LAPSUS. In: International conference on Desertification, Murcia, Spain, 16 - 18 September, 2009. - Murcia, Spain : Ediciones de la universidad de Murcia, Advances in studies on desertification, 2009-09-16/ 2009-09-18
- Buis E, Veldkamp A (2008) Modelling dynamic water redistribution patterns in arid catchments in the Negev Desert of Israel. *Earth Surface Processes and Landforms* **33**, 107-122.

- Claessens L, Heuvelink GBM, Schoorl JM, Veldkamp A (2005) DEM resolution effects on shallow landslide hazard and soil redistribution modelling. *Earth Surface Processes and Landforms* **30**, 461-477.
- Claessens L, Lowe DJ, Hayward BW, Schaap BF, Schoorl JM, Veldkamp A (2006a) Reconstructing high-magnitude/low-frequency landslide events based on soil redistribution modelling and a Late-Holocene sediment record from New Zealand. *Geomorphology* **74**, 29-49.
- Claessens L, Verburg PH, Schoorl JM, Veldkamp A (2006b), Contribution of topographical based landslide hazard Modelling to the analysis of the spatial distribution and ecology of Kauri (*Agathis australis*): *Landscape Ecology* **21**, 63-76.
- Claessens L, Schoorl JM, Veldkamp A (2007a) Modelling the location of shallow landslides and their effects on landscape dynamics in large watersheds: an application for Northern New Zealand. *Geomorphology* **87**(1-2), 16-27.
- Claessens L, Knapen A, Kitutu MG, Poesen J, Deckers JA (2007b) Modelling landslide hazard, soil redistribution and sediment yield of landslides on the Ugandan footslopes of Mount Elgon. *Geomorphology* **90**, 23-35.
- Claessens L, JM Schoorl, Verburg PH, Geraedts L, Veldkamp A (2009) Modelling interactions and feedback mechanisms between land use change and landscape processes. *Agriculture, Ecosystems and Environment* **129**, 157-170.
- Claessens L, Stoorvogel JJ, Antle JM, (in prep). Exploring the Impacts of Field Interactions on an Integrated Assessment of Terraced Crop Systems in the Peruvian Andes. Submitted to *Journal of Land Use Science*.
- Coevert-Keesstra SD, Schoorl JM, Temme AJAM (2009) Modelling daily sediment yield from a meso-scale catchment, a case study in SW Poland. In: International conference on Desertification, Advances in studies on desertification, Murcia, Spain, 16-18 September 2009. - Murcia, Spain : Ediciones de la universidad de Murcia, Congreso internacional sobre desertificacion en memoria del professor John B. Thornes, 2009-09-16/ 2009-09-18
- Hailelassie A, Priess J, Veldkamp E, Teketay D, Lesschen JP (2005) Assessment of soil nutrient depletion and its spatial variability on smallholders' mixed farming systems in Ethiopia using partial versus full nutrient balances. *Agriculture, Ecosystems and Environment* **108**, 1-16.
- Hailelassie A, Priess JA, Veldkamp E, Lesschen JP (2006) Smallholders' soil fertility management in the Central Highlands of Ethiopia: implications for nutrient stocks, balances and sustainability of agroecosystems. *Nutrient Cycling in Agroecosystems* **75**, 135-146.
- Hailelassie A, Priess JA, Veldkamp E, Lesschen JP (2007) Nutrient flows and balances at the field and farm scale: Exploring effects of land-use strategies and access to resources. *Agricultural Systems* **94** (2), 459-470.
- Heuvelink GBM, Schoorl JM, Veldkamp A, Pennock DJ (2006) Space-time Kalman filtering of soil redistribution. *Geoderma* **133**, 124-137.
- Lesschen JP, Asiamah RD, Gicheru P, Kante S, Stoorvogel JJ, Smaling EMA (2005) Scaling Soil Nutrient Balances - Enabling mesoscale approaches for African realities. *FAO Fertilizer and Plant Nutrition Bulletin* 15, FAO, Rome.
- Lesschen JP; Stoorvogel JJ; Smaling EMA, Heuvelink GBM; Veldkamp A (2007) A spatially explicit methodology to quantify soil nutrient balances and their uncertainties at the national level *Nutrient Cycling in Agroecosystems* **78**, 111 - 131.
- Lesschen JP, Schoorl JM, Cammeraat LH (2009). Modelling runoff and erosion for a semi-arid catchment using a multi-scale approach based on hydrological connectivity. *Geomorphology* **109** (3-4), 174-183.
- Roy RN, Misra RV, Lesschen JP, Smaling EM (2004) 'Assessment of soil nutrient balances: Approaches and Methodologies'. *Fertilizer and Plant Nutrition Bulletin* 14. (FAO: Rome).
- Schoorl JM, Boix Fayos C, de Meijer RJ, van der Graaf ER, Veldkamp A (2004). The 137Cs technique on steep Mediterranean slopes (Part 2): landscape evolution and model calibration. *Catena* **57**, 35-54.
- Schoorl JM, Veldkamp A, Bouma J (2002) Modelling water and soil redistribution in a dynamic landscape context. *Soil. Sci. Soc. Am. J.* **66**, 1610-1619.
- Schoorl JM, Veldkamp A (2001) Linking land use and landscape process modelling: a case study for the Alora region (South Spain). *Agric.Ecosyst.Environ* **85**, 281-292.
- Schoorl JM, Sonneveld MPW, Veldkamp A (2000) Three-dimensional landscape process modelling: the

- effect of DEM resolution. *Earth Surf.Proc.Landforms* **25**, 1025-1034.
- Schoorl JM, Claessens L, Lopez Ulloa M, de Koning GHJ, Veldkamp A (2006) Geomorphological Analysis and Scenario Modelling in the Noboa – Pajan Area, Manabi Province, Ecuador. *Zeitschrift Fur Geomorfologie* **145**, 105 - 118.
- Schoorl JM, Veldkamp A (2006) Multi-Scale Soil-Landscape Process Modeling. In ‘Environmental Soil-Landscape Modeling: Geographic Information Technologies and Pedometrics’. (Ed S Grunwald) pp. 417 – 435. (Boca Raton, FL, CRC press: Taylor and Francis Group).
- Schoorl JM, Chang KT, Chiu YJ, Veldkamp A (2009) Landscape Dynamics: Calibrating landscape process modelling with Caesium-137 data, separating water driven erosion from landslides? In ‘Proceedings of The International Conference in Commemoration of the 10th Anniversary of the 1999 Chi-Chi Earthquake. Sessions on Land Dynamics in Mountainous Watersheds: Typhoons, Landslides, and Land Use, Taipei, Taiwan, 17-21 September 2009’.
- Sonneveld MPW, Schoorl JM, Veldkamp A (2006) Evaluating The Fate Of Phosphorus In Apparent Homogeneous Landscapes Using a High-Resolution DEM. *Geoderma* **133**, 32 - 42.
- Stoorvogel JJ, Temme AJAM, Antle JM, Claessens L, Schoorl JM (2009) A novel site-specific methodology to assess the supply curve of environmental services. In ‘Proceedings of the Conference on Integrated Assessment of Agriculture and Sustainable Development; Setting the Agenda for Science and Policy (AgSAP 2009), Egmond aan Zee, The Netherlands, 10-12 March 2009’. (Wageningen, The Netherlands: 2009-03-10/ 2009-03-12)
- Temme AJAM, Schoorl JM, Veldkamp A (2006) Algorithm for dealing with depressions in dynamic landscape evolution models. *Comp.Geosci* **32**, 452 - 461.
- Temme AJAM, Veldkamp A (2009) Multi-process Late Quaternary landscape evolution modelling reveals lags in climate response over small spatial scales. *Earth Surface Processes and Landforms* **34**, 573-589.
- Temme AJAM, Heuvelink GBM, Schoorl JM, Claessens L (2009a) Chapter 5: Geostatistical simulation and error propagation in geomorphometry. In: *Geomorphometry: Concepts, Software, Applications*. (Eds T Hengl, HI Reuter) pp. 121 - 140. (Elsevier) (*Developments in Soil Science* **33**)
- Temme AJAM; Baartman JEM; Schoorl JM (2009b) Can uncertain landscape evolution models discriminate between landscape responses to stable and changing future climate? A millennial-scale test. *Global and Planetary Change* **69**, 48 - 58.
- Viveen W, Maas GJ, Schoorl JM (2009) Preliminary investigation of the spatial erosion and sedimentation patterns and processes in the catchment of German – Dutch Vecht river: potentials for recovery of natural river dynamics. (NL: Een verkenning van ruimtelijke erosie- en sedimentatieprocessen in het stroomgebied van de Nederlands-Duitse Vecht; Potenties voor herstel van natuurlijke rivierdynamiek). Wageningen, Alterra, Alterra Report 1939. pp 33.
- Temme AJAM, Claessens L, Veldkamp A, Schoorl JM (In prep) Evaluating choices in multi-process landscape evolution models. Submitted to *Geomorphology*.
- Viveen W, Schoorl JM, Veldkamp A, Vidal Romani JR (2009) Quaternary tectonics, sea level and climate change: the case of the river Miño. In: Livro de resumos, VII Reuniao do Quaternário Ibérico: O futuro do ambiente da Península Iberica: as lições do passado geológico recente, Portugal, Algarve, 5-9 October 2009. - Faro, Portugal : CIMA, 2009-10-05/ 2009-10-09