

High prevalence of nasal MRSA carriage in slaughterhouse workers in contact with live pigs in The Netherlands

B. A. G. L. VAN CLEEF^{1,2*}, E. M. BROENS^{1,3}, A. VOSS⁴, X. W. HUIJSDENS¹,
L. ZÜCHNER⁵, B. H. B. VAN BENTHEM¹, J. A. J. W. KLUYTMANS²,
M. N. MULDER¹ AND A. W. VAN DE GIESSEN¹

¹ RIVM National Institute for Public Health and the Environment, Centre for Infectious Disease Control Netherlands, Bilthoven, The Netherlands

² VU University Medical Centre, Department of Medical Microbiology and Infection Prevention, Amsterdam, The Netherlands

³ Quantitative Veterinary Epidemiology, Wageningen Institute of Animal Sciences (WIAS), Wageningen University, Wageningen, The Netherlands

⁴ Canisius-Wilhelmina Hospital, Department of Medical Microbiology and Infection Control, Nijmegen, The Netherlands

⁵ Food and Consumer Product Safety Authority (VWA) Region East, Zutphen, The Netherlands

(Accepted 15 January 2009; first published online 9 February 2010)

SUMMARY

Livestock-associated MRSA has been found in various animals, livestock farmers and retail meat. This study aimed to determine the prevalence and determinants of nasal MRSA carriage in pig slaughterhouse workers. Three large pig slaughterhouses in The Netherlands were studied in 2008 using human and environmental samples. The overall prevalence of nasal MRSA carriage in employees of pig slaughterhouses was 5·6% (14/249) (95% CI 3·4–9·2) and working with live pigs was the single most important factor for being MRSA positive (OR 38·2, $P < 0\cdot0001$). At the start of the day MRSA was only found in environmental samples from the lairages (10/12), whereas at the end of the day MRSA was found in the lairages (11/12), the dirty (5/12) and clean (3/12) areas and green offal (1/3). The MRSA status of the environmental samples correlated well with the MRSA status of humans working in these sections ($r = 0\cdot75$). In conclusion, a high prevalence of nasal MRSA carriage was found in pig-slaughterhouse workers, and working with live pigs is the most important risk factor. Exact transmission routes from animals to humans remain to be elucidated in order to enable application of targeted preventive measures.

Key words: Abattoirs, cross-sectional studies, domestic animals, humans, methicillin-resistant *Staphylococcus aureus*.

INTRODUCTION

Since 2003, a distinct clone of methicillin-resistant *Staphylococcus aureus* (MRSA), related to the

livestock reservoir has emerged in the human population [1]. As this clone was found to be non-typable (NT) by pulsed-field gel electrophoresis using the *Sma*I restriction enzyme, it was originally called NT-MRSA [2, 3]. Multi-locus sequence typing revealed that all strains belonged to the clonal complex 398 (CC398) [4]. At present, it is clear that people who have frequent contact with pigs or veal calves

* Author for correspondence: Drs B. A. G. L. van Cleef, Epidemiology and Surveillance Unit, National Institute for Public Health and the Environment, PO Box 1, 3720 BA Bilthoven, The Netherlands.
(Email: brigitte.van.cleef@rivm.nl)

have extremely high MRSA CC398 carriage rates compared to national community prevalences (25–35% vs. 0.1% in The Netherlands) [5–8].

As a result of the elevated prevalences in this specific population, the ‘search and destroy’ policy in The Netherlands was adapted; persons in contact with live pigs and veal calves are added to the high-risk group and should be screened for MRSA upon hospital admission [9]. As a consequence, the number of MRSA CC398-carrying patients found in The Netherlands increased dramatically to nearly 30% of all newly detected MRSA strains in 2007 [10], and 42% in 2008 [11]. The proportion of MRSA in *S. aureus* nosocomial infections remained very low (<2%), compared to other countries [12].

In a recent survey by the Food and Consumer Product Safety Authority in the Netherlands (VWA) MRSA was found in 11% of retail meat (with a minimum MRSA prevalence of 2% in game and a maximum of 35% in turkey) [13]. Other studies also found MRSA in retail meat, in varying percentages (2.5% [14], 19% [15], 0.7% [16], 5% [17], 0% [18] and 17%, R. de Jonge, J. E. Verdier and A. H. Havelaar, unpublished observations).

In animal husbandry-dense areas, the majority of newly identified human MRSA carriers concerns this livestock-associated MRSA [19], and recently, the first hospital outbreaks of CC398 have been reported [20, 21]. Meanwhile, serious invasive infections due to CC398 have been observed [22–27]. Therefore, the emergence of this new livestock-associated clone poses a potential public health risk that warrants close monitoring.

The high prevalence of MRSA in meat products and in people working with livestock raises the question whether slaughterhouse workers, who are in contact with pigs (dead or alive) and meat products, are also at risk. Therefore, we performed a cross-sectional survey on nasal MRSA CC398 carriage in employees of pig slaughterhouses, and on the occurrence of MRSA in different slaughterhouse sections.

METHODS

Study population, questionnaires and human sampling

Three pig slaughterhouses were enrolled in the study on the basis of voluntary participation, from a complete list of 10 large pig slaughterhouses in The Netherlands. All were located in the south and the east of the country, in areas with a high pig density. By using a structured questionnaire,

slaughterhouse-specific information was collected, e.g. number of employees, slaughterhouse capacity, specifics on lairages and the production process, information on microbiological contamination of the carcasses and working benches and hygiene measures.

Slaughterhouse workers were enrolled in the survey based on voluntary participation. A written consent was obtained from each participant. The survey contained questions on age, gender, country of birth, recent antibiotic use, job description, working in more than one section of the slaughterhouse (rotation), wearing plastic gloves, living on a livestock farm, and contact with family members working in healthcare or in livestock farming. Slaughterhouse workers were divided in three different categories according to their activities: contact with live pigs, dead pigs or other. When subjects indicated that they worked in more than one section, they were included in the category with the most intense contact with live animals.

Nasal swabs (Venturi Transystem, Copan Innovation, Italy) were taken from workers in order to determine the presence of MRSA. This study was approved by the Medical Ethical Committee of the University Hospital Utrecht (file no. 08/050).

Environmental sampling

To determine the MRSA status of the different slaughterhouse sections, environmental wipe samples were taken from surfaces in each section (Fig. 1) at the beginning and at the end of the working day using Sodibox wipes (Raisio Diagnostics B.V. Nieuwerkerk aan den IJssel, The Netherlands). Sections of the slaughterhouse were divided in two different categories according to the cleanliness of the animal/carcass: dirty or clean areas. In the dirty area, the carcass surface is cleaned by scalding, depilation and singeing. In the clean area, the carcass is eviscerated and processed into meat products.

Microbiological methods

Nasal swabs were incubated in Mueller–Hinton enrichment broth (Becton Dickinson, USA) with 6.5% NaCl, for 18–48 h at 35 °C. Then 10 µl of the broth was plated onto a MRSA-ID culture plate (bioMérieux, France), and incubated overnight at 35 °C. Suspect (green) colonies were identified as *S. aureus* by a latex agglutination test (Staphaurex Plus; Murex Diagnostics Ltd, UK) and tested for cefoxitin sensitivity by the disc diffusion method [28]. The obtained MRSA isolates were subsequently stored at –80 °C.




Fig. 1. Schematic representation of the sections of the production chain (dotted lines) in a pig slaughterhouse. The shaded area represents sections where live pigs are located (dirty area). Each human figure represents about 10 persons, circled persons are not actual slaughterhouse employees (livestock transport workers and official veterinarians and auxiliaries).

Environmental sample wipes were soaked in 100 ml Mueller–Hinton enrichment broth with 6.5% NaCl and incubated for 18 h at 37 °C. Next, 1 ml of the broth was transferred into 9 ml Phenol Red mannitol broth with 5 mg/ml ceftizoxime and 75 mg/ml aztreonam (bioMérieux) and incubated for 18 h at 37 °C. Subsequently, 10 µl of the suspension was transferred onto a Columbia agar plate with 5% sheep blood. In parallel, Brilliance MRSA culture plates (Oxoid, UK) were inoculated with 10 µl suspension and incubated for 18 h at 37 °C. Colonies were subcultured until pure.

Confirmation of the isolates was done by a multiplex PCR specific for *S. aureus* [29], the *mecA* gene [30], and the Panton–Valentine leucocidin (PVL) toxin genes [31]. Isolates were defined as MRSA on the basis of their *mecA* gene presence. Staphylococcal protein A (*spa*) typing was conducted according to Harmsen *et al.* [32]. On all MRSA-positive environmental and human samples, antimicrobial susceptibility was tested using the Vitek system (bioMérieux SA, France) according to the manufacturer's instructions.

Sample size and statistical analysis

The prevalence of MRSA nasal carriage in the general population in The Netherlands was assumed to be

<0.5%. A nasal carriage rate of $\geq 2\%$ in slaughterhouse workers was considered as a significant increase. The required sample size was calculated as 450 subjects ($\alpha = 0.05$, $\beta = 0.10$).

Prevalence of MRSA in slaughterhouse workers was calculated as a percentage of the total amount of samples in general and specified per category and job description. Wilson confidence intervals (CI) were calculated. Univariable exact logistic regression was performed using SAS, version 9.1 [33]. Odds ratios (OR) were determined by comparing different categories and job descriptions within those categories. In order to calculate the association between the human and environmental samples and because of the skewed distributions of the percentages of positive persons and environmental samples per section, Spearman's rank correlation was used.

RESULTS

Slaughterhouse characteristics

In the three selected slaughterhouses, the total number of employees varied between 80 and 260. The total number of slaughtered pigs per day varied between 3800 and 5000, all pigs originated from farms in The Netherlands. In one slaughterhouse, cattle were

Table 1. Prevalence of nasal MRSA carriage in slaughterhouse workers

Contact with pigs	Function	Total	MRSA	Percentage	95% CI
Live pigs	Livestock transport worker	41	9	22.0	12.0–36.7
	Official veterinarian + auxiliary	13	2	15.4	4.3–42.2
	Lairage worker	32	2	6.3	1.7–20.1
	Dirty area worker	7	1	14.3	2.6–51.3
Dead pigs*		127	0	0.0	0.0–2.9
Other†		29	0	0.0	0.0–11.7
Total		249	14	5.6	3.4–9.2

CI, Confidence interval (data from three slaughterhouses combined).

* Clean area worker, carcass cooling and cutting plant worker, green offal worker, meat hygiene inspector, quality assurance worker.

† Administrative and technical personnel.

slaughtered as well, but in separate rooms in the same building.

Humans

Of the total of 497 slaughterhouse workers 195 (39.2%) agreed to participate. An additional 41 livestock transport workers and 13 official veterinarians and auxiliaries (i.e. persons from the VWA, who monitor and assist the meat hygiene inspectors) were included, yielding a total of 249 study subjects, including 16 female participants. Mean age was 43 years (range 19–73 years), and the mean working week was 41 h (range 7–80 h).

We found an overall nasal MRSA prevalence of 5.6% in slaughterhouse workers (14/249, Table 1). MRSA carriage was found exclusively in persons having contact with live pigs (15.1%), compared to subjects not working with live pigs (0.0%, OR 38.2, Table 2).

Nine of the 41 (22%) livestock transport workers were MRSA positive, as well as 2/13 (15%) veterinarians and auxiliaries. In total, 3/195 (1.5%, 95% CI 0.5–4.4%) employees of slaughterhouses (excluding livestock transport workers and official veterinarians and auxiliaries) were MRSA positive; these were all working in the dirty area of the slaughterhouse. No specific slaughterhouse function proved to be a significant risk factor, when comparing different activities within the clean and the dirty areas. Twenty-three persons indicated working in both dirty and clean areas and only one of these was found MRSA-positive.

Regarding potential determinants and confounders, no significant difference in persons with and without MRSA was found (Table 2). Furthermore, no

significant differences in MRSA prevalence in humans between slaughterhouses were found.

Environment

At the start of the day MRSA was only found in environmental samples from the lairages (10/12) (Table 3, Fig. 1). At the end of the day MRSA was found in the lairages (11/12), the dirty (5/12) and clean (3/12) areas and green offal (1/3). Spearman's correlation coefficient, a measure for the correlation between MRSA status of the environmental samples and the humans working in these areas, is 0.75 ($P=0.002$). The squared correlation ($0.75 \times 0.75 = 0.56$) gives the coefficient of determination; 56% of variance in percentage of positive persons can be explained by environmental contamination.

Spa typing and antimicrobial susceptibility testing

In total, 14 human and 32 environmental MRSA strains were collected. The predominant *spa* type was t011 in both human subjects (11/14) and environmental samples (21/32). *Spa* type t108 was only found once in a human nasal sample, and also once in an environmental sample from the corresponding slaughterhouse. An additional 10 environmental isolates from the other slaughterhouses were typed as t108. *Spa* type t571 was only found once in environmental samples, and t034 and t1451 were found only once in humans, not in environmental samples of the corresponding slaughterhouse. From two environmental samples two different *spa* types were isolated, in both cases t011 and t108. PVL-positive strains were not found.

Table 2. *Univariable exact logistic regression analysis*

Characteristic	Total	MRSA	Percentage	OR	95% CI	P value
Female gender	16	0	0.0	Ref.		
Male gender	233	14	6.0	1.4	0.2–∞	0.77
Born abroad	60	1	1.7	0.2	0.0–1.6	0.22
Living on livestock farm	24	3	12.5	2.8	0.5–11.7	0.28
Recent antibiotic use	28	3	10.7	2.3	0.4–9.5	0.40
Contact with family members in healthcare or livestock farming	47	3	6.4	1.2	0.2–4.7	1.00
Working with live pigs	93	14	15.1	38.2	6.3–∞	<0.0001
Rotation	59	3	5.1	0.9	0.3–3.5	1.00
Always wearing plastic gloves	53	2	3.8	Ref.		
Sometimes wearing plastic gloves	76	6	7.9	2.2	0.4–22.9	0.57
Never wearing plastic gloves	113	6	5.3	1.4	0.2–14.9	1.00

OR, Odds ratio; CI, confidence interval; ref. reference category.

Boldface values belong to characteristics that are significantly related to MRSA, when comparing the presence of the relevant factor *vs.* the absence of it.

Table 3. *MRSA in environmental samples taken at start and end of working day*

Pigs	Department	Start of the day			End of the day		
		Total	MRSA	Percentage	Total	MRSA	Percentage
Live	Lairage	12	10	83.3	12	11	91.7
	Dirty area	12	0	0.0	12	5	41.7
Dead	Clean area	12	0	0.0	12	3	25.0
	Carcass cooling	12	0	0.0	12	0	0.0
	Cutting plant	8	0	0.0	8	0	0.0
	Green offal	3	0	0.0	3	1	30.0

Data from three slaughterhouses combined.

Antimicrobial susceptibility testing revealed that all MRSA isolates from humans and the environment are resistant against tetracycline (Table 4), and 19/46 isolates show combined erythromycin and clindamycin resistance. Furthermore, all isolates are sensitive for mupirocin and vancomycin (only human isolates tested). *Spa* type t108 appears to have less combined erythromycin + clindamycin resistance (0/11 = 0.0%) than t011 (17/32 = 53.1%, $P = 0.002$). No clear difference in resistance pattern between the human and environmental isolates was determined.

DISCUSSION

To our knowledge, this is the first study on the prevalence of nasal MRSA in pig slaughterhouse workers. Working with live pigs is the most important determinant for nasal CC398 carriage, justifying the present hospital infection control guidelines in The

Netherlands, which indicate that contact with live pigs is a risk factor for MRSA carriage. Working with dead pigs does not seem to be a risk factor for MRSA carriage.

The prevalence of 15.1% in persons working with live pigs is comparable to data found elsewhere, e.g. 26% and 14% in pig farmers and 12.5% in veterinarians attending an international pig health convention [1, 5, 34]. A low prevalence was found in Danish veterinarians (3.9%) [35], but higher nasal prevalences were found in German pig farmers on MRSA-positive farms (86%), German pig veterinarians (45%) and USA pig farmers (45%) [36, 37].

The overall MRSA prevalence in all subjects in the current study is 5.6%, which is significantly higher than the general population prevalence reported in The Netherlands (0.1%) [7, 8, 38]. The higher prevalence in livestock transport workers compared to lairage workers might be explained by the less intense

Table 4. Antimicrobial susceptibility profiles of all human and environmental MRSA isolates

Antimicrobial	Human (n=14)		Environmental (n=32)	
	Resistant	Percentage	Resistant	Percentage
Tetracycline	14	100.0	32	100.0
Erythromycin	8	57.1	12	37.5
Clindamycin	8	57.1	12	37.5
Gentamicin	1	7.1	11	34.4
Ciprofloxacin	0	0.0	6	18.8
Trimethoprim/ sulfamethoxazole	3	21.4	1	3.1
Rifampicin	0	0.0	0	0.0
Fusidic acid	0	0.0	0	0.0
Linezolid	0	0.0	0	0.0
Mupirocin	0	0.0	0	0.0
Tobramycin	1	7.1	n.t.	
Vancomycin	0	0.0	n.t.	
Nitrofurantoin	0	0.0	n.t.	
Neomycin	n.t.		1	3.1
Amikacin	n.t.		0	0.0

n.t., Not tested.

physical contact with pigs by lairage workers, who often use sticks to herd the animals. Transport workers earmark all animals at pick up and often herd the animals with their bare hands. Second, high-pressure spray cleaning of the truck may result in formation of MRSA aerosols, which can be inhaled by the transport worker. Insight into these mechanisms may give more information on the transmission route of MRSA.

During the day MRSA accumulates, particularly in the first stages of the production process, which predominantly deals with live pigs. Since pigs were loaded into the lairages at night, the lairages were not clean at the time of sample collection at the beginning of the day. Moreover, the lairages are cleaned every day, but not disinfected.

There is a significant association between the presence of MRSA in different sections, and the percentage of MRSA-positive persons working in these relevant sections. It is possible that acquisition of MRSA occurs through contaminated surfaces [39]. However, presence of MRSA on different surfaces does not necessarily imply that there is an increased risk of human MRSA acquisition via the environment: where the lairages have a high percentage of MRSA-positive samples at the end of the day (92%), a relatively low percentage of lairage workers had acquired the bacterium (6.3%). It is plausible that animals spread MRSA to both humans and the environment, and human acquisition of MRSA seems

to be more likely by contact with MRSA-positive animals than through environments with MRSA in dust or aerosols.

All *spa* types found in our study were previously confirmed as belonging to the CC398 livestock-associated MRSA clone [40]. The most predominant *spa* types in both human and environmental isolates were t011 and t108, which is in accord with previous studies in pigs and pig farmers [1, 4, 5, 22, 41, 42]. The subject with t034 was an official veterinarian and the *spa* type t1451 came from a livestock transport worker, these persons often have more animal contacts than in the slaughterhouse alone. Antimicrobial susceptibility, in particular tetracycline resistance was comparable to profiles found in other studies for livestock-associated MRSA [2, 5, 22].

The prevalence of MRSA found in retail meat in other studies is considerable, the prevalence of MRSA found in employees of pig slaughterhouses in this study is low. The role of slaughterhouse employees in transmitting MRSA to the meat products thus does not seem to be large. Especially as persons working with meat products were all negative in this study. This finding is in accord with an unpublished study (R. de Jonge, J. E. Verdier and A. H. Havelaar, unpublished observations), where none of 101 employees from the cold-meat processing industry and institutional kitchens carried MRSA. It is probable that another transmission route to retail meat is

involved here. Contamination of meat with MRSA by the environment (surfaces) and/or equipment, or from animals to carcasses/meat products is more likely to occur. This kind of cross-contamination has already been demonstrated for *Salmonella* spp. in pig slaughterhouses [43].

Our study has a few limitations. As with every questionnaire, survey recall bias, selection bias, and language bias may have occurred. Next, the low number of slaughterhouses visited ($n=3$) yields little power to find significant differences between slaughterhouses. Nevertheless, we assume that these results are representative for all Dutch pig slaughterhouses, because the working conditions in all pig slaughterhouses in The Netherlands are comparable due to automation and the strict legislation on hygiene and animal handling. Despite a smaller sample size than calculated beforehand, the number of subjects is still sufficient to confirm previous findings on the risk of acquiring MRSA for people in contact with live pigs. Possibly more risk factors could be found if the number of slaughterhouse workers was larger, e.g. country of birth, recent antibiotic use, amount of hours worked per week, and contact with healthcare. Furthermore, no pigs were sampled in our study, but in a previous study on MRSA at Dutch slaughterhouses MRSA was detected in 81% of the Dutch slaughter batches and 39% of the individual pigs [2]. Environmental samples are considered to be a good proxy for animal MRSA carriage, concerning the association found between environmental and animal samples in other studies (OR 27.5, $\kappa=0.68$) [44]. Longitudinal information on duration of MRSA carriage and the possibility of transient colonization is not yet available; this will be our group's next study subject.

In conclusion, nasal MRSA CC398 is found in pig slaughterhouse workers in significantly higher percentages than the general population prevalence in The Netherlands. It is found exclusively in persons working with live pigs. In addition to contact with live pigs, environmental contamination might also play a role in the acquisition of MRSA, but exact transmission routes from animals to humans remain to be elucidated in order to enable application of targeted preventive measures.

ACKNOWLEDGEMENTS

We thank the three slaughterhouses and in particular their quality assurance workers for their participation and help in the data collection in this study.

DECLARATION OF INTEREST

None.

REFERENCES

1. Voss A, *et al.* Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerging Infectious Diseases* 2005; **11**: 1965–1966.
2. de Neeling AJ, *et al.* High prevalence of methicillin resistant *Staphylococcus aureus* in pigs. *Veterinary Microbiology* 2007; **122**: 366–372.
3. Bens CC, Voss A, Klaassen CH. Presence of a novel DNA methylation enzyme in methicillin-resistant *Staphylococcus aureus* isolates associated with pig farming leads to uninterpretable results in standard pulsed-field gel electrophoresis analysis. *Journal of Clinical Microbiology* 2006; **44**: 1875–1876.
4. Huijsdens XW, *et al.* Community-acquired MRSA and pig-farming. *Annals of Clinical Microbiology and Antimicrobials* 2006; **5**: 26.
5. Broek van den IVF, *et al.* Methicillin-resistant *Staphylococcus aureus* in people living and working in pig farms. *Epidemiology and Infection* 2008; **137**: 700–708.
6. Graveland H, *et al.* Methicillin resistant *Staphylococcus aureus* (MRSA) in veal calf farmers and veal calves in the Netherlands. ASM Conference on Antimicrobial Resistance in Zoonotic Bacteria and Foodborne Pathogens, 2008. Copenhagen, Denmark: American Society for Microbiology, 2008.
7. Wertheim HF, *et al.* Low prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) at hospital admission in the Netherlands: the value of search and destroy and restrictive antibiotic use. *Journal of Hospital Infection* 2004; **56**: 321–325.
8. Dutch Working Party on Antibiotic Policy (SWAB). Nethmap 2008 – consumption of antimicrobial agents and antimicrobial resistance among medically important bacteria in the Netherlands (www.swab.nl). Accessed 24 January 2010.
9. Dutch Working Party on Infection Prevention (WIP). (<http://www.wip.nl>). Accessed 27 November 2009.
10. Haenen A, *et al.* Surveillance of MRSA in the Netherlands in 2007: an increasing trend of livestock-related MRSA [in Dutch]. *Infectieziekten Bulletin* 2009; **20**: 138–145.
11. National Institute of Public Health and the Environment. MRSA site (<http://www.rivm.nl/MRSA>). Accessed 27 November 2009.
12. The European Antimicrobial Resistance Surveillance System. (<http://www.rivm.nl/earss>). Accessed 27 November 2009.
13. de Boer E, *et al.* Prevalence of methicillin-resistant *Staphylococcus aureus* in meat. *International Journal of Food Microbiology* 2009; **134**: 52–56.
14. van Loo IH, *et al.* Methicillin-resistant *Staphylococcus aureus* in meat products, the Netherlands. *Emerging Infectious Diseases* 2007; **13**: 1753–1755.

15. **Lin J, et al.** *Staphylococcus aureus* isolated from pork and chicken carcasses in Taiwan: Prevalence and antimicrobial susceptibility. *Journal of Food Protection* 2009; **72**: 608–611.
16. **Pereira V, et al.** Characterization for enterotoxin production, virulence factors, and antibiotic susceptibility of *Staphylococcus aureus* isolates from various foods in Portugal. *Food Microbiology* 2009; **26**: 278–282.
17. **Pu S, Han F, Ge B.** Isolation and characterization of methicillin-resistant *Staphylococcus aureus* strains from Louisiana retail meats. *Applied and Environmental Microbiology* 2009; **75**: 265–267.
18. **Lee do K, et al.** New antimicrobial drug resistance and epidemiological typing patterns of staphylococci from clinical isolates and raw meats. *Archives of Pharmacological Research* 2008; **31**: 1016–1022.
19. **van Rijen MM, Van Keulen PH, Kluytmans JA.** Increase in a Dutch hospital of methicillin-resistant *Staphylococcus aureus* related to animal farming. *Clinical Infectious Diseases* 2008; **46**: 261–263.
20. **Wulf MW, et al.** First outbreak of methicillin-resistant *Staphylococcus aureus* ST398 in a Dutch hospital, June 2007. *Eurosurveillance* 2008; **13**.
21. **Fanoy E, et al.** An outbreak of non-typeable MRSA within a residential care facility. *Eurosurveillance* 2009; **14**.
22. **van Loo I, et al.** Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans. *Emerging Infectious Diseases* 2007; **13**: 1834–1839.
23. **Ekkelenkamp MB, et al.** Endocarditis due to methicillin-resistant *Staphylococcus aureus* originating from pigs [in Dutch]. *Nederlands Tijdschrift voor Geneeskunde* 2006; **150**: 2442–2447.
24. **Declercq P, et al.** Complicated community-acquired soft tissue infection by MRSA from porcine origin. *Infection* 2008; **36**: 590–592.
25. **Robicsek A, et al.** Prediction of methicillin-resistant *Staphylococcus aureus* involvement in disease sites by concomitant nasal sampling. *Journal of Clinical Microbiology* 2008; **46**: 588–592.
26. **Witte W, et al.** Methicillin-resistant *Staphylococcus aureus* ST398 in humans and animals, Central Europe. *Emerging Infectious Diseases* 2007; **13**: 255–258.
27. **Lewis HC, et al.** Pigs as source of methicillin-resistant *Staphylococcus aureus* CC398 infections in humans, Denmark. *Emerging Infectious Diseases* 2008; **14**: 1383–1389.
28. **CLSI.** Performance standards for antimicrobial susceptibility testing, 17th informational supplement. CLSI document M100-S17. Clinical and Laboratory Standards Institute, Wayne, PA, 2007.
29. **Martineau F, et al.** Species-specific and ubiquitous-DNA-based assays for rapid identification of *Staphylococcus aureus*. *Journal of Clinical Microbiology* 1998; **36**: 618–623.
30. **de Neeling AJ, et al.** Resistance of staphylococci in the Netherlands: surveillance by an electronic network during 1989–1995. *Journal of Antimicrobial Chemotherapy* 1998; **41**: 93–101.
31. **Lina G, et al.** Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clinical Infectious Diseases* 1999; **29**: 1128–1132.
32. **Harmsen D, et al.** Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. *Journal of Clinical Microbiology* 2003; **41**: 5442–5448.
33. **SAS.** SAS 9.1 (version SAS 9.1.3 service pack 3). SAS Institute Inc., Cary, NC, USA.
34. **Wulf MW, et al.** Prevalence of methicillin-resistant *Staphylococcus aureus* among veterinarians: An international study. *Clinical Microbiology and Infection* 2008; **14**: 29–34.
35. **Moodley A, et al.** High risk for nasal carriage of methicillin-resistant *Staphylococcus aureus* among Danish veterinary practitioners. *Scandinavian Journal of Work, Environment and Health* 2008; **34**: 151–157.
36. **Cuny C, et al.** Nasal colonization of humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without exposure to pigs. *PLoS ONE* 2009; **4**: e6800.
37. **Smith TC, et al.** Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS ONE* 2009; **4**: e4258.
38. **Donker G, Stobberingh E.** Is MRSA lurking everywhere? [in Dutch]. *Huisarts en wetenschap* 2008; **51**: 113.
39. **Boyce JM.** Environmental contamination makes an important contribution to hospital infection. *Journal of Hospital Infection* 2007; **65** (Suppl. 2): 50–54.
40. **Huijsdens XW, et al.** Molecular characterisation of PFGE non-typable methicillin-resistant *Staphylococcus aureus* in the Netherlands, 2007. *Eurosurveillance* 2009; **14**.
41. **Armand-Lefevre L, Ruimy R, Andreumont A.** Clonal comparison of *Staphylococcus aureus* isolates from healthy pig farmers, human controls, and pigs. *Emerging Infectious Diseases* 2005; **11**: 711–714.
42. **van Belkum A, et al.** Methicillin-resistant and -susceptible *Staphylococcus aureus* sequence type 398 in pigs and humans. *Emerging Infectious Diseases* 2008; **14**: 479–483.
43. **Prendergast DM, et al.** Prevalence and numbers of *Salmonella* spp. and enterobacteriaceae on pork cuts in abattoirs in the Republic of Ireland. *Journal of Applied Microbiology* 2008; **105**: 1209–1219.
44. **Broens EM, et al.** Prevalence study and risk factor analysis of NT-MRSA in pigs in the Netherlands. ASM Conference on Antimicrobial Resistance in Zoonotic Bacteria and Foodborne Pathogens, 2008. Copenhagen, Denmark: American Society for Microbiology, 2008.