

Cyanobacteria blooms cannot be controlled by Effective Microorganisms (EM[®]) from mud- or Bokashi-balls

Miquel Lurling · Yora Tolman ·
Frank van Oosterhout

Published online: 7 March 2010

© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract In controlled experiments, the ability of “Effective Microorganisms (EM, in the form of mudballs or Bokashi-balls)” was tested for clearing waters from cyanobacteria. We found suspensions of EM-mudballs up to 1 g l^{-1} to be ineffective in reducing cyanobacterial growth. In all controls and EM-mudball treatments up to 1 g l^{-1} the cyanobacterial chlorophyll-*a* (Chl-*a*) concentrations increased within 4 weeks from ≈ 120 to $325\text{--}435 \mu\text{g l}^{-1}$. When pieces of EM-mudballs (42.5 g) were added to 25-l lake water with cyanobacteria, no decrease of cyanobacteria as compared to untreated controls was observed. In contrast, after 4 weeks cyanobacterial Chl-*a* concentrations were significantly higher in EM-mudball treatments ($52 \mu\text{g l}^{-1}$) than in controls ($20 \mu\text{g l}^{-1}$). Only when suspensions with extremely

high EM-mudball concentrations were applied (i.e., 5 and 10 g l^{-1}), exceeding the recommended concentrations by orders of magnitude, cyanobacterial growth was inhibited and a bloom forming concentration was reduced strongly. In these high dosing treatments, the oxygen concentration dropped initially to very low levels of 1.8 g l^{-1} . This was most probably through forcing strong light limitation on the cyanobacteria caused by the high amount of clay and subsequent high turbidity of the water. Hence, this study yields no support for the hypothesis that EM is effective in preventing cyanobacterial proliferation or in terminating blooms. We consider EM products to be ineffective because they neither permanently bind nor remove phosphorus from eutroficated systems, they have no inhibiting effect on cyanobacteria, and they could even be an extra source of nutrients.

Guest editors: M. Meerhoff, M. Beklioglu, R. Burks, F. García-Rodríguez, N. Mazzeo & B. Moss / Structure and Function of World Shallow Lakes: Proceedings from the 6th Shallow Lakes Congress, held in Punta del Este, Uruguay, 23–28 November, 2008

M. Lurling (✉) · Y. Tolman · F. van Oosterhout
Aquatic Ecology & Water Quality Management Group,
Department of Environmental Sciences,
Wageningen University, P.O. Box 47, 6700 AA
Wageningen, The Netherlands
e-mail: miquel.lurling@wur.nl

Y. Tolman
Waterboard Delfland, P.O. Box 3061, 2061 DB Delft,
The Netherlands

Keywords Biological control · Eutrophication ·
Lake restoration · *Microcystis aeruginosa*

Introduction

Anthropogenic nutrient over-enrichment of surface waters is the primary cause of cyanobacterial blooms and nuisance scum's (Fogg, 1969; Reynolds & Walsby, 1975; Reynolds, 1987; Paerl, 1988, 2008). The massive blooms are a serious environmental problem and pose a risk to human health, as blooms

may contain a variety of very potent toxins, and may produce stomach-turning odors and may cause high turbidity, anoxia, fish kills, and food web alterations (Codd et al., 2005; Dittmann & Wiegand, 2006; Paerl, 2008; Paerl & Huisman, 2008, 2009). Climate change is expected to aggravate hazardous blooms favoring cyanobacterial dominance in a wide range of aquatic ecosystems (Paerl & Huisman, 2008, 2009). This expectation is underpinned by the coincidence of the two hottest summers in Europe, 2003 and 2006, since recording started (Luterbacher et al., 2004; Rebetz et al., 2009), with major cyanobacterial nuisance.

During the 2006 heat waves, at least 114 mostly shallow lakes and ponds in The Netherlands suffered heavily from cyanobacterial blooms. One of the seriously affected sites was Lake Gooimeer, a 2,576 ha shallow lake in The Netherlands, which includes a harbor (Almere-Haven) and a bathing area at the northern part of the lake. From August 2006 until the end of October 2006, a major cyanobacterial bloom caused fish kills and foul odors. As a consequence of this bloom, a swimming ban followed, the harbor was closed and the start and finish of the European Championship Triathlon had to be relocated to elsewhere. Following the 2006 heat-waves, many, mostly commercial, parties claimed to have Columbus's egg to alleviate these undesirable cyanobacteria blooms. One of the most heavily promoted products was "Effective-Microorganisms" (EM[®]) in so-called "mudballs" or "Bokashi-balls". They are based on the concept of Higa (1998) that addition of EM[®] would change the microbial community towards dominance of beneficial species, while suppressing harmful bacteria. Subsequently, EM-mudballs are claimed to wipe out cyanobacterial blooms, thereby making the infested waters clear again (<http://www.emvereniging.nl> accessed on June 28, 2008). The manufacturers intended to apply 12,000 EM-mudballs in Almere-Haven (The Netherlands). However, the authorities forbid the application of EM-mudballs because they contain heavy metals such as mercury as well as nutrients (Rijkswaterstaat, 2007). Moreover, the statements and claims about effectiveness were based on anecdotal evidence rather than on scientifically defensible evidence. In this study, we tested the claim that EM-mudballs would suppress and remove cyanobacteria from infested water in controlled experiments.

Materials and methods

EM-mudballs and phosphorus release

The EM-mudballs were obtained from Agriton BV (Noordwolde, The Netherlands) and belonged to a batch of 12,000 EM-mudballs that were made with the intention of applying in Almere-Haven (The Netherlands). Almere-Haven consists of two harbors, each one has the size of 4 ha (mean depth 1.9 m, max depth 3 m) that had suffered from major *Microcystis* blooms in 2006. The EM-mudballs had an average fresh weight of ± 300 g (A. de Puisselaar, B. V. Agriton, personal communication). The suggested dose was 1 ball per m², which makes maximum in situ concentrations of 0.1–0.3 g l⁻¹.

To get an estimate of the potential release of phosphate from EM-mudball material, 0.5 g EM-mudball material was brought into 100 ml nanopure water in triplicate. Three additional Erlenmeyer's only contained 100 ml nanopure water. The Erlenmeyer's were closed with Parafilm and placed for 48 h in an incubator in darkness at 22°C, and continuous orbital shaking (200 rpm). After this, the material was centrifuged for 5 min at 3,000 rpm, followed by filtration over a 0.45- μ m membrane filter. The filtrates were analyzed on soluble P as molybdate reactive phosphorus (Murphy & Riley, 1962) using a SKALAR auto-analyzer.

Cyanobacterial scum material was harvested from Almere-Haven (The Netherlands) on August 16th and 28th 2007. The collected material was counted under a NIKON light microscope using a Burkner Chamber and subjected to fluorescence analysis for estimating the contribution of major algal groups, i.e., cyanobacteria, chlorophytes, and diatoms to the overall chlorophyll-*a* (Chl-*a*) concentration using a PHYTO-PAM (Lürling & Verschoor, 2003). It revealed that the field material consisted of >95% *M. aeruginosa* (mostly present in large colonies) and some *M. flos-aquae*, *Anabaena* sp. and 2.5% chlorophytes. *Pseudooanabaena* sp. and *Navicula* sp. were present on the mucus of *Microcystis* colonies, but diatoms did not show up in the PHYTO-PAM signal.

Effect of EM-mudball material on cyanobacteria in artificial medium

In the first experiment, cyanobacterial scum material was inoculated in 400-ml beakers that contained

250 ml suspensions of artificial algal growth medium (WC) medium (Lüring & Beekman, 1999) and EM-mudball material. The EM-mudball was tested in the following concentration range: 0 (untreated control), 0.01, 0.1, 0.25, 0.5, 1.0, 5.0 and 10 g l⁻¹. Five replicates per concentration were used. The initial phytoplankton concentration was 127 (±8) µg l⁻¹ Chl-*a*, which consisted of 95% cyanobacteria (121 ± 6 µg l⁻¹). The Chl-*a* and the Photosystem II efficiency (Φ_{PSII}) were measured using a PHYTO-PAM phytoplankton analyzer (Heinz Walz GmbH, Effeltrich, Germany). One of features of the PHYTO-PAM phytoplankton analyzer is that it uses four different excitation wavelengths, which allows a separation between cyanobacteria, green algae, and diatoms (Kolbowski & Schreiber, 1995; Heinz Walz GmbH, 1999; Lüring & Verschoor, 2003).

The beakers were placed at random in a Gallenkamp ORBI-SAFE Netwise Orbital Incubator at 20°C, in 25 rpm and in a 18:6 h light:dark rhythm. The daily light:dark cycle was programmed so that the light intensity increased gradually to a maximum of 130 µmol quanta m⁻² s⁻¹ and subsequently decreased again to darkness, which resulted in a daily average light intensity of ~57 µmol photons m⁻² s⁻¹. Initially, and after 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, and 42 days samples were taken for analyses of Chl-*a* (and the share of cyanobacterial based Chl-*a*), pH (using a WTW-pH320 meter), conductivity (using a WTW-Cond 3300i meter), oxygen concentration and saturation (using an Oxyguard oxygen meter). Nano-pure water was added to compensate for evaporation losses. The water quality variables (conductivity, pH, oxygen concentration, and saturation) and Chl-*a* concentrations (and the share of cyanobacterial based Chl-*a*) were statistically analyzed running repeated measures ANOVA in the tool pack SPSS version 16.0.1. When the ANOVA indicated significant differences a Tukey post-hoc comparison test was run to distinguish means that were significantly different ($P < 0.05$).

Effect of EM-mudball on cyanobacteria from Almere-Haven

On August 28th 2007, 220 l water with cyanobacteria was collected in the harbor of Almere-Haven and transported immediately to the laboratory. Here, each of eight aquaria was filled with 25 l of the water. The

aquaria were placed in a temperature controlled water bath at 20°C and illuminated from above with 175 µmol photons m⁻² s⁻¹ in a 16:8 h light–dark cycle. The next day, four aquaria received an equal piece of EM-mudball (42.5 ± 2.2 g) each (treatments), whereas the four others were used as controls.

Initially and after 1, 2, 3, 4, 6, 8, 9, 10, 13, 15, and 28 days samples were taken for analysis of Chl-*a* and Φ_{PSII}, turbidity, pH, conductivity, oxygen concentration, and saturation as in the previous experiment. The temperature was measured with each of the above-mentioned meters. The evaporation losses were compensated with adding distilled water. The aquaria were sampled after mixing and resuspension of sedimented material.

Water quality variables (turbidity, temperature, conductivity, pH, oxygen concentration, and saturation) and phytoplankton characteristics (Chl-*a* and Φ_{PSII}) were statistically analyzed as in the previous experiment.

Results

EM-mudballs and phosphorus release

The phosphate concentrations in filtrates of EM-mudball suspensions (5 g l⁻¹) were 822 (±33) µg P l⁻¹ and significantly higher (t test: $t = 42.9$; $P < 0.001$) than those in the controls, which were 6 (±1) µg P l⁻¹. Hence, under the conditions employed in this study (48 h at 22°C in darkness at 200 rpm) EM-mudballs released 162 (±6) µg P g⁻¹ EM mudball.

Effect of EM-mudball material on cyanobacteria in artificial medium

In all controls and EM-mudball treatments up to 1 g l⁻¹ the cyanobacteria Chl-*a* concentration increased within 4 weeks from ≈120 µg l⁻¹ to 325–435 µg l⁻¹ (Fig. 1). At very high EM-mudball concentrations (5 and 10 g l⁻¹) cyanobacteria Chl-*a* concentration decreased from ≈120 µg l⁻¹ to on average 8 and 5 µg l⁻¹, respectively (Fig. 1). The repeated measure ANOVA indicated a significant time effect ($F_{14,434} = 84.4$; $P < 0.001$), a significant EM-mudball concentration effect ($F_{7,31} = 53.5$; $P < 0.001$) and a significant EM-mudball × time




Fig. 1 Total chlorophyll-*a* concentrations ($\mu\text{g l}^{-1}$, open bars) and cyanobacteria chlorophyll-*a* concentrations ($\mu\text{g l}^{-1}$, filled gray bars) in phytoplankton from the lake (inoculated scum from Almere-Haven area, i.e., mainly colonial *M. aeruginosa*)

in controls (0 g l^{-1}) and suspensions with different concentrations EM-mudball material ($0.01\text{--}10 \text{ g l}^{-1}$). Error bars indicate 1 SD ($n = 5$)

interaction ($F_{98,434} = 6.65$; $P < 0.001$). A Tukey post-hoc comparison test revealed three homogeneous groups that were significantly different from each other: (1) 0, 0.01, 0.1, 0.25 and $0.5 \text{ g EM-mudball l}^{-1}$, (2) 0.1, 0.25, 0.5 and 1.0 g l^{-1} , and (3) 5 and 10 g l^{-1} .


In as much as the collected and inoculated scum material from Almere-Haven was not exclusively comprised of cyanobacteria, but also contained some chlorophytes, over the course of the experiment gradually more green algae were detected (Fig. 1). During the course of the experiment, in controls and

EM-mudball treatments up to 1 g l^{-1} total Chl-*a* concentrations (cyanobacteria + eukaryotic algae) had increased from $127 \mu\text{g l}^{-1}$ to maximum values between 590 and $975 \mu\text{g l}^{-1}$ (Fig. 1). In contrast, high dosage of 5 and $10 \text{ g EM-mudball l}^{-1}$ reduced the initial Chl-*a* concentration of $127 \mu\text{g l}^{-1}$ to on average 49 and $37 \mu\text{g l}^{-1}$, respectively (Fig. 1). The repeated measure ANOVA indicated a significant time effect ($F_{14,434} = 90.6$; $P < 0.001$), a significant EM-mudball concentration effect ($F_{7,31} = 52.5$; $p < 0.001$) and a significant EM-mudball \times time

interaction ($F_{98,434} = 5.93$; $P < 0.001$). A Tukey post-hoc comparison test revealed four homogeneous groups that were significantly different from each other: (1) 0, 0.01, 0.1 and $0.5 \text{ g EM-mudball l}^{-1}$, (2) 0.01, 0.1, 0.25, and 0.5 g l^{-1} , (3) 0.25, 0.5 and 1 g l^{-1} , and (4) 5 and 10 g l^{-1} .

A repeated measure ANOVA revealed a significant EM-mudball concentration effect on Φ_{PSII} ($F_{7,31} = 58.7$; $P < 0.001$). The Φ_{PSII} was on average between 0.38 and 0.41 in controls and EM-mudball treatments up to 1 g l^{-1} (Fig. 2). In the 5 and

Fig. 2 The Photosystem II efficiency for cyanobacteria (inoculated scum from Almere-Haven area, i.e., mainly colonial *M. aeruginosa*) exposed in beakers to different concentrations EM-mudball material ($0.01\text{--}10 \text{ g l}^{-1}$). Error bars indicate 1 SD ($n = 5$)



10 g l⁻¹ treatments, Φ_{PSII} was on average 0.31 and 0.16, and showed considerable variation among replicate (Fig. 2). Tukey's test showed that Φ_{PSII} in controls and EM-mudball treatments up to 1 g l⁻¹ were similar, while the 5 and 10 g l⁻¹ treatments were significantly lower and also significantly different from each other.

Water quality variables were significantly influenced by the addition of different concentrations of EM-mudball material (Fig. 3). Suspensions with higher EM-mudball material showed rapid and stronger oxygen depletion followed by a gradual increase over the course of the experiment (Fig. 3A). Patterns for oxygen saturation were similar to those of oxygen concentration. The repeated measures ANOVA revealed significant time effects (oxygen concentration: $F_{14,434} = 183.7$; $P < 0.001$, saturation: $F_{14,434} = 87.9$; $P < 0.001$), significant EM-mudball concentration effects (oxygen concentration: $F_{7,31} = 525.2$; $P < 0.001$, saturation: $F_{7,31} = 622.4$;

$P < 0.001$), and significant time \times EM-mudball interactions (oxygen concentration: $F_{98,434} = 19.5$; $P < 0.001$, saturation: $F_{98,434} = 23.8$; $P < 0.001$). Tukey's post-hoc comparisons revealed similar homogeneous groups for both oxygen concentration and saturation: (1) 0 and 0.01 g l⁻¹, (2) 0.1 and 0.25 g l⁻¹, (3) 0.25 and 0.5 g l⁻¹, (4) 1 g l⁻¹, (5) 5 g l⁻¹, and (6) 10 g l⁻¹.

Adding EM-mudball material caused significant turbidity in the water ($F_{5,24} = 197.1$; $P < 0.001$). When turbidity was measured directly after the EM-mudball suspensions had been made, mean values were 77, 81, 108, 246, 324, 443, <1,000 and <1,000 NTU for controls and increasing EM-mudball concentrations, respectively (Fig. 3B). The majority of the suspended matter settled within 3 days and after 2–3 weeks, the increase in turbidity paralleled the increase in Chl-*a* (see Fig. 1).

No major differences in pH of the various incubations were observed (Fig. 3C). Nonetheless,




Fig. 3 Dissolved oxygen concentration (**A**; mg l⁻¹), the turbidity (**B**; NTU), the pH (**C**), and the conductivity (**D**; $\mu\text{S cm}^{-1}$) in beakers that contained 250 ml suspensions of cyanobacterial bloom material collected from the field, artificial growth medium

(WC-medium) and EM-mudball material included doses of 0, 0.01, 0.1, 0.25, 0.5, 1.0, 5.0, and 10 g l⁻¹. Error bars indicate 1 SD ($n = 5$)

the repeated measures ANOVA indicate a significant time effect ($F_{14,434} = 233.1$; $P < 0.001$), a significant EM-mudball effect ($F_{7,31} = 39.0$; $P < 0.001$), and a significant time \times EM-mudball interaction ($F_{98,434} = 9.14$; $P < 0.001$). Three homogeneous groups were distinguished by a Tukey test in the order from higher to lower pH: (1) 0–0.5 g l⁻¹, (2) 0.1–1 g l⁻¹, and (3) 5 and 10 g l⁻¹.

The conductivity of the water was affected by the EM-mudball material where higher concentrations of EM-mudball caused higher conductivity (Fig. 3D). The repeated measures ANOVA indicate a significant time effect ($F_{13,403} = 27.3$; $P < 0.001$), a significant EM-mudball effect ($F_{7,31} = 124.5$; $P < 0.001$), and a significant time \times EM-mudball interaction ($F_{91,403} = 2.50$; $P < 0.001$). Four homogeneous groups were distinguished by a Tukey test that in the order from lower to higher conductivity were: (1) 0–0.25 g l⁻¹, (2) 0, 0.25, 0.5, and 1 g l⁻¹, (3) 5 g l⁻¹, and (4) 10 g l⁻¹.

Effect of EM-mudball on cyanobacteria from Almere-Haven

The water from the Lake contained 113 $\mu\text{g l}^{-1}$ of Chl-*a* of which 96 $\mu\text{g l}^{-1}$ was from cyanobacteria. After 1 day in the laboratory, these concentrations had increased to 141 and 122 $\mu\text{g l}^{-1}$, respectively (Fig. 4). Then Chl-*a* concentrations both in treated and untreated aquaria decreased in a similar way for some days, stabilized for 1 week and started to increase again, but only in the treated aquaria (Fig. 4). Averaged over time there was no difference between controls and treatments in the total Chl-*a* concentrations ($F_{1,6} = 0.10$; $P = 0.761$) and the cyanobacteria Chl-*a* ($F_{1,6} = 0.13$; $P = 0.731$). However, after 28 days, total Chl-*a* ($t = 3.72$; $P = 0.010$) and the cyanobacterial Chl-*a* concentrations ($t = 6.00$; $P = 0.001$) were significantly higher in the treatments than in the controls (Fig. 4). Total Chl-*a* concentrations were 100 (± 21) $\mu\text{g l}^{-1}$ and 47 (± 19) $\mu\text{g l}^{-1}$ in treatments and controls, respectively. After 28 days, cyanobacterial Chl-*a* concentrations were 53 (± 7) $\mu\text{g l}^{-1}$ and 20 (± 8) $\mu\text{g l}^{-1}$ in treatments and controls, respectively. This might be caused by the release of phosphate as a 48 h incubation (in darkness, at 22°C) of EM-mudball material revealed a release of 162 (± 6) $\mu\text{g PO}_4\text{-P g}^{-1}$ EM mudball ($N = 3$).

In EM-mudball treatments, despite initial high productivity, within a few days a sharp drop in




Fig. 4 Cyanobacteria chlorophyll-*a* concentrations ($\mu\text{g l}^{-1}$) (A) and the total chlorophyll-*a* concentrations ($\mu\text{g l}^{-1}$) (B) in aquaria containing 25 l lake water from Almere-Haven without (*open symbols*, control) and with a part of EM-mudball (42.5 ± 2.2 g; *filled symbols*). Error bars indicate 1 SD ($n = 4$)

oxygen concentrations and saturation was observed (Fig. 5). After 9 days, oxygen concentrations in controls and treatments resembled each other. Averaged over the entire experimental period, oxygen concentrations ($F_{1,6} = 423.3$; $P < 0.001$) and saturation ($F_{1,6} = 361.0$; $P < 0.001$) were significantly higher in controls than in the EM-mudball treatments (Fig. 5). The pH appeared significantly lower in the EM-mudball treatments than in controls ($F_{1,6} = 130.5$; $P < 0.001$). Conductivity was significantly higher ($F_{1,6} = 58.8$; $P < 0.001$) in the EM-mudball treatments than in controls (Fig. 5). Turbidity ($F_{1,6} = 5.39$; $P = 0.059$) and temperature ($F_{1,6} = 1.40$; $P = 0.281$) were similar in controls and EM-mudball treated aquaria (Fig. 5).

The Φ_{PSII} varied between 0.4 and 0.55, but there was no difference ($F_{1,6} = 4.22$; $P = 0.086$) in Φ_{PSII} of cyanobacteria in controls and EM-mudball treated aquaria (Fig. 6).




Fig. 5 Dissolved oxygen concentration (A; mg l⁻¹) and saturation (B; %), the pH (C), the temperature (D; °C), the conductivity (E; μS cm⁻¹) and the turbidity (F; NTU) in aquaria

that contained 25 l from Almere-Haven without (*open symbols, control*) and with a part of an EM-mudball (42.5 ± 2.2 g; *filled symbols*). Error bars indicate 1 SD (n = 4)

Discussion

The results of this study clearly showed that within the 4–6 weeks experimental periods EM-mudballs were ineffective in suppressing and removing cyanobacteria from infested water, both in artificial medium and lake water. Only in suspensions with very high concentrations of EM-mudball material (i.e., 5 and 10 g l⁻¹), the phytoplankton biomass was

reduced strongly. This, however, can be attributed to initially very high turbidity of the water causing strong light inhibition, attachment to mud and burial of *Microcystis* (Verspagen et al., 2006). Moreover, it is not advisable using such high concentrations of 5 and 10 g l⁻¹ in situ as it would result in many additional centimeters of sediment, strong oxygen consumption which might be hazardous to many aquatic animals especially fish fauna, may cause




Fig. 6 The Photosystem II efficiency for cyanobacteria in aquaria containing lake water from Almere-Haven without (*open symbols*, control) and with a part of an EM-mudball (42.5 ± 2.2 g; *filled symbols*). Error bars indicate 1 SD ($n = 4$)

release of nutrients and the input of heavy metals (Rijkswaterstaat, 2007). Also when parts of EM-mudballs were dropped in lake water, resembling actual applications, the mudballs gradually disintegrated without having the claimed effect on the cyanobacteria. This experiment showed that after 4 weeks EM mudballs, as compared to the controls, actually promoted cyanobacterial growth rather than inhibiting it. In as much as the dosage was even higher than has been suggested being effective, the results are not in favor of the claim that EM-mudballs are effective in suppressing and removing cyanobacteria from infested water.

The underlying idea in adding EM-mudballs to a water body might be promotion of the bacterial flora as bacterial competitive superiority in terms of phosphorus uptake has been described well (e.g. Currie & Kalf, 1984a, b). However, in eutrophic lakes bacteria might be P and C limited leading to stronger increase in algal abundance than bacterial abundance (Lean & White, 1983; Currie, 1990). Bacteria are on the menu of grazers, such as heterotrophic nanoflagellates (Sheldon et al., 1986), while the success of cyanobacteria is often explained from their grazing resistance (Lampert, 1987). The grazing activity of heterotrophic nanoflagellates might promote the release of phosphate from the P-rich bacteria by several hundreds of percents (Bloem et al., 1988). Consequently, phytoplankton might flourish in freshwater systems despite bacteria often dominate the P uptake, which is corroborated

by controlled experiments revealing that bacterivorous nanoflagellates could prevent a strong competition for phosphate between algae and bacteria (Pengerud et al., 1987).

The manufacturers of the EM-mudballs claimed that the ‘Effective Micro-organisms’ would “consume the phosphates and eat away the cyanobacteria” (<http://www.emvereniging.nl/index2.php/weblog/C11/P34>). If the EM had been effective in removing phosphate, a sharp decline in the Φ_{PSII} of the cyanobacteria should have been observed (Wykoff et al., 1998). However, the Φ_{PSII} revealed the cyanobacteria were in good physiological state. Moreover, in artificial medium the cyanobacteria and algae expressed good growth even in high doses of EM-mudball suspensions, while in aquaria with EM-mudball material cyanobacteria remained present and reached higher concentrations than the controls. These results not only show that cyanobacteria were not eaten away by the so-called EM, but also suggest that EM-material might even further aggravate a cyanobacteria bloom through the release of phosphate. A short term (48 h) experiment revealed a release of phosphate of about $162 \mu\text{g P g}^{-1}$ EM mudball. In the 8 ha harbor area of Almere-Haven, based on the leachate experiment, the intended application of 12,000 EM-mudballs would then have led to an increase of water column P by $4 \mu\text{g l}^{-1}$, or even more through bacterial mineralization of mudball material on the sediment (Kairesalo et al., 1995). Further, as application of EM-mudballs cause a drop in oxygen concentration, release of phosphate direct from the sediment or associated bacteria is expected (Hupfer & Lewandowski, 2008). Hence, this study provided no evidence for the manufacturers claims.

Although it has been asserted that EM-cocktails contain about 80 species of micro-organisms, such as photosynthetic bacteria, lactic acid bacteria, actinomycetes, yeasts and fermenting fungi (Higa, 1998), such claims could not be confirmed by independent research (Van Egeraat, 1998; Van Vliet et al., 2006). The addition of EM to slurry manure had no effects on the composition of the bacterial community present in the slurry manure; EM had no effect on the chemical composition of the manure and no effect on grass production (Van Vliet et al., 2006). In experiments on agar plates, Van Egeraat (1998) found no effect of micro-organisms in EM-1[®] on the growth of eight test bacteria (species belonging to the

genera *Azotobacter*, *Rhizobium*, *Pseudomonas*, *Bacillus*, *Streptomyces*, *Mycobacterium*, *Serratia* and *Escherichia*), one yeast (*Saccharomyces*) and one fungus (*Penicillium*). Therefore, and because EM-products contain common micro-organisms already present in the environment, we doubt that an EM-application will have the potential to change the microflora in situ.

Manufacturers recommend dosages of 1 EM-mudball per square-meter or 1 ball per 1 to 10 m³ of water (<http://www.eminfo.nl/HuisEnTuinBrochNLweb.pdf>; www.em-natuurlijk-actief.com/NL-ned/HuisEnTuinBrochNLweb.pdf; <http://www.emamerica.com/>; Agriton BV). The total surface area of the Almere harbors is 8 ha, for which 12,000 mudballs in this case 0.15 mudball per square meter was considered sufficient by the manufacturers. It should, however, be taken into consideration that the Almere-Haven area is part of a large (2,576 ha) shallow lake (Gooimeer), where the majority of the *Microcystis* bloom is being produced outside the harbor area. The water quality in the Almere-Haven area is strongly influenced by the Gooimeer, and mitigating the cyanobacterial nuisance in the harbor area seems only to be possible by strong nutrient reduction in the large lake Gooimeer (Burger et al., 2007). With massive blooms and scum being produced on the lake and driven into the harbor area by wind action, the suggested EM-application in the harbor would have been of no use.

The Centre for Water Management (Ministry of Transport, Public Works and Water Management, Directorate, General for Public Works and Water Management, Lelystad, The Netherlands) has estimated a total of 0.25 gram mercury in the 12,000 mudballs, or about 21 µg per mudball (Rijkswaterstaat, 2007). Completely mixed through the water column in the harbor this would have yielded about 1.6 ng mercury l⁻¹. Although this concentration is below the maximum allowable concentration of 1.2 µg mercury l⁻¹ (RIVM, 2008), in regulation 2455/2001/EC of the European Union, mercury has been pointed out as a priority dangerous compound of which discharge into the water is no longer allowed (Rijkswaterstaat, 2007). As EM-products (EM-mudballs or EM-suspensions) might contain such undesirable compounds, might vary in quality (Van Vliet et al., 2006), will not permanently fix or remove phosphorus from eutroficated systems, have no inhibiting effect on cyanobacteria, and could even be a

source of nutrients, we consider EM non-effective in mitigating eutrophication as such and blooms of cyanobacteria in specific.

The primary cause of cyanobacterial blooms is an over-enrichment of the water with nutrients, in which phosphorus plays a key role (Schindler, 1977; Schindler et al., 2008). Hence, in reducing eutrophication, source-oriented measures principally targeting the phosphorus inflow and internal loading remain essential.

Acknowledgments We wish to thank Wendy Beekman, Dr. Carol Soares (AEW), and Dr. Hans Ruiter (Centre for Water Management) and Drs. Gert Butijn, and Ton Vrieze (Directie IJsselmeergebied) for fruitful discussions. We thank Mr. A. A. de Puisselaar (Agriton, B. V., Noordwolde, The Netherlands) for delivery of the EM-mudballs. This study was supported by the Ministry of Transport, Public Works and Water Management, Directorate, General for Public Works and Water Management, Directie IJsselmeergebied.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Bloem, J., M. Starink, M.-J. B. Bar-Gilissen & T. E. Cappenberg, 1988. Protozoan grazing, bacterial activity, and mineralization in two-stage continuous cultures. *Applied Environmental Microbiology* 54: 3113–3121.
- Burger, D., H. Los & S. Groot, 2007. Bestrijding van blauwalgen in de havenkom Almere-Haven. Rapport Q4328.00, WL|Delft Hydraulics: 50 pp (in Dutch).
- Codd, G. A., L. F. Morrison & J. S. Metcalf, 2005. Cyanobacterial toxins: risk management for health protection. *Toxicology and Applied Pharmacology* 203: 264–272.
- Currie, D. J., 1990. Large-scale variability and interactions among phytoplankton, bacterioplankton, and phosphorus. *Limnology and Oceanography* 35: 1437–1455.
- Currie, D. J. & J. Kallf, 1984a. A comparison of the abilities of freshwater algae and bacteria to acquire and retain phosphorus. *Limnology and Oceanography* 29: 298–310.
- Currie, D. J. & J. Kallf, 1984b. The relative importance of bacterioplankton and phytoplankton in phosphorus uptake in freshwater. *Limnology and Oceanography* 29: 311–321.
- Dittmann, E. & C. Wiegand, 2006. Cyanobacterial toxins – occurrence, biosynthesis and impact on human affairs. *Molecular Nutrition Food Research* 50: 7–17.
- Fogg, G. E., 1969. The physiology of an algal nuisance. *Proceedings of the Royal Society London B* 173: 175–189.
- Heinz Walz GmbH, 1999. *Phytoplankton Analyzer PHYTO-PAM. System Components and Principles of Operation*, 1st ed. <http://www.walz.com>.

- Higa, T., 1998. Effective Micro-organisms. For Sustainable Agriculture and Healthy Environment. Jan van Arkel, Utrecht: 191 pp (in Dutch).
- Hupfer, M. & J. Lewandowski, 2008. Oxygen controls the phosphorus release from lake sediments – a long-lasting paradigm in limnology. *International Review Hydrobiology* 93: 415–432.
- Kairesalo, T., L. Tuominen, H. Hartikainen & K. Rankinen, 1995. The role of bacteria in the nutrient exchange between sediment and water in a flow-through system. *Microbial Ecology* 29: 129–144.
- Kolbowski, J. & U. Schreiber, 1995. Computer-controlled phytoplankton analyzer based on a 4-wavelength PAM Chl fluorometer. In Mathis, P. (ed.), *Photosynthesis: From Light to Biosphere*, Vol. V. Kluwer Academic Publishers, Dordrecht: 825–828.
- Lampert, W., 1987. Feeding and nutrition in *Daphnia*. In Peters, R. H. & R. De Bernardi (eds), *Daphnia*. *Memorie dell'Istituto Italiano di Idrobiologia* 45:143–192.
- Lean, D. R. S. & E. White, 1983. Chemical and radio-tracer measurement of phosphorus uptake by lake plankton. *Canadian Journal of Fisheries and Aquatic Sciences* 40: 147–155.
- Lürling, M. & W. Beekman, 1999. Grazer-induced defenses in *Scenedesmus* (Chlorococcales; Chlorophyceae): coenobium and spine formation. *Phycologia* 38: 368–376.
- Lürling, M. & A. M. Verschoor, 2003. F_0 -spectra of chlorophyll fluorescence for the determination of zooplankton grazing. *Hydrobiologia* 491: 145–157.
- Luterbacher, J., D. Dietrich, E. Xoplaki, M. Grosjean & H. Wanner, 2004. European seasonal and annual temperature variability, trends, and extremes since 1500. *Science* 303: 1499–1503.
- Murphy, J. & J. P. Riley, 1962. A modified single solution method for determination of phosphate in natural waters. *Analytica Chimica Acta* 27: 31–36.
- Paerl, H. W., 1988. Nuisance phytoplankton blooms in coastal, estuarine, and inland waters. *Limnology and Oceanography* 33: 823–847.
- Paerl, H. W., 2008. Nutrient and other environmental controls of harmful cyanobacterial blooms along the freshwater-marine continuum. In Hudnell, K. E. (ed.), *Cyanobacterial Harmful Algal Blooms: State of the Science and Research Needs*, Chapter 10. *Advances in Experimental Medicine and Biology* 619: 217–237.
- Paerl, H. W. & J. Huisman, 2008. Blooms like it hot. *Science* 320: 57–58.
- Paerl, H. W. & J. Huisman, 2009. Climate change: a catalyst for global expansion of harmful cyanobacterial blooms. *Environmental Microbiology Reports* 1: 27–37.
- Pengerud, B., E. F. Skjoldal & T. F. Thingstad, 1987. The reciprocal interaction between degradation of glucose and ecosystem structure. *Studies in mixed chemostat cultures of marine bacteria, algae, and bacterivorous nanoflagellates*. *Marine Ecology Progress Series* 35: 111–117.
- Rebetez, M., O. Dupont & M. Giroud, 2009. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 95: 1–7.
- Reynolds, C. S., 1987. Cyanobacterial water-blooms. *Advances in Botanical Research* 13: 67–143.
- Reynolds, C. S. & A. E. Walsby, 1975. Water blooms. *Biological Reviews of the Cambridge Philosophical Society* 50: 437–481.
- Rijkswaterstaat, 2007. Advies Beheer-en bestrijdingsmaatregelen Blauwalg: 19 pp (in Dutch).
- RIVM, 2008. Kwik en kwikverbindingen. http://www.rivm.nl/rvs/stiffen/prio/totale_prior_stoffenlijst.jsp (in Dutch).
- Schindler, D. W., 1977. Evolution of phosphorus limitation in lakes: natural mechanisms compensate for deficiencies of nitrogen and carbon in eutrophied lakes. *Science* 195: 260–262.
- Schindler, D. W., R. E. Hecky, D. L. Findlay, M. P. Stainton, B. R. Parker, M. J. Paterson, K. G. Beaty, M. Lyng & S. E. M. Kasian, 2008. Eutrophication of lakes cannot be controlled by reducing nitrogen input: results of a 37-year whole-ecosystem experiment. *Proceedings of the National Academy of Sciences USA* 105: 11254–11258.
- Sheldon, R. W., P. Nival & F. Rassoulzadegan, 1986. An experimental investigation of a flagellate-ciliate-copepod food-chain with some observations relevant to the linear biomass hypothesis. *Limnology and Oceanography* 31: 184–188.
- Van Egeraat, A. W. S. M., 1998. EM1 and the Growth of Microorganisms. *Micro-organisms in EM1*. Wageningen University, The Netherlands (in Dutch).
- Van Vliet, P. C. J., J. Bloem & R. G. M. de Goede, 2006. Microbial diversity, nitrogen loss and grass production after addition of Effective Micro-organisms® (EM) to slurry manure. *Applied Soil Ecology* 32: 188–198.
- Verspagen, J. M. H., P. M. Visser & J. Huisman, 2006. Aggregation with clay causes sedimentation of the buoyant cyanobacterium *Microcystis*. *Aquatic Microbial Ecology* 44: 165–174.
- Wykoff, D. D., J. P. Davies, A. Melis & A. R. Grossman, 1998. The regulation of photosynthetic electron transport during nutrient deprivation in *Chlamydomonas reinhardtii*. *Plant Physiology* 117: 129–139.