

Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands

A. P. Schrier-Uijl · A. J. Veraart ·
P. A. Leffelaar · F. Berendse · E. M. Veenendaal

Received: 8 January 2010 / Accepted: 26 March 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Shallow fresh water bodies in peat areas are important contributors to greenhouse gas fluxes to the atmosphere. In this study we determined the magnitude of CH₄ and CO₂ fluxes from 12 water bodies in Dutch wetlands during the summer season and studied the factors that might regulate emissions of CH₄ and CO₂ from these lakes and ditches. The lakes and ditches acted as CO₂ and CH₄ sources of emissions to the atmosphere; the fluxes from the ditches were significantly larger than the fluxes from the lakes. The mean greenhouse gas flux from ditches and lakes amounted to 129.1 ± 8.2 (mean \pm SE) and 61.5 ± 7.1 mg m⁻² h⁻¹ for CO₂ and 33.7 ± 9.3 and 3.9 ± 1.6 mg m⁻² h⁻¹ for CH₄, respectively. In most water bodies CH₄ was the dominant greenhouse gas in terms of warming potential. Trophic status of

the water and the sediment was an important factor regulating emissions. By using multiple linear regression 87% of the variation in CH₄ could be explained by PO₄³⁻ concentration in the sediment and Fe²⁺ concentration in the water, and 89% of the CO₂ flux could be explained by depth, EC and pH of the water. Decreasing the nutrient loads and input of organic substrates to ditches and lakes by for example reducing application of fertilizers and manure within the catchments and decreasing upward seepage of nutrient rich water from the surrounding area will likely reduce summer emissions of CO₂ and CH₄ from these water bodies.

Keywords Biogeochemistry · Drainage ditches · Greenhouse gas emission · Nutrients · Peatlands · Temperate lakes

A. P. Schrier-Uijl (✉) · F. Berendse · E. M. Veenendaal
Nature Conservation and Plant Ecology Group,
Wageningen University, Droevendaalse steeg 3a,
6708 PD Wageningen, The Netherlands
e-mail: arina.schrier@wur.nl

A. J. Veraart
Department of Aquatic Ecology and Water Quality
Management, Wageningen University, P.O. Box 47,
6700 AA Wageningen, The Netherlands

P. A. Leffelaar
Plant Production Systems Group, Wageningen University,
P.O. Box 430, 6700 AK Wageningen, The Netherlands

Introduction

Freshwater bodies such as ditches, streams, wetlands and lakes contribute appreciably to the processing of carbon and its transport to the atmosphere (e.g. Bastviken et al. 2004; Walter et al. 2006; Wang et al. 2006). It has been estimated that lakes annually emit 8–48 Tg methane (CH₄), which is 6–16% of the global natural CH₄ emissions (Bastviken et al. 2004; St. Louis et al. 2000), and 513 Tg C carbon dioxide (CO₂) (Cole et al. 1994). Saarnio et al. (2009) have

estimated that large lakes alone account for 24% of all wetland CH_4 emissions in Europe. It has been shown that small water bodies also significantly contribute to the landscape-scale CH_4 budgets in wetland regions (e.g. Schrier-Uijl et al. 2009a, b; Juutinen et al. 2009; Repo et al. 2007; Walter et al. 2007; Roulet and Moore 1995). Yet though it is likely that both lakes with organic-rich sediment and also eutrophic ditches contribute especially significantly to regional greenhouse gas balances, they are poorly studied and very little is known about their underlying biogeochemical processes (Saarnio et al. 2009).

CO_2 is produced by respiration in sediments and throughout the water column and can also be a product of biological processes in the sediment (Fig. 1). As CO_2 is highly soluble, high concentrations can accumulate near the sediment/water interface, which results in oversaturation and release to the atmosphere. It has been suggested that the transport of dissolved organic carbon (DOC) from terrestrial environments is an important source of carbon in aquatic environments. If this is the case, lakes in organic-rich peatlands have larger CO_2 fluxes than lakes in mineral catchments (Rantakari and Kortelainen 2005; Huttunen et al. 2002).

CH_4 emission is the balance of two counteracting processes: methanogenesis in anoxic conditions and the oxidation of the generated CH_4 (Minkinen and Laine 2006; Bastviken et al. 2002), (Fig. 1). CH_4 is a major product of carbon metabolism in lakes; its production depends on the availability of alternative electron acceptors such as O_2 , NO_3^- , Fe^{3+} and SO_4^{2-}

(van Bodegom and Scholten 2001). After these electron acceptors have been used up, CH_4 production becomes the dominating degradation process of organic matter and is the terminal microbial process in the anaerobic degradation of organic matter. The CH_4 travels from the sediment through the water column to the atmosphere and on the way it can be oxidised into CO_2 (Whiting and Chanton 2001). Most of the CH_4 that remains unoxidised will be emitted by diffusive flux to the atmosphere.

The underlying microbial processes affecting CO_2 and CH_4 production and emission are regulated by variables such as sediment and water temperature, oxygen availability, organic matter availability and composition, sediment and water chemistry, the presence of electron acceptors (redox conditions), pH, electrical conductivity (EC) and factors such as water depth and lake size (e.g. Stadmark and Leonardson 2005; Juutinen et al. 2009; Repo et al. 2007; Frei et al. 2006; Loeb et al. 2007; Casper et al. 2003).

Most of the freshwater lakes in the Netherlands are in peat areas, are very shallow (<2 m), and were created by large-scale dredging and removal of peat during the early seventeenth century (Gulati and van Donk 2002). They vary in area, depth, hydrology and physico-chemical characteristics, but most of them are eutrophic, due to the application of fertilisers and manure within their catchments, the oxidation of peat and the upward seepage of nutrient-rich water from the surrounding area. Drainage since the Middle Ages has resulted in the typical landscape of narrow fields separated by drainage ditches (Fig. 2).




Fig. 1 Simplified illustration of CO_2 and CH_4 dynamics in water bodies, with *OM* organic matter



Fig. 2 The typical Dutch peat area landscape of lakes and narrow fields separated by drainage ditches (TOP10vector Product information 2007)

Shallow fresh water bodies are not very well understood in terms of their greenhouse gas emissions and have not been incorporated in previous regional or global greenhouse gas budgets. The emission from these water bodies is probably high and poses an international rather than a domestic problem because in many lowland regions of Europe agriculture continues to contribute appreciably to the nutrient loading of lakes and ditches (Gulati and van Donk 2002; Lamers et al. 1998).

About 16% of the total area (41,864 km²) of the Netherlands is covered by water, mostly classified as wetland (Gulati and van Donk 2002) and with over 300,000 km of drainage ditches. Much of this wetland is in peat areas. It is very important to quantify how these wetlands contribute to the greenhouse gas balance and which factors regulate the emission. In this study we focus on the high-emitting temperate lakes and drainage ditches in peat areas in the Netherlands and on many variables that can alter the emission of CH₄ and CO₂. The two aims of this study were (1) to quantify CH₄ and CO₂ fluxes from shallow lakes and drainage ditches in the Netherlands during a 3-week period in the summer season and (2) to identify the factors that regulate the emissions of CH₄ and CO₂ from lakes and ditches.

Materials and methods

Study sites

Measurements were performed in a 3-week period between June 16th and July 6th in the summer of 2009 in 5 shallow fresh water lakes and 14 drainage ditches at 7 locations in peat areas in the Netherlands (Fig. 3).

The 5 lakes are located in peat areas in the Netherlands and differ in trophic status (de Haan et al. 1993) and depth (Table 1). L4 and L5 are located in the east of the Netherlands where the subsoil consists of mesotrophic to oligotrophic sedge-peat overlying sand. The other lakes are located in the southwest of the Netherlands where the subsoil consists of eutrophic to mesotrophic reed-sedge peat and alder carr peat.

Drainage ditches at 7 locations in different peat areas in the Netherlands were sampled. They differed in trophic status and water depth (Table 2). At each




Fig. 3 Geographical distribution of the 5 sampled lakes (L1–L5) and 14 sampled drainage ditches (at 7 locations) (D1–D7) in peat areas in the Netherlands

location 2 connected ditches were sampled and because there were no significant differences between them related to water quality they were treated as 1 location in the analyses. All the drainage ditches sampled contained some aquatic vegetation.

Measurements

Flux measurements and calculation of fluxes

Detailed measurements of CH₄ emission and CO₂ emission were performed with floating chambers from a dinghy at different locations in the lakes and drainage ditches. We measured the emissions from each lake on two different days. On each of these days we measured at three different locations per lake, and repeated the measurements five times at each location. This yielded 30 measurements per lake. Each ditch was sampled on 1 day in the 3-week period, with 8 replicates per ditch. This yielded 16 measurements per location in the two connected ditches. All measurements were performed between 10.30 and 14.30 h. Data quality was assessed and outliers resulting from disturbances were removed from the dataset. Emissions of CH₄, CO₂ and N₂O were determined using a closed dark chamber method and a Photo Acoustic Field Gas Monitor (INNOVA 1412 sn, 710-113, ENMO services, Belgium) connected to a PVC chamber by Teflon tubing (e.g. van

Table 1 Characteristics of the lakes sampled

Lake	Abbreviation	Trophic status	Lake size (ha)	Average depth (m)
Reeuwijkse plas	L1	Eutrophic	927	2.1
Vinkeveense plas	L2	Eutrophic	1,079	2.4
Nieuwkoopse plas	L3	Eutrophic	676	2.5
Belterwiede	L4	Mesotrophic to eutrophic	613	1.8
Schutsloterwiede	L5	Mesotrophic to eutrophic	141	1.2

Table 2 Characteristics of the drainage ditches

Drainage ditch	Abbreviation	Trophic status	Ditch width (m)	Mean water depth (m)
Oukoop	D1	Eutrophic	8	0.25
Stein	D2	Mesotrophic	6	0.28
Horstermeer	D3	Eutrophic	3	0.90
Drie berken	D4	Mesotrophic	3	0.87
Koole	D5	Mesotrophic	3	0.80
Sint Jans	D6	Eutrophic	2	0.28
Doosje	D7	Mesotrophic	6	0.43

Huissteden et al. 2005; Hendriks et al. 2007). Fluxes of N₂O appeared to be too low to detect with the gas analyzer, therefore the N₂O flux measurements were not included in the analyses. Samples were taken from the headspace of this closed cylindrical dark chamber (30 cm diameter, 25 cm height). Gas samples were taken every minute during a 5-min period and every single measurement was checked on linearity of the build up of the gas concentration in the chamber. This check eliminated about 30% of the measurements. The slope dC/dt of the gas concentration curve at time $t = 0$ was estimated using linear regression (e.g. van Huissteden et al. 2005; Schrier-Uijl et al. 2009b). A small fan was installed in the chamber to homogenise the inside air and a water lock was used to control pressure in the chamber. We used a floater to place the chamber onto the water surface, carefully avoiding the effect of pressure differences and the disturbance of the water surface (for details, see Schrier-Uijl et al. 2009a). Since the gas monitor software does not compensate fully for cross-interference of CO₂ and water vapour at high concentrations, air was led through glass tubes filled with silica gel and soda lime before it entered the gas analyser, to remove water vapour. To cross-validate the chamber-based measurements, we also performed eddy covariance measurements on L1 at the same time and location and compared these with the chamber measurements

within the footprint of the system. The eddy covariance system was located along a boardwalk in L1 and the footprint of the mast was on the lake. Within this footprint chamber measurements were performed on the lake during a period of 4 h. The two independent methods had previously been compared at different temporal scales in a heterogeneous landscape of fields and ditches (Schrier-Uijl et al. 2009b; Kroon et al. 2007).

Variables measured

At each lake and drainage ditch we measured water temperature and pH at two depths (10 and 30 cm and at 25 cm depth in D1 and D2), dissolved oxygen at 10 cm intervals from the water surface to the sediment surface, and the EC at 10 cm depth. Oxygen, pH, temperature and EC were measured with an HQ multiprobe with a luminescent dissolved oxygen sensor (Hach Company, Loveland, Colorado, USA). The variables investigated in the ditches were the dissolved CH₄ concentrations at the water surface, the middle of the water column and in the water immediately above the sediment. Samples for dissolved methane analysis were taken using an airtight 20 ml glass syringe at three depths in the water column: at the sediment surface, at the water surface, and at a depth half-way in between. The water samples were

transferred into airtight glass Exetainers[®] (Labco, high Wycombe, UK) containing 120 μl ZnCl_2 to halt biological processes; to prevent air bubbles being trapped in these vials they were filled to overflowing before being capped. The samples were stored in water at 20°C until analysis. Dissolved methane was measured by membrane inlet mass spectrometry (MIMS) (Lloyd and Scott 1983) using an OmniStar[™] Gas Analysis System (Pfeiffer Vacuum, Asslar, Germany), equipped with a quadrupole QMS 200 mass spectrometer with a Channeltron detector (Burle Industries). The MS was operated by Quadstar 32-bit software for data acquisition. The sample was pumped through a water bath at 20°C before passing through silicon membrane tubing in which gases were released to the MS. An inlet as described by Kana (1994) was used for the analysis, but without using a cryotrap, as this would have frozen out the methane. Instead, to prevent confounding effects of water vapour, the inlet at the MS side was heated to 180°C. Methane was measured at mass to charge ration (m/z) of 15, as a pre-calibration experiment had shown that this gave the most reliable results. Concentrations of methane were calculated by comparing the ion current at m/z 15 of the sample to the ion current at m/z 15 of air-saturated water at 20°C.

The water in each lake was sampled at three locations with 3 replicates (mixed sample). The water in each ditch was sampled at two locations with 3 replicates (mixed sample). Undisturbed sediment samples were taken from the sediment top layer (upper 10 cm) by means of a plastic cup perforated with holes 2 cm apart at the end of a length-adjustable pipe.

Two of the three water samples were filtered immediately with a Whatman 0.45 μm cellulose membrane filter (Whatman International Ltd, Maidstone, England); the third sample was not filtered. All samples were transported in coolers and stored frozen (−20°C) until analyses. The unfiltered water samples were analysed for organic matter (OM) content, %C, %N, Chlorophyll-*a* content, total N and total P; the filtered samples were analysed for $\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , SO_4^{2-} , Fe^{2+} and PO_4^{3-} using a SAN^{plus} autoanalyzer (Skalar Analytical, Breda, the Netherlands). Dissolved organic carbon (DOC) and dissolved inorganic carbon (DIC) were measured in filtered samples using a carbon analyser. Total N and total P were measured using a SAN^{plus} auto analyser with laser destructor. All these samples were measured in

duplicate. Chlorophyll-*a* content in unfiltered samples from the microcosms was measured using a phyto-PAM fluorometer (Heinz Waltz GmbH, Effeltrich, Germany). For the sediment samples a CaCl_2 extraction was used to obtain the available $\text{PO}_4\text{-P}$, $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$, and an ammonium oxalate extraction was used to obtain the active form of Fe.

Data analysis

Correlations between the measured variables and fluxes of CO_2 and CH_4 were first tested by using Pearson correlation analysis. Data were tested for normality. We used stepwise, multiple linear regression analyses to quantify the relationships between environmental variables and fluxes of CO_2 and CH_4 (SPSS 15.0). The variables that significantly enhanced the emissions of CO_2 and CH_4 were selected and were used to build regression models. Differences in the fluxes and variables between and within lakes and ditches were tested using one-way ANOVA (SPSS 15.0).

Results

Climatic variables

During the sampling period the mean day air temperatures ranged from 15 to 25°C, the average temperature at the surface of the water bodies studied ranged from 19.2 to 25.4°C and the wind speed at 3 m above water level ranged from 2.1 to 4.5 m s^{-1} (Table 3).

Characteristics of lakes and drainage ditches

The lakes and drainage ditches were humic, shallow and nutrient-rich. The sediment in D6 and D7 had the lowest organic matter content because these two ditches are located in an area with shallow peat on sand. The EC in all the lakes and ditches sampled ranged from 269–866 $\mu\text{S cm}^{-1}$; the pH ranged from 6.8–9.0, with the highest values in the lakes (Table 4).

Emissions to the atmosphere

Lakes and drainage ditches studied acted as sources of CO_2 and CH_4 emissions to the atmosphere

Table 3 Average (mean \pm SD) water temperatures at 10 cm depth (T_{10} in $^{\circ}\text{C}$), at 30 cm depth (T_{30} in $^{\circ}\text{C}$) and average wind speed (U in m s^{-1}) during

	Mean T_{10}	Mean T_{30}	Mean U
D1	20.1 (\pm 0.1)	19.5 (\pm 0.9)	4.1
D2	25.4 (\pm 2.5)	25.2 (\pm 2.4)	3
D3	19.2 (\pm 0.6)	18.4 (\pm 0.4)	2.7
D4	25 (\pm 0.5)	25.8 (\pm 0.3)	3
D5	22.3 (\pm 0.3)	22 (\pm 0.1)	4.6
D6	20.2 (\pm 0.7)	19.1 (\pm 1.4)	2.1
D7	22.8 (\pm 0.2)	21.8 (\pm 1.2)	3.9
L1	23.6 (\pm 3.0)	23.7 (\pm 0.5)	3
L2	21.4 (\pm 1.1)	20.3 (\pm 0.5)	3.6
L3	23.3 (\pm 1.1)	23.2 (\pm 1.1)	3
L4	23.2 (\pm 1.4)	22.7 (\pm 1.2)	3.9
L5	25 (\pm 1.2)	24.9 (\pm 0.6)	3.9

For the location codes see Tables 1 and 2

(Figs. 4, 5), except for L1 where a small uptake of CO_2 was measured. The mean release of both gases to the atmosphere was significantly higher from the ditches than from the lakes ($P < 0.001$).

The contribution of CO_2 emission compared to CH_4 emission in terms of warming potential is given in Fig. 6, where CH_4 fluxes have been transformed to CO_2 equivalents (CH_4 is 23 times as potent as CO_2).

Lakes

The emission of CO_2 from the lakes ($n = 93$) ranged from -6.0 to $123.9 \text{ mg m}^{-2} \text{ h}^{-1}$ and CH_4 emission ($n = 96$) ranged from 1.4 – $18.1 \text{ mg m}^{-2} \text{ h}^{-1}$. The CO_2 fluxes from L1 were significantly lower than those from the other lakes ($P < 0.01$); the CO_2 fluxes from L2 were significantly higher than those from L3, L1 and L5 ($P < 0.05$). The highest CH_4 emission was measured from L5 and the lowest from L2, but the differences were not significant. The lakes acted as sources of emissions of both gases, except for L1 that acted as a very small sink for CO_2 . In terms of warming potential, in 3 lakes the dominant emitted greenhouse gas was CH_4 and in 2 lakes it was CO_2 (Fig. 6).

Ditches

The emission of CO_2 from the drainage ditches ($n = 80$) ranged from $69.6 \text{ mg m}^{-2} \text{ h}^{-1}$ to $199.0 \text{ mg m}^{-2} \text{ h}^{-1}$ and CH_4 emission ($n = 79$) ranged from 1.2 to $39.3 \text{ mg m}^{-2} \text{ h}^{-1}$. The CO_2 emission from D3 was significantly higher than the fluxes from D7, D1 and D5. The highest CH_4 emission was measured from D4 and the lowest from D7, but the CH_4 fluxes did not differ significantly because there was great variability among the ditches. In all ditches except D6 and D7, the dominant greenhouse gas in terms of

Table 4 General characteristics of the lakes and ditches sampled, with standard deviations for lake/ditch depth, EC, pH and percentage organic matter in the sediment

Site	Lake size (ha) or Ditch width (m)	Sediment type	Mean depth (cm)	Mean EC ($\mu\text{S cm}^{-1}$)	Mean pH	Organic matter sediment (% dry weight)
L1	927	Peat	209.2 (\pm 42)	508.7 (\pm 53)	9.0 (\pm 0.3)	46.63 (\pm 9.5)
L2	1079	Peat	240.0 (\pm 42)	866.0 (\pm 13)	8.4 (\pm 0.1)	36.89 (\pm 12.9)
L3	676	Peat	253.3 (\pm 71)	392.5 (\pm 2)	8.2 (\pm 0.1)	63.08 (\pm 14.1)
L4	613	Peat on sand	178.0 (\pm 27)	398.0 (\pm 9)	8.4 (\pm 0.1)	58.72 (\pm 3.4)
L5	141	Peat on sand	120.0 (\pm 21)	393.0 (\pm 15)	8.4 (\pm 0.2)	59.81 (\pm 5.8)
D1	8	Peat	25.0 (\pm 3)	523.0 (\pm 38)	9.0 (\pm 0.0)	55.94 (\pm 3.0)
D2	6	Peat	27.5 (\pm 9)	269.3 (\pm 46)	6.8 (\pm 0.3)	57.37 (\pm 7.9)
D3	3	Peat	90.0 (\pm 14)	662.0 (\pm 48)	7.0 (\pm 0.4)	37.77 (\pm 3.7)
D4	3	Peat	86.7 (\pm 8)	386.7 (\pm 2)	7.2 (\pm 0.3)	74.22 (\pm 0.8)
D5	3	Peat	80.0 (\pm 17)	387.0 (\pm 2)	7.7 (\pm 0.1)	73.89 (\pm 0.8)
D6	2	Peat on sand	27.5 (\pm 3)	350.3 (\pm 26)	7.2 (\pm 0.3)	9.53 (\pm 3.1)
D7	6	Peat on sand	42.5 (\pm 3)	395.5 (\pm 6)	8.1 (\pm 0.3)	2.65 (\pm 2.4)

Lake area was determined from Top10vector maps: TDN (2006)

Fig. 4 Mean CO₂ fluxes (mg m⁻² h⁻¹) and their standard deviations in the sampling period for the ditches (D1–D7) and lakes (L1–L5) at different locations in peat areas in the Netherlands. Positive flux values indicate release from the water to the atmosphere




Fig. 5 Mean CH₄ fluxes (mg m⁻² h⁻¹) with their standard deviations for the ditches (D1–D7) and lakes (L1–L5) at different locations in peat areas in the Netherlands. Ditches were sampled on 1 day ($n = 16$ per location) in the period 16 June–6 July 2009; Lakes were sampled during 2 days in the same period ($n = 24$ per location). Positive flux values indicate release from the water to the atmosphere





Fig. 6 Contribution to greenhouse warming of CO₂ emission (dark grey) compared to CH₄ emission (light grey) in lakes and ditches, given in CO₂ equivalents



warming potential was CH₄. All ditches acted as sources of emissions of both gases.

Cross-validation

Large-scale CH₄ flux measurements by eddy covariance were performed on one of the lakes (L1) to cross-validate flux values from this homogeneous landscape on a diurnal base. A cross-validation of chamber based CH₄ and CO₂ values and eddy covariance based values

is also performed earlier for a more heterogeneous peat area (Schrier-Uijl et al. 2009b). Details for the used eddy covariance instruments have been reported in Veenendaal et al. 2007 and Kroon et al. 2007. In this study, CH₄ fluxes within the footprint of the eddy covariance system were 5.8 ± 3.26 (mean \pm SD, $n = 24$) measured by chambers compared to 4.6 ± 1.3 measured by eddy covariance over a 4-h period. It would be of great interest in the future to also use eddy covariance to capture temporal variability of

greenhouse gas fluxes (CH_4 and CO_2) from water bodies and to explain more of the measured variability.

Dissolved oxygen and dissolved CH_4

Typical vertical profiles of oxygen saturation during the measurements are shown in Fig. 7.

On average, the lakes had a higher O_2 saturation than ditches. In both types of waterbody, oxygen saturation decreased only slightly at the top of the water column, which suggests that there was hardly any respiration by aquatic organisms. Deeper in the water column the oxygen saturation fell rapidly to values close to 0% just above the sediment. Of the lakes, L2 had the highest O_2 saturation throughout the profile, and of the ditches D3 and D6 had the lowest O_2 saturation.

Dissolved CH_4 concentrations were measured at three depths: at the top and middle of the water column and just above the ditch sediments. In all the ditches the dissolved CH_4 concentrations increased with depth (Fig. 8).

Concentrations of dissolved CH_4 ($\mu\text{g l}^{-1}$) in the water of ditches at the top, middle and bottom of the water column. The y-axis shows the depth (cm).

D1 and D3 had a high dissolved CH_4 concentration and also a high CH_4 emission (Fig. 5). None of the following variables correlated significantly with the dissolved CH_4 in the ditch water immediately above the sediment or with the difference between dissolved CH_4 concentration at the water surface and

at the sediment surface: nutrient content (NO_3^- , NH_4^+ , Fe, PO_4^{3-}); sediment oxygen demand (SOD); O_2 saturation of the water, organic matter content (% organic matter, %N, %C); amount of green algae and plants. We did not find any significant correlation between dissolved CH_4 concentration at the water surface and CH_4 release to the atmosphere. The oxygen saturation at the sediment surface correlated negatively with CH_4 emission to the atmosphere ($P = 0.065$).

The variables measured and their correlation with CH_4 and CO_2 emission

Climate, depth, EC and pH

Climatic conditions in the 3-week sampling period were stable. No significant correlation was found between the CO_2 and CH_4 fluxes and the temperature or the wind velocity. Neither the depth of water in the ditches (range 0.28–0.90 m) or the depth of water in the lakes (range 1.20–2.53 m) correlated significantly with CO_2 or CH_4 release to the atmosphere, although the deepest lakes tended to have the lowest CH_4 and CO_2 emissions. A positive correlation was found between EC and CO_2 flux and a significant negative correlation was found between CO_2 emission and the pH of the water ($r = -0.81$; $P = 0.001$). Though the correlation between CH_4 emission and pH was also negative, it was not significant ($r = -0.23$; $P = 0.41$).

Fig. 7 Oxygen saturation (%) in the water columns of lakes (right) and drainage ditches (left) at different depths in the water column at the time of the measurements





Fig. 8 Concentrations of dissolved CH_4 ($\mu\text{g l}^{-1}$) in the water of ditches at the top, middle and bottom of the water column. The y-axis shows the depth of the ditch (cm)



Nutrients and organic matter in water and sediment

The percentage of N measured in the lake sediments was significantly positively correlated with the release of CO₂ ($P < 0.01$); in ditches, the %N and the %OM measured in the sediments were significantly positively correlated with the release of CH₄ ($P = 0.02$ and $P = 0.05$, respectively). The lowest organic matter contents of the sediments were found in D6, D7 and L2 (Table 4), which had the lowest CH₄ fluxes (Fig. 5). In this study neither the DIC nor the DOC correlated significantly with the CO₂ flux or the CH₄ flux.

The ammonium (NH₄⁺) concentration in the water ranged from 0.1 to 478.6 µg NH₄-N l⁻¹; the highest concentrations were found in D3, D6 and L2. Ammonium concentration correlated positively with CO₂ emission ($r = 0.67$; $P < 0.05$). The NO₃⁻ concentration in the water was around 0 mg N l⁻¹—except for L2 and L4, where the mean concentrations were 0.43 and 0.12 mg N l⁻¹, respectively. In the sediment of the lakes and ditches the NH₄⁺ concentrations ranged from 12.3 to 478.1 mg/kg dry weight with the highest concentrations in D4 (324.2 mg kg⁻¹ dry weight) and D5 (478.1 mg kg⁻¹ dry weight). The NO₃-N concentration ranged from 0.0 to 3.55 mg kg⁻¹ dry weight, with the highest concentration in D4 and the lowest in L4. The only lake with high NO₃ concentrations in the water and sediment was L2: it was also the only lake where measurable N₂O emissions were observed (0.163 mg m⁻² h⁻¹,

$n = 23$). See Table 5 for the concentrations of NH₄⁺ and table 6 for the concentrations of NO₃-N.

A weak negative correlation was found between the SO₄²⁻ concentrations in lake water (range 17.8–53.4 mg l⁻¹) and ditch water (range 3.4–47.1 mg l⁻¹) and CO₂ and CH₄ fluxes ($r = -0.43$, $P = 0.16$; $r = -0.1$, $P = 0.81$). The water of D3 had the lowest SO₄²⁻ concentrations of all the lakes and ditches sampled; this ditch had a high Fe concentration in its water and sediment (Table 6), which suggests the binding of SO₄²⁻ to iron. Overall, the SO₄²⁻ concentrations did not significantly differ between the ditches and lakes. The Pearson correlations between CO₂, CH₄ and Fe²⁺ in the water were weakly positive; the Fe²⁺ concentrations we measured ranged from 33.6 to 1,032 µg l⁻¹, with an average of 301.6 µg l⁻¹. Methane emission correlated significantly positively with the PO₄³⁻ concentration of the sediments of the lakes and ditches ($r = 0.77$, $P = 0.81$). The PO₄³⁻ concentration of the sediments correlated positively both with the Fe concentration and the SO₄²⁻ concentration of water. The total P concentration in the water correlated positively with CO₂ emission, indicating the high availability of organic substrates. The nutrient concentrations are given in Tables 5 and 6.

Multiple regression analyses

Multiple regression with stepwise elimination of variables showed that for summer CH₄ fluxes the

Table 5 Chemical composition of the water of lakes and ditches with means for sulphate (SO₄²⁻), iron (Fe²⁺), phosphate (PO₄³⁻), ammonium (NH₄⁺), nitrate (NO₃⁻) and total P

Location	SO ₄ ²⁻ (mg l ⁻¹)	Fe ²⁺ (µg l ⁻¹)	PO ₄ ³⁻ (µg P l ⁻¹)	NH ₄ ⁺ (µg N l ⁻¹)	NO ₃ ⁻ (mg N l ⁻¹)	Ptot (µg P l ⁻¹)
D1	47.1	324.3	98	13.8	0.00	153.4
D2	16.6	864.5	33	0.1	0.00	66.3
D3	3.4	1032.0	291	478.6	0.00	515.7
D4	21.6	58.5	11	8.8	0.00	18.3
D5	22.1	26.7	24	9.2	0.00	15.0
D6	7.1	470.5	240	166.4	0.00	201.0
D7	19.2	146.7	45	3.8	0.00	15.8
L1	34.8	48	53	12.2	0.00	11.3
L2	53.4	59.7	11	76	0.42	14.0
L3	21.0	31.7	25	11.5	0.00	10.3
L4	18.2	113.5	10	15.8	0.12	12.6
L5	16.9	640.0	18	6.1	0.00	25.4

For location codes see Tables 1 and 2

Table 6 Means for Fe, PO₄-P, NH₄-N, NO₃-N in the top10 cm of the bottom sediments of the sampled lakes and drainage ditches

Location	Fe (mg kg ⁻¹)	PO ₄ -P (mg kg ⁻¹)	NH ₄ -N (mg kg ⁻¹)	NO ₃ -N (mg kg ⁻¹)
D1	2.7	8.6	154.9	0.4
D2	3.0	0.6	215.4	0.7
D3	7.0	0.3	187.8	0.9
D4	6	20.7	478.1	3.6
D5	3.1	8.8	323.6	0.5
D6	2.0	0.2	51.9	0.2
D7	0.8	0.1	12.3	0.1
L1	1.5	2.6	106.0	0.4
L2	1.7	3.3	8.0	1.2
L3	0.5	5.2	123.7	0.3
L4	5.7	0.5	86.7	0
L5	na	na	na	na

Table 7 Statistical details of the CH₄ and the CO₂ multiple regression models with PO₄ concentration in the sediment, Fe concentration in the water, depth, EC and pH as explanatory

variables: R square of the model, significance of the model, Pearson correlations and significance of the separate variables


	R ² model	P model	Pearson corr.			P parameter		
			PO ₄ sediment	Fe _{water}		PO ₄ sediment	Fe _{water}	
CH ₄ model	0.87	0.000	0.77	0.19		0.000	0.003	
			Depth	EC	pH	Depth	EC	pH
CO ₂ model	0.89	0.000	-0.62	0.17	-0.81	0.06	0.004	0.001

PO₄³⁻ concentrations in the sediment and the Fe²⁺ concentrations in the water explained 87% of the variation when Eq. 1 was used:

$$F_{\text{CH}_4} = -3.482 + 2.183[\text{PO}_4]_{\text{sediment}} + 22.116[\text{FE}]_{\text{water}} \quad (1)$$

where F_{CH_4} is the CH₄ flux (mg m⁻² h⁻¹), $[\text{PO}_4]_{\text{sediment}}$ is the PO₄-P concentration in the sediment (mg kg⁻¹) and $[\text{FE}]_{\text{water}}$ is the concentration of Fe²⁺ in the water (mg l⁻¹). Table 7 presents statistical details of the model. Fig. 9 shows the measured CH₄ fluxes versus the CH₄ fluxes in the sampled lakes and ditches, modelled by means of Eq. 1. By performing regression analyses with ditches only, the fit of the regression improved to R² = 0.94.

For CO₂, a regression model with mean depth of lake or ditch water and the EC and pH of the water as independent variables explained up to 89% of the variation in summer CO₂ emission at the water-atmosphere interface when Eq. 2 was used:

**Fig. 9** Measured CH₄ fluxes versus modelled CH₄ fluxes (mg m⁻² h⁻¹) based on a multiple regression model with PO₄-P concentration in the sediment and Fe²⁺ concentration in the water as explanatory variables. The grey line is the 1:1 line

$$F_{\text{CO}_2} = 477.359 - 0.213\text{depth} + 0.171\text{EC} - 54.4\text{pH} \quad (2)$$

where F_{CO_2} is the CO₂ flux, depth is the mean depth of the water in the sampled lake or ditch (cm), EC is the mean electrical conductivity and pH is the mean pH in the sampled lakes and ditches. Figure 10 shows




Fig. 10 Measured CO₂ flux versus modelled CO₂ flux (mg m⁻² h⁻¹) based on a multiple regression model with mean depth of water, EC and pH as explanatory variables. The grey line is the 1:1 line

the measured CO₂ fluxes in lakes and ditches versus the modelled CO₂ fluxes by means of Eq. 2. Table 7 presents statistical details of the model.

Regression analyses on the data from the ditches only improved the predictive power of the regression to $R^2 = 0.91$.

Discussion

In this study we determined the magnitude of CH₄ and CO₂ fluxes from 12 water bodies in Dutch wetlands during a 3-week period in the summer season and studied the factors that might regulate emissions of CH₄ and CO₂ from these lakes and ditches. During this period the lakes and ditches acted as CO₂ and CH₄ sources of emissions to the atmosphere; the fluxes from the ditches were significantly larger. One lake (L1) was in equilibrium with the atmosphere in terms

of CO₂ emission. Kosten et al. 2010 found that <10% of lakes worldwide are in equilibrium with the atmosphere in terms of $p\text{CO}_2$, and they found that most other lakes are CO₂ sources. Compared with other studies, the lake emissions founding our study were in the intermediate to high range (see Table 8 for CH₄ fluxes). For example, Rantakari and Kortelainen (2008) found CO₂ fluxes in the range 7.48–11.5 mg m⁻² h⁻¹ in 37 boreal Finnish lakes. The average CH₄ emission from our drainage ditches was higher than the lake fluxes found in other studies (Table 7). As the CH₄ emissions measured by the gas analyser within the footprint area of an eddy covariance system at location L2 agreed within the uncertainty limits with the EC system, we are confident that our measurement technique provided reliable flux estimates that are applicable to larger areas (Kroon et al. 2007; Schrier-Uijl et al. 2009b).

The temporal variability of emissions of CH₄ and CO₂ from water bodies is normally found to be related to temperature and wind velocity when measuring over longer time spans (e.g. Stadmark and Leonardson 2005; Frei et al. 2006; Hendriks et al. 2007; Repo et al. 2007; Schrier-Uijl et al. 2009a, b; Kroon et al., in press). However, a large part of the variability of fluxes cannot be explained by temperature or wind velocity only. Our results refer to data collected during summer, a period in which around 70% of the annual ditch emissions are generated. The study did not last long enough to include seasonal patterns of CH₄ and CO₂ production and emission. Diurnal stratification and mixing due to day–night temperature differences may bias flux estimates if the

Table 8 Comparison between the CH₄ emission rates in this study and the CH₄ emission rates reported in other studies

Reference	System	Location	Sampling period	Flux CH ₄ (mg m ⁻² h ⁻¹)
Guerin and Abril 2007	Tropical lake	French Guiana	Late spring	4 ± 4.7
Juutinen et al. 2009	30 Eutrophic Boreal lakes	Finland	All seasons	Median of 0.137
Stadmark and Leonardson 2005	3 Ponds	South Sweden	Summer	10
Huttunen et al. 2002	Boreal lakes	Finland	Summer	1.0
Bastviken et al. 2004	11 Lakes	North America	Summer 2000	Range 0.15–3.2
Repo et al. 2007	3 Boreal lakes	Siberia	Summer	0.34
Present research	5 Temperate lakes	Netherlands	Early summer 2009	1.4–18.1, mean 5.0
Present research	7 Drainage ditches	Netherlands	Early summer 2009	1.2–39.3, mean 18.8

Mean CH₄ emission rates are in mg CH₄ m⁻² h⁻¹. The period of sampling and the location are given

For units: mg CH₄ m⁻² h⁻¹ refer to mg CH₄ emitted per m² of water area per hour

only measurements available are from the daytime (Repo et al. 2007). In Schrier-Uijl et al. (2009a, b) the diurnal variation of CH_4 fluxes over an area with fields and ditches was tested in October/November 2006. After correction for temperature dependency, the emission of CH_4 did not differ significantly between day and night. Nevertheless, there could be diurnal variation of fluxes from water bodies, because less oxygen will be produced at night, which will result in a lower redox potential and higher CH_4 production. In addition, less CO_2 will be taken up by aquatic plants at night, because then they are not photosynthesising. These effects should be considered when estimating annual fluxes from water bodies by using continuous measurements such as eddy covariance. In our study, only diffusive fluxes of CO_2 and CH_4 were measured; however, ebullition can also contribute to the emission of CH_4 from water bodies (Walter et al. 2006; Walter et al. 2007). While sampling ditches D5 and D4, we observed ebullition, so it is possible that we underestimated the release of CH_4 fluxes. In a summer study done by Repo et al. (2007) in Siberian water bodies, ebullition was observed in two of the three lakes sampled (depth <1.5 m) and accounted for 19–37% and 11–40% of the total CH_4 emissions from these two lakes.

The fact that the lakes and ditches acted as sources for CH_4 and CO_2 indicates that CO_2 production exceeded CO_2 uptake during photosynthesis by plants and that CH_4 production exceeded CH_4 oxidation. Our observation that the deeper, mostly less eutrophic lakes with low EC had the smallest fluxes agrees with findings reported for lakes in the boreal zone in Finland (depth range 3.8–26.5 m) (Juutinen et al. 2009). Deeper water bodies usually have less degradable organic matter and more oxidation of CH_4 than shallow lakes, because the transport pathway is longer (e.g. Borges et al. 2004). The EC, which is an indicator of trophic status in fresh water lakes, and the depth of the water body were two of the three significant predictors in the regression analyses for CO_2 fluxes.

The pH correlated negatively with emissions from both gases, yet at lower pH values (pH < 7) the correlations are usually positive (e.g. Inubushi et al. 2005). CO_2 enters the water as a result of the biological processes of organic carbon degradation and respiration by plants. In our ecosystems it is likely that through uptake of CO_2 by plants during the

day in the growing season, HCO_3^- is transformed to CO_2 , causing the HCO_3^- concentration to decline and diminishing the buffering effect. This reduced buffering effect can result in pH values above 9.0 and in a negative relation between CO_2 flux and pH. Incorporating pH in the regression equation for CO_2 significantly improved the equation's predictive power. In peat soils in temperate areas the optimum pH for methanogenesis is between 5.5 and 7.0, which explains the slightly negative correlation we found between CH_4 emission and pH (Le Mer and Roger 2001).

Water turbulence due to wind can increase mixing of oxygen in the water. This is illustrated by the higher oxygen concentration throughout the water column of L2, which had been subjected to high wind speeds on the day before sampling. The high O_2 saturation in L2 corresponded with higher CO_2 fluxes and very low CH_4 fluxes, illustrating the oxidation of CH_4 to CO_2 . The opposite can be seen in D3, D6 and L5, where the oxygen concentrations at the water surface were low and the CH_4 fluxes were high, illustrating the low turnover of CH_4 carbon to CO_2 . The fast decrease in dissolved CH_4 from the sediments to the water surface in D3 and D1 indicates that most of the dissolved CH_4 is oxidised during transport or passes through the water column and escapes to the atmosphere very quickly. As found in other studies, dissolved CH_4 poorly predicted the diffusive fluxes at the water–air interface (e.g. Huttunen et al. 2006; Juutinen et al. 2009). The factors responsible for this finding could be variation in duration of storage, release of CH_4 to the atmosphere and complex processes during transport of CH_4 through the water column (e.g. Kankaala et al. 2003).

The input of organic matter as a substrate in the lake or drainage ditch system increases the availability of substrates and this can increase the production of CO_2 and CH_4 (e.g. Casper 1992) and increases the possibility of minimising the competition for electron donors between methanogenesis and other anaerobic processes (Scholten et al. 2002; Scheid et al. 2003). As long as O_2 reaches the sediments, it will act as the primary oxidant of organic matter.

Permanently anaerobic conditions in the sediment may hamper nitrification of NH_4^+ to NO_3^- , but stimulate denitrification of NO_3^- to N_2 by microorganisms, leading to a high NH_4^+ to NO_3^- ratio, as was found in this study. Also, the greater availability

of NH_4^+ compared to NO_3^- suggests the occurrence of dissimilatory reduction of NO_3^- to NH_4^+ (DNRA) under anaerobic conditions in these ditches. DNRA is likely to occur in the organic sediments that we sampled, as this process usually occurs at high carbon inputs (Burgin and Hamilton 2007). As our ditch systems did not contain much aquatic plant biomass, it is unlikely that the NO_3^- uptake by plants and algae was influential in the ditches. Other possible sources of NH_4^+ in the water could be cation exchange of adsorbed NH_4^+ by Fe^{2+} (but this only occurs at very high Fe^{2+} concentrations: Loeb et al. 2007), and leaching through groundwater from surrounding, managed agricultural areas. NH_4^+ inhibits methanotrophy and therefore may reduce CH_4 oxidation and increase its emission (Conrad and Rothfuss 1991), which may explain the positive correlation between the NH_4^+ and CH_4 fluxes in our study. The positive correlation of NH_4^+ with CO_2 emission is in line with the findings of other studies.

Our finding is that the two most significant predictors of CH_4 fluxes were the PO_4^{3-} concentration in the sediment of lakes and ditches and the Fe^{2+} concentration in the water of lakes and ditches. In anaerobic sediments, Fe^{3+} will be reduced to Fe^{2+} . At the sediment–water interface some of the Fe^{2+} will be oxidized to Fe^{3+} (how much depends on the oxygen concentration just above the sediment) and some of this will be released into the water. Thus a high concentration of Fe^{2+} in the water is related to anaerobic conditions. Both Fe concentration and SO_4^{2-} concentration correlate with PO_4^{3-} availability at the sediment–water interface. The PO_4^{3-} in sediments is bound to Fe^{3+} and when the Fe^{3+} is reduced to Fe^{2+} , PO_4^{3-} will be released to the water (e.g. Smolders et al. 2006; Smolders and Roelofs 1993).

Overall, a higher trophic status was positively correlated with summer emissions of CO_2 and CH_4 , while the depth of the water and the pH were inversely correlated with CO_2 emission. It is therefore likely that decreasing the inputs of organic matter and nutrients (for example, by changing the management of the surrounding areas) will reduce emissions and that this effect will be strongest in drainage ditches.

Much of the uncertainty in flux estimates is due to temporal variation. So, also diurnal, seasonal, annual and inter annual variability must be studied in more detail to get insight in climatic responses, extreme

drought/rainfall events and the influence of management in the surrounding catchments. In this respect, there is a need for long-term, continuous measurements of emissions (e.g. by eddy covariance).

Conclusion

The current study focused on emissions from temperate, shallow lakes ($n = 5$) and drainage ditches ($n = 14$) in agricultural peat areas in the Netherlands. It was found that in general, both these types of waterbodies are important sources of CO_2 and CH_4 . The ditches had significantly higher CO_2 and CH_4 fluxes than the lakes. Trophic status was an important indicator of the magnitude of fluxes. 87% of the variation in the summer fluxes of CH_4 could be explained by PO_4^{3-} in the sediment and Fe^{2+} concentration in the water, and 89% of the CO_2 flux could be explained by water depth, EC and pH. Our results can be used to refine greenhouse gas emission inventories and to ascertain possible ways of reducing the release of CO_2 and CH_4 from water bodies to the atmosphere. Decreasing the nutrient loads and input of organic substrates to ditches and lakes will likely reduce summer emissions of CO_2 and CH_4 from these water bodies.

Acknowledgements Many thanks to Roos Loeb and Joop Harmsen for their critical comments on an earlier version of this manuscript, and to Jan van Walsem, Frits Gilissen and John Beijer for their assistance in the field and in the laboratory. We are also grateful to the State Forestry Service for the use of their boats and to Arjan Hensen and Petra Kroon who carried out the eddy covariance measurements at L1. Joy Burrough advised on the English. This study was funded by Wageningen University, The Province of North Holland, CarboEurope-IP and the Klimaat voor Ruimte Project (BSIK).

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Bastviken D, Ejlertsson J, Tranvik L (2002) Measurement of methane oxidation in lakes: a comparison of methods. *Environ Sci Tech* 36:3354–3361
- Bastviken D, Cole J, Pace M et al (2004) Methane emissions from lakes: dependence of lake characteristics two

- regional assessments and a global estimate. *Glob Biogeochem Cycles* 18:GB4009. doi: [10.1029/2004GB002238](https://doi.org/10.1029/2004GB002238)
- Borges AV, Delille B, Schiettecatte LS et al (2004) Gas transfer velocities of CO₂ in three European estuaries (Randers Fjord, Scheldt, and Thames). *Limnol Oceanogr* 49(5):1630–1641
- Burgin AJ, Hamilton SK (2007) Have we overemphasized the role of denitrification in aquatic ecosystems? A review of nitrate removal pathways. *Front Ecol Environ* 5(2):89–96
- Casper P (1992) Methane production in lakes of different trophic state. *Arch Hydrobiol Beih Ergebn Limnol* 37:149–154
- Casper P, Chan OC, Furtrado ALS et al (2003) Methane in an acidic bog lake: the influence of peat in the catchment on the biogeochemistry of methane. *Aquat Sci* 65:36–46. doi: [1015-1621/03/010036-11](https://doi.org/10.1015-1621/03/010036-11)
- Cole JJ, Caraco NF, Kling GW et al (1994) Carbon-dioxide supersaturation in the surface waters of lakes. *Science* 265(5178):1568–1570
- Conrad R, Rothfuss F (1991) Methane oxidation in the soil surface-layer of a flooded rice field and the effect of ammonium. *Biol Fert Soils* 12(1):28–32
- De Haan H, van Liere L, Klapwijk SP et al (1993) The structure and function of fen lakes in relation to water table management in The Netherlands. *Hydrobiologia* 265:1–3. doi:[10.1007/BF00007266](https://doi.org/10.1007/BF00007266)
- Frei M, Razzak MA, Hossain MM et al (2006) Methane emissions and related physicochemical soil and water parameters in rice-fish systems in Bangladesh. *Agric Ecosyst Environ* 120:391–398. doi:[10/1016/j.agee](https://doi.org/10/1016/j.agee)
- Guerin F, Abril G (2007) Significance of pelagic aerobic methane oxidation in the methane and carbon budget of a tropical reservoir. *J Geophys Res Biogeosci* 112(G3). Art. number G03006
- Gulati R, van Donk E (2002) Lakes in the Netherlands their origin eutrophication and restoration: state-of-the-art review. *Hydrobiologia* 478:73–106
- Hendriks DMD, van Huissteden J, Dolman AJ et al (2007) The full greenhouse gas balance of an abandoned peat meadow. *Biogeosci Discuss* 4:277–316
- Huttunen JT, Väisänen T, Heikkinen M et al (2002) Exchange of CO₂ CH₄ and N₂O between the atmosphere and two northern boreal ponds with catchments dominated by peatlands or forests. *Plant Soil* 242:137–146
- Huttunen JT, Väisänen T, Hellsten SK et al (2006) Methane fluxes at the sediment-water interface in some boreal lakes and reservoirs. *Boreal Environ Res* 11:27–34. ISSN:1239-6095
- Inubushi K, Otake S, Furukawa Y et al (2005) Factors influencing methane emission from peat soils: comparison of tropical and temperate wetlands. *Nutr Cycl Agroecosyst* 71:93–99
- Juutinen S, Rantakari M, Kortelaine P et al (2009) Methane dynamics in different boreal lake types. *Biogeosciences* 6:209–223
- Kana TM (1994) Membrane inlet mass spectrometer for rapid high-precision determination of N₂ O₂ and Air in environmental water samples. *Anal Chem* 66:4166
- Kankaala P, Taipale S, Nykanen H et al (2003) Oxidation, efflux, and isotopic fractionation of methane during autumnal turnover in a polyhumic, boreal lake. *J Geophys Res Biogeosci* 112:G2 G02003
- Kosten S, Roland F, Da Motta Marques DML, Van Nes EH, Mazzeo N, Sternberg LDSL, Scheffer M, Cole JJ (2010) Climate-dependent CO₂ emissions from lakes. *Glob Biogeochem Cycles*. doi:[10.1029/2009GB003618](https://doi.org/10.1029/2009GB003618)
- Kroon PS, Hensen A, Zahniser MS et al (2007) Suitability of quantum cascade laser spectrometry for CH₄ and N₂O eddy covariance measurements. *Biogeosciences* 4:715–728
- Kroon PS, Schrier-Uijl AP, Hensen A et al (in press) Annual balances of CH₄ and N₂O from a managed fen meadow using eddy covariance flux measurements. *Eur J Soil Sci*
- Lamers PM, van Roozendaal ME, Roelofs GM (1998) Acidification of freshwater wetlands: combined effects of non-airborne sulfur pollution and desiccation. *Water Air Soil Pollut* 105:95–106
- Le mer J, Roger P (2001) Production oxidation emission and consumption of methane by soils: a review. *Eur J Soil Biol* 37(1):25–50
- Lloyd D, Scott RI (1983) Direct measurement of dissolved gases in microbiological systems using membrane inlet mass spectrometry. *J Microbiol Methods* 1:313–328
- Loeb R, van Daalen E, Lamers LPM et al (2007) How soil characteristics and water quality influence the biogeochemical response to flooding in riverine wetlands. *Biogeochemistry* 85:289–302. doi:[10.1007/s1-533-007-9135-x](https://doi.org/10.1007/s1-533-007-9135-x)
- Minkinen K, Laine J (2006) Vegetation heterogeneity and ditches create spatial variability in methane emissions from peatlands drained for forestry. *Plant Soil* 285:289–304
- Rantakari M, Kortelainen P (2005) Interannual variation and climatic regulation of the CO₂ emission from large boreal lakes. *Glob Change Biol* 11:1368–1380
- Rantakari M, Kortelainen P (2008) Controls of organic and inorganic carbon in randomly selected Boreal lakes in varied catchments. *Biogeochemistry* 91(2–3):151–162
- Repo ME, Huttunen JT, Naumov AV et al (2007) Release of CO₂ and CH₄ from small wetland lakes in western Siberia. *Tellus* 59:788–796. doi:[10.1111/j.1600-0889.2007.00301.x](https://doi.org/10.1111/j.1600-0889.2007.00301.x)
- Roulet NT, Moore TR (1995) The effect of forestry drainage practices on the emission of methane from northern peatlands. *Can J For Res* 25(3):491–499
- Saarnio S, Winiwater W, Leitão J (2009) Methane release from wetlands and watercourses in Europe. *Atmos Environ* 43:1421–1429. doi:[10.1016/j.atmosenv.2008.04.007](https://doi.org/10.1016/j.atmosenv.2008.04.007)
- Scheid D, Stubner S, Conrad R (2003) Effects of nitrate- and sulfate-amendment on the methanogenic populations in rice root incubations. *Fems Microbiol Ecol* 43(3):309–315
- Scholten JCM, Conrad R, Stams AJM (2002) Effect of 2-bromo-ethane sulfonate molybdate and chloroform on acetate consumption by methanogenic and sulfate-reducing populations in freshwater sediment. *Fems Microbiol Ecol* 32(1):35–42
- Schrier-Uijl AP, Kroon PS, Leffelaar PA et al (2009a) Methane emissions in two drained peat agro-ecosystems with high and low agricultural intensity. *Plant Soil*. doi:[10.1007/s11104-009-0180-12009](https://doi.org/10.1007/s11104-009-0180-12009)
- Schrier-Uijl AP, Kroon PS, Hensen A et al (2009b) Comparison of chamber and eddy covariance based CO₂ and CH₄ emission estimates in a heterogeneous grass ecosystem on

- peat. *Agric For Meteorol.* doi:[10.1016/j.agrformet.2009.11.007](https://doi.org/10.1016/j.agrformet.2009.11.007)
- Smolders AJP, Roelofs JGM (1993) Sulphate-mediated iron limitation and eutrophication in aquatic ecosystems. *Aquat Bot* 46:247–253
- Smolders AJP, Lamers LPM, Lucassen ECHET et al (2006) Internal eutrophication: how it works and what to do about it—a review. *Chem Ecol* 22(2):93–111
- St. Louis VL, Duchemin CAE, Rudd JWM et al (2000) Reservoir surfaces as sources of greenhouse gases to the atmosphere: a global estimate. *Bioscience* 50:766–775. doi:[10.1641/0006-3568\(2000\)050\[0766:RSASOG](https://doi.org/10.1641/0006-3568(2000)050[0766:RSASOG)
- Stadmark J, Leonardson L (2005) Emissions of greenhouse gases from ponds constructed for nitrogen removal. *Ecol Eng* 25:542–551. doi:[10.1016/j.ecoleng.2005.07.004](https://doi.org/10.1016/j.ecoleng.2005.07.004)
- TOP10vector Product information (in Dutch). Topografische Dienst Kadaster Emmen the Netherlands. Available at <http://www.tdn.nl/> (verified 6 Dec 2007)
- van Bodegom PM, Scholten JCM (2001) Microbial processes of CH₄ production in a rice paddy soil: model and experimental validation. *Geochemica et Cosmochimica Acta* 65(13):2055–2066
- van Huissteden J, Maximov TC, Dolman AJ (2005) High methane flux from an arctic floodplain (indigirka lowlands eastern siberia). *J Geophys Res Biogeosci* 110. Art number G02018
- Veenendaal EM, Kolle O, Leffelaar PA et al (2007) CO₂ exchange and carbon balance in two grassland sites on eutrophic drained peat soils. *Biogeosciences* 4:1027–1040
- Walter KM, Zimov SA, Chanton JP (2006) Methane bubbling from Siberian thaw lakes as a positive feedback to climate warming. *Nature* 443:71–75. doi:[10.1038/nature05040](https://doi.org/10.1038/nature05040)
- Walter KM, Smith LC, Chapin FS (2007) Methane bubbling from northern lakes: present and future contributions to the global methane budget. *Phil Trans R Soc* 365:1657–1676
- Wang HJ, Lu JW, Wang WD (2006) Methane fluxes from the littoral zone of hyper eutrophic Taihu Lake China. *J Geophys Res* 111:D17109. doi:[10.1029/2005JD006864](https://doi.org/10.1029/2005JD006864)
- Whiting GJ, Chanton JP (2001) Greenhouse carbon balance of wetlands: methane emissions versus carbon sequestration. *Tellus* 53B:521–528. ISSN 0280-6509 2001