

This is a "Post-Print" accepted manuscript, which has been published in
"Foodborne Pathogens and Disease".

Please cite this publication as follows:

Islam, M.A., Mondol, A.S., Azmi, I.J., De Boer, E., Beumer, R.R., Zwietering, M.H., Heuvelink, A.E., Talukder, K.A. (2010) Occurrence and characterization of Shiga toxin-producing *Escherichia coli* in raw meat, raw milk and street vended juices in Bangladesh. *Foodborne Pathogens and Disease* 7 (11), 1381-1385.

You can download the published version at:

<http://dx.doi.org/10.1089/fpd.2010.0569>

1 **Occurrence and characterization of Shiga toxin-producing *Escherichia coli***
2 **in raw meat, raw milk and street vended juices in Bangladesh**

3
4 Mohammad A. Islam^{1,2,3}, Abdus S. Mondol¹, Ishrat J. Azmi¹, Enne de boer², Rijkelt R.
5 Beumer³, Marcel H. Zwietering³, Annet E. Heuvelink² and Kaiser A. Talukder¹

6
7 ¹ *Enteric and Food Microbiology Laboratory, International Centre for Diarrhoeal Disease*
8 *Research, Bangladesh, GPO Box-128, Dhaka-1000, Bangladesh.*

9 ² *Food and Consumer Product Safety Authority, P.O. Box 202, 7200 AE Zutphen, The*
10 *Netherlands.*

11 ³ *Laboratory of Food Microbiology, Wageningen University, 6700 EV Wageningen*
12 *The Netherlands.*

13
14 **Running title: Shiga toxin-producing *Escherichia coli* in Bangladesh**

15
16 Contact information for the corresponding author:

17 *Dr. Mohammad A. Islam*

18 *Enteric and Food Microbiology Laboratory*

19 *ICDDR,B*

20 *GPO Box-128, Dhaka-1000, Bangladesh*

21 *Tel.: +88 2 8860523-32*

22 *Fax: +88 2 8812529 or 88 2 8823116.*

23 *E-mail address: maislam@icddr.org*

24

25

1 **Abstract**

2

3 The major objective of this study was to investigate the prevalence of Shiga toxin (Stx)-
4 producing *Escherichia coli* (STEC) in different types of food samples and to compare their
5 genetic relatedness with STEC strains isolated previously from animal sources in Bangladesh.
6 We investigated a total of 213 food samples, including 90 raw meat samples collected from
7 retail butcher shops, 20 raw milk samples from domestic cattle, and 103 fresh juice samples
8 from street vendors in Dhaka city. We found that more than 68% ($n = 62$) of the raw meat
9 samples were positive for the *stx* gene(s); 34% ($n = 21$) of buffalo meats and 66% ($n = 41$) of
10 beef. Approximately 10% ($n = 2$) of the raw milk and 8% ($n = 8$) of the fresh juice samples
11 were positive for *stx*. We isolated STEC O157 from seven meat samples (7.8%) of which two
12 were from buffalo meats and five from beef, and no other STEC serotypes could be isolated.
13 We could not isolate STEC from any of the *stx*-positive raw milk and juice samples. The
14 STEC O157 isolates from raw meats were positive for the *stx*₂, *eae*, *katP*, *etpD* and
15 enterohemorrhagic *E. coli hly* (*hly*_{EHEC}) virulence genes, and belonged to three different
16 phage types: 8 (14.3%), 31 (42.8%), and 32 (42.8%). Pulsed-field gel electrophoresis (PFGE)
17 typing revealed six distinct patterns among seven isolates of STEC O157 suggesting a
18 heterogeneous clonal diversity. Of the six PFGE patterns, one was identical and other two
19 were $\geq 90\%$ related to PFGE patterns of STEC O157 strains isolated previously from animal
20 feces indicating that raw meats are readily contaminated with fecal materials. This study
21 represents the first survey of STEC in the food chain in Bangladesh.

22

23 *Key words:* Shiga toxin-producing *Escherichia coli*, raw meat, raw milk, street vended juice

24

25

1 **Introduction**

2

3 Shiga toxin-producing *Escherichia coli* (STEC) organisms, also called verocytotoxin
4 (VT)-producing *E. coli* (VTEC), are one of the most important groups of foodborne
5 pathogens (Karmali, 1989; Paton and Paton, 1998). They are a major cause of gastroenteritis
6 that may be complicated by hemorrhagic colitis (HC) or the hemolytic-uremic syndrome
7 (HUS), which is the main cause of acute renal failure in children. STEC are a serologically
8 diverse group of zoonotic pathogens. Serotype O157:H7 has been reported as the
9 predominant type worldwide (Tarr et al., 2005). Since its identification as a pathogen in 1982,
10 STEC O157:H7 has been the cause of a series of outbreaks in Canada, Japan, the United
11 Kingdom, the United States and other countries (Karmali, 1989; Paton and Paton, 1998;
12 Willshaw et al., 2001). However, more than 200 STEC serotypes have been reported and
13 more than 100 have been linked with human infection (Eklund et al., 2001).

14 The pathogenicity of STEC strains is associated with various virulence factors. The main
15 factor is the capacity to form two potent phage-encoded cytotoxins called Shiga toxins (Stx1,
16 Stx2 and their variants) or verocytotoxins (VT1 and VT2) (Paton and Paton, 1998). Another
17 virulence factor associated with clinical STEC isolates is the protein intimin, encoded by the
18 *eae* gene and responsible for the intimate attachment of STEC strains to the intestinal
19 epithelial cells and causing the formation of attaching and effacing (A/E) lesions in the
20 intestinal mucosa (Kaper et al., 1998). Enterohemolysin, also called enterohemorrhagic *E.*
21 *coli* (EHEC) hemolysin has also been identified as a possible STEC virulence factor (Beutin
22 et al., 1989).

23 STEC represent the only pathogenic group of *E. coli* that has a definite zoonotic origin,
24 although not all the STEC strains have been demonstrated to cause disease in humans. STEC
25 strains rarely cause disease in animals, and ruminants are recognized as their main natural

1 reservoir. Cattle are considered to be the major animal source of STEC that are virulent to
2 humans, in particular STEC O157 (Caprioli et al., 2005). STEC O157 and other serotypes
3 associated with human infections have also frequently been isolated from the intestinal
4 content of other ruminant species, including sheep, goat, water buffalo, and wild ruminants.
5 Contamination of carcasses with STEC usually occurs during slaughter and subsequent
6 processing through fecal material originating directly or indirectly from the rectal-cecal area
7 (Erickson and Doyle, 2007). Dairy products (milk, cheese, cream) associated with infection
8 have included those that are unpasteurized, have had a pasteurization failure or have been
9 contaminated post-pasteurization. Ready-to-eat foods have also been associated with
10 infection, particularly cooked meats contaminated by raw materials during processing, in
11 catering establishments, at retail sale, and in the home. In developed countries, STEC O157
12 outbreaks associated with beef have caused concern among public health workers, clinicians,
13 and the public, prompting major changes in clinical and laboratory practice, meat production,
14 and food preparation (Mahon et al., 1997).

15 Like many of the developing and underdeveloped countries, the hygienic conditions in
16 Bangladesh are severely compromised, and living with domestic animals within the same
17 premises is a common practice among Bangladeshi population. In our previous study we
18 found that the prevalence of STEC in slaughter animals was 24% and 43% of these STEC
19 isolates belonged to serogroup O157 (Islam et al., 2008). Although this high prevalence of
20 STEC in animal reservoirs has little impact on public health considering the lack of STEC
21 associated infections among Bangladeshi population (Islam et al., 2007), it is important to
22 know the occurrence and transmission of these organisms in the human food chain.

23 Since the incidence of STEC in Bangladeshi foods has remained unknown, the study being
24 reported here examined the occurrence of STEC in different types of food sample, with the
25 isolates further characterized according to serotype, phage type, virulence gene profile, and

1 pulsed-field gel electrophoresis (PFGE) type. In addition, we compared the PFGE types of
2 the food isolates with that of a collection of STEC strains isolated at adjacent period from
3 animal feces in Bangladesh to find out clonal relatedness.

4 5 **Materials and methods**

6 *Sampling*

7 We collected randomly 90 raw meat samples (60 beef and 30 buffalo) from July 2006 to
8 January 2007 from 15 retail butcher shops located in five different areas (3 from each area)
9 in Dhaka city. From each butcher shop, we collected 6 meat samples at two different time
10 points (3 at one time point) during the sampling period. In the same period, we randomly
11 collected raw milk samples from 20 dairy cows domesticated at 20 different households
12 located in two suburbs (10 from each area) of Dhaka city. In addition, we collected 103
13 freshly prepared juice samples, including mixed herbal juice ($n = 49$), sugar cane juice ($n =$
14 40), wood apple juice ($n = 7$) and some other types of juice (three milk shake, one lemon, two
15 mix fruit, and one pineapple) from street vendors who operate in the main streets of 10
16 different locations in the Dhaka city. Only one juice sample was collected at one time point
17 from each vendor. Each sample (minimum of 25 g or 25 ml) was aseptically transferred to a
18 sterile plastic bag, placed in an ice box, and transported to the laboratory within 2 h. We
19 started microbiological investigation of the samples within 4 h of collection.

20 21 *Isolation of STEC O157*

22 For meat samples, a 25-g portion, and for raw milk and juice samples, a 25-ml portion
23 was added to 225 ml of modified tryptone soy broth (Oxoid Ltd., Basingstoke, United
24 Kingdom). After homogenization in a stomacher for 1 min, the samples were incubated for
25 18-20 h at 37°C. STEC O157 strains were isolated as described previously (Islam et al.,

1 2006), by using the immunomagnetic separation (IMS) technique and presumptive isolates
2 were confirmed by an agglutination test using *E. coli* O157 latex agglutination kit (Murex
3 Biotech Ltd., Dartford, Kent, United Kingdom) and a PCR for the *rfb* (O-antigen-encoding)
4 region of *E. coli* O157 (*rfbE*_{O157}) (Paton and Paton, 1998).

5

6 *Isolation procedure for STEC non-O157*

7 The overnight enrichment cultures were screened by PCR for the presence of *stx*₁ and *stx*₂
8 using the primers as described earlier (Islam et al., 2007). DNA was extracted from 1 ml of
9 broth culture by thermal cell lysis using Chelex-100 resin (Bio-Rad Laboratories Ltd.,
10 Richmond, Calif.) and 5 µl of the DNA extract was used in the PCR (Malorny et al., 2003).
11 Multiplex PCR for *stx*₁ and *stx*₂ was carried out in a PTC-200 peltier thermal cycler (Bio-
12 Rad) using the program as described previously (Islam et al., 2007). DNA extracted from *E.*
13 *coli* O157:H7 strain NCTC 12079 and *E. coli* strain ATCC 25922 were used as positive and
14 negative controls, respectively in each PCR reaction. Broth cultures that yielded positive PCR
15 results for *stx*₁ and/or *stx*₂ were streaked onto tryptone bile x-glucuronide (TBX) agar (Oxoid)
16 supplemented with cefixime (20 mg/l), cefsulodin (3 mg/l) and vancomycin (30 mg/l) (CCV-
17 TBX). Single colonies, including different morphological types were transferred from the
18 CCV-TBX and plated onto Luria agar (Difco, BD diagnostics, USA) to create a grid pattern
19 of 96 colonies (12×8). The plates were incubated overnight at 37°C. The isolates were
20 subjected to PCR for the detection of *stx*₁ and *stx*₂ as described above. The number of PCR
21 reactions was reduced to 20 by pooling the colonies per row and per column in 500 µl sterile
22 distilled water. DNA was extracted by boiling the suspensions for 10 min and 5 µl of the
23 supernatant was used in the PCR. The isolates with positive PCR results were identified as *E.*
24 *coli* by an API 20E test (bioMérieux, Marcy l'Etoile, France).

25

1 *PCR detection of virulence factors*

2 PCR for detecting both chromosome (e.g., *stx*₁, *stx*₂, and *eae*)- and plasmid (e.g., *hly*_{EHEC},
3 *katP*, and *etpD*)-encoded virulence genes was performed as described earlier (Islam et al.,
4 2007) in a total volume of 25 µl containing 2.5 µl of 10× Fast Start buffer with 1.5 mM
5 MgCl₂ (Applied Biosystems, Weiterstadt, Germany), 200 µM concentration of each
6 deoxynucleoside triphosphate, a 0.2 pmol of each primer, and 1 U of Fast start *Taq* DNA
7 polymerase (Roche, Mannheim, Germany) using a PTC-200 peltier thermal cycler (Bio-Rad).

8

9 *Phage typing*

10 Isolates confirmed to be STEC O157 were phage typed at the laboratory of Enteric
11 Pathogens, Central Public Health Laboratory, London, United Kingdom.

12

13 *PFGE*

14 PFGE was performed following the standardized protocol developed by PulseNet for *E.*
15 *coli* O157:H7 (Ribot et al., 2006). Analysis of the TIFF images was carried out by the
16 BioNumerics software (Applied Maths, Belgium) using the dice coefficient and unweighted-
17 pair group method using average linkages to generate dendrograms with 1.0% tolerance
18 values. Normalized PFGE patterns (bundle files) of the food isolates were compared with that
19 of the animal isolates from our previous studies (Islam et al., 2008) in order to find out the
20 clonal relationships.

21

22 **Results and discussion**

23 *Occurrence of STEC in food samples*

24 Of the 90 meat samples tested in this study, we found that 62 (69%) were positive for *stx*
25 gene(s); 21 samples of buffalo meats, and 41 samples of beef (Table 1). STEC O157 strains

1 were isolated from seven samples (7.8%); two samples of buffalo meats, and five samples of
2 beef (Table 1). No STEC non-O157 strains were isolated. During the processing of carcasses,
3 meat can easily become contaminated with animal feces if there are unhygienic conditions,
4 lack of good processing practices and a lack of knowledge of the butchers on basic hygiene.
5 These features are commonly observed in butcher shops in Bangladesh where the same
6 premise is being used to slaughter the animal, to process carcasses and as vending point. Our
7 result correlates with data reported from neighboring countries, including India where STEC
8 O157 have been identified as one of the predominant serogroups from buffalo meats
9 (Hazarika et al., 2007). STEC O157 strains isolated from raw minced beef samples (9%; $n =$
10 22) (Dutta et al., 2000), beef surface swabs (3.7%; $n = 27$) and milk samples (2.4%; $n = 81$)
11 (Manna et al., 2006) in India. Of the 20 raw milk samples, two (10%) were positive for *stx*
12 gene(s) (Table 1). Of the 103 fresh juice samples, eight (8%) were positive *stx* gene(s); six
13 samples of mixed herbal juices, and two samples of wood apple juices (Table 1). However,
14 no STEC could be isolated from *stx*-positive samples of raw milk and street vended juices.
15 This might be due to the presence of a very small number of organisms present in the sample
16 that were insensitive to the culture method used in the study. Another reason might be the
17 presence of free *stx* phages in the sample that turned out to be positive by PCR for *stx* genes.

18

19 *Characterization of STEC O157 isolates*

20 STEC O157 strains that were isolated from raw meat samples were further characterized.
21 All isolates were positive for the *stx*₂ gene only. There is considerable epidemiological
22 evidence to indicate that STEC O157 isolates producing Stx₂ are more commonly associated
23 with serious disease than isolates producing Stx₁ or Stx₁ and Stx₂ (Law, 2000). In addition to
24 *stx*₂ gene, all isolates in the study were positive for *eae*, *etpD*, *katP* and *hly*_{EHEC} genes, which
25 accentuate their capability to cause human infection.

1 The strains could be grouped into three different phage types (PT): PT 31 ($n = 4$), PT 32
2 ($n = 2$), and PT 8 ($n = 1$). Phage types 31 and 32 have been commonly found among STEC
3 O157 isolates obtained from both human and non-human sources (food and animal feces)
4 from different countries suggesting a foodborne transmission of the organism (Khakria et al.,
5 1990, Mannix et al., 2007). In our previous study we found that these two phage types (PT 31
6 and PT 32) were the predominant PTs among STEC O157 strains isolated from animal feces
7 (Islam et al., 2008). Phage type 8 has been found frequently among STEC O157 strains from
8 humans and bovine in many European countries, including Spain, Belgium, Finland,
9 Germany, Italy, England and Scotland (Khakria et al., 1990). We also found this phage type
10 in our previous study in a STEC O157 strain isolated from goat feces. Therefore, it is more
11 likely that raw meat products are often contaminated with animal feces in the butcher shop
12 environment.

13 PFGE analysis of the seven STEC O157 isolates revealed six distinct patterns, suggesting
14 a genetically heterogeneous group. We compared these patterns with a panel of PFGE pattern
15 that comprises 44 STEC O157 strains isolated previously from feces of cow, buffalo and
16 goats from slaughterhouses in Dhaka city (Fig. 1). One strain (CM52) isolated from beef had
17 an identical PFGE pattern with that of a strain isolated from buffalo feces (M139).
18 Interestingly, both the beef and fecal strains were isolated from the same slaughterhouse (data
19 not shown). Another beef isolate (CM2) had a PFGE pattern that was 92% related to that of a
20 cluster of strains isolated from cow (C8) and goat feces (G72 and G85). Two strains, one
21 from beef (CM56) and one from buffalo meat (MM28) belonged to the same clonal type (Fig.
22 1) and this PFGE pattern was 90% related to that of a strain isolated previously from cow
23 feces (C21). Among the six PFGE patterns obtained from meat isolates, 3 (50%) were
24 identical/closely ($\geq 90\%$) related to that of the isolates from animal feces. However, this result
25 was not concordant with the phage typing results. Overall results of the study indicate that

1 meats are readily contaminated with fecal materials containing STEC O157 in the butcher
2 shops due to unhygienic practices of carcass processing.

3

4 **Conclusion**

5 To our knowledge, this is the first survey to establish the prevalence of Shiga toxin-
6 producing *E. coli* in various food items in Bangladesh. Despite frequent contamination of raw
7 meat products by potentially virulent *E. coli* O157, to date, no clinical data about impact on
8 human health are available in Bangladesh. Improved surveillance and hygienic conditions
9 along with the development of control measures are therefore highly recommended to
10 minimize contamination of raw meat products by potentially virulent *E. coli* O157.

11

12 **Acknowledgments**

13 This research was funded by the International Centre for Diarrhoeal Disease Research,
14 Bangladesh (ICDDR,B) and by the Netherlands Foundation for the Advancement of Tropical
15 Research (NWO-WOTRO) grant (Award number WB 93-415). ICDDR,B acknowledges
16 with gratitude the commitment of the Netherlands Foundation for the Advancement of
17 Tropical Research (NWO-WOTRO) grant to the Centre's research efforts. ICDDR,B also
18 gratefully acknowledges these donors who provide unrestricted support to the Centre's
19 research efforts: Australian International Development Agency (AusAID), Government of
20 Bangladesh, Canadian International Development Agency (CIDA), The Kingdom of Saudi
21 Arabia (KSA), Government of the Netherlands, Government of Sri Lanka, Swedish
22 International Development Cooperative Agency (SIDA), Swiss Development Cooperation
23 (SDC) and Department for International Development, UK (DFID).

24

25

1 **References**

2

3 Beutin L, Montenegro MA, Orskov I, *et al.* Close association of verotoxin (Shiga-like toxin)
4 production with enterohemolysin production in strains of *Escherichia coli*. J Clin
5 Microbiol 1989;**27**:2559–2564.

6 Caprioli A, Morabito S, Brugère H, *et al.* Enterohaemorrhagic *Escherichia coli*: emerging
7 issues on virulence and modes of transmission. Vet Res 2005;**36**:289-311.

8 Dutta S, Deb A, Chattopadhyay UK, *et al.* Isolation of Shiga toxin-producing *Escherichia*
9 *coli* including O157:H7 strains from dairy cattle and beef samples marketed in
10 Calcutta, India. J Med Microbiol 2000;**49**:765-767.

11 Eklund M, Scheutz F, and Siitonen A. Clinical isolates of non-O157 Shiga toxin-producing
12 *Escherichia coli*: serotypes, virulence characteristics, and molecular profiles of strains
13 of the same serotype. J Clin Microbiol 2001;**39**:2829-2834.

14 Erickson MC and Doyle MP. Food as a vehicle for transmission of Shiga toxin-producing
15 *Escherichia coli*. J Food Prot 2007;**70**:2426-2449.

16 Hazarika RA, Singh DK, Kapoor KN, *et al.* Verotoxic *Escherichia coli* (STEC) from beef
17 and its products. Indian J Exp Biol 2007;**45**:207-211.

18 Islam MA, Heuvelink AE, Talukder KA, *et al.* Evaluation of immunomagnetic separation and
19 PCR for the detection of *Escherichia coli* O157 in animal feces and meats. J. Food
20 Prot. 2006;**69**:2865-2869.

21 Islam MA, Heuvelink AE, de Boer E, *et al.* Shiga toxin-producing *Escherichia coli* isolated
22 from patients with diarrhoea in Bangladesh. J Med Microbiol 2007;**56**:380-385.

23 Islam MA, Mondol AS, de Boer E, *et al.* Prevalence and genetic characterization of Shiga
24 toxin-producing *Escherichia coli* isolated from slaughtered animals in Bangladesh.
25 Appl Environ Microbiol 2008;**74**:5414-5421.

1 Kaper JB, Elliott S, Sperandio V, *et al.* Attaching and effacing intestinal histopathology and
2 the locus of enterocyte effacement. In: *Escherichia coli* O157:H7 and other Shiga
3 toxin-producing *E. coli* strains. Kaper JB, and O'Brien AD (eds.). Washington DC:
4 ASM press, 1998, pp. 163-182.

5 Karmali MA. Infection by verocytotoxin-producing *Escherichia coli*. Clin Microbiol Rev
6 1989;**2**:5-38.

7 Khakria R, Duck D and Lior H. Extended phage-typing scheme for *Escherichia coli*
8 O157:H7. Epidemiol Infect 1990;**105**:511-520.

9 Law D. Virulence factors of *Escherichia coli* O157 and other Shiga toxin-producing *E. coli*. J
10 Appl Microbiol. 2000;**88**:729-45.

11 Mahon BE, Griffin PM, Mead PS, *et al.* Hemolytic uremic syndrome surveillance to monitor
12 trends in infection with *Escherichia coli* O157:H7 and other shiga toxin-producing *E.*
13 *coli*. Emerg Infect Dis 1997;**3**:409-412.

14 Malorny B, Hoorfar J, Hugas M, *et al.* 2003. Interlaboratory diagnostic accuracy of
15 *Salmonella* specific PCR-based method. Int J Food Microbiol 2003;**89**:241-249.

16 Manna SK, Brahmane MP, Manna C, *et al.* Occurrence, virulence characteristics and
17 antimicrobial resistance of *Escherichia coli* O157 in slaughtered cattle and diarrhoeic
18 calves in West Bengal, India. Lett Appl Microbiol 2006;**43**:405–409.

19 Mannix M, Whyte D, McNamara E, *et al.* Large outbreak of *E. coli* O157 in 2005, Ireland.
20 Eur Surveill 2007;**12**:article 3.

21 Paton JC and Paton AW. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia*
22 *coli* infections. Clin Microbiol Rev 1998;**11**:450-479.

23 Ribot EM, Fair MA, Gautom R, *et al.* Standardization of pulsed-field gel electrophoresis
24 protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for
25 PulseNet. Foodborne Pathog Dis 2006;**3**:59-67.

- 1 Tarr PI, Gordon CA and Chandler WL. Shiga-toxin-producing *Escherichia coli* and
2 haemolytic uraemic syndrome. Lancet 2005;**365**:1073-1086.
- 3 Willshaw GA, Cheasty T, Smith HR, *et al.* Verocytotoxin-producing *Escherichia coli*
4 (VTEC) O157 and other VTEC from human infections in England and Wales: 1995-
5 1998. J Med Microbiol 2001;**50**:135-142.
- 6

1 TABLE 1. PCR DETECTION OF *STX* GENE(S) IN BROTH CULTURES FROM FOOD
 2 SAMPLES AND ISOLATION OF SHIGA TOXIN-PRODUCING *E. COLI* O157 AND
 3 NON-O157 BY SUBCULTURING ONTO SELECTIVE AGAR

<i>Sources (n = 213)</i>	<i>No. of PCR positive (%)</i>				<i>No. of Isolates (%)</i>		
	<i>Only stx₁</i>	<i>Only stx₂</i>	<i>stx₁₊₂</i>	<i>Total</i>	<i>E. coli O157^a</i>	<i>Non-O157</i>	<i>Total</i>
Meat (90)	1 (1.1)	9 (10.0)	52 (57.8)	62 (68.9)	7 (7.8)	0	7 (7.8)
Buffalo (30)	0	2 (6.7)	19 (63.3)	21 (70)	2 (6.7)	0	2 (6.7)
Beef (60)	1 (1.7)	7 (11.7)	33 (55)	41 (68.3)	5 (8.3)	0	5 (8.3)
Raw milk (20)	1 (5)	1 (5)	0	2 (10)	0	0	0
Fresh juices (103)	2 (1.9)	4 (3.9)	2 (1.9)	8 (7.8)	0	0	0
Mixed herbal (49)	2 (4)	2 (4)	2 (4)	6 (12)	0	0	0
Sugarcane (40)	0	0	0	0	0	0	0
Wood apple (7)	0	2 (28.6)	0 (0)	2 (28.6)	0	0	0
Others (7)	0	0	0	0	0	0	0

4 ^aBy using immunomagnetic separation.

5

6

1 **Legend to figure**

2 **FIG. 1.** Dendrogram generated by Bionumeric software, showing distance calculated by the
3 dice similarity index of PFGE *XbaI* profiles for STEC O157 isolates isolated from raw meats
4 ($n = 7$) and animal feces ($n = 44$). Box with dotted line indicated profiles of raw meat
5 isolates. The degree of similarity (%) is shown on the scale.

6

7

1 **FIG.1.**

