

Providing Seasonal-to-Interannual Climate Information for Risk Management and Decision-making

L. Goddard^{a,*}, Y. Aitchellouche^b, W. Baethgen^a, M. Dettinger^c, R. Graham^d, P. Hayman^c,
M. Kadi^f, R. Martínez^g and H. Meinke^h with additional contributions by E. Conrad^a

^aInternational Research Institute for Climate and Society, Columbia, United States

^bInternational Federation of Red Cross and Red Crescent Societies, Dakar, Senegal

^cUnited States Geological Survey, Scripps Institution of Oceanography, La Jolla, United States

^dMet Office, Exeter, United Kingdom

^ePrimary Industries and Resources of South Australia/South Australia Research and Development Institute, Australia

^fAfrican Centre of Meteorological Application for Development (ACMAD), Niamey, Niger

^gInternational Research Center on El Niño (CIIFEN), Guayaquil, Ecuador

^hWageningen University and Research Centre, Wageningen, The Netherlands

Abstract

Much has been learned in the interpretation and use of climate information since the 1997/1998 El Niño event that garnered so much attention. Seasonal-to-interannual forecasts are now produced around the world. However, mismatches in their scales, specificity or communication (of forecast content and uncertainties) with decision-maker needs still hinder their use. More work is needed to improve a) the utility of models, b) access to observational and model/forecast data, c) understanding and communication of the opportunities and limitations of forecasts, and d) methods by which decision systems use climate predictions – both through modifications of decision systems and more tailored forecast information. This white paper discusses these issues and recent advances in providing climate information needed in effective climate risk management. The paper advocates for the establishment and/or strengthening of “chains of experts and communications” to better enable appropriate and effective dissemination, assimilation and further use and application of climate information.

Keywords: Information and education; climate services for health; health monitoring and surveillance

1. Introduction

The history of physically based seasonal climate forecasts is relatively short and strongly linked to the ability to predict sea-surface temperatures (SST) in the El Niño region. The first physically based model forecast of equatorial Pacific Ocean temperatures was produced only in the mid-1980s [1]. The growing ability to predict El Niño led to a cascade of efforts for developing and improving the seasonal climate forecasts and attempting to make those useful to society. El Niño is the overall dominant influence in regional climate variability worldwide, though other modes of sea-surface temperature variability can be more important in some regions [2].

In developing a white paper for understanding and predicting seasonal-to-interannual climate variability from a user perspective or from a producer perspective, what is immediately clear is how blurred the boundary between these communities has become over the past decade. A user may be a decision-maker acting individually or as part of a collective. A user may also be a translator of information regarding climate variability or its associated impacts such that the information can be used by decision-makers. Similarly, a producer may be the climate scientist running a dynamical model of the climate on a large computer. A producer may be the sectoral scientist who takes the information from the climate model and feeds it through a hydrology model or crop model. Or, the producer may be that same translator referred to above who modifies the initial forecast information into a more usable format for the policymaker or decision-maker. In some places the term “producer” may even mean “producer of agricultural products” (a farmer), highlighting the confusion this terminology can cause.

The blurring of boundaries between these communities began when it became clear that effective climate risk management could not be accomplished in isolated communities. It has been an important realization, but much work remains. This white paper is written from the perspective of users, but not those who have been traditionally targeted as end-users. The user’s voice here is from the translators – those who build on the scientific advances in climate modelling and diagnostics, those who design decision systems for resource management and those who are participating in the conversation regarding climate risk management and the prospects of seasonal prediction. The following discussion concerns the advances that are necessary in providing climate information for effective climate risk management at seasonal-to-interannual timescales. These efforts constitute the wealth of research, communication and application that has been attempting to bridge the gap between the traditional provider and user communities.

The paper begins with the historical evolution of developing and using climate information for management and decisions. The discussion then progresses through the current infrastructure of accessing climate forecast information, the gaps between the

*Corresponding author. Tel: +1-845-6804430
E-mail address: goddard@iri.columbia.edu

operational provision of forecasts and their use, considerations for increasing the value of forecast information, some of the lessons learned and concludes with our view of the way forward. Climate information refers in this paper to both current and historical observations-based data and to predictions of future climate conditions, with particular focus on seasonal-to-interannual variations. Most of the attention here will be upon the user's perspective on forecasts, but those forecasts become valuable primarily in context of past climate variations and, where known, the past performances of the forecasts. Also, although this paper focuses on seasonal-to-interannual climate variability, the context set by the background climate, which may be slowly varying, is well recognized. The successful future of climate risk and resource management depends critically on the reliability of the entire chain of information that the international community is working so hard to forge.

2. The historical situation

Until the last decade, plans and decisions that needed climate information often followed an approach based on the long-term means of relevant climate variables. For example, maize in a given region was sown at a certain date because the combination of rainfall and temperatures in the following 4–6 months for that region was, on average, the most favorable for the crop growth and development. Plans for distributing water in multi-purpose reservoirs (hydroelectricity, irrigation and human consumption) were established with lead times of several months, based on the mean values of the precipitation for the entire year (and in some cases also based on the current situation, for example, snowpack). Health institutions based their action plans for infectious disease outbreaks in a given area in consideration of the long-term average temperatures and rainfall of that area.

This approach to management is not a general truth, however. It is not uncommon for farmers to use environmental observables to guide their actions. Soil moisture availability, for example, might suggest when to plant what. In particular, resource-poor farmers in semi-arid regions are excellent, intuitive risk managers. Of course they are also conservative: they do not plan for the average season, but for the poor season (low plant densities, no or low inputs), so they ensure their survival, but never manage to make a real profit because they miss out in the good years.

Interestingly, the probability that an entire year will behave as an average year (12 months of average rainfall) is virtually zero. Moreover, by definition, the probability that the rainfall of two subsequent trimesters will fall in the central (normal) tercile, is less than 10 per cent. Still, and up to the 1990s, planning and decisions in many climate-dependent activities could only be based on these very unlikely "average" or "normal" years.

Where mean conditions have not been used as the default climate forecast, resource and risk managers have relied, and often continue to rely, heavily on observed conditions at the time that decisions need to be made for the basis of their forecasts of future eventualities. For example, it has been (and remains) common for managers to base decisions and forecasts of future water supplies solely on observed snowpack and soil moisture for all but the shortest-term, multiday problems [3]. Observed conditions on the ground have been more reliable and more immediately relevant to the decisions to be made than other climate science resources for many applications.

Efforts to make these improvements have been prompted, in part, by a growing understanding of the effects of global scale climate phenomena like the El Niño–Southern Oscillation (ENSO) on the climates, resources and hazards of many regions, and a growing expectation that improvements in climate forecasts may provide a basis for decision-making that is not currently being exploited. Generally speaking, climate forecasts are forecasts of those climate system variations that reflect predictable responses to predictable changes in slowly varying boundary conditions like sea-surface temperatures and radiative imbalances in the Earth's energy budget. Not all possible sources of climate forecast skill have been identified or exploited, but boundary condition contributors may include a variety of large-scale air–sea connections [4][5][6][7], snow and sea-ice patterns [8][9] and soil moisture and vegetation [10]. Long-term radiative imbalances associated with human-caused emissions of greenhouse gases into the atmosphere have been a focus of much attention on even longer timescales [11].

Within the past decade, however, climate scientists have begun to identify potential improvements in long-lead, seasons-to-years-ahead climate forecasts [12][13][14] and to link them with resource models [15][16] or statistical distributions of parameters relevant to management [17][18] to improve the immediacy and, in some cases, the reliability of the climate forecasts for use in management decisions. Consequently, research institutes such as the International Research Institute for Climate and Society (IRI), invested huge efforts to provide synthesized climate forecasts based on the inputs from the international modelling community, and started supplying them to climate information providers (National Weather Services, Regional Climate Centres and Specialized Meteorological Centres) and to key socio-economic sectors (agriculture, health, water resources and disaster prevention and reduction). The premise of these efforts was that supplying the best possible seasonal forecasts would immediately result in better decisions and more effective planning activities in those sectors. Efforts were thus concentrated in investing increased efforts in the dynamical models and statistical methods that resulted in forecasts with better skill.

The initial reaction in the different sectoral communities was extremely optimistic: the new seasonal climate forecasts were viewed as tools that would assist these communities to cope better with the immense challenges posed by climate variability on their activities. Planning and decisions in activities that depend on, or are affected by, climate would now be better informed.

However, this initial optimistic environment was shortly followed by frustration in both the climate science community and the socio-economic sectors since expectations from neither group were fulfilled. Excellent achievements were obtained in the climate science community for supplying seasonal forecasts that were continuously improving. Many studies demonstrated the potential value of incorporating this information into the decision-making of different sectors [19][20][21][22][23][24]. However, there was little or no evidence that the generated information was effectively being embedded in the policies, planning or decision-making within the sectors. On the other hand, the socio-economic sectors started receiving vast amounts of information resulting from the seasonal forecasts but in most cases could not find ways to incorporate it in a useful manner for their routine activities.

3. The current situation

3.1 Forecast producers and their products

Producers of seasonal forecasts may be broadly categorized according to the geographical coverage addressed. Forecasts with global coverage often require the use of dynamical prediction models (an exception would be predictions based on ENSO teleconnections with skill at the global scale (Stone and others, 1996 [25]) and producers are limited to centres with capacity for numerical seasonal climate prediction (typically some National Meteorological or Hydrometeorological Services [NMSs] and research centres). Forecasts with competitive skill may also be produced with relatively inexpensive statistical models for some regions, and such models have been developed by a host of producers from individual researchers to NMSs for their country or region. Finally there are centres and activities that synthesize global and regional forecasts from the various local and international sources, and from both dynamical and statistical methods, into consolidated outlooks for a region, specifically the Regional Climate Centres (RCCs) and Regional Climate Outlook Forums (RCOFs) [26]. This latter consolidation activity also takes place on a national level at some NMSs [27].

3.1.1 Forecasts with global coverage

In recent years there has been substantial progress in coordinating and disseminating the output from centres producing forecasts with global coverage using dynamical prediction methods. The aim has been to increase the accessibility and usability of the information to NMSs, RCCs and RCOFs, and to make available the benefits of multi-model ensemble combinations [28]. To coordinate convergence among the forecast centres, the World Meteorological Organization (WMO) has defined standards in real-time forecast output and hindcast validation. Centres adhering to the criteria may apply for designation as WMO Global Producing Centres of Long-range Forecasts (GPCs). Eleven such GPCs have now been designated. A significant further boost to the coordination of GPC output occurred with WMO designation, in 2009, of a Lead Centre for Long-range Forecast Multi-model Ensembles (LC-LRFMME), jointly hosted by the Korea Meteorological Agency (KMA) and the NOAA National Centers for Environmental Prediction (NCEP). The LC-LRFMME has a range of functions, with two central themes:

- (a) To provide a single portal from which users (RCCs, RCOFs, NMSs and GPCs) can access GPC forecast output in common formats;
- (b) To promote research into, and to generate and provide multi-model products from, the GPC forecasts.

The single portal and uniform presentation introduced by the LC-LRFMME greatly enhance the accessibility and usability of the GPC products, and provide RCCs, RCOFs and NMSs with validated model output from a range of models for use, together with other inputs, in generating a consolidated forecast for their region or country.

In line with WMO criteria, prediction products typically take the form of anomalies in 3-month-mean quantities with a range of 6 months to 1 year ahead. Predicted variables made available include 2m temperature, precipitation, sea-surface temperature, mean sea-level pressure, 850 hPa temperature and 500 hPa temperature. Anomalies are usually expressed both deterministically, using the mean of the prediction ensemble, and in terms of probabilities for categories (typically tercile categories of the model climatology), based on the ensemble distribution. Some centres also produce and make available forecasts of extremes, but in general these products are not well developed. In these cases extremes forecasts are typically presented as probabilities for outer quintile categories, rather than risk of floods, drought, heatwaves or cold spells. Examples of forecast products presently available from producing centres may be viewed at http://www.wmo.int/pages/prog/wcp/wcasp/clips/producers_forecasts.html.

A key question faced in creating or using a consolidated forecast from various prediction inputs is: what are the relative skills of the inputs and of the final product? For this reason GPC accreditation requires provision of hindcast validation according to a WMO defined set of diagnostics and procedures, the Standard Verification System for Long-range Forecasts (SVSLRF). Such diagnostics are typically available on the Websites of the producing centres. In addition, SVSLRF diagnostics from designated GPCs are available in common format on the WMO Lead Centre for the SVSLRF (co-hosted by the Australian Bureau of Meteorology and the Meteorological Service of Canada – <http://www.bom.gov.au/wmo/lrfvs/>). The SVSLRF information provides a basis for making qualitative judgments on appropriate weights to be given to the GPC products. However, further international cooperation is required to make the validation information more accessible and useable. For example, adherence to a common hindcast period to be used by all GPCs for generating validation diagnostics would be ideal. It must be acknowledged, however, that given the current embryonic state of seasonal forecast systems at many centres, such convergence may take some time to achieve.

The importance of the current role of expert judgment in interpreting both the model predictions and the hindcast skill evaluations cannot be overestimated. Models are not perfect and often require correction, or at least interpretation. Although success with ENSO prediction is encouraging, it is well known that models do not currently represent well many other important modes of intra-seasonal-to-interannual variability (for example, the Madden–Julian Oscillation, the North Atlantic Oscillation or the Indian monsoon), and such shortcomings need to be accounted for by forecasters. Hindcast skill assessments also require careful interpretation. Although very useful, most hindcast evaluations represent “average” skill assessed over many years, and as such they provide little information on the credibility of predicted signals in the context of the unique SST forcing in any particular year, for example, when ENSO forcing is strong [13].

The model predictions provided are intended for use by climate professionals (such as NMSs, RCOFs and RCCs) to help in construction of consolidated forecasts for users. Although the model predictions are often freely available on the Internet, at least in terms of visualized products, data from the current and past forecasts are typically neither freely available nor easily accessible. The free availability of maps leads to issues concerning the potential for misinterpretation or confusion in cases when some model predictions conflict with an official consolidated forecast. However, the lack of easily available data reduces the actual use of many of these predictions.

3.1.2 Forecasts with regional coverage

Examples of consolidated regional forecasts are those of the Regional Climate Outlook Forums currently held in Africa and Latin America for some regions with spatial and temporal coherency in rainfall seasonality [26]. Regional Climate Outlook Forums have not played a big role in south and South-East Asia. The RCOFs are not permanent entities, but are convened ahead of the main rainy seasons (for example, in May for west Africa, and in February for the east African short rainy seasons and in August for the long rainy season). Key objectives of the RCOFs are development of a consensus prediction for the region and dissemination of the forecast to application sectors including agriculture, water resources, energy, health and media. The RCOFs also provide unique opportunities to form alliances between forecast users and producers, for users to feed back requirements to the producers and for institutional capacity-building. The consensus forecast provides a broad-scale outlook for the region, which may be further elaborated to national scales (with more targeted dissemination) by the NMSs of the regions. To deal with geographical variations in the forecast signals, the region is divided into zones, and each zone assigned numerical probabilities for tercile categories of rainfall. (See Figure 1.) Analysis of 10 years of real-time consensus seasonal predictions from regional forums convened in Africa shows that, despite some shortcomings, the forecasts have evidence of skill [29].

The procedure used to generate the consensus involves a blending of all available evidence, which may include examination of the principal mode or modes of climate forcing likely to operate over the forecast period and to influence the region (for example, the status of ENSO and/or regionally relevant SST anomalies); the prediction skill of the climate models used; and the current predictions from dynamical and empirical models. Typically, statistical predictions for individual countries – based on historical links with global SST patterns – form a key component in the consensus. The forecasts are blended where discrepancies occur at national boundaries and may be modified in light of interpretation from regional experts. As is often found with consensus methods, the resulting blended forecasts tend to overestimate probabilities for the average category reducing the usefulness for applications. (See Chidzambwa and Mason [29].)

Input from predictions from the dynamical modelling centres also plays an important role, and standardization of this input, through the WMO GPC criteria, has assisted in its use. Standardized GPC products from the LC-LRFMME were first introduced to the RCOF process at the twenty-third Greater Horn of Africa Climate Outlook Forum (GHACOF 23, Mombasa, Kenya, 2–4 March 2009), coordinated by the Intergovernmental Authority on Development Climate Prediction and Applications Centre (ICPAC), and are being introduced as key prediction tools available to all RCOFs.




Figure 1. PRESAO-11 RCOF consensus forecast issued 21 May 2008

Vertically stacked numbers show predicted probabilities (top to bottom) of above-normal, near-normal and below-normal rainfall over west Africa (latitude and longitude shown on axes), relative to the 1971–2000 climatology period. Each zone indicates the area over which the overlaid 3-category probabilities apply.

A fully satisfactory objective procedure for optimal blending of all the available information has yet to be agreed upon within the RCOFs – and this was one of the development needs identified at the RCOF Review 2008 (Arusha, United Republic of Tanzania, November 2008; WMO report and position papers, in preparation). Currently, dynamical model predictions are used to inform qualitative adjustments to the consensus forecast. However, objective use of the dynamical forecasts has been initiated at the West Africa Forum as part of an initiative known as PRESAO-Second Generation. The PRESAO initiative is coordinated by the African Centre of Meteorological Application for Development (ACMAD). Using model hindcasts and observational records for each country, dynamical model predictions are calibrated for each country in the region. (See section 5.1.2 on toolboxes.) The calibration process uses similar statistical tools used in developing the statistical prediction models (the IRI Climate Prediction Tool [CPT] is one of the software packages used), and skill scores for statistical methods and different GPC models can be generated for each country. This allows the RCOF users to gain an appreciation of the relative strengths and weaknesses of statistical and dynamical predictions.

The resulting national forecasts should contain a degree of common modulation consistent with large-scale signals, and thus should require less ad hoc blending when forming the regional consensus. Similarly, in western South America, the International Research Centre on El Niño (CIIFEN) has organized training activities to improve the capabilities of the Meteorological Services to use statistical and dynamic methods to produce downscaled seasonal forecasts at both regional and national levels. With less need for manual blending, the regional forecasts require less deliberation at the RCOF meeting. For example, using more objective calibration and combination techniques, CIIFEN has established institutional agreements with the Meteorological Services of six countries of the region to produce a unified seasonal climate forecast that is updated every month.

3.2 Forecast applications

The types of seasonal forecast applications products thus far developed are quite varied and depend, among other things, on the available prediction skill and the target audience. A full review of the status of forecast applications is beyond the scope of this paper. The following sections illustrate some example applications typical of current capabilities.

3.2.1 General applications: advisory statements

An important and widespread application is the provision of advice on seasonal prospects to government, non-governmental organizations (NGOs), the public and the media. Such advice often takes the form of written statements (Figure 2). Because of current imperfections in predictions, such applications are ideally generated using consolidated forecast products developed through expert consensus. However, given the wide availability of seasonal forecasts on the Internet, it is likely that statements may in some cases be constructed through relatively inexpert interpretation of limited inputs (for example, the interpretation of output from a single model, which may have limited skill). This situation needs to be addressed through continued education and through better channelling of forecast information through a global climate service structure (for example, GPCs to RCCs/RCOFs to National Meteorological and Hydrological Services [NMHSS] to users).

- | | |
|----|--|
| 1. | The probability of rainfall deficit is very low in the subregion. The probability of rainfall less than normal is equal to 0.20–0.15 and 0.20 in Zones I, II and III, respectively |
| 2. | A high probability of higher-than-normal rainfall in Zone I and II, and near-normal rainfall in Zone III (probability of 0.45–0.50 and 0.50, respectively) |
| 3. | In this regard, strengthening the Early Warning Systems in place for community protection (flooding risks), vegetation protection (risks from locust invasion) and public health protection (likely severe malaria epidemics and other water-borne diseases) is recommended. |

Figure 2. Statements/advice issued by ACMAD on 21 May 2008 regarding prospects for the July–September 2008 west Africa season. These statements were based on the consensus forecast product shown in Figure 1. Note: heavy rain and flooding in July caused damage and some fatalities in Benin, Burkina Faso, Chad, the Gambia, Côte d'Ivoire, Niger, Nigeria and Togo.

Forecast statements are delivered to the users in a variety of ways, and delivery may cascade through different levels of producer and user. In the case of RCOFs the first opportunity for communication is at the Forum itself. The rainfall forecast might be used with other inputs to develop regional basin streamflow predictions and food security assessments, which are then cascaded to relevant users. Disaster management applications were prominent following the 2008 PRESAO-11 Forum (Figures 1 and 2 above). Representatives of the International Federation of Red Cross and Red Crescent Societies (IFRC) participated in the Forum, forming the kind of user–producer partnership that leads to good understanding of the forecast and its implications. On the basis of this new understanding, IFRC launched an appeal to raise funds for flood disaster management; pre-positioned emergency stock (such as drinking water and sanitation equipment); secured staff visas in advance of entering countries at risk; and initiated a community dialogue to increase preparedness to act on output from Early Warning Systems.

In their response to the 2008 PRESAO forecast, IFRC adopted a "no-regrets" approach, in which the probabilities for above-average rainfall were considered sufficient to assume a deterministic forecast of above average. This was a pioneering action of IFRC to prepare for climate impacts rather than wait and respond to an existing impact. Prior to the IFRC proactive response to the RCOF product, IFRC staff underwent considerable capacity-building through education or training on the causes of climate variability, the interpretation of probabilistic forecasts and the opportunities and limitations of those forecasts. The IFRC also carefully monitored the observed conditions. However, it would be interesting to speculate what the agency's response would be to the next forecast, if the high probability event (floods) had not occurred. Credibility is a rather fickle commodity, even when probabilities are specified correctly.

The cascade of forecast information through different levels of producers and users represents an important complement to dissemination of the forecast through media channels. The media play an important role in disseminating top level messages through a wide variety of channels, but cannot be expected to offer advice or enter into dialogue. Moreover, there is still much work to do in enhancing media understanding of the probability forecasts. This is discussed further in Section 6.2.

3.2.2 Applications tailored to specific sectors

Developing tailored seasonal forecast products that are useful enough and specific enough to find real-world applications requires persistent collaborations to co-evolve the uses of, and confidence in, the climate products. The direct use of dynamical forecasts, unfiltered by human and statistical corrections, is not yet realistic with the current state-of-the-art models. Some examples illustrating

3.2.2.1 *Water resource management examples*

Being less sensitive compared to many other applications and to some types of seasonal prediction errors, predictions of reservoir inflow represent something of an opportunity for applications. For large catchments (covering several model gridpoints), sensitivity to modest position errors in predicted rainfall anomalies is less critical than, for example, in some agricultural applications where local detail may be required. If the predictand of interest is total inflow over a long period (say, a 4- or 5-month rainfall season), the sensitivity to modest temporal errors is also reduced. In addition, for older reservoirs, long historical records of inflow are often available allowing relationships between predicted rainfall and observed inflow to be developed and used to make real-time predictions of inflow.

A notable example of the use of model-based climate predictions in reservoir management has been documented by Georgakakos et al. [30] for the northern California region. The Integrated Forecast and Reservoir Management (INFORM) project aims to directly integrate ensemble forecasts from the United States National Centers for Environmental Prediction Global Forecast System and Climate Forecast System into decision-making tools already used by federal and state managers of some of the region's largest reservoirs. The same team has developed similar tools and evaluations on the Nile [31] and in Panama [32] in service of a number of very different decision-making environments.

In the Philippines, efforts by IRI in partnership with the National Water Resources Board (NWRB) and the national meteorological service, Philippine Atmospheric Geophysical and Astronomical Services Administration (PAGASA), have led to the integration of seasonal climate forecasts of reservoir inflow into a water allocation model for the Angat Reservoir. Angat provides 97 per cent of the water supply for metro Manila, irrigation water for rice and hydropower. Seasonal forecasts help decision-makers at the NWRB to better assess options for distributing water across these multiple uses. While PAGASA had already been providing general precipitation forecast information to NWRB, this effort helped PAGASA build the capacity to use statistical downscaling techniques to produce seasonal forecasts of reservoir inflow. The existing NWRB reservoir model was revised to include these forecasts, with an interface that enables users to visualize the uncertainty of projected inflow for a given set of water allocations [33].

Objective seasonal reservoir prediction systems based on dynamical model predictions have also been developed for west Africa. Meteo France has developed a system to predict flow in the Senegal River. The system is used in management of the Manantali dam in Mali [34]. Improved management from application of the prediction system has optimized hydroelectricity generation and also enabled a guarantee that artificial flooding for flood recession farming can be achieved in three out of four years (compared to once in five years if no forecast information is used). The Meteorological Office Hadley Centre, working with the Volta River Authority, has developed a system to predict water volume inflow into Lake Volta, Ghana. The predictions are used to aid management of hydroelectricity generation at the plant at Akosombo, which provides approximately 50 per cent of Ghana's electricity [35].

Full integration of the climate forecasts with other information impacting on decision strategies is not easily won. Without collaborative involvement of climate scientists and decision-makers in developing the management tools, there is a danger that tailored forecasts become no more than an elaboration of the original seasonal climate forecast, leaving the user little better off in understanding how to actually apply the information. The INFORM project has involved years of development of both the reservoir management tools and the technologies to obtain and incorporate climate forecasts in the operational setting. More importantly, it has required years of collaboration between the researchers and reservoir managers to define the working constraints and procedures that such a system would need before being used operationally. At this time, the INFORM system is functioning in an operational mode in parallel with the existing decision-making procedures so that the decision-making agencies can themselves determine whether the system is safe and beneficial. There is a need to share experience more widely to help bring a fully integrated approach to the various developing applications in this sector.

3.2.2.2 *Agricultural decision-making example*

The International Research Centre on El Niño implemented a geographic information system for western South America that can incorporate seasonal forecasts as a layer of information relevant to dynamic risk of several regional crops types (Figure 3). The agricultural decision tool addresses the vulnerability of designated crops spatially according to the area of intervention of every country. Additional information was added such as layers of exposure to different climate hazard levels. Resiliency levels were estimated on social, economic, political and institutional parameters; land use characterization; and water retention capacity, including topography and texture, among other factors.

The combination of various information layers allows estimation of vulnerability of specific crops, and the influence of each layer can be weighted according to the region and crop in order to assemble dynamic layers from seasonal forecasts of precipitation and maximum and minimum temperatures. This allows dynamic agro-climatic risk maps to be generated for each crop. The system has been validated in every country in close coordination with expert teams from the NMHSs. Finally, the tool is able to generate maps which are updated following every forecast providing three-month risk scenarios and if monthly climate forecasts are available, every one or two months. Users can visualize the vulnerability layer, forecast layer and the associated risk for the next season represented in a simple colour scale display next to the map. The system is available through <http://ac.ciifen-int.org/sig-agroclimatico/> and is becoming operational in each NMHS in western South America.




Figure 3. Dynamic Climate–Agriculture Risk Maps generated on the basis of seasonal forecasts for pilot areas in six countries in western South America

3.2.2.3 Health – malaria example

Of the current applications in the health sector, malaria prediction in Africa is probably the best developed, although still fairly rudimentary. In malaria-prone regions, incidence is positively correlated with seasonal rainfall totals. Thomson et al. [24] demonstrated the potential for using dynamical seasonal predictions to predict malaria incidence in Southern Africa. Three centres participating in the European Union project DEMETER – the Met Office, the European Centre for Medium-Range Weather Forecasts (ECMWF) and Meteo France – demonstrated that ensemble mean November–February rainfall anomalies from three dynamical models show spatial distributions that are very similar to observed distributions when composited for the five years with highest malaria incidence and the five years with lowest incidence. Multi-model output from the three centres – constituting the EURO-SIP multi-model – has since been used as input to Malaria Outlook Forums in Botswana. Together with careful monitoring of the population immunity, the environment and the climate, the climate forecasts are the basis of a Malaria Early Warning System, and have been used effectively by the Ministry of Health–Botswana, in coordination with district authorities, the World Health Organization and regional NGOs.

3.2.2.4 Other sectors

Climate information and forecasts are being used to help forecast fire activity in peatland areas of Central Kalimantan, Indonesia. Research by IRI, Bogor Agriculture University and CARE Indonesia revealed that rainfall during the June–October dry season – when most fires are started – is critical in determining the severity of fire activity. A simple statistical forecast using the observed NINO4 sea-surface temperature index can provide decision-makers with 1–2 months early warning of the likely severity of the fire season. Forecasts of NINO4 can increase lead time further. The government of Central Kalimantan province has recognized the potential to use this seasonal forecast to take early action to help reduce fire activity in high-risk years, and a 2008 regulation on fire incorporated the use of this early warning system to help determine when burning to clear land would be too risky [36].

Additional examples from other sectors include the direct use of dynamical models in seasonal prediction of tropical storm activity [37][38], which is potentially of great value to the insurance sector. Predictions of crop yield have also been the subject of much research interest (for example, *Tellus 57A*, DEMETER special issue). However, experience from the user viewpoint is currently very limited.

4. Gaps between the provision of climate information and its use

Seasonal forecasts are more widely available than they were 10 years ago, and the dialogue between producers of information, researchers and different categories of decision-makers has been enhanced by the Regional Climate Outlook Forums. However, gaps still exist between information provided and information desired. Providers of information do not always understand the users' needs or that the seasonal predictions may not be understood by their possible users. Providers of information also may not understand the kinds of information that decision-makers can act on or appreciate the organizational and decision-making contexts in which potential users operate. Decision-makers need to be made aware of the opportunities and limitations of the information being offered and have credible demonstration of use and benefit. In addition, those who would use the information should be made aware that some of the desired forecast information may not be scientifically feasible at this time, or ever, due to limitations in prediction tools or the inherent uncertainty in the climate system. This awareness on all sides of seasonal climate forecasts requires continuous and receptive dialogue, as well as continued effort on all sides to improve the provision and use of the forecast information.

The common or perceived gaps between provision of climate information and its use primarily concern:

- (a) A lack of the spatial, temporal and element specificity, given the typical format of seasonal totals, applicable to relatively large areas, for just mean temperature and rainfall totals;
- (b) The way in which information is communicated, such as timing, content, phrasing or language, and even the means of delivery, all of which can influence the effectiveness of uptake.

(See http://www.bom.gov.au/climate/pi-cpp/training/nms/Criticism_SCO_Comments.doc.)

As a consequence, in recent years, much research also has been devoted to developing assessment methods to identify, foster and design improved and sustained connections and services linking the climate community with users. The aim of these efforts has been to support and advance the co-discovery and co-evolution of climate services and applications [39]. Recent efforts have centred around the concepts of climate services and decision support, which from the climate information producers' side is now understood to be a threefold process: first, involvement of the generation of useful climate information and predictions; second, translation of that information into forms useful for decision-makers; and third, dissemination and communication of the results [3]. All three of these processes require regular and intense collaboration with the potential users and decision-makers, if the decision supports are to be successful and are to provide better understanding of decision-making and the human dimensions of uses of climate information. These issues are expanded below with some strategies for improvement.

4.1 Spatial and temporal scale

The skill of Global Climate Model (GCM)-based seasonal climate forecasts tends to increase with increasing spatial aggregation relative to a single GCM grid cell [40]. For example, over North-east Brazil, where regional predictability is uniformly high, forecast accuracy was shown to decrease for smaller averaging regions. Interpretation of a grid box as representing a single station within it reduces skill further. This should be expected as local realization of weather and climate averages out on the larger scale. Locally specific information therefore has greater uncertainty. One option is translation of larger-scale forecasts to local scale, done either statistically or dynamically, that appropriately considers how the quantitative information, particularly its uncertainty, changes at that scale and at that particular location. Of course, such detailed information also requires sufficient observational records in order to be meaningful.

The temporal resolution of seasonal forecasts is mostly a 3-month seasonal mean. Often decision-makers are more concerned with a particular month or how the characteristics of the weather within the seasonal climate, such as dry spells, start of the rainy season or hot/cold spells may be different in the coming season. It is the change in characteristics – changing persistence or magnitude of weather events – that leads to the seasonal mean variability. There is encouraging research to suggest that this may be possible in some cases; for example, Robertson et al. [41] find that the frequency of rainfall within a season exhibits higher predictability than the seasonal total rainfall. Similarly, Lo et al. [42] find a strong ENSO influence on the start of the northern Australian wet season, which is an important decision variable for the cattle industry. However, information on higher temporal characteristics of the seasonal climate is not routinely provided by forecast providers. Yet, for some regions the Madden–Julian Oscillation (MJO)-based forecasts are showing considerable promise at this intra-seasonal timescale [43]. These are examples of tailored products most appropriately developed by regional or national climate centres in collaboration with local users and decision-makers.

Dynamical or statistical models can be employed to downscale the coarse output of global climate prediction models spatially and temporally to meet the needs for local information. Some examples can be found in the special issue of *Tellus* (2005) and by Sun et al. [44][45]. A clear benefit of dynamical downscaling over statistical downscaling has yet to be demonstrated, although dynamical downscaling experiments can reveal more information about processes of the climate and its variability. Regardless of the downscaling approach, it is necessary to demonstrate and document the ability to predict climate at scales relevant to applications or decisions, such as local values of temperature and precipitation and/or the temporal characteristics of the seasonal climate variables in the specific instances. It is also important to document to what extent prediction quality is improved or degraded relative to the coarser input; higher resolution information does not necessarily guarantee improved quality of the information. In other words, interpretation of large-scale predictions to local scale involves considerably more than blind use of models. Care must be exercised when developing such information. Some detail may never be possible to predict. Users must find ways of getting the most out of the level of predicted detail that is feasible – just as producers should strive to increase reliable detail where possible.

Regional Climate Outlook Forums can play a significant role in closing this gap by motivating participating countries to improve national capacity for seasonal forecasts, including downscaling. A better climate service for decision-making must ensure that NMHSs be able to respond to local users, often by providing locally relevant information. Applications that draw on such information encourage the private sector, government and other stakeholders to invest in the improvement of the forecast capacities.

4.2 Forecast variables and their specificity

Seasonal forecasts are typically provided as probabilistic outlooks for precipitation totals and monthly mean temperature, and these variables have been cited as useful indicators in many applications. However, decision-makers often desire more detailed prediction information as described above (number of warm days, rainy days, onset date, length of season, possible long dry spell within the season and locally specific information). Decision-makers may also require prediction of environmental variables impacted by climate variability that are connected more directly to climate risk and resource management, such as streamflow, crop yield and mosquito distributions. In addition to being variables that may not be included in dynamical models, they span a range of space and timescales: streamflow is an integrator of many rain events (in time and space) and hence tends to strengthen a signal; the date of the last or first frost is very local and a weather event that will be very difficult to forecast; many extreme events are sub-grid-scale and very challenging; crop yields and mosquito distributions are influenced by non-climate effects and are sensitive to starting conditions. The prediction of seasonal climate-related variables that are closer to many aspects of applications problems are being actively researched. Also, programs are being developed by different centres (for example, the PRESAO-SG at ACMAD) to use available science and data to produce forecasts for these elements.

No forecast is complete without a description of its uncertainty. Due to imperfect dynamical models and inherent unpredictability of the land surface–ocean–atmosphere continuum, climate forecasts are necessarily probabilistic. Uncertainties generally increase with forecast lead time and vary with weather situation and location. The methods of generating the forecast probabilities vary among producers. If the dynamical models (note that statistical predictions typically provide an estimate of uncertainty that is determined by training the prediction on past data. Careful attention is required to make sure that such hindcasts of even statistical models are generated independently of the prediction target(s)) were perfect, and, for example, 70 per cent of the ensemble members had warmer than average temperature, it would be possible to forecast a 70 per cent chance that it will be warmer than average. However, models are not perfect, and calibration of model probabilities based on past probabilistic performance is often required to produce reliable probabilities [46][47]. Reliability is one desirable attribute of probabilistic forecasts [48]. In view of the preceding example, for a reliable forecast, over a large number of cases predicting a 70 per cent likelihood for a warmer-than-average season, it should turn out to be warmer than average in 70 per cent of those cases.

To help users appreciate the necessarily probabilistic nature of seasonal climate forecasts, the forecasts should:

- (a) Be accompanied by performance statistics of the previous forecasts (or hindcasts);
- (b) Be tailored to the needs of RCCs and RCOFs and National Centres for Ocean Forecasting (NCOFs) who may add further regional or national tailoring before delivery by extension services to local users such as individual farmers in local language and including local knowledge whenever possible.

The temptation to interpret a probabilistic forecast deterministically, or as an answer with no uncertainty, is to be avoided. Government or private sector decision-makers always work with risk and uncertainty; uncertainty is part of the decision-making process. Risk, by definition, is probabilistic. Furthermore, even if a precise deterministic forecast could be issued, it would not necessarily result in a correct decision.

4.3 Communication

Making quantitative and intelligent use of probability forecasts requires an effective chain of communication between information providers and users. As discussed above, forecasts must be presented probabilistically. But are they perceived as such by local users? Do users assimilate this probabilistic information easily? Where the level of climate literacy is low, the potential for misinterpretation and inappropriate use of the forecast is likely to be high. If the climate information is not provided and communicated adequately, the whole system is ineffective, and to some extent useless or damaging.

Language, phrasing or context is often an impediment to proper interpretation of climate information. Often translation could be made by equating the likely climate shifts in terms of environmental indicators. Evidence of the use of indigenous forecasting methods to predict seasonal climate variations have been discovered in several regions [49]. Different communities may use different indicators such as observations of clouds; wind or lightning; behaviour of livestock, wildlife or local flora; or even the appearance of certain stars [50]. Unfortunately, few studies have been done to connect the physical climate system to these environmental indicators, making it difficult for climate scientists to capitalize on the possible connections. Beyond tradition, indigenous climate forecasting has been appealing because it is made for tailored elements and focuses on those crucial elements needed by the community (for example onset, length of the season, timing of the rains) at the desired spatial resolution and provided in the understood local language.

Even if indigenous forecasts are not competing, technical jargon may compromise the value of the forecast for applications. For example, the terms "normal," "average," and "climatology" are used too casually [51]. Efforts are made by producers to use clear and simple language, for example, by including a summary for non-technical users and explaining the main factors and forecast tools. In some cases a glossary may be provided to explain terminology. (See: <http://www.cpc.ncep.noaa.gov/products/predictions/90day/seasglossary.html>.) Such details are also available within the process of RCOF/NCOF to those who can attend. However, this supplementary information is often insufficient or inaccessible to users in remote areas with little access to modern IT facilities.

Many would-be users of climate forecast information rely on the radio. This again emphasizes the importance of a well formed, and well informed, chain of communication. In one example of providing forecasts over the radio, new and easy-to-use technology (RANET system) was introduced in 2000 for dissemination of forecasts through an organized system to local communities in local language. RANET is a collaborative effort of ACMAD, NOAA, many National Hydrometeorological Services, non-government organizations, and communities. These varied partners come together to make weather, water, and climate information available to

communications technology such as EUMETcast from the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT), are being implemented and tested for dissemination of climate information. However, there is still considerable need to improve the effectiveness of communication of the climate information, by using multiple channels, including press, cellular phones, satellite television and the Internet. Some alliances with local media, who may be open to broadcasting seasonal predictions free of charge, have been positive. Such alliances require coordination, but they can then ensure sustainability of information delivery.

The interaction between providers and users has been historically low within the NMHSs and in the scientific climate community. Along the chain of information, the RCOFs play an essential role in bridging the communication gap. The RCOFs and NCOFs can provide a venue for the NMHSs to build strategic alliances that can sustain an interaction with the key stakeholders. Such alliances are especially important in cases where such outreach cannot be supported within the NMHSs. The private sector and the media must also assume responsibility and become part of long-term solutions, avoiding the tendency to focus on climate information only when disasters occur.

Clearly, the media also play a key role in the climate information chain. In many RCOFs, media professionals are major partners in the discussions with both researchers/producers and users. (See also Section 5.) Examples of media misinterpretation abound and can result from genuine misunderstanding or from a desire for eye-catching headlines. For example, a prediction by the Meteorological Office (United Kingdom) of a 2 in 3 chance of below-median (1971–2000) European temperatures for winter 2005–2006 [27] was widely interpreted by the United Kingdom media as implying a very cold winter – even though the forecast itself contained no information on severity (for the United Kingdom, two-category precision was the best that prediction skill would allow). To alleviate the risk of media misinterpretation, many forecast centres employ media experts in the crafting of the forecast statement, following the reasoning that – as headlines will be written – it is better that the forecasters (rather than journalists) make the first draft. Press briefings are also used by many centres to help convey the correct message to the media. While it is helpful to train the media regarding seasonal climate forecasts, such forecasts do not often make for exciting headlines. Thus, the main goal with respect to the media should be to ensure, at the least, that they understand the best that science can do about seasonal forecasts, and the dangers of the miscommunication of that information.

4.4 ENSO as a case study

El Niño is used here as a case study to demonstrate the degree of evolution in climate information systems. Although El Niño 1997/1998, was an extraordinary event and the prediction was successfully done before the middle of 1997, many countries did not react or take the necessary actions to cope with the expected climate impacts. One of the main reasons for this was the multiple sources of information. Within western South America, official statements from the national level were ignored by the media and more attention was given to the global agencies. However, while the global statement indicated the evolution of El Niño, the manifestation of El Niño through climate impacts were not evident in much of north-western South America until very late 1997 due mainly to the seasonal cycle. The lack of confidence in the national institutions plus the chaotic dissemination of information from multiple, and often non-qualified, sources led some governments to ignore the best ENSO prediction to date. By the time that El Niño climate impacts were manifest, several Latin American countries were effectively taken by surprise and impacted negatively due to limited planning.

Ten years after the El Niño 1997/1998, the climate information management regarding ENSO has become critical. When the media, governments and users are informed about ENSO, however, they retain a fixed idea of the 1997/1998 impacts, without possibility of variation. In the first decade of the twenty-first century, moderate to weak ENSO events have not led to the same dramatic impacts at the local level in western South America that were experienced in 1997/1998. Although the climate forecasts have been consistent with that outcome, the final result is that ENSO predictions are becoming useless in some countries due to the disconnect between Niño Region 3.4 evolution and the expectation of local ENSO effects in each country. The International Research Centre on El Niño supported a survey in Latin America about what ENSO is for each country. The finding was that the number of ENSO definitions is close to the number of countries involved in the survey. One conclusion was that for Latin America, communicating just ENSO as a forecast can be a particularly ambiguous and confusing way to communicate climate information. Several countries are now more concerned with improving their capacity to consider the wider range of factors that can influence the climate forecasts over the region and concentrating more effort on communicating those forecasts.

5. Techniques for increasing the value of climate forecast information

As discussed in the previous section, several gaps exist between the information typically provided in seasonal forecasts and that needed for climate risk management and decision-making. Some of these gaps can be addressed by the scientific community, which includes climate and sectoral specialists. Others must be addressed by the larger integrated process that enables actions to be taken and realized in the presence of forecasts of the climate and its impacts.

5.1 Technical efforts

5.1.1 Translating climate forecasts into more relevant variables

Operational climate risk management requires knowledge about the likely consequences of the climate. Variables of interest for decision-makers often differ from what climate forecasts routinely provide. Instead of seasonal rainfall forecasts, decision-makers often require quantities such as crop yields [52] or return on investment [53], provided probabilistically. Such probabilistic representation of decision variables helps risk managers to conduct rapid assessments of management options. In some cases, this might even mean bypassing climate forecasts and going straight to the decision-relevant variable, often by using several models. The term "model" here refers to any simplified representation of a system that enables the investigation of the properties of that system and allows prediction of future outcomes.

In agriculture, modelling can be used to evaluate the efficacy of potential innovations (such as climate forecasts) at various levels of integration – from genetic engineering, to phenotype expression, to crop and cropping system management, to regional governance and policymaking. At the field to farm level, models are already used for operational risk management [54] and for the design of more resilient farm business [55], while they have also become indispensable in plant sciences to understand and predict the complexity of biological systems [56][57]. Likewise, public as well as private sector policy decisions are increasingly informed by the design of and output from simulation models [58][59][60] including the development of new insurance products [61][62]. Connecting agricultural models with climate models remains a substantial challenge for the use of General Circulation Models (GCMs) for operational risk management, but it is not insurmountable [63]. The most pressing issue that should be addressed is designing scientifically and statistically robust methods for providing GCM output at the appropriate spatial and temporal scales: while most agricultural models operate at a point scale using historical daily station data as input, GCM output is generally provided for large grid cells with vastly different attributes to the climate data for which the agricultural models were developed. In addition, methods that would readily allow one to obtain reliable probability distributions of forecast variables must be available. This is an essential condition before the information can be used by biophysical models. Although statistical tools exist for providing calibrated probabilistic information at the right scales [63][64], agreed protocols for implementation are lacking.

5.1.2 *Toolboxes*

Providing information probabilistically is absolutely essential for responsible forecast provision. This is emphasized in a WMO report [65] that explicitly states that only probabilistic forecast systems should be considered for risk management. The report lists four key forecast system attributes: a) consistency (whether the forecasts correspond with the forecaster's judgment); b) quality (whether the forecasts correspond with the observations consistent with its issuance); c) relevancy (whether what is forecast is of concern to the user); and d) value (whether the forecasts are/can be beneficial when used). Forecast quality, often referred to as "skill", encompasses a wide range of statistical properties and is an essential precondition before a forecast can become valuable. Because dynamical models contain notable deficiencies, the quality of dynamically based predictions can often be improved through the use of statistical post-processing. Freely available software such as the Climate Predictability Tool (CPT) (<http://iri.columbia.edu/climate/tools/cpt>) is one example of a toolbox that can spatially and probabilistically recalibrate predictions from a dynamical model, generally improving the quality of the prediction at a given spatial resolution. The CPT can also be used to produce statistical forecasts based on current or recent observed conditions that can serve as a comparison to the dynamical prediction. And the CPT can be used to statistically downscale coarse model predictions, assuming a sufficient history of observations exists at the desired scale.

5.1.3 *Historical climate records as a tool for decision-making*

The analogue year approach – still often held as the benchmark to be beaten by GCM-based methods – has been successfully used in some regions based on a range of different climate indicators such as the Southern Oscillation Index (SOI) [25]; SST or Real-time Multivariate MJO predictions based on upper level wind; and outgoing long-wave radiation anomalies [43]. The advantage to such an approach is that the climate information consists of past observations, so the implications can be readily connected with decision models, and no supercomputer is needed. Some of the obvious problems include difficulties in dealing with short record length and low station density, particularly in developing countries. Non-stationarity due to decadal/multi-decadal climate variability and climate change can also severely limit the usefulness of analogue approaches.

5.2 *Chain(s) of communication enabling the use of climate information*

Decision-making in sectors such as agriculture, health or natural resources consists of individual choices to which collective action at local level is often the appropriate response [66]. It takes place within specific social and institutional settings that provide the framework within which actions have to be taken. It also takes place within a global setting, where global and national policies influence choices and actions. Decision-makers require quantitative information to supplement their already existing empirical knowledge that is based on years of experience, expert judgment, insight and intuition. Hence, decision-making is highly context-specific and scale-specific [67]. Climate science and forecasting might be able to add to this process, but it should never try to replace it.

The following discussion relates largely to decision-making in agriculture and resource management, two of the most climate sensitive sectors in our economy. While the lessons learned from these examples can be generalized, the sectoral context is critical for the success of the forecast. For illustration purposes, we retained this context.

Here we ask the question: why is the available climate information often not used and embraced by decision-makers? Lack of "ownership" of the information by the intended end-users is clearly one issue and has led to a growing acceptance among climate scientists that they must move out of their disciplinary confines and engage in a process of continued, shared learning and joint problem solving [68][69].

More efficient and effective policies would, at least in principle, result from common approaches and technologies, and would provide decision-makers at all levels and scales (from farm and agribusiness to policy) with more objective, faster and lower-cost information. All stakeholder groups could then objectively compare options, evaluate choices and assess policy or management consequences. Analytical approaches, such as systems modelling, facilitate this process and help stakeholders to identify and choose between inevitable trade-offs along the path to sustainable development [70].

Cash et al. [71] and Cash and Buizer [72] argue that translating climate information into real life action requires three essential components – salience, credibility and legitimacy. Salience relates to the perceived relevance of climate information: does the system provide information that these users think they need, in a form and at a time that they can use it? Credibility addresses the perceived technical quality of information: does the system provide information that is perceived to be valid, accurate, tested, or, more generally, at least as likely to be "true" as alternative views? Legitimacy concerns the perception that the system has the interests of

Decision-makers usually manage risk holistically and often intuitively [73][74], while climate science tends to provide specific, detailed and generally technical information. This can create a perceived lack of science relevance that can be overcome by embedding technological approaches within context-specific, multi-stakeholder dialogues. Such participatory processes can translate scientific information into real life action by paying attention to salience, credibility and legitimacy as proposed by Cash et al. [71]. They contain at least two critical elements that need to be reconciled:

- (a) A participatory process of negotiating, building trust and creating knowledge;
- (b) A solid scientific and technical basis that can be used for such knowledge creation.

So far much of the emphasis has been on the latter. We suggest taking a complementary approach: rather than using climate science as the starting point for this process (supply), we need to critically appraise stakeholders' problems or decisions and – in a participatory approach – jointly identify and outline the specific needs for trans-disciplinary scientific input. Through this process we can then better match technical options with socio-economically feasible solutions. Here we review some examples of these technologies and suggest ways how they are – or might be – applied.

5.3 A human-centric approach to the definition of forecast value

A forecast has no value unless someone uses the information, acts upon it and subsequently achieves a better outcome than they would have achieved otherwise [70]. This is quite different from the largely technical definitions of forecast quality that has so far dominated forecast assessments [48][65]. While it is important to establish forecast quality for the basic credibility of the scientific approach on which forecasts are based, the efforts necessary to estimate forecast value extend much beyond the climate science community. A real impact on decision-making requires a more human-centric definition. We therefore define “forecast value” as “the features or characteristics of a forecast product and/or forecast service that enables action and satisfies identified and agreed needs of the user community”.

This definition, although it cannot be determined by the climate community alone, makes the concept of forecast value tangible and enables useful, meaningful and measurable improvements in climate science by:

- (a) Highlighting aspects of the forecast system that matter to users through a dialogue about the role and value of science in decision-making;
- (b) Enabling action on the feedback received from users by designing forecast products that comply with user requirements, making forecasts more relevant and useful for society;
- (c) Explicitly considering different relevant temporal and spatial scales when designing new forecast products.

A better understanding of the importance of the timing of climate information can affect its uptake. Providing climate information in forms, and at times, that support decisions requires significant understanding of the decisions that must be made and the way that climate information can enter those decisions. For example, one useful approach has been collaborative development of decision calendars [75][76] that identify regularly occurring moments when, predictably, existing or needed climate products would be of maximum use in, for example, decisions about sowing crops or releasing water from reservoirs. Developing such information requires persistent collaborations with the potential users, because the full range of ways that climate influences many management systems is poorly known; the producers of climate information need to learn the needs of the users, and the users need to learn the capabilities of the producers.

Howden et al. [77] suggest a reflective risk management loop that clearly articulates the type of information needed for successful change management in agriculture and natural ecosystems. At the centre of this loop are three interacting domains – environment; production; and norms and values. At each step a stakeholder dialogue ensures that the needs of these domains are aligned with the scientific knowledge to be created. The following steps form part of this loop:

- (1) Understand the existing system, and scope possible changes to values;
- (2) Identify likely core issues and decision criteria, and clarify who, what and when;
- (3) Assess (climate) impacts and trends, including the uncertainties;
- (4) Evaluate impacts;
- (5) Assess the adaptation options, and their broader consequences and links;
- (6) Design and evaluate implementation options.

In cases where the evaluation of the final step is positive, changes to the environment, to production practices or to the norms and values (which include policy) will be implemented and the loop starts again, this time with the modified system. In contrast, should the evaluation fail, no action takes place, but the loop continues.

Climate information is required for step 3, while step 5 determines the type of climate information that is needed for better risk management (for example, onset dates for the wet season). The following example used this framework.

Profitability of the cattle industry in northern Australia is directly related to the amount of live weight gain that can be achieved before early monsoonal rains make the transport and marketing of cattle impossible. Cattle producers therefore have to balance their decisions: selling stock too early results in lower weight and reduced profits, while waiting too long can prevent the selling of cattle altogether as access becomes impossible after a certain amount of rainfall. Until recently there was no reliable forecast that cattle producers could use to help in their decision-making. Existing forecasts of monsoon onset, based on meteorological definitions, are meaningless to them because the onset of the monsoon occurs long after the build-up of the wet season has started and transport routes have become impassable. This motivated Lo et al. [42] to develop an ENSO-based statistical forecast system that is tailored to the needs of the cattle industry in northern Australia. Their system probabilistically predicts the wet season onset date, defined on the accumulation of rainfall to a threshold determined in consultation with the cattle industry. Consistent with earlier studies, the interannual variability of the onset dates is shown to be well related to the immediately preceding July-August Southern Oscillation Index. The probability that onset will occur later than the climatological mean date is predicted using logistic regression. When assessed using cross-validated hindcasts, the skill of the predictions exceeds that of climatological forecasts in the majority of locations in northern Australia. At times of strong anomalies in the July-August SOI, the forecasts are reliably emphatic.

5.4 The need for ongoing climate science and prediction research for improving the quality and value of climate information

Efforts toward interpretation and tailoring of climate predictions can improve the quality and usability of the information, but much research and development are still needed on the development of models and observing systems that are essential to those forecasts. Not all possible sources of climate forecast skill have been identified or exploited. Even for those contributors that the scientific community recognizes as contributing to forecast skill, certain processes may be inadequately represented in models, or poorly understood through observations. Examples include land–biosphere–atmosphere interactions and the predictability due to the cryosphere in seasonal-to-interannual predictions. Modern climate forecast models strive to capture predictive influences from all these conditions [78][79] but forecast skills have grown only grudgingly [80][81]. Improvements in our ability to forecast ENSO and its impacts [82], along with whatever other long-lead predictability, less exploited climate modes may contribute, remain the focus of attempts to improve climate forecast skills at large scales. Improved models will be necessary to realize such improvements. Deficiencies in nearly all dynamical models in representing certain key physical processes, such as tropical convection and aspects of the tropical upper ocean dynamics, suggest strongly that room for improvement exists.

At still longer timescales, predictability is more difficult to come by. Recognition during the past decade of the comparable role of inter-decadal climate variability with seasonal to interannual variability in many areas of the globe has improved our understanding of climate variations overall. However, the difficulties of predicting climate beyond a year have thus far limited the predictive value of that recognition. The climate system is, for most part, too chaotic and internally variable to support most useful, longer-lead forecasts of climate variability with current technologies [83]. Again considerable research is required to develop models that appropriately capture the relevant processes and data assimilation systems that can incorporate the observed state of the climate system that is essential to initialize decadal scale predictions.

As an example of innovative research into the generation and improvement of the utility of climate predictions, methods for determining the quality and value of operational climate forecast products in very specific, and often local, applications and decisions have been developed [51]. These assessments not only address whether forecasts or other information are good enough for inclusion in decision-making but, because nearly all decisions already depend on imperfect information about a wide range of stressors and conflicts, also inform users about how much to trust or value the climate information relative to many other information types and sources. This latter step has been recognized as the more important aspect of climate information valuation in most real-world applications.

6. Lessons learned

This section provides a synthesis of lessons learned in RCOFs around the globe after the RCOF review process held in Arusha, United Republic of Tanzania, in November 2008 [84]. It is based on practical experiences in each region and sets up a good reference for further recommendations.

6.1 Learning about the users

Regional Climate Outlook Forums provide face-to-face contact. A Forum is not a regular meeting; it is a mutual learning process where users learn about the nature, quality and value of forecasts, become better acquainted with terminology and, more importantly, understand the limitations of climate predictions. On the other side, forecast providers learn about user perception of their information such as their interpretation of a coloured map of the terciles. The Forums have provided the opportunity for the climate community to learn more about the users' profiles and with this knowledge, develop ways to tailor and communicate climate information. Language, native phraseology, ancestral climate knowledge, local culture and the intuition of the colour scales are all part of the elements used to better communicate [85].

Additionally, the RCOFs have allowed for the identification of key contacts, or relevant actors in the region, who have the capabilities to disseminate, share or transfer information to other contacts for a positive socio-economic effect. These contacts maximize dissemination efforts and reduce the risk that climate information goes unused within a decision-making process. Key contacts must include government officials who provide advice and/or ensure that climate forecasts reach the authority or decision-maker. The list of key contacts should be a widely accessible document that is continuously checked and updated.

6.2 Involving the media

Climate forecasts generally produce high expectations from the media. This situation could be good or bad depending on how a press conference, a personal interview or a newspaper report is handled. Even after several training workshops for journalists in some regions, the results may not be satisfactory. However, it has been demonstrated that efficiency and effectiveness can be obtained by

sectoral specialists in the fields of water, agriculture, health, energy and disasters who are trained in the interpretation and applications of climate information, including both recent observations and predictions, as these experts are already communicating other types of information to users (prices, markets and epidemics, for example).

Press, radio and TV require different formats to communicate the same forecast, and an additional effort to prepare and tailor products according to the interested medium can be of value.

Some RCOFs have included the media contacts as participants. They play the roles and responsibilities of key contacts and assist NMHSs with dissemination. Special media sessions have facilitated good communication in some RCOFs. This approach requires regularity and quality of climate forecast bulletins as well as the media interest, which usually comes from the relevance of the provided information for their audience [86].

Climate forecasts in some regions receive specific spaces in press, magazines and other media without any cost, an arrangement that provides more visibility to the information. In these cases, the media benefit is likely to have been demonstrated.

Press conferences could be the best or the worst way to disseminate climate information. Several regions have experienced the negative impact of an ill-handled press conference. The entire effort of producing the forecast, carefully reviewing the statement and other elements could be wasted with a single misstatement of the official spokesperson, generating instead misinformation and lack of credibility with the user community. Experience shows that it is best to disseminate an accurate statement of the forecast that can inform the media, but that is still concise, friendly and explicit.

6.3 Changing the language

The best way to ensure a communication channel with users is to make the wording of forecasts simple. This requires an additional, but worthwhile, effort from climate information providers that usually helps to expand the user community. In such cases, some technical and complex details are omitted leading to an understandable message that can trigger a response from users. Good experiences have been noted in traditional communities where the climate information has been reworded using some native phraseology rather than technical jargon. The level of positive reception is evidenced by the further demand for regular information.

6.4 Receiving and assimilating user feedback

The connections along the chain of information are enhanced when the perceptions of users are integrated in the communication process. In some regions the RCOFs have been replicated as national outlook forums providing the space to receive the user feedback. An effective process involves listening to users' comments, analyzing them and then integrating them into the communication strategy. For example, CIIFEN receives positive reactions when users see some of their suggestions incorporated in climate bulletins. As a consequence, the relationship with users gets stronger and the use and communication of the climate information improves. Climate information providers should be flexible enough to make information available in different formats, wording and graphic design.

6.5 Involving the private sector

One of the best practices in the delivery of climate information to other users, such as the private sector, has been to transform their passive role as listeners into that of actors. This has been achieved when the users are requested to participate in the dissemination process. After understanding the potential benefits of climate forecasts, private sector partners become key actors. Funding required for sustainability is increasingly provided by the private sector, making their involvement very important in future climate information strategy, especially in developing countries.

6.6 Getting positive responses from governments

An important segment of users is composed of government authorities. At the national level, the NMHSs are important liaisons with relevant contacts with the government. Although it is not an easy process, the identification of experts who can advise authorities about the application of climate forecasts can be highly effective. The initial target for engaging governments may not be the top authorities (Ministers, for example); it will likely be their technical advisors. Involvement of national authorities from different sectors such as civil protection, planning, agriculture, health, energy, infrastructure, water resources, banking and finance and science and technology, among others, results in powerful connections that facilitate effective responses to climate information and services at the national level. A good practice to implement where possible is the invitation to participate in RCOFs via videoconference. Relevant authorities or decision-makers get involved, participate in discussions and gain access to RCOF conclusions at a very limited cost.

6.7 Customizing climate products

Special bulletins for specific industries or sectors are a "best practice" with evident results. This requires additional effort to know user needs and to be able to reflect them in the product. The era of standard weather or climate bulletins is over. Among the many challenges for the meteorological community in its effort to support sustainable economic growth with better climate services, the development of more customized information through more regular interactions with the users is one of the most important. When done properly, the communication process is quite effective at improving user response to climate information.

6.8 Getting involved in user activities

Users organize several meetings and activities where members of the climate community, mainly the NMHSs, are invited. Positive response to these invitations builds a stronger relationship, allowing climate information providers to listen and learn about

users and their problems, some of which may be handled with climate information. It is important and very rewarding to be involved in users' activities as much as possible to lay the foundation for better application and dissemination of the climate information.

6.9 *Generating trust from users*

To build more trust in the user community, a set of actions are needed. Regular face-to-face contact and personalized e-mails or phone calls encourage the user to integrate climate information in his day-to-day climate-sensitive activities. User trust also relies on the presentation of climate forecasts and the articulating of uncertainties and limitations. Users require climate information in spite of the limitations, but these must be informed properly and with transparency [87]. Such individualized interaction with the user community is often not possible by the limited staff in NMHSs or even regional climate centres. This again highlights the need and the value of the chain of information and the role that local authorities – whether boundary organizations, sectoral specialists, government officials or the media – can serve in building a strong climate service network.

6.10 *Demonstrating the effectiveness of climate applications*

The best way to convince users and to involve government authorities, media, private sector and others is to demonstrate the benefit of incorporating climate information into their efforts. This can be done through pilot projects that document the process and the outcomes over time. Once the results are evident, additional support will come from partners who become more motivated to scale up pilot projects to other geographic areas and/or development sectors. More pilot demonstrations are needed to further document, test and exchange best practices of user-provider interactions.

7. **Summary and the way forward**

Many of the factors that contributed to the under-utilization of the available science-based seasonal climate forecasts were discussed in previous sections of this article. The lessons learned strongly suggest that the way forward needs a cultural change in the interaction of the climate science community and the users. A first element of this needed change consists of considering the demand side as the starting point and is the main focus of this interaction, as opposed to using a supply-oriented approach [88][89]. When the main focus is on the demand side (users), climate information and climate products become just one of the many types of information that feed the decision-making and planning processes in the different socio-economic sectors [54]. Just like the many other types of information needed for decisions and planning, climate products must be available in formats that are understandable, with spatial resolution that makes them usable, and must be provided with adequate lead times to make them actionable.

A second lesson learned is that the sectoral stakeholders and the climate scientists need to approach the ongoing efforts in research and applications of seasonal climate forecasts as a work in progress. Up to the 1990s climate-related decisions could only be made on the exclusive basis of the highly unlikely “average” climate conditions described above. The last few years have witnessed significant advances in the ability of the climate science community to provide probabilistic information on the expected climate conditions for the next season, and that information is potentially very useful to assist the decision-making and planning activities in the different sectors. Thus, the current capabilities of seasonal climate forecasts allowed stakeholders to evolve from having nothing to assist decisions and planning, to having something. Climate scientists continue to invest efforts to improve their something by enhancing the skill of the forecasts, increasing their spatial resolution, and by improving the ability to forecast climate variables that are more relevant for the different sectors than the total seasonal means (for example, the onset of the wet season, the frequency and duration of dry spells, the timing of cold spells, dust density or the probability of extreme events).

A thrust is now needed to clearly understand the climate-related problems of the sectoral stakeholders, and to translate the climate information at various scales into sectoral products and information that can be directly embedded into decisions and policies. This translation requires that the climate information is communicated using formats and language that can be readily understood, provided at lead times and temporal/spatial resolution that make it actionable and that it is explicitly and strongly linked to relevant sectoral information. For example, there is a need to develop methods that combine good monitoring systems with climate forecasts that produce streamflow forecasts, crop yield outlooks and infectious disease alerts. There is a consequent need to enhance the local capacities to develop, establish and operate information and decision support systems that use climate-related products but also sectoral information and simulation tools.

Effectively communicating climate information and embedding it in decision-making, planning and policies requires establishing chains of information. It is unrealistic to expect that an organization that provides seasonal climate forecasts, whether a National Weather Service, a Regional Climate Centre or a Specialized Meteorological Centre, will be able to include experts in all the socio-economic sectors that are relevant in any given country or region. A much more effective strategy consists of taking advantage of the structures and institutional arrangements that already exist in the different sectors [90]. For example, the agricultural sector of developed and developing countries includes technical advisers of NGOs, private companies and public institutions who continually interact with farmers. The interaction with the farmer community is holistic in nature, that is, advice is provided in issues that embrace the whole farming operation such as fertilizer use, crop and livestock management practices, pest control, markets and commercialization. This holistic interaction forces advisers to learn and use adequate communication strategies and language, and results in establishing and enhancing their trust within the farmer communities. The communication of relevant climate information is therefore much more effective when it is carried out through these advisors. Consequently, efforts are needed to improve a two-way communication between the providers of climate information and this type of intermediary. Improving this communication includes developing the capacity of advisers to understand the climate information and products (including their limitations), and translating it into information and products that are understandable and usable for the farmers. It also includes improving the capacity of the climate service providers to better understand the types and formats of information and products that are relevant to the agricultural sector.

Similar two-way communication is needed between the climate science community and the seasonal climate providers (National Weather Services, Regional Climate Centres or Specialized Meteorological Centres) as part of the chain of information.

example, understanding the causes of climate variability. Such research is essential for providing a scientific basis for claiming predictability of seasonal climate. This type of research, as well as work targeted to create or improve coupled climate models, is grounded in typical academic and scientific institutions. It must be endorsed by publication of innovative methods in leading peer-reviewed scientific journals, and it can be completely disconnected from the demands of socio-economic sectors. Scientific groups can ensure the socio-economic relevance of the scientific advances by capitalizing on the scientific outcomes and developing applications and products that are helpful for intermediary organizations (climate service providers and/or boundary organizations directly linked to socio-economic sectors). Here again, a two-way communication has proven to be effective in shaping aspects of the research agendas in academic institutions to facilitate a smooth transition from basic science to applied research.

The ideal set of components of these chains of information varies for the different settings, regions, socio-economic sectors, scales and other factors. However, the successful cases always include strong two-way communications between the different components of the information chain, from the end-users, through a set of boundary organizations to the institutions conducting basic research.

The last few years have also witnessed increased awareness in the impacts of observed and expected changes in climate at longer timescales (climate change). Accordingly, there is a huge demand from stakeholders acting at different spatial scales (global, regional or local) and in different socio-economic sectors (agriculture, food security, health, water and disasters) for climate information across a continuum of temporal scales from days through seasons to decades. For example, the International Federation of the Red Cross and Red Crescent Societies and the United Nations World Food Programme require seasonal climate forecasts to improve their planning and disaster risk management strategies, but they also need climate information at much shorter timescales (from monitoring current conditions to forecasting a few days ahead) to improve their immediate preparedness and response to climate-related disasters. On the other hand, governments and development agencies (such as the World Bank and United Nations Development Programme) are increasingly demanding information at longer timescales. In the developing country context, the longer-term climate information that is considered most actionable and therefore is being most intensively demanded is the information for the next one to three decades (near term climate change).

Consequently, efforts to improve the provision, communication and effective application of climate information and products at seasonal timescales must be carried out in connection with the demands on other temporal scales. According to the Intergovernmental Panel on Climate Change (IPCC) Working Group II (2007), adaptation assessments benefit from linking future changes in climate to past and present variability. The new seasonal climate forecast needs to be conceived as part of the portfolio of climate relevant information that is increasingly needed by the stakeholders who act at different geographic scales and in the different socio-economic sectors.

8. Conclusions and recommendations

Conclusions:

- (a) The importance and visibility of climate information systems (CIS) has risen dramatically in the last few years, a trend that is likely to continue. Climate information systems provide information that is relevant for climate-related risk management and decision-making. Climate forecasts are important components of CIS, but the utility of CIS goes much further than just forecasting.
- (b) The awareness of seasonal-to-interannual climate forecasts has also increased considerably since the late 1990s, due in large part to Regional Climate Outlook Forums and intense media coverage of the 1997/1998 El Niño event. However, much more effort must be invested in demonstrating use and increasing utility of these forecasts.
- (c) Increased use and benefit of seasonal-to-interannual forecasts will occur only with appropriate interpretation or tailoring of climate predictions, particularly in the case of dynamical model predictions. Much work is required to further develop and link seasonal-to-interannual prediction models with application models (for example, crop yield prediction).
- (d) Effective CIS must involve all actors and not just the National Meteorological and Hydrological Services.
- (e) A better climate service for decision-making must ensure that NMHSs and local climate services be able to respond to local users, often by providing locally relevant information, and those services must be supported even as local needs may vary from year to year.
- (f) A culture change is required to build a chain of communication that realizes the benefits of advances in climate predictions to society. The chain must target decision-makers responsible for national infrastructures and welfare, and should include climate intermediaries and NMHSs, sectoral scientists, government, business sectors, media and others.
- (g) Involving the relevant sectoral scientists and decision-makers as collaborating partners very early in the process is critical to ensuring relevance, trust and ownership of climate-related decision systems.
- (h) Using CIS that focus on managing climate variability is an important means of preparing for the near future of climate change.

Recommendations:

- (a) The climate science community should:

- (1) Advocate for wider consideration of climate information in climate-related decision-making and risk management;
 - (2) Promote training on communication of climate information for NMHSs and stakeholders;
 - (3) Encourage governments to invest in diverse dissemination structures;
 - (4) Encourage the implementation of more demonstrations through pilot projects;
 - (5) Request more government support for the participation of NMHSs and stakeholders in RCOFs;
 - (6) Promote mechanisms for advances in climate science to support existing networks/responsibilities by enhancing understanding and technical capacity for better decision systems and climate risk management;
 - (7) Establish and maintain regular, tailored climate bulletins to meet specific user requirements;
 - (8) Document and disseminate the pitfalls, benefits and success stories of climate products at national, regional and global levels;
 - (9) Encourage research and technology transfer of methods to tailor climate predictions/projections;
 - (10) Encourage open access to data from both observations and dynamical models for present and past conditions;
 - (11) Promote investments at international and national levels to improve intermediary structures for climate information.
- (b) Global Producing Centres, RCCs and NMHSs should maintain and manage a network of key contacts through science collaboration, staff exchanges, regular visits, e-mails, phone calls, local workshops and videoconferences.
- (c) On-going investments must be placed in continuing to improve dynamical climate systems, including models, data assimilation systems and ensemble techniques.

References

- [1] Cane, M.A., S.E. Zebiak and S.C. Dolan, Experimental forecasts of El Niño. *Nature*. 321 (1986) 827.
- [2] Folland, C.K., J. Owen, M.N. Ward and A. Colman, Prediction of seasonal rainfall in the Sahel region using empirical and dynamical methods. *Journal of Forecasting*. 10 (1991) 21.
- [3] Beller-Simms, N., H. Ingram, D. Feldman, N. Mantua, K.L. Jacobs and A.M. Waple (eds), Decision-Support Experiments and Evaluations Using Seasonal-to-Interannual Forecasts and Observational Data: A Focus on Water Resources, U.S. Climate Change Science Program, Synthesis and Assessment Product 5.3, Asheville, NOAA National Climatic Data Center. 2008.
- [4] Redmond, K.T. and R.W. Koch, Surface climate and stream flow variability in the western United States and their relationship to large-scale circulation indices, *Water Resources Research*. 27 (1991) 2381.
- [5] Mantua, N.J. and S.R. Hare, Y. Zhang, J.M. Wallace and R.C. Francis, A Pacific interdecadal climate oscillation with impacts on salmon production, *Bulletin of American Meteorological Society*. 78 (1997) 1069
- [6] Enfield, D.B. and L. Cid-Serrano, Projecting the risk of future climate shifts, *International Journal of Climatology*. 26 (2006) 885.
- [7] Hoerling, M.P. and A. Kumar, A perfect ocean for drought, *Science*. 299 (2003) 691.
- [8] Cohen, J. and D. Entekhabi, Eurasian snow cover variability and northern hemisphere climate predictability, *Geophysical Research Letters*. 26 (1999) 345.
- [9] Clark, M.P. and M.C. Serreze, Effects of variations in East Asian snow cover on modulating atmospheric circulation over the North Pacific Ocean, *Journal of Climate*. 13 (2000) 3700.
- [10] Koster, R.D. and M.J. Suarez, Soil moisture memory in climate models, *Journal of Hydrometeorology*. 2 (2001) 558.
- [11] Intergovernmental Panel on Climate Change (IPCC), Summary for Policymakers, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller, eds). Cambridge and New York, Cambridge University Press, 2007.
- [12] Krishnamurti, T.N. C.M. Kishtawal, T. LaRow, D. Bachiochi, Z. Zhang, C.E. Williford, S. Gadgil and S. Surendran, Multi-model superensemble forecasts for weather and seasonal climate, *Journal of Climate*. 13 (2000) 4196.

- [14] Zheng, F. J. Zhu, R.-H. Zhang and G. Zhou, Improved ENSO forecasts by assimilating sea surface temperature observations into an intermediate coupled model, *Advances in Atmospheric Sciences*. 23 (2006) 615.
- [15] Kim, J. N.L. Miller, J.D. Farrara and S.-Y. Hong, A seasonal precipitation and stream flow hindcast and prediction study in the western United States during the 1997/98 winter season using a dynamic downscaling system, *Journal of Hydrometeorology*. 1(4) (2000) 311.
- [16] Kyriakidis, P.C. N.L. Miller and J. Kim, Uncertainty propagation of regional climate model precipitation forecasts to hydrologic impact assessment, *Journal of Hydrometeorology*. 2 (2001) 140.
- [17] Dettinger, M.D. D.R. Cayan and K.T. Redmond, United States stream flow probabilities based on forecasted La Niña, winter-spring 2000, *Experimental Long-Lead Forecast Bulletin*, Center for Land-Ocean-Atmosphere Studies, 8 (1999) 57.
- [18] Sankarasubramanian, A. and U. Lall, Flood quantiles in a changing climate: Seasonal forecasts and causal relations, *Water Resources Research*. 39 (2003) 1134.
- [19] Hansen, J.W. Realizing the potential benefits of climate prediction to agriculture: Issues, approaches, challenges. *Agricultural Systems*. 74 (2002) 309.
- [20] Cabrera, V.E. D. Letson and G. Podestá, The value of the climate information when farm programs matter, *Agricultural Systems*. 93 (2007) 25.
- [21] McIntosh, P.C. M.J. Pook, J.S. Risbey, S.N. Lisson and M. Rebbeck, Seasonal climate forecasts for agriculture: Towards better understanding and value, *Field Crops Research*. 104 (2007) 130.
- [22] Hansen, J.W. A. Mishra, K.P.C. Rao, M. Indeje and R. K. Ngugi, Potential value of GCM-based seasonal rainfall forecasts for maize management in semi-arid Kenya. *Agricultural Systems*. 101 (2009) 80.
- [23] Hammer, G.L. J.W. Hansen, J.G. Phillips, J.W. Mjelde, H.S.J. Hill, A. Love and A. Potgieter, Advances in application of climate prediction in agriculture *Agricultural Systems*. 70 (2001) 515.
- [24] Thomson, M.C. F.J. Doblus-Reyes, S.J. Mason, R. Hagedorn, S.J. Connor, T. Phindela, A.P. Morse and T.N. Palmer, Malaria early warnings based on seasonal climate forecasts from multi-model ensembles, *Nature*. 439 (2006) 576.
- [25] Stone, R.C. G.L. Hammer and T. Marcussen, Prediction of global rainfall probabilities using phases of the Southern Oscillation Index, *Nature*. 384 (1996) 252.
- [26] Ogallo, L., P. Bessemoulin, J.-P. Ceron, S. Mason and S.J. Connor, Adapting to climate variability and change: The Climate Outlook Forum process, *WMO Bulletin*, 57 (2008) 93.
- [27] Graham, R.J. C. Gordon, M.R. Huddleston, M. Davey, W. Norton, A. Colman, A.A. Scaife, A. Brookshaw, B. Ingleby, P. McLean, S. Cusack, E. McCallum, W. Elliot, K. Groves, D. Cotgrove and D. Robinson, The 2005/06 winter in Europe and the United Kingdom: Part 1 – How the Met Office forecast was produced and communicated, *Weather*. 61 (2006) 327.
- [28] Hagedorn, R., F.J. Doblus-Reyes and T.N. Palmer, The rationale behind the success of multi-model ensembles in seasonal forecasting – 1. Basic concept, *Tellus*. 57 (2005) 219.
- [29] Chidzambwa, S. and S.J. Mason, Report of the valuation of Regional Climate Outlook Forecasts for Africa during the Period 1997–2007, ACMAD Technical Report, Niamey, ACMAD, 2008.
- [30] Georgakakos, K.P. N.E. Graham, T.M. Carpenter, A.P. Georgakakos and H. Yao, Integrating climate-hydrology forecasts and multi-objective reservoir management for Northern California, *Eos Transactions American Geophysical Union*. 86 (2005) 122.
- [31] Georgakakos, A.P. Decision-support systems for integrated water resources management with an application to the Nile basin, in: *Topics on System Analysis and Integrated Water Resources Management* (A. Castelletti and R. Soncini-Sessa, (eds), Amsterdam, Elsevier, 2006.
- [32] Graham, N.E., K.P. Georgakakos, C. Vargas and M. Echevers, Simulating the Value of El Niño Forecasts for the Panama Canal, *Advances in Water Research*. 29 (2007) 1667.
- [33] Brown, C. E. Conrad, A. Sankarasubramanian, S. Someshwar and D. Elazegui, The use of seasonal climate forecasts within a shared reservoir system: The case of Angat Reservoir, the Philippines, in: *Climate Change Adaptation in the Water Sector* (F. Ludwig, P. Kabat, H. van Shaik and M. van der Valk, (eds), London, Earthscan, 2009.
- [34] Julie, A. and J.P. Céron, Seasonal forecasting in West Africa: A strategic partnership for a sustainable development of a cross boundary river catchment, in: *Elements for Life* (Journal. Griffiths, eds.), London, Tudor Rose and WMO, 2007.
- [35] Graham, R.J. Personal communication. (2009).

- [37] Vitart, F. Seasonal forecasting of tropical storm frequency using a multi-model ensemble, *Quarterly Journal of Royal Meteorological Society*. 132 (2006) 647.
- [38] Vitart, F. M.R. Huddleston, M. Déqué, D. Peake, T.N. Palmer, T.N. Stockdale, M.K. Davey, S. Ineson and A. Weisheimer, Dynamically-Based Seasonal forecasts of Atlantic Tropical -Storm Activity issued in June by EUROSIP, *Geophysical Research Letters*. 34 (2007) L16815.
- [39] Miles, E.L. A.K. Snover, L.C. Whiteley Binder, E.S. Sarachik, P.W. Mote and N.J. Mantua, An approach to designing a national climate service. *Proceedings of the National Academy of Sciences*. 103 (2006) 19616.
- [40] Gong, X.A.G. Barnston and M.N. Ward, The effect of spatial aggregation on the skill of seasonal rainfall forecasts, *Journal of Climate*. 16 (2003) 3059.
- [41] Robertson, A.W.V. Moron and Y. Swarinoto, Seasonal predictability of daily rainfall statistics over Indramayu district, Indonesia, *International Journal of Climatology*. 29 (2009) 1449.
- [42] Lo, F. M.C. Wheeler, H. Meinke and A. Donald, Probabilistic forecasts of the onset of the north Australian wet season, *Monthly Weather Review*. 135 (2007) 3506.
- [43] Wheeler, M.C. H.H. Hendon, S. Cleland, H. Meinke and A. Donald, Impacts of the Madden-Julian Oscillation on Australian rainfall and circulation, *Journal of Climate*. 22 (2009) 1482.
- [44] Sun, L. D.F. Moncunill, H. Li, A.D. Moura, F.A.S. Filho and S.E. Zebiak, An operational dynamical downscaling prediction system for Nordeste Brazil and the 2002-04 real-time forecast validation, *Journal of Climate*. 19 (2007) 1990.
- [45] Sun, L. H. Li, M.N. Ward, and D. Moncunill: Climate variability and corn yields in semi-arid Ceará, Brazil, *Journal of Applied Meteorology and Climatology*. 46 (2007) 226.
- [46] Robertson, A.W. U. Lall, S.E. Zebiak and L. Goddard, Optimal combination of multiple atmospheric GCM ensembles for seasonal prediction, *Monthly Weather Review*. 132 (2004) 2732.
- [47] Stephenson, D.B. C.A.S. Coelho, F.J. Doblas-Reyes and M. Balmaseda, Forecast assimilation: A unified framework for the combination of multi-model weather and climate predictions, *Tellus*. 57 (2005) 253.
- [48] Wilks, D.S. *Statistical Methods in the Atmospheric Sciences*, New York, Academic Press, 1995.
- [49] Roncoli, C. K. Ingram and P. Kirshen, Reading the rains: Local knowledge and rainfall forecasting in Burkina Faso, *Society and Nature Research*. 15 (2002) 409.
- [50] Orlove, B.S. J.H.C. Chiang and M.A. Cane, Forecasting Andean rainfall and crop yield from the influence of El Niño of Pleiades visibility, *Nature*. 403 (2000) 68.
- [51] Hartmann, H.C. T.C. Pagano, S. Sorooshian and R. Bales, Confidence builders: Evaluating seasonal climate forecasts from user perspectives, *Bulletin of American Meteorological Society*. 83 (2002) 683.
- [52] Meinke, H. R.C. Stone and G.L. Hammer, Using SOI phases to forecast climatic risk to peanut production: A case study for northern Australia, *International Journal of Climatology*. 16 (1996) 783.
- [53] Twomlow, S. F.T. Mugabe, M. Mwale, R. Delve, D. Nanja, P. Carberry and M. Howden, Building adaptive capacity to cope with increasing vulnerability due to climatic change in Africa - A new approach, *Physics and Chemistry of the Earth*. 33 (2008) 780.
- [54] Meinke, H. and R.C. Stone, Seasonal and inter-annual climate forecasting: The new tool for increasing preparedness to climate variability and change in agricultural planning and operations, *Climatic Change*. 70 (2005) 221.
- [55] Rodriguez, D., P. DeVoi and H. Cox, Helping practical wisdom: A niche for whole farm systems modelling, in: *Proceedings of the First Farming Systems Design Conference*, Catania, Italy, 2007.
- [56] Hammer, G. M. Cooper, F. Tardieu, S. Welch, B. Walch, F. van Eeuwijk F., S. Chapman and D. Podlich, Models for navigating biological complexity in breeding improved crop plants, *Trends in Plant Science*. 11 (2006) 587.
- [57] Yin, X. and P.C. Struik, Applying modelling experiences from the past to shape crop systems biology: The need to converge crop physiology and functional genomics, *New Phytologist*. 179 (2008) 629.
- [58] Kokic, P. R. Nelson, H. Meinke, A. Potgieter and J. Carter, From rainfall to farm incomes – transforming advice for managing climate risk in Australia. Part I: Development and testing of a bioeconomic modelling system, *Australian Journal of Agricultural Research*. 58 (2007) 993.
- [59] Nelson, R. P. Kokic and H. Meinke, From rainfall to farm incomes – transforming policy advice for managing climate risk in Australia, Part II: Forecasting farm incomes, *Australian Journal of Agricultural Research*. 58 (2007) 1004.

- [60] van Ittersum, M.K. J. Wolf and H.H. van Laar, eds, Proceedings of the Conference on Integrated Assessment of Agriculture and Sustainable Development: Setting the Agenda for Science and Policy (AgSAP 2009). Egmond aan Zee, The Netherlands, 10–12 March 2009. Wageningen, Wageningen University and Research Centre, 2009.
- [61] Barrett, C.B. B.J. Barnett, M.R. Carter, S. Chantarat, J.W. Hansen, A.G. Mude, D.E. Osgood, J.R. Skees, C.G. Turvey and M.N. Ward, Poverty Traps and Climate Risk: Limitations and Opportunities of Index-Based Risk Financing, IRI Tech. Rep. No. 07-03. Palisades, International Research Institute for Climate and Society, 2007.
- [62] Hertzler, G. Adapting to climate change and managing climate risks by using real options, *Australian Journal of Agricultural Research*. 58 (2007) 985.
- [63] Hansen, J.W. M.V.K. Sivakumar and B.C. Bates, eds, Advances in applying climate prediction to agriculture, *CR Special 16 Climate Research*. 33 (2006) 1.
- [64] Maia, A.H.M. and H. Meinke, Quantifying climate-related risks and uncertainties using Cox regression models. *CLIVAR Exchanges*. 47 (2008) 23. <http://www.clivar.org/publications/exchanges/exchanges.php>.
- [65] World Meteorological Organization (WMO), Proceedings of the meeting of the CCI OPAG3 expert team on verification, 2005. http://www.wmo.int/pages/prog/wcp/wcasp/meetings/documents/ETVerification_FinalReport_February2005.pdf.
- [66] Adger, W.N. Scales of governance and environmental justice for adaptation and mitigation of climate change, *Journal of International Development*. 13 (2001) 921.
- [67] Adger, W.N. N.W. Arnella and E.L. Tompkins, Successful adaptation to climate change across scales, *Global Environmental Change*. 15 (2005) 77.
- [68] Glantz, M.H. *Climate Affairs: A Primer*, Washington, Island Press, 2003.
- [69] Glantz, M.H. (ed.), *Usable science 9: El Niño early warning for sustainable development in Pacific Rim countries and islands*. Report of workshop held 13–16 September 2004 in the Galapagos Islands, Ecuador. Boulder, ISSE/NCAR, 2005.
- [70] Hammer, G.L.N. Nicholls and C. Mitchell, eds, *Applications of Seasonal Climate Forecasting in Agricultural and Natural Ecosystems – The Australian Experience*, Dordrecht, Kluwer Academic Publishers, 2000.
- [71] Meinke, H. R. Nelson, P. Kokic, R. Stone, R. Selvaraju and W. Baethgen, Actionable climate knowledge – from analysis to synthesis, *Climate Research*. 33 (2006) 101.
- [72] Cash, D.W. W.C. Clark, F. Alcock, N.M. Dickson, N. Eckley, D.H. Guston, J. Jäger and R.B. Mitchell R.B., Knowledge systems for sustainable development, *Proceedings of the National Academy of Sciences*. 100 (2003) 8086.
- [73] Cash, D.W. and J. Buizer, Knowledge–action systems for seasonal to interannual climate forecasting: Summary of a workshop, Roundtable on Science and Technology for Sustainability, Policy and Global Affairs, Washington, The National Academies Press, 2005. <http://books.nap.edu/catalog/11204.html>.
- [74] Schwartz, B. and K. Sharpe, Practical wisdom: Aristotle meets positive psychology, *Journal of Happiness Studies*. 7 (2006) 377.
- [75] Pulwarty, R.S. and T.S. Melis, Climate extremes and adaptive management on the Colorado River – Lessons from the 1997–98 ENSO event, *Journal of Environmental Management*. 63 (2001) 307.
- [76] Corringham, T.W. A.L. Westerling and B.J. Morehouse, Exploring use of climate information in wildland fire management: A decision calendar study, *Journal of Forestry*. 106 (2008) 71.
- [77] Howden, S.M. J.-F. Soussana, F.N. Tubiello, N. Chhetri, M. Dunlop and H. Meinke, Adapting agriculture to climate change, *Proceedings of the National Academy of Sciences*. 104 (2007) 19691.
- [78] Arakawa, A. Future development of general circulation models, in: *General Circulation Model Development* (D. Randall, ed.), San Diego, Academic Press, 2000.
- [79] Goddard, L. and M.P. Hoerling, Practices for seasonal-to-interannual climate predictions, *US CLIVAR Variations*. 4 (2006) 1.
- [80] Barnston, A.G. H.M. van den Dool, D.R. Rodenhuis, C.R. Ropelewski, V.E. Kousky, E.A. O'Lenic, R.E. Livezey, S.E. Zebiak, M.A. Cane, T.P. Barnett, N.E. Graham, M. Ji and A. Leetmaa, Long-lead seasonal forecasts—Where do we stand?. *Bulletin of American Meteorological Society*. 75 (1994) 2097.
- [81] Livezey, R.E. and M.M. Timofeyeva, The first decade of long-lead U.S. seasonal forecasts: Insights from a skill analysis, *Bulletin of American Meteorological Society*. 89 (2008) 843.
- [82] Mason, S.J. L. Goddard, N.E. Graham, E. Yulaeva, L. Sun and P.A. Arkin, The IRI seasonal climate prediction system and the 1997/98 El Niño event. *Bulletin of American Meteorological Society*. 80 (1999) 1853.

- [83] Smith, D.M. S. Cusack, A.W. Colman, C.K. Folland, G.R. Harris and J.M. Murphy, Improved surface temperature prediction from the coming decade from a global climate model, *Science*. 317 (2007) 796.
- [84] Martinez, R. Position Paper on Users Liaison, An International expert review meeting on Regional Climate Outlook Forums, WMO, 2008.
- [85] Power, S. N. Plummer and P. Alford, Making climate model forecast more useful, *Australain Journal of Agricultural Research*. 58 (2007) 945.
- [86] Hansen, J. W. Baethgen, D. Osgood, P. Ceccato and R. Kinuthia, Innovations in climate risk management: Protecting and building rural livelihoods in a variable and changing climate, *Journal of Semi-Arid Tropical Agricultural Research*. 4 (2007) 1. (published online at <http://www.icrisat.org/Journal/specialproject.htm>).
- [87] Brown, C. and J.W. Hansen, Agricultural Water Management and Climate Risk. Report to the Bill and Melinda Gates Foundation, IRI Tech. Rep. No. 08-01. Palisades, International Research Institute for Climate and Society, 2008.
- [88] Lemos, M.C. T.J. Finan, R.W. Fox, D.R. Nelson and J. Tucker, The use of seasonal climate forecasting in policymaking: Lessons from Northeast Brazil, *Climatic Change*. 55 (2002) 479.
- [89] Ziervogel, G. Targeting seasonal climate forecasts for integration into household level decisions: The case of smallholder farmers in Lesotho, *Geographical Journal*. 170 (2004) 6.
- [90] Meinke, H. W.E. Baethgen, P.S. Carberry, M. Donatelli, G.L. Hammer, R. Selvaraju and C.O. Stöckle, Increasing profits and reducing risks in crop production using participatory systems simulation approaches, *Agricultural Systems*. 70 (2001) 493.