

ORAL PRESENTATION

Open Access

# Sugar-fermenting yeast as an organic source of carbon dioxide to attract the malaria mosquito *Anopheles gambiae* s.s.

Renate C Smallegange<sup>1\*</sup>, Wolfgang H Schmied<sup>1</sup>, Karel J van Roey<sup>1</sup>, Niels O Verhulst<sup>1</sup>, Jeroen Spitzen<sup>1</sup>, Wolfgang R Mukabana<sup>2,3</sup>, Willem Takken<sup>1</sup>

*From Parasite to Prevention: Advances in the understanding of malaria*  
Edinburgh, UK. 20-22 October 2010

## Background

Carbon dioxide (CO<sub>2</sub>) plays an important role in the host-seeking process of opportunistic, zoophilic and anthropophilic mosquito species and is therefore commonly added to mosquito sampling tools. The African malaria vector *Anopheles gambiae* Giles *sensu stricto* is attracted to human volatiles augmented by CO<sub>2</sub>. We investigated whether CO<sub>2</sub>, usually supplied from gas cylinders acquired from commercial industry, could be replaced by CO<sub>2</sub> derived from fermenting yeast (yeast-produced CO<sub>2</sub>).

## Methods

Trapping experiments were conducted in the laboratory, semi-field and field, with *An. gambiae* s.s. as the target species. MM-X traps were baited with volatiles produced by yeast-sugar solutions, prepared in bottles. Catches were compared with traps baited with industrial CO<sub>2</sub>. The additional effect of human odours was also examined.

## Results

Traps baited with yeast-produced CO<sub>2</sub> caught significantly more mosquitoes than unbaited traps and also significantly more than traps baited with industrial CO<sub>2</sub>, both in the laboratory and semi-field. Adding yeast-produced CO<sub>2</sub> to traps baited with human odour significantly increased trap catches. During the field trials, traps baited with yeast-produced CO<sub>2</sub> caught similar numbers of *An. arabiensis* Patton as traps baited with

industrial CO<sub>2</sub>. Addition of human odour increased trap catches.

## Conclusions

We conclude that yeast-produced CO<sub>2</sub> can effectively replace industrial CO<sub>2</sub> for sampling of *An. gambiae* s.s.. This will significantly reduce costs and allow sustainable mass application of odour-baited devices for mosquito sampling in remote areas.

## Acknowledgements

This study was funded by a grant from the Foundation for the National Institutes of Health (NIH) through the Grand Challenges in Global Health Initiative (GCGH#121).

## Author details

<sup>1</sup>Laboratory of Entomology, Wageningen University, P.O. Box 8031, 6700 EH, Wageningen, The Netherlands. <sup>2</sup>International Centre of Insect Physiology and Ecology, P.O.Box 30772 - 00100, GPO, Nairobi, Kenya. <sup>3</sup>School of Biological Sciences, University of Nairobi, P.O. Box 30197 - 00100 GPO, Nairobi, Kenya.

Published: 20 October 2010

doi:10.1186/1475-2875-9-S2-O29

Cite this article as: Smallegange et al.: Sugar-fermenting yeast as an organic source of carbon dioxide to attract the malaria mosquito *Anopheles gambiae* s.s. *Malaria Journal* 2010 **9**(Suppl 2):O29.

<sup>1</sup>Laboratory of Entomology, Wageningen University, P.O. Box 8031, 6700 EH, Wageningen, The Netherlands

Full list of author information is available at the end of the article