

The invasion of an introduced predator, Nile perch (*Lates niloticus*, L.) in Lake Victoria (East Africa): chronology and causes

Kees (P.C.) Goudswaard · Frans Witte ·
Egid F. B. Katunzi

Received: 15 May 2006 / Accepted: 17 November 2006 / Published online: 23 December 2006
© Springer Science+Business Media B.V. 2006

Abstract Nile perch, a large predatory fish, was introduced into Lake Victoria in 1954. The upsurge of Nile perch in Lake Victoria was first observed in the Nyanza Gulf, Kenya, in 1979. In Ugandan waters this occurred 2–3 years later and in the Tanzanian Mwanza Gulf 4–5 years later. At the beginning of the upsurge in the Mwanza Gulf in 1983/1984 only sub-adult and adult fishes were found. The first juveniles appeared in 1985, suggesting that the initial increase of Nile perch was mainly caused by migration of sub-adults and adults. Shortly after the onset of trawl fishery in the area in 1973, haplochromines in the Mwanza Gulf started to decline. The final disappearance of the haplochromines, in 1987, only occurred after the Nile perch boom, and despite the abandoning of the haplochromine fishery in 1986. We hypothesize that the decline of haplochromines

decreased predation on and competition with juvenile Nile perch and then facilitated survival of these juveniles. Consequently the immigration of sub-adult and adult Nile perch in an area may have paved the way for successful recruitment. Over-exploitation of haplochromine cichlids in the 1970s in the Nyanza Gulf, where the Nile perch upsurge was first observed, may have played a similar role.

Keywords Species-introduction · Colonization · Extinction · Predation · *Caridina nilotica* · Haplochromine cichlids

Introduction

Deliberate introductions of animals to areas outside their natural distribution, has taken place for centuries (Williamson 1996). However, within Africa, the transfer of fish species from one water system into another is a relatively recent phenomenon. Most species were transferred between the 1920s and 1970s with a peak (67 species) in the 1950s (Welcomme 1988). Worldwide, at least 237 fish species were transferred, of which the fate of 72 has been documented. Of these, 25 were harmful for the indigenous fauna and/or the local environment (Beverton 1992). The introduction of Nile perch, *Lates niloticus* L, into Lake Victoria is considered to belong to this last group

Kees (P.C.) Goudswaard · F. Witte (✉)
Institute of Biology, University of Leiden,
P.O. Box 9516, 2300 RA Leiden, The Netherlands
e-mail: F.Witte@Biology.leidenuniv.nl

Present Address:
Kees (P.C.) Goudswaard
Institute for Marine Resources and Ecosystem
Studies, Wageningen IMARES, P.O. Box 77, 4400
AB Yerseke, The Netherlands

E. F. B. Katunzi
Tanzania Fisheries Research Institute (TAFIRI),
P.O. Box 475, Mwanza, Tanzania

(Barel et al. 1985; Ogutu-Ohwayo 1990a; Kaufman 1992; Witte et al. 1992a, b).

Nile perch were unofficially introduced into Lake Victoria in August 1954, when individuals from Lake Albert were released at Jinja, Uganda (Amaras 1986). In May 1960, local fishermen reported the first Nile perch from gillnet catches near Jinja. Seven others were reported in the same year. The captured fishes were sub-adults with a size range of 28–43 cm (Hamblyn 1961).

After heavy debates (Fryer 1960; Anderson 1961; Pringle 2005), the first official introductions of Nile perch took place in May 1962 and September 1963 at Entebbe, Uganda. They originated from Lake Albert and comprised, respectively, 35 fishes between 16 and 43 cm, and 339 fingerlings (Gee 1964). In the Nyanza Gulf, Kenya, eight Nile perch from Lake Turkana were released in 1963 (Arunga 1981; Pringle 2005). During a bottom trawl survey in 1969/1970 Nile perch was caught lake wide in low densities (Kudhongania and Cordone 1974b). In the 1980s the species suddenly increased strongly, but not simultaneously, throughout the lake.

The effects of the Nile perch boom on the indigenous fauna of Lake Victoria, has been discussed in several publications (Barel et al. 1985; Hughes 1986; Acere 1988; Harrisson et al. 1989; Ogutu-Ohwayo 1990a; Kudhongania et al. 1992; Witte et al. 1992a, b; Goldschmidt et al. 1993; Bundy and Pitcher 1995; Kudhongania and Chitamwebwa 1995; Goudswaard and Witte 1997; Goudswaard et al. 2002a, b). Remarkably, the process by which Nile perch established itself successfully in Lake Victoria is an underexposed aspect in the whole discussion. In this paper we present data on Nile perch densities, length frequency distributions, the appearance of recruits, and the decline of the haplochromines during successive periods at different localities in Lake Victoria. Although the data sets from several areas are incomplete, to our knowledge they are the best available. With the aid of these data, we make an attempt to elucidate the process of colonization of Lake Victoria by Nile perch.

Material and techniques

In the Tanzanian waters of Lake Victoria fishing was done with MV Kiboko (105 hp, bottom trawl


head rope 18 m, codend mesh 20 mm), which made trawl shots of half an hour. The sampling period ran from October 1984 until July 1990. Most frequently fished was the Mwanza Gulf, where five stations with a soft mud bottom and a depth range from 1.5 m in the south to 35 m in the north were sampled monthly (Fig. 1). The sixth station at a depth of 57 m (Fig. 1) was fished until the end of 1986. Quarterly samples were taken at: (i) Kagera—an open water area in the western part of the lake with a mud/clay bottom. Samples were collected at depths ranging from 4 m in the west to 48 m in the east. The shallow areas (<10 m) were restricted to Chigoga and Kemono Bay as well as the area near Bukoba harbour. (ii) Speke Gulf—a large bay in the south-east corner of the lake with a soft muddy bottom and a depth ranging from 2 m in the east to 34 m in the west. (iii) The Emin-Pasha Gulf in the southwest corner of the lake was surveyed in 1985 and 1986. The area is up to 18 m deep and has a mud and sandy bottom. Directly north of this gulf the lake floor deepens to 48 m.

Information was also gathered from other vessels like: MV Mdiria (170 hp, trawl head rope 25 m, codend mesh 20 mm and 90 mm) fishing in the Mwanza Gulf and incidentally elsewhere; four boats of the Nyanza Fishing and Processing Company (NFPC) (170 hp, trawl head rope 21 m, codend mesh 90 mm) fishing in the Emin Pasha Gulf from November 1975 to August 1976; three trawlers (75 hp, head rope 17 m, codend mesh 70 mm) from a commercial company operating in the Speke Gulf. These boats started fishing in 1986 and records were available up to June 1990. For commercial reasons Nile perch was separated by the crew in small (<35 cm TL) and big (>35 cm) fishes. Data in this paper from other ships than MV Kiboko are based on catch records of the crews.

Catch records of Nile perch and haplochromines were standardized to kg h^{-1} for all vessels. Trawl data from Ugandan and Kenyan waters were obtained from literature (Table 1). The total length (TL) of Nile perch was measured and rounded to the nearest cm below.

Catch data were tested for normal distribution with a Kolmogorov-Smirnov Test. Most Nile perch catches were not normally distributed. Therefore, we used Mann-Whitney *U*-tests to test for differences in catch sizes between years

Fig. 1 Map of Lake Victoria with main fishing areas discussed in the present study. Hatched areas in the Mwanza Gulf were fished by MV Mdiria. Asterisks in the Mwanza Gulf were sampling stations of MV Kiboko. The hatched area in the Speke Gulf was fished by MV Mdiria and MV Kiboko. The double hatched area in the Speke Gulf was also fished by commercial trawlers. The hatched areas in the Emin Pasha Gulf and Kagera area were fished by MV Kiboko and before 1984 occasionally by MV Mdiria and trawlers of the NFPC (for details see “Material and techniques” section)



and for differences in catch sizes of different size groups. Sequential Bonferroni corrections were applied in case of repeated use of data. A correlation between mean monthly catch rates of Nile perch by commercial trawlers in the Speke Gulf and months was calculated. As mean monthly catch rates were normally distributed, the Pearson correlation test was used. All tests were done in SPSS10.0 for Windows.

Results

Nile perch appearance in the Tanzanian waters

In the early 1970s, Nile perch catches in the Mwanza Gulf were low (Fig. 2a, b). A slow

increase was observed in the second half of the 1970s and the beginning of the 1980s. In 1983 a strong increase was noticed (Fig. 2a, b).

Most Nile perch caught in the Mwanza Gulf between 1977 and 1983 were adults or sub-adults (personal observation). The same was observed in bottom trawl catches from 1982/1983 (Fig. 3a) and from the last quarter of 1984 (Fig. 3b). In 1987 small fishes (1–20 cm) were abundant in the catches (Fig. 3c). Detailed observations on the increase of small Nile perch were collected at the station of 12–16 m deep in the Mwanza Gulf. In 1984 there were less small (1–10 cm TL) than medium and large sized Nile perch (21–50 cm and >50 cm; Mann–Whitney *U*-test, $P \leq 0.002$ in each case). The number of small sized Nile perch started to increase in 1985. In

Table 1 Mean catch rates (kg h⁻¹) of Nile perch and haplochromines in bottom trawl catches made in the Ugandan waters, the Nyanza Gulf (Kenya) and in four different areas in Tanzania

Year	Uganda		Kenya		Tanzania							
			Nyanza Gulf		Kagera		Speke Gulf		Mwanza Gulf		Emin Pasha Gulf	
	Hap.	Np.	Hap.	Np.	Hap.	Np.	Hap.	Np.	Hap.	Np.	Hap.	Np.
1969	327	+	440	1	610	+	-	-	548	0	-	-
1973	-	-	-	-	-	-	-	-	1235	0	-	-
1974	-	-	149	6	-	-	1294	0	1376	0	-	-
1975	-	-	-	-	308	0	1025	0	1365	1	-	-
1976	-	-	-	-	-	-	-	-	1915	1	-	-
1977	-	-	-	-	490	0	-	-	1165	1	540	+
1978	-	-	-	-	-	-	-	-	1095	1	-	-
1979	-	-	59	*46	-	-	1005	0	1182	1	-	-
1980	-	-	-	-	-	-	-	-	709	4	-	-
1981	392	4	0	169	-	-	-	-	789	4	-	-
1982	298	45	0	169	-	-	-	-	682	4	-	-
1983	265	54	-	-	-	-	288	101	629	19	-	-
1984	114	140	-	-	-	-	-	-	290	82	-	-
1985	18	235	-	-	38	*146	-	-	326	*103	108	*71
1986	0	80	9	273	0	259	-	-	93	174	0	76
1987	-	-	-	-	0	244	-	-	4	203	-	-
1988	-	-	0	145	0	199	0	*224	0	115	-	-
1989	-	-	0	-	0	118	0	119	0	107	-	-
1990	-	-	-	-	-	-	0	163	1	133	-	-

Nile perch catches for the first time reaching values ≥ 45 kg h⁻¹ (an arbitrarily chosen density) are given in bold. *indicates the first year that juvenile Nile perch ≤ 10 cm were observed; +indicates presence of low densities of Nile perch, without precise data for the records used. All 1969 data are from Kudhongania and Cordone (1974a, b). The Nyanza Gulf data are from Benda (1981), Okemwa (1981), Fisheries Department Nairobi Kenya (1988) and Ogari and Asila (1990). Data from Ugandan waters are from Acere (1988). Tanzanian data from 1973 till 1983 are from MV Mdiria and from 1984–1990 from MV Kiboko. Hauls made in 1969 and in Ugandan waters were made with trawlers ranging from 105–180 hp, utilizing conventional bottom trawls with head ropes of 18–25 m and 19 and 38 mm codend mesh. In the Nyanza Gulf, part of the catches was made with a 85 hp trawler, towing a net with a 13.7 m head rope. In this area mainly a 38 mm codend mesh size was used. Hap., Haplochromines, Np., *Lates niloticus*


1986 and 1987 densities of 1–10 cm Nile perches were significantly higher than in each of the previous years (Fig. 4a; Mann–Whitney *U*-test, $P = 0.000$ in each case). In the following years the densities fluctuated around the same level. From 1987 onwards the densities of 1–10 cm class Nile perch were significantly higher than the densities of all other classes (Mann–Whitney *U*-test, P values ranging from 0.000 to 0.027), with the exception of 1990, when the densities of the 21–50 and the 1–10 cm class did not differ significantly ($P = 0.055$).

In the Emin Pasha Gulf Nile perch catches showed a similar pattern as in the Mwanza Gulf (Fig. 4b). In 1973 and 1974 MV Mdiria, with a 60 mm codend mesh, for 74 h caught no Nile perch. Between November 1975 and August 1976, bottom trawlers of NFPC caught 6 large Nile

perch with a total weight of 172 kg in 850 h. In 1977 MV Mdiria caught 28 kg Nile perch (possibly one individual) in 12 h trawling (2.3 kg h⁻¹), using either a 20 or a 90 mm codend mesh. In June 1985 MV Kiboko caught 71 kg Nile perch h⁻¹ in the Emin Pasha Gulf (Table 1), with relatively few juveniles (1–10 cm TL, Fig. 4b). The densities of these juveniles did not differ significantly from those of other length classes. In August 1986 the numbers of juveniles had increased (Mann–Whitney *U*-test, $P = 0.000$) and were higher than those of large sized Nile perch (Mann–Whitney *U*-test, $P = 0.000$).

The Kagera area in 1985 and 1986 revealed similar densities of juvenile Nile perch as found in the Mwanza Gulf in the same years (Fig. 4a, c). The same held for the densities in the Emin Pasha Gulf in October 1985 and August 1986 (Fig. 4b).

Fig. 2 Catch rates of Nile perch and haplochromines by bottom trawlers in the Mwanza Gulf. **(a)** 90 mm codend mesh during the period 1973 till 1987 (3875 h trawling); **(b)** 20 mm codend mesh during the period 1973–1990. The data from 1984 till 1990 (543 h trawling) for the 20 mm net are from MV Kiboko, all other data from MV Mdiria (2176 h trawling)



The average monthly catches of large Nile perch in commercial trawlers in the Speke Gulf, decreased over time (Fig. 5a, Pearson correlation test, $R = -0.513$, $P = 0.000$, $N = 58$), while the catches of small Nile perch increased (Fig. 5b, Pearson correlation test, $R = 0.812$, $P = 0.000$, $N = 58$).

Nile perch upsurge in Kenyan and Ugandan waters of Lake Victoria


In the Kenyan and Ugandan waters, a strong increase of Nile perch occurred earlier than in the Mwanza Gulf (Table 1). A Nile perch catch rate of 46 kg h⁻¹ was observed in the Nyanza Gulf in 1979. Values ≥ 45 kg h⁻¹ were first observed in the Ugandan waters in 1982 and in the Mwanza Gulf in 1984. Catches above 100 kg h⁻¹ were reached

1 or 2 years later at each of these stations. Juveniles (<10 cm) were present in the Nyanza Gulf in 1979, while they appeared in 1985 for the first time in the Mwanza Gulf.

Nile perch and haplochromines

The catches of haplochromine cichlids in the Mwanza Gulf declined from the onset of their exploitation with bottom trawlers in 1973 (Fig. 2a, b). Large haplochromines (mainly piscivores > 17 cm SL) declined much faster than the smaller ones and by 1980 they had disappeared from the catches (Fig. 2a). The small species also declined, but at a much slower rate (Fig. 2b). Commercial exploitation of haplochromines ended in 1986. The trawl survey by MV Kiboko at five different depth zones in the Mwanza Gulf

Fig. 3 Length frequency distribution of Nile perch in the Mwanza Gulf. **(a)** October 1982–April 1983, ($N = 67$); **(b)** October–December 1984, ($N = 434$); **(c)** January–December 1987 ($N = 277,487$). Catches from 1982–1983 were made with NFPC trawlers, those from 1984 and 1987 with MV Kiboko. Note log scale of the Y-axes



showed an increase in Nile perch catches and a concomitant decline of the haplochromines (Table 1, Fig. 6). This observation was also noted at other places in the lake (Table 1, Fig. 6). Between November 1975 and August 1976 NFPC trawlers with 90 mm codend mesh fished in the Emin Pasha Gulf for 850 h and caught on average $21.9 \pm 7.9 \text{ kg h}^{-1}$ large-sized haplochromines. In 1977 MV Mdiria made two hauls with 20 mm codend mesh, which contained an average catch of haplochromines of 540 kg h^{-1} (Table 1). Nile perch were also caught in this period (see above). In June 1985, both haplochromines and Nile perch were present in bottom trawl catches with MV Kiboko (Table 1, Fig. 6). Large haplochromines were rare. In October 1985 the stock of

haplochromines in the 11–20 m deep zone had vanished (Fig. 6), while in the shallow water (1–10 m deep) only a fraction of the original stock was still present. In August 1986 catches of haplochromines had been reduced to a few individuals in both depth zones, resulting in weights that are not visible in Fig. 6.

In the Kagera area, we found haplochromines in all depth zones in the second quarter of 1985. After 1985 at Chigoga- and Kemonondo Bay as well as near Bukoba harbour (all <10 m deep), we collected very small quantities ($<<1 \text{ kg}$) of haplochromines in 3.5 h trawling. In the 11–20 m and 21–48 m zones that were fished 16.5 and 299 h, respectively, hardly any haplochromines were caught.




Fig. 4 Development of the number of Nile perches over time, in different size classes. **(a)** Mwanza Gulf (12–16 m deep) 1984–1990; **(b)** Emin Pasha Gulf (4–18 m deep)

June 1985–August 1986; **(c)** Kagera area (6–48 m deep) 1985–1989. *Note* log scale of the Y-axes. Numbers above X-axes indicate the number of trawl shots

Discussion

The colonization process

Around 400 Nile perch from Lake Albert (339 fingerlings and 35 fishes of 16–43 cm) were introduced into Lake Victoria (Gee 1964; Amaras 1986; Pringle 2005). Additionally, eight fishes of unknown size were introduced from Lake Turkana in 1963 (Gee 1964; Arunga 1981; Pringle 2005). Apart from *Lates niloticus*, each of these lakes harbours a deep-water *Lates* species, *L. macrophthalmus* Worthington 1929 and *L. longispinis* Worthington 1932, respectively. Allozyme data indicate that Nile perch of Lake Victoria were mainly *Lates niloticus* from Lake Albert, although there were indications for some genetic contribution of *L. macrophthalmus* (Hauser et al. 1998).


Although Nile perch was introduced in 1954, it was first reported from a catch of a fisherman in Uganda in May 1960 (Gee 1964). Several others were caught later that year. The eight fishes recorded in 1960 from the northeast corner of the lake, were 28–43 cm long (Hamblyn 1961). They

were probably offspring of those introduced in 1954, as in 6 years time the introduced individuals would have attained a much greater length (e.g., Hughes 1992). According to Gee (1964) a breeding population existed in the lake in 1962/1963, probably centered in Hannington Bay and down the Buvumu Channel.

According to Pringle (2005), the first Nile perch reported from Tanzanian waters was caught near Mwanza in October 1961. Other reports suggest that Nile perch was landed for the first time from Tanzanian waters of the lake in August 1963 at Musoma (Kudhongania and Cordone 1974a). During a bottom trawl survey in 1969–1970 Nile perch was rarely caught in Tanzanian waters. Catches constituted approximately 0.01 kg h⁻¹, while the catches in Uganda and Kenya were 0.48 and 1.12 kg h⁻¹, respectively (Kudhongania and Cordone 1974a, b).

In the Nyanza Gulf the average catch rate of Nile perch was 46 kg h⁻¹ as early as 1979. These catches contained all stages from early juveniles to adults. In contrast, in the Mwanza Gulf Nile perch catches above 45 kg h⁻¹ were not observed

Fig. 5 Length frequency distribution of Nile perch caught by commercial trawlers in the Speke Gulf, based on 12 539 h of bottom trawling during the period 1986–1990. **(a)** large Nile perch > 35 cm TL; **(b)** small Nile perch < 35 cm TL




before 1984, and juveniles were first caught in 1985 (Table 1). Up to 1985, all Nile perch found in the Mwanza Gulf apparently had been born elsewhere. The length frequency distribution of Nile perch from trawl catches in 1983 and 1984 (Fig. 3) indicates that the strong increase in number of Nile perch in the Mwanza Gulf in 1983–1984 was caused by sub-adult fishes. It is most likely that these fishes arrived by immigration. In Ugandan waters and in the Speke Gulf, Nile perch catches around 45 kg h⁻¹ were observed 1–2 years earlier than in the Mwanza Gulf, and 3 years later than in the Nyanza Gulf (Table 1). These observations suggest that Nile perch spread from the Nyanza Gulf towards the North and the South. Zaret and Paine (1973) described the spread of the introduced predator *Cichla ocellaris* in Lake Gatun in Panama. They

note that ‘the fish have not diffused haphazardly throughout the lake, but have moved as a wave, with a leading edge composed of sub-adults’. Though there are many gaps, our data suggest a similar process for Nile perch in Lake Victoria. A similar migration of sub-adults of an introduced predator fish, *Parachanna obscura* (Steindachner, 1879), was noticed in the upper course of the Zambian Congo River. Between 1996 and 1999 a migration from a local nucleus at the Luapula River towards Lake Mweru was observed (PCG unpublished data).

Although allozyme electrophoresis provided some indication for genetic differentiation of Nile perch within Lake Victoria (Hauser et al. 1998), preliminary results of a tagging program in the Mwanza region revealed that Nile perch was able to cover 50 km per week (Ligtvoet and Mkumbo

Fig. 6 Average catch rates (kg h^{-1}) of Nile perch and haplochromines in bottom trawls according to depth zone in different years. **(a)** Mwanza Gulf; **(b)** Emin Pasha Gulf; **(c)** Kagera area. Numbers under X-axes indicate the number of trawl shots. Numbers after years indicate the quarters of the year that catches were made



1990). This is well within the range that Nile perch could cover based on a cruising speed fish of 1–2 body lengths s^{-1} (Videler 1993), and implies that Nile perch was able to cover the distance from Kenyan waters to the Mwanza Gulf in a few months. However, after the Nile perch became

abundant in the Nyanza Gulf in Kenya in 1979, it took 4 years before the sub-adult Nile perch “wave” reached the Mwanza Gulf. This might indicate that there was no need for these fishes to migrate in large numbers at that time. Nile perch boomed in the Ugandan and the Tanzanian waters

after haplochromines had been depleted in the Nyanza Gulf (Table 1). Consequently, it is possible that local resource depletion caused more fish to disperse away from their natal site.

Decline of haplochromines

Evidence has been given by Ogutu-Ohwayo (1990b) and by Witte et al. (1992a, b) that the decline and extinction of several hundreds of haplochromine species was due to predation by Nile perch. Other causes for the decline of the haplochromines were local over-exploitation (Witte and Goudswaard 1985; Acere 1988; Harrison et al 1989; Bundy and Pitcher 1995; Kudhongania and Chitamwebwa 1995) and eutrophication, leading to oxygen deficiencies (Hecky 1993; Hecky et al. 1994; Verschuren et al. 2002; but see Witte et al. 2005) and to increased turbidity (Seehausen et al. 1997).

Exploitation of a virgin stock generally leads to a stock reduction and changes in species composition. Larger species, which mature above the 50% retention size of the net, are generally more sensitive to over-exploitation than small sized species. In Lake Malawi commercial trawl fishery on haplochromine cichlids resulted in a change in dominance from medium and large sized species to small sized species (Turner et al. 1995). The same was observed in the Mwanza Gulf, where the large species had vanished in the catches with the 90 mm codend by 1979 (Fig. 2b). The catch per unit of effort (in kg h⁻¹ trawling) also declined for the small sized haplochromines. However, for the following reasons, the steep decline in the Mwanza Gulf, after 1984 cannot be attributed to fishing only: (1) After the trawl fishery in the Mwanza Gulf was stopped in 1986, the decline of the haplochromines continued (Table 1); (2) In areas where fishing pressure for haplochromines was low, e.g., Emin Pasha Gulf and Kagera area, they vanished as well (Table 1); (3) In all areas the disappearance of the haplochromines was concomitant with a strong increase of the Nile perch, thus suggesting a major impact of this predator (Table 1).

With respect to eutrophication, especially increased turbidity may have had a strong impact, as cichlids are highly dependent on vision for

feeding and reproduction (Seehausen et al. 1997, 2003; Seehausen and Van Alphen 1998; Witte et al. 2005). Because the Nile perch upsurge in Lake Victoria and the increase of eutrophication both occurred in about the same period (Verschuren et al. 2002), it is difficult to establish the relative impact of these phenomena separately. Nevertheless, as indicated above, predation by Nile perch was a major factor.

Large Nile perch as a pioneer for juvenile Nile perch?

It is remarkable that after the first release of Nile perch in Lake Victoria it took more than 25 years before a sudden increase of its population took place, resulting in its dominance in the demersal fish stock. The initial abundance of haplochromine cichlids suggests that food cannot have been a limiting factor for the piscivorous Nile perch between its introduction and the early 1980s. Therefore, we assume that other factors have played a role. An important factor could have been the survival of very young Nile perch and the food abundance for these fishes. Although we have no evidence from stomach contents, it is likely that the abundant haplochromines fed upon eggs and larvae of Nile perch, whenever available. Moreover, the haplochromines were probably competing with juvenile Nile perch for zooplankton and insect larvae. During the colonization process the adult and sub-adult Nile perch fed heavily upon haplochromine cichlids (Hughes 1986; Ogutu-Ohwayo 1990a, b; Ligtoet et al. 1990) and reduced their numbers to extremely low levels, thus enhancing the chance of survival of eggs and larvae of Nile perch. Further, it has been suggested that the reduction of haplochromine cichlids also reduced the predation on juvenile shrimps, *Caridina nilotica* (Roux, 1833) (Goudswaard et al. 2006). Apart from zooplankton and insect larvae, juvenile shrimps also appeared to be an important food item for small (<10 cm) Nile perch (Goudswaard et al. 2006; Katunzi et al. 2006). Both shrimps and insect larvae increased after the disappearance of the haplochromines (Kaufman 1992; Mbahinzireki 1992; Goldschmidt et al. 1993; Goudswaard et al. 2006), which were their predators. In this

situation adult Nile perch facilitated the food availability for juvenile Nile perch by eradicating the former competitors of their juveniles.

If the foregoing is correct, then Nile perch initially could have been successful in reproduction only at places where the haplochromine stock was reduced. Before the Nile perch upsurge this must have been an area where the reduction of haplochromines was caused by other factors than predation by Nile perch. The heavily exploited Nyanza Gulf in Kenya might have acted as such a place, as already in the early 1970s over-exploitation of haplochromines was reported for this area (Marten 1979).

Nile perch after the boom

After the disappearance of the haplochromines and the establishment of a large stock of shrimps, the Nile perch became a resident species of which successive year classes were present. Juvenile Nile perches became one of the main food sources of adult Nile perch >50 cm (Hughes 1986, 1992, Ogari and Dadzie 1988; Ogutu-Ohwayo 1990b; Mkumbo and Ligvoet 1992; Katunzi et al. 2006). In the Speke Gulf the amount of large Nile perch in trawl catches declined (Fig. 5a). As Nile perch fishery mainly aimed at large fishes, fishing may have caused this reduction. It has been observed in Kenyan waters that size at 50% maturity has declined from 102 cm for females and 74 cm for males before 1986 (Asila and Ogari 1988) to 55 cm for males and 70 cm for females around 1990 (Ogari and Asila 1992).

The explanation of the simultaneous increase of small (<35 cm) individuals is more complex. It might be the combined result of: (1) Growth of the recruits that were born in the area after the initial invasion of the sub-adults; (2) A relatively high predation pressure on juvenile Nile perch by adult Nile perch in the early years when the densities of large Nile perch were still high.

Conclusions

Several hypotheses for the Nile perch boom have been published. For instance, Hecky

(1993) suggested that the increasing turbidity and deoxygenation, caused by the eutrophication, broke down the complexly balanced food web in Lake Victoria, resulting in increased abundances of insect larvae and the shrimp, *Caridina nilotica*, that favored survival of juvenile Nile perch. Verschuren et al. (2002) suggested that deepwater oxygen loss caused the deepwater haplochromines to migrate to shallower depth ranges where they augmented the food base for Nile perch. However, he correctly adds that there is no evidence that characteristic deepwater haplochromines migrated to shallow areas, and that densities of haplochromines at depths of more than 30 m used to be significantly lower than in shallow habitats.

We hypothesize the following steps in the colonization process of Nile perch in Lake Victoria. (1) Initially, over-fishing of haplochromines in the Nyanza Gulf facilitated a local recruitment of juvenile Nile perch due to a decrease of predation by and competition with haplochromines. (2) Colonization of other areas in the lake took place at different moments. In each of these areas the first step in the process comprised the invasion of sub-adults from neighbouring areas, where haplochromines had been depleted. (3) Subsequently, the local haplochromine stock vanished, and both juvenile Nile perch and juvenile shrimps had a higher survival rate than before. (4) As a consequence a strong recruitment of Nile perch occurred in the area.

Thus, the pioneering sub-adult Nile perch created, by eradication of the haplochromines, a changed species composition in favour of juvenile Nile perch.

Acknowledgements Thanks go to the staff of the Tanzania Fisheries Research Institute (TAFIRI) and the Freshwater Fisheries Training Institute Nyegezi. The crew of MV Kiboko collected many data under difficult circumstances. J.J. Videler and C.D.N. Barel are thanked for their comments on the draft of this paper. The Haplochromis Ecology Surevy Team (HEST) was financially supported by the section for Research and Technology of the Netherlands Minister of Development Co-operation, by the Netherlands Foundation for the Advancement of Tropical Research (WOTRO) grants W87-129, W87-161, W87-189 and W84-488 and, by the Schure-Beijerinck-Popping Fund.

References

- Acere TO (1988) Recent trends in the fisheries of Lake Victoria (Uganda, Northern part). In: CIFA. Report of the 4th session of the sub-committee for the development and management of the fisheries of Lake Victoria. 6–10 April 1987. Kisumu. FAO fisheries report 338, pp 36–45
- Amaras JO (1986) Editorial letter in Newspaper “The Standard”, July 22, Nairobi, Kenya
- Anderson AM (1961) Further observations concerning the proposed introduction of Nile Perch in Lake Victoria. East Afr Agric Forest J 24:195–201
- Arunga JO (1981) A case study of the Lake Victoria Nile perch *Lates niloticus* (Mbuta) fishery In: Proceedings of the workshop of the Kenya Marine and Fisheries Research Institute on Aquatic resources of Kenya, 13–19 July 1981, Mombasa Kenya, pp 167–183
- Asila AA, Ogari J (1988) Growth parameters and mortality rates of Nile perch (*Lates niloticus*), estimated from length frequency data in the Nyanza Gulf (Lake Victoria). In: Venema SC, Christensen JM, Pauly D (eds) Contributions to tropical fisheries biology. FAO fisheries report 389, pp 272–287
- Barel CDN, Dorit R, Greenwood PH et al (1985) Destruction of fisheries in Africa’s lakes. Nature 315:19–20
- Benda RS (1981) A comparison of bottom trawl catches in the Kenya waters of Lake Victoria. J Fish Biol 18:609–613
- Beverton RJH (1992) Fish resources: threats and protection. Netherlands J Zool 42:39–175
- Bundy A, Pitcher TJ (1995) An analysis of species changes in Lake Victoria: did the Nile perch act alone? In: Pitcher TJ, Hart PJB (eds) The impact of species changes in African lakes. Chapman & Hall, London
- Fisheries Department Kenya (1988) State of Lake Victoria Fisheries, Kenya. In: CIFA. Report of the 4th session of the Sub-Committee for the development and management of the fisheries of Lake Victoria. FAO fisheries report 388, pp 20–28
- Fryer G (1960) Concerning the proposed introduction of Nile perch into Lake Victoria. East Afr Agric Forest J 25:267–270
- Gee JM (1964) Nile perch investigation. EAFFRO Annual Report 1962/63:14–24
- Goldschmidt T, Witte F, Wanink JH (1993) Cascading effects of the introduced Nile perch on the detritivorous/phytoplanktivorous species in the sub littoral areas of Lake Victoria. Conserv Biol 7:686–700
- Goudswaard PC, Witte F (1997) The catfish fauna of Lake Victoria after the Nile perch upsurge. Environm Biol Fishes 49:21–43
- Goudswaard PC, Witte F, Katunzi EFB (2002a) The tilapiine fish stock of Lake Victoria before and after the Nile perch upsurge. J Fish Biol 60:838–856
- Goudswaard PC, Witte F, Chapman LJ (2002b) Decline of the African lungfish *Protopterus aethiopicus* in Lake Victoria (East Africa). Afr J Ecol 40:42–52
- Goudswaard PC, Witte F, Wanink JH (2006) The Shrimp *Caridina nilotica* in Lake Victoria (East Africa), before and after the Nile perch increase. Hydrobiologia 563:31–44
- Hamblyn EL (1961) The Nile perch project. EAFFRO Annual Report 1960:26–32
- Harrison K, Crimmen O, Travers R et al (1989) Balancing the scales of Lake Victoria. Biologist 36:189–191
- Hauser L, Carvalho GR, Pitcher TJ et al (1998) Genetic affinities of an introduced predator: Nile perch in Lake Victoria, East Africa. Mol Ecol 7:849–857
- Hecky RE (1993) The eutrophication of Lake Victoria. Internationale Vereinigung für Theoretische und Angewandte Limnologie. Verhandlungen 25:39–48
- Hecky RE, Bugenyi FWB, Ochumba P et al (1994) Deoxygenation of the deep water of Lake Victoria, East Africa. Limnol Oceanogr 39:1476–1481
- Hughes NF (1986) Changes in the feeding biology of the Nile perch, *Lates niloticus*, in Lake Victoria since its introduction in 1960, and its impact on the native fish community of the Nyanza Gulf. J Fish Biol 29:541–548
- Hughes NF (1992) Nile perch, *Lates niloticus*, predation on the freshwater prawn, *Caridina nilotica*, in the Nyanza Gulf of Lake Victoria, East Africa. Environm Biol Fishes 33:307–309
- Katunzi EFB, Van Densen WLT, Wanink JH et al. (2006) Spatial and temporal patterns in the feeding habits of juvenile Nile perch in the Mwanza Gulf of Lake Victoria. Hydrobiologia 256:121–133
- Kaufman L (1992) Catastrophic change in species-rich freshwater ecosystems, the lessons of Lake Victoria. BioScience 42:846–848
- Kudhongania AW, Cordone AJ (1974a) Past trends, present stocks and possible future state of the fisheries of the Tanzanian part of the Lake Victoria. African J Trop Hydrobiol Fish 3:167–181
- Kudhongania AW, Cordone AJ (1974b) Batho-spatial distribution pattern and biomass estimate of the major demersal fishes in Lake Victoria. Afr J Trop Hydrobiol Fish 3:15–31
- Kudhongania AW, Twongo T, Ogotu-Ohwayo R (1992) Impact of the Nile perch on the fisheries of Lake Victoria and Kyoga. Hydrobiologia 232: 1–10
- Kudhongania AW, Chitamwebwa DBR (1995) Impact of environmental change, species introductions and ecological interactions on the fish stocks of Lake Victoria. In: Pitcher TJ, Hart PJB (eds). The impact of species changes in African lakes. Chapman & Hall, London: 19–32
- Ligtvoet W, Mkumbo OC (1990) Synopsis of ecological and fishery research on Nile perch (*Lates niloticus*) in Lake Victoria, conducted by HEST/TAFIRI. pp 35–74. In: CIFA. Report of the 5th session of the Sub-Committee for the Development and Management of the Fisheries in Lake Victoria, 12–24 Sept. 1989, Mwanza. FAO fisheries report 430
- Marten GG (1979) Impact of fishing on the inshore fishery of Lake Victoria (East Africa). J Fish Res Board Canada 36:891–900

- Mbahinzireki GB (1992) Distribution and abundance of two dipteran aquatic larvae in northern Lake Victoria. *FAO fisheries report* 475:39
- Mkumbo OC, Ligvoet W (1992) Changes in the diet of Nile perch, *Lates niloticus* (L), in the Mwanza Gulf of Lake Victoria. *Hydrobiologia* 232:79–84
- Ogari J, Asila A (1990) The state of Lake Victoria fisheries, Kenya waters. In: CIFA. Report of the 5th session of the Sub-Committee for the development and management of the fisheries of Lake Victoria. *FAO fisheries report* 430, pp 18–23
- Ogari J, Asila A (1992) Status of the fisheries of Lake Victoria Kenya Sector. In: CIFA. Report of the 6th session of the sub-committee for the development and management of the fisheries in Lake Victoria, 10–13 Feb. 1992, Jinja. *FAO fisheries report* 475, pp 15–23
- Ogari J, Dadzie S (1988) The food of Nile perch, *Lates niloticus* (L.), after the disappearance of the haplochromine cichlids in the Nyanza Gulf of Lake Victoria (Kenya). *J Fish Biol* 32:571–577
- Ogutu-Ohwayo R (1990a) The decline of the native fishes of lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus*, and the Nile tilapia, *Oreochromis niloticus*. *Environm Biol Fish* 27:81–96
- Ogutu-Ohwayo R (1990b) Changes in the prey ingested and the variations in the Nile perch and other fish stocks of Lake Victoria (Uganda). *J Fish Biol* 37:55–63
- Okemwa EN (1981) Changes in fish species composition of Nyanza Gulf of Lake Victoria. In: Proceedings of the workshop of the Kenya Marine and Fisheries Research institute on aquatic resources of Kenya, 13–19 July 1981, Mombasa Kenya, pp 138–156
- Pringle (2005) The origins of the Nile perch in Lake Victoria. *BioScience* 55:780–787
- Seehausen O, Van Alphen JJM, Witte F (1997) Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* 277:1808–1811
- Seehausen O, Van Alphen JJM (1998) Effect of the male coloration on female mate choice in closely related Lake Victoria cichlids (Haplochromis nyererei complex). *Behav Ecol Sociobiol* 42:1–8
- Seehausen O, Van Alphen JJM, Witte F (2003) Implications of eutrophication for fish vision, behavioural ecology, and species coexistence. In: Crisman TL, Chapman LJ, Chapman CA et al (eds) *Conservation, Ecology and Management of African fresh waters*. University Press of Florida, Gainesville
- Turner GF, Tweddle D, Makwinja RD (1995) Changes in the demersal cichlid communities as a result of trawling in southern Lake Malawi. In: Pitcher TJ, Hart PJB (eds) *The impact of species change in African lakes*. Chapman & Hall, London, pp 397–412
- Videler JJ (1993) *Fish swimming*. Chapman & Hall, London, 260 pp
- Verschuren D, Johnson TC, Kling HJ, Edgington DN, Laevitt PR, Brown ET, Talbot MR, Hecky RE (2002) History and timing of human impact on Lake Victoria, East Africa. *Proc R Soc London* 269:289–294
- Welcomme RL (1988) International introductions of inland aquatic species. *FAO fisheries technical paper* 249
- Williamson M (1996) *Biological invasions*. Chapman & Hall, London. 244 pp
- Witte F, Goudswaard PC (1985) Aspects of the haplochromine fishery in southern Lake Victoria. In: CIFA. Report of the 3rd Session of the Sub-Committee for the Development and Management of the Fisheries in Lake Victoria, 4–5 October 1984, Jinja. *FAO fisheries report* 335, pp 81–88
- Witte F, Goldschmidt T, Wanink JH et al (1992a) The destruction of an endemic species flock Quantitative data on the decline of the haplochromine cichlids of Lake Victoria. *Environm Biol Fish* 34:1–28
- Witte F, Goldschmidt T, Goudswaard PC et al (1992b) Species extinction and concomitant ecological changes in Lake Victoria. *Netherlands J Zool* 42:214–232
- Witte F, Wanink JH, Rutjes HA et al (2005) Eutrophication and the fish fauna of Lake Victoria. In: Vikram Reddy M (ed) *Restoration and management of tropical eutrophic lakes*. Science Publishers Inc, Enfield (NH), USA, pp 301–338
- Zaret TM, Paine RT (1973) Species introduction in a tropical Lake. *Science* 182:449–455