

A fast ethanol assay to detect seed deterioration

Jan Kodde¹, Wayne T. Buckley², Corine C. de Groot³, Marie Retiere¹, Ana Marcela Viquez Zamora⁴ and Steven P.C. Groot^{1*}

¹Plant Research International B.V., PO Box 619, 6700AP Wageningen, The Netherlands; ²Agriculture and Agri-Food Canada, Brandon Research Centre, PO Box 1000A, R.R. #3 Brandon, Manitoba R7A 5Y3, Canada; ³Bejo Zaden BV., PO Box 50, 1749ZH Warmenhuizen, The Netherlands; ⁴Wageningen University, Horticultural Production Chains Group, PO Box 630, 6700AP Wageningen, The Netherlands

(Received 4 July 2011; accepted after revision 11 September 2011)

Abstract

The most common way to test seed quality is to use a simple and reliable but time- and space-consuming germination test. In this paper we present a fast and simple method to analyse cabbage seed deterioration by measuring ethanol production from partially imbibed seeds. The method uses a modified breath analyser and is simple compared to gas chromatographic or enzymatic procedures. A modified method using elevated temperatures (40°C instead of 20°C) shortened the assay time and improved its sensitivity. The analysis showed an inverse correlation between ethanol production and seed quality (e.g. the final percentages or speed of germination and the number of normal seedlings). The increase in ethanol production was observed when cabbage seeds were deteriorated by storage under ambient conditions or hot water treatments, both of which reduced the number of normal seedlings. Premature seeds produced more ethanol upon imbibition than mature seeds. Ethanol production occurred simultaneously with oxygen consumption, indicating that lack of oxygen is not the major trigger for ethanol production.

Keywords: *Brassica*, breath analyser, ethanol, fermentation, mitochondria, seed deterioration, seed quality

Introduction

Seed quality with respect to germination and seedling establishment is important for seed producers and farmers. This quality is influenced by various factors,

including seed production conditions, harvesting, cleaning, sorting, seed treatments and seed health. In this paper we use the term 'quality' to refer to germination characteristics including the percentage or speed of germination, and the percentage of normal seedlings.

Premature harvest of seeds is known to have a negative effect on seed germination and tolerance to adverse conditions, e.g. hot water treatments (Jalink *et al.*, 1998; Groot *et al.*, 2006). There is a need for a fast test to determine the quality of seeds to optimize the settings for seed sorting based on maturity or hot water treatments in the seed industry.

Seeds may accumulate damage during storage or treatments. The degree of the deterioration varies through the amounts of stress imposed by these processes. This deterioration may result in reduced germination and seedling emergence, which is not necessarily easy to predict. Matthews and Khajeh-Hosseini (2006, 2007) showed a clear negative correlation between the mean time required for radicle protrusion and the average, final shoot length or the mean time for emergence from soil, by comparing ten maize seed lots. These results were explained by possible, prolonged periods of metabolic repair of the damage. Analysis of the degree of damage might predict germination performance of a seed lot. Mitochondria are known to be among the primary sites of deterioration, for instance during seed ageing (Bewley and Black, 1994).

Upon imbibition metabolic processes are re-initiated, including transcription, translation and enzyme activation. All these processes require the production of energy, generated by aerobic or anaerobic respiration. During aerobic respiration pyruvate is catabolized in the citric acid cycle, which proceeds in the mitochondria. This citric acid cycle generates NADH and FADH₂ which are oxidized in the electron transport chain. During electron transport a proton gradient is established over the inner

*Correspondence
Fax: + 31 317 480587
Email: steven.groot@wur.nl

mitochondrial membrane, which drives the production of ATP via ATPase. Glycolysis, which takes place in the cytosol, is the primary source of ATP in the absence of functional mitochondria. During glycolysis glucose is converted into two pyruvate molecules and NAD^+ is reduced to NADH which is oxidized to NAD^+ in functional mitochondria (Fan *et al.*, 1997; Rasmusson *et al.*, 2004).

Although lack of oxygen (e.g. due to submergence) is often the major reason for fermentation to occur, it may also take place under aerobic conditions. For example, during pollen tube growth both aerobic respiration and fermentation occur (Bucher *et al.*, 1995).

During imbibition in pea seeds, it takes some time for the mitochondria to reconstitute to a morphologically 'mature' structure with a folded inner membrane (Benamar *et al.*, 2003). When mitochondria are not functional, an alternative mechanism of regenerating NAD^+ is needed. This is provided by the conversion of pyruvate to ethanol or lactic acid during fermentation (Smith and Rees, 1979; Feireira de Sousa and Sodek, 2002).

Many plant species are known to have the ability to produce ethanol (Cossins and Beevers, 1963). Elevated levels of ethanol and its precursor, acetaldehyde, were detected in soybean seeds 30 min after the beginning of imbibition (Woodstock and Taylorson, 1981). The soybean seeds were thought unable to generate adequate energy (ATP) via aerobic respiration early in imbibition and employed anaerobic respiration, or fermentation, as an alternative mechanism of generating energy.

Among many factors affecting seed performance is the damage that accumulates in the seed during drying and storage, for instance that to DNA and cell, including mitochondrial, membranes (Bewley and Black, 1994). When mitochondria are non-functional due to the damage, initiation of anaerobic respiration is expected to deliver the energy for metabolic processes, at least for repair. Indeed, a relation between ethanol production and seed deterioration, as evidenced by a decline in germination or seedling growth, has been suggested (Perry, 1978; Woodstock and Taylorson, 1981; Taylor *et al.*, 1999; Kataki and Taylor, 2001; Bicanic *et al.*, 2003; Akimoto *et al.*, 2004; Rutzke *et al.*, 2008). In those studies seed deterioration was imposed by natural or conditional ageing of the seeds and ethanol production was measured using gas chromatography, immobilized enzyme technology or laser-based photo-acoustic spectroscopy. For routine analyses of multiple samples, these analytical methods are relatively slow and costly. Buckley and co-workers (Buckley *et al.*, 2006; Buckley and Huang, 2011) introduced a simplified method of measuring seed-produced ethanol with a modified breath analyser. Using the latter method, we studied ethanol

production by imbibed cabbage seeds of different maturity, after hot water treatments or natural ageing in the laboratory.

Materials and methods

Seed samples

Seeds of cabbage (*Brassica oleracea* L.) from a recently harvested commercial seed lot (cv. Stanton F1, Savoy winter cabbage) and an approximately 15-year-old, commercial seed lot (cv. Bartolo F1, white cabbage) were sorted into six fractions differing in maturity based on chlorophyll fluorescence (ScanMaster I, Satake, Houston, Texas, USA) (Jalink *et al.*, 1998). To test the effect of hot water treatments on seed germination and ethanol production, seeds of the highest maturity fraction (Bartolo, maturity fraction 6) were submerged for different periods of time (15, 30, 60 and 90 min) in tap water at 55°C. The seeds were then cooled with running tap water, blotted on filter paper and dried to 6% seed moisture (fresh weight basis) by incubating them for at least 3 d under circulating air at 32% relative humidity (RH) and 20°C. A relative humidity of 32% was maintained with a saturated solution of CaCl_2 . Seeds that were rinsed under tap water and dried, but not incubated in water at 55°C were used as a control.

Natural ageing was also investigated. A 9-month-old cabbage seed lot (cv. Amtrak F1, white cabbage) was stored under ambient conditions on a laboratory bench at about 20°C and 40–50% RH. At various times, subsamples were transferred to –20°C to suspend ageing. About 31 months after the start of the experiment the seeds were retrieved from the –20°C storage and subjected to a germination test and analysis of ethanol production.

Germination tests

Germination tests were performed using a Jacobsen germination table (Karma, 1968) with two or four replicates of 50 seeds on filter paper. To maintain high humidity around the seeds an inverted plastic funnel with a hole in the conical end was placed over each filter paper with seeds. Two different germination assays were performed. In assay 1, seeds were incubated at 20°C with 8 h light ($40\text{--}65\ \mu\text{mol m}^{-2}\text{s}^{-1}$)/16 h dark. Seeds were scored as germinated when the radicle had protruded. Radicle emergence was counted daily to determine the speed of germination. Using GERMINATOR software (Joosen *et al.*, 2010), a curve-fitting program designed for the analysis of germination data, time required to reach 50% of the final germination (T_{50}) was calculated.

The germination data were fitted to a four-parameter Hill function by iteration using the Solver tool (Microsoft Excel). The resulting curve formula was used to calculate germination parameters including the final percentage, speed and uniformity of germination. After 30 d, seedling quality, total germination percentage and percentage of normal seedlings were recorded. Normal seedlings were evaluated as described in the rulebook of the International Seed Testing Association (ISTA, 1999). In assay 2, seeds were germinated under continuous light ($40\text{--}65\ \mu\text{mol m}^{-2}\text{s}^{-1}$) with an alternating temperature of 20°C (16 h) and 30°C (8 h). As an indicator of seed quality, the number of normal seedlings established after 5 d was also counted. Final germination was determined after 7 d in this assay.

Ethanol assay

Five hundred milligrams of seeds (treated, sorted or control) equilibrated at 32% RH were placed in a 20 ml glass vial. Water was added to achieve the desired seed moisture percentage, taking into account the initial seed moisture content of 6%. The vials were sealed with aluminium crimp caps with 3-mm-thick butyl rubber and Teflon septa immediately after adding water. The vials containing wetted seeds were incubated at either 20°C or 40°C . Ethanol concentration in the headspace was measured at different time points using the modified breath analyser (Alcotest 6810, Dräger safety AG & Co. KGaA, Lübeck, Germany). To avoid temporary cooling, sample measurement outside the incubator was performed within 5 s.

The original breath analyser (Dräger 6810) was modified by replacing the breath sampling port with a Luer fitting and by making adjustments to the software. A disposable hypodermic needle (18 gauge, 5 cm long) was used to collect gas samples from the vials. The needle was shortened to approximately 2.5 cm and cut half through approximately 1.5 cm from the Luer hub to allow for pressure equilibration in the vial during its insertion. The device collected a subsample of approximately 0.3 ml from the headspace for every measurement. The analytical range of the breath analyser was $10\text{--}2500\ \mu\text{g}$ ethanol per litre of gas mixture. The breath analyser was calibrated annually by the supplier.

Oxygen and carbon dioxide measurements

In a separate experiment, oxygen and carbon dioxide were measured in the headspace of 20 ml vials containing 500 mg of either the most mature or the least mature fractions of the Bartolo and Stanton seed lots. At the start of the experiment water was added to a final moisture content of 30%. The vials were then

incubated at 40°C . After 6 h, oxygen and carbon dioxide were determined in 1 ml samples of headspace atmosphere by gas chromatography (Chrompack CP-2002 Micro-GC, Chrompack, Middelburg, The Netherlands). A Molsieve 5A PLOT column (Restec, Bellefonte, Pennsylvania, USA) was used for oxygen and a HayeSep A column (Valco Instruments Company Inc., Houston, Texas, USA) for carbon dioxide. The data were analysed using the Maestro II software (Gerstel GmbH & Co. KG, Mülheim an der Ruhr, Germany).

Results

Hot water treatments

To optimize the ethanol assay, seed samples of different quality were produced using four durations (15, 30, 60 and 90 min) of a hot water treatment at 55°C . The hot water treatments were applied to mature Bartolo seeds and the subsequent effect on seed quality was determined using germination assay 1 with four replicates. Hot water treatment for 15 min delayed germination (i.e. increased T_{50}), but did not affect total germination of maturity fraction 6 (the most mature) of the Bartolo seed lot (Fig. 1). However, the frequency of normal seedlings declined. Longer hot water treatments decreased total radicle emergence, increased T_{50} and strongly decreased the frequency of normal seedlings. After treatment for 1 h radicle protrusion failed in all seeds.

Ethanol production from the hot water treated, deteriorated seeds was analysed after equilibrating them to five different seed moisture contents (20, 30, 40, 50 and 60%) and incubating them at 20°C (24 h) or 40°C (6 h). The samples showed a marked variation in




Figure 1. Effect of hot water treatments (55°C ; 15, 30, 60 and 90 min) on germination performance of cv. Bartolo seeds (fraction 6: the most mature). Total germination percentage after 30 d (□), the percentage of normal seedlings (■) and the time to reach 50% of the final germination percentage (T_{50} , ●) are shown (mean \pm SE, $n = 4$).




Figure 2. Effect of duration of hot water (55°C) treatment and seed moisture content (%) on headspace ethanol levels above 500 mg samples of cv. Bartolo seeds in 20 ml vials. Ethanol was measured after 24 h incubation at 20°C (a) or 6 h incubation at 40°C (b).

ethanol production (Fig. 2a). Control seeds (only kept under running tap water for a short time) and seeds treated with hot water for 15 min did not produce ethanol at 20°C, whereas 30–90 min hot water treatments resulted in ethanol production (Fig. 2a). The highest ethanol production at 20°C was observed from the seeds treated with hot water for 60 min and incubated at 40% moisture at both 20°C and 40°C. Headspace ethanol concentrations were much higher in seed samples incubated at 40°C than those at 20°C (Fig. 2a and b). When no (control: 0 min) or only a short (15 min) hot water treatment was given, less ethanol was produced from seeds with 40% moisture than from those with 30, 50 or 60% moisture. This local minimum was reproducible in repetitive experiments (data not shown). Tests at other incubation temperatures (30, 35, 45, 50 and 60°C) confirmed that 40°C was the most efficient temperature to detect differential ethanol production from hot water treated cabbage seeds.

Seed maturity

The second set of experiments was performed on cabbage seeds varying in maturity. The seed quality of six maturity fractions, which were sorted on the basis of their chlorophyll fluorescence (i.e. decreasing chlorophyll indicating increasing maturity), was analysed using germination assay 1, with four replicates. The frequency of normal seedlings gradually increased from fraction 1 (the least mature, 69%) to fraction 6 (the most mature, 93%) (Fig. 3), while total germination was affected only in fraction 1 (the least mature seeds). After incubation at 20°C (30% moisture) for 24 h, headspace ethanol was detectable only from fraction 1 (the least mature) (Fig. 3). When seeds were

incubated at 40°C for 6 h, much higher ethanol levels were found for seeds across all maturity fractions. Headspace ethanol levels decreased with increasing seed maturity and showed a negative correlation with the frequency of normal seedlings ($r = -0.97$).

Oxygen and carbon dioxide

The previous experiment was partly repeated and extended with maturity fractions 1 and 6 from the same Bartolo seed lot and from another cultivar (Stanton), using germination assay 1 with two




Figure 3. Seeds (cv. Bartolo) were sorted into six fractions differing in maturity based on their chlorophyll content, with fraction 1 (the highest chlorophyll content, i.e. the least mature) through fraction 6 (the lowest chlorophyll content, i.e. the most mature). The percentages of total germination and normal seedlings (mean \pm SE, $n = 4$) were determined according to germination analysis 1. Headspace ethanol levels (500 mg samples, 20 ml vials and 30% final seed moisture) were measured after 24 h incubation at 20°C (▲) or 6 h incubation at 40°C (●).

replicates. The most mature fraction 6 performed better, in terms of the percentages of germination and normal seedlings, than the least mature fraction 1 for both seed lots (Fig. 4). Headspace ethanol, carbon dioxide and oxygen levels of fractions 1 and 6 of Bartolo or Stanton seeds were determined by gas chromatography after 6 h incubation at 40°C with a 30% moisture content. Ethanol and carbon dioxide levels were higher while oxygen was significantly lower for fraction 1 (the least mature) compared to fraction 6 (the most mature) for both cultivars (Fig. 4) (ANOVA, $P = 0.02$). Oxygen levels never dropped below 18% during the assay.

Natural ageing

Quality variation was also obtained from naturally aged cabbage seeds (cv. Amtrak). Storage of the cabbage seeds at the laboratory bench resulted in a gradual decline in germination performance

(germination assay 2, with two replicates) (Fig. 5). The number of normal seedlings after 5 d (germination assay 2: early evaluation) declined from 92 to 58% over a period of 31 months. The decline was less obvious (from 94 to 82%) when evaluation was performed after 7 d. When samples from these aged seeds were incubated at 20°C with 30% moisture for 24 h, headspace ethanol was detected from seeds stored for 19 months or longer (Fig. 5). Headspace ethanol levels were much higher when incubated at 40°C (3.5 h). Ethanol production was already detectable from seeds stored for 4.8 months when the 40°C incubation was used for ethanol assay. Prolonged (24 h) incubation at 40°C allowed ethanol detection from all samples, while ethanol production from seeds aged for 17 months or more reached the maximum detection limit.

The ethanol levels measured at 40°C after 3.5 h were negatively correlated ($r = -0.92$) with the number of normal seedlings after 5 d. This correlation was slightly less ($r = -0.87$) when the ethanol assay was performed




Figure 4. Effect of seed maturity on germination performance (germination assay 1) and respiration. Seeds of two cultivars Bartolo (B) and Stanton (S) differing in maturity were sorted based on their chlorophyll content into six fractions, fraction 1 (the highest chlorophyll content, i.e. the least mature) through fraction 6 (the lowest chlorophyll content, i.e. the most mature). The control was water. The percentages of final germination and normal seedlings are shown in (a). Headspace ethanol (b), oxygen (c) and carbon dioxide (d) levels were determined in the headspace after 6 h incubation at 30% seed moisture and 40°C. Values are means \pm two data points.




Figure 5. Effect of seed ageing on ethanol production. Cabbage seeds (cv. Amtrak) were naturally aged at about 20°C on a laboratory bench. Seed quality was assessed by the percentage of normal seedlings after 5 d (■) and total germination after 7 d (□) (mean \pm two data points). Headspace ethanol was measured above 500 mg seed samples at 30% moisture in 20 ml vials after 24 h incubation (▲) at 20°C or after 3.5 h (○) or 24 h (●) incubation at 40°C.

at 20°C for 24 h. The amount of ethanol produced at 40°C after 3.5 h was almost linearly correlated ($r = 0.98$) with ageing time at the laboratory bench.

Discussion

Field emergence of seeds is an important aspect of crop establishment. Seed companies and growers are seeking reliable and rapid methods to predict field performance of their seeds. Field performance is a complex trait related to a collection of physiological and genetic factors, all of which cannot be measured in a single assay. Most likely, multiple tests will be needed to provide a comprehensive evaluation and reliable prediction of field performance.

Delouche and Baskin (1973) proposed a sequence of events during seed deterioration which leads to loss of germination and field performance. The first event in this model is membrane degradation. An ideal test, therefore, would be one that could measure membrane deterioration (McDonald, 1980).

As ethanol production most likely is initiated or enhanced by the loss of mitochondrial membrane integrity, an ethanol assay has good potential as a test to examine the level of deterioration in seeds. While ethanol can be quantified with research-oriented, analytical equipment, the assay has to be simple for practical purposes. In the present work we demonstrated that variation in cabbage seed quality can be analysed by measuring ethanol production with a hand-held breath analyser (Buckley *et al.*, 2006; Buckley and Huang, 2011).

The sensitivity of the assay was improved by incubating the seeds at 40°C instead of the moderate temperatures of 20 or 25°C used in previous studies (Perry, 1978; Woodstock and Taylorson, 1981; Taylor *et al.*, 1999; Kataki and Taylor, 2001; Bicanic *et al.*, 2003; Akimoto *et al.*, 2004; Rutzke *et al.*, 2008; Buckley and Buckley, 2009; Buckley and Huang, 2011). The advantage of using 40°C for ethanol assay was also well exemplified by the analysis of different fractions of seeds differing in maturity (Fig. 3) and the naturally aged seeds with different quality (Fig. 5).

Our results also identified some precautions necessary for the reliable application of this method. The ethanol levels at 40°C after prolonged incubation (24 h) of the aged seeds exceeded those commonly measured with the breath analyser (e.g. human breath analysis by law enforcement). This potential problem can be addressed by a newer version of the analyser, which is dedicated particularly for use in agricultural research and application. The maximum ethanol detection limit of the modified breath analyser could be 9990 $\mu\text{g l}^{-1}$ which exceeds the previous detection limit of 2500 $\mu\text{g l}^{-1}$ (Burkhard Stock, Dräger safety AG&CO KGaA, pers. comm.).

In general, incubation at 40°C yielded a substantial amount of headspace ethanol within hours and enabled the entire assay to be run within a single working day. Higher temperatures are not recommended because discrimination of differences in seed quality was not improved (data not shown).

The increase of ethanol levels in the headspace at higher temperatures can be explained, in part, by the effect of temperature on the partitioning of ethanol in

an air/water system. Elevated metabolism in seeds could additionally explain higher headspace ethanol levels at 40°C compared to those at 20°C. Pyruvate decarboxylase is one of the regulatory enzymes in the conversion of pyruvate to ethanol (Davies *et al.*, 1974). The activity of this enzyme in wheat seeds is about five times greater at 40°C than at 20°C (Singer and Pensky, 1952). It is also possible that incubation at 40°C induced elevated ethanol production by additional seed deterioration, because 40°C is above the maximum germination temperature for cabbage seeds (Elson *et al.*, 1992). Furthermore, immature or deteriorated seeds may be more sensitive to high temperature stress than high-quality seeds. The elevated headspace ethanol observed at 40°C may be due to a combination of physical and biological factors. During the assay oxygen levels did not drop below 18% (Fig. 4), indicating that ethanol production was not due to lack of oxygen.

A moisture content of 30% seems optimal for determining ethanol production from cabbage seeds (Fig. 2). Also, this is below the minimal moisture content (40%) required for radicle protrusion (data not shown). This agrees with previous research (Rutzke *et al.*, 2008), in which the quality differences of artificially aged cabbage seed could be resolved better at 30% than at 60% moisture.

Hot water treatment of seeds is performed to reduce seed-borne diseases (McGee, 1995). In practice, it is important to find a balance between effectiveness against pathogens and seed damage. Seed lots may differ in their sensitivity to hot water treatments, and therefore test treatments should be performed to determine the optimal conditions (Groot *et al.*, 2006, 2008). However, test treatments are time-consuming, and a rapid assay to determine potential negative effects on seed germination performance is needed. The ethanol assay developed here may offer such a rapid assay.

Acknowledgements

Dräger safety AG&CO KGaA (Germany) is acknowledged for providing modified breath analysers. The research presented here has been funded by the Netherlands Ministry of Economic affairs, Agriculture and Innovation in the frame of supporting organic agriculture through research.

References

Akimoto, T., Cho, S., Yoshida, H., Furuta, H. and Esashi, Y. (2004) Involvement of acetaldehyde in seed deterioration of some recalcitrant woody species through the acceleration of aerobic respiration. *Plant and Cell Physiology* **45**, 201–210.

- Benamar, A., Tallon, C. and Macherel, D. (2003) Membrane integrity and oxidative properties of mitochondria isolated from imbibing pea seeds after priming or accelerated ageing. *Seed Science Research* **13**, 35–45.
- Bewley, J.D. and Black, M. (1994) *Seeds: Physiology of development and germination* (2nd edition). New York, Plenum Press.
- Bicanic, D., Persijn, S., Taylor, A., Cozijnsen, J., van Veldhuijzen, D., Lenssen, G. and Wegh, H. (2003) Detection of ethanol and acetaldehyde released from cabbage seeds of different quality: laser photoacoustic spectroscopy versus FTIR and headspace gas chromatography. *Review of Scientific Instruments* **74**, 690–693.
- Bucher, M., Brander, K.A., Sbicego, S., Mandel, T. and Kuhlemeier, C. (1995) Aerobic fermentation in tobacco pollen. *Plant Molecular Biology* **28**, 739–750.
- Buckley, W.T. and Buckley, K.E. (2009) Low-molecular-weight volatile indicators of canola seed deterioration. *Seed Science and Technology* **37**, 676–690.
- Buckley, W.T. and Huang, J. (2011) An ethanol-based seed vigour assay for canola. *Seed Science and Technology* **39**, 510–526.
- Buckley W.T., Irvine R.B., Buckley K.E. and Elliott R.H. (2006) New, 24-h seed vigour assays for canola. The ASA–CSSA–SSSA International Annual Meetings (12–16 November 2006), Indianapolis. Available at website <http://a-c-s.confex.com/crops/2006am/techprogram/P26853.HTM> (accessed 27 October 2010).
- Cossins, E.A. and Beever, H. (1963) Ethanol metabolism in plant tissues. *Plant Physiology* **38**, 375–380.
- Davies, D.D., Grego, S. and Kenworthy, P. (1974) The control of the production of lactate and ethanol by higher plants. *Planta* **118**, 297–310.
- Delouche, J.C. and Baskin, C.C. (1973) Accelerated aging techniques for predicting the relative storability of seed lots. *Seed Science and Technology* **1**, 427–452.
- Elson, M.K., Morse, R.D., Wolf, D.D. and Vaughan, D.H. (1992) High-temperature inhibition of seed germination and seedling emergence of broccoli. *HortTechnology* **2**, 417–419.
- Fan, T.W.M., Higashi, R.M., Frenkiel, T.A. and Lane, A.N. (1997) Anaerobic nitrate and ammonium metabolism in flood-tolerant rice coleoptiles. *Journal of Experimental Botany* **48**, 1655–1666.
- Feireira de Sousa, C.A. and Sodek, L. (2002) The metabolic response of plants to oxygen deficiency. *Brazilian Journal of Plant Physiology* **14**, 83–94.
- Groot, S.P.C., Birnbaum, Y., Rop, N., Jalink, H., Forsberg, G., Kromphardt, C., Werner, S. and Koch, E. (2006) Effect of seed maturity on sensitivity of seeds towards physical sanitation treatments. *Seed Science and Technology* **34**, 403–413.
- Groot, S.P.C., Birnbaum, Y., Kromphardt, C., Forsberg, G., Rop, N. and Werner, S. (2008) Effect of the activation of germination processes on the sensitivity of seeds towards physical sanitation treatments. *Seed Science and Technology* **36**, 609–620.
- ISTA (1999) International Seed Testing Association. International rules for seed testing. *Seed Science and Technology* **27**.
- Jalink, H., van der Schoor, R., Frandas, A., van Pijlen, J.G. and Bino, R.J. (1998) Chlorophyll fluorescence of *Brassica oleracea* seeds as a non-destructive marker for

- seed maturity and seed performance. *Seed Science Research* **8**, 437–443.
- Joosen, R.V.L., Kodde, J., Willems, L.A.J., Ligterink, W., Van der Plas, L.H.W. and Hilhorst, H.W.M.** (2010) GERMINATOR: a software package for high-throughput scoring and curve fitting of Arabidopsis seed germination. *The Plant Journal* **62**, 148–159.
- Karma, S.K.** (1968) Effect of different distances between water level and seed bed on Jacobsen apparatus on the germination of *Pinus silvestris* L. seed. *Studia Forestalia Suecica* **65**, 1–17.
- Kataki, P.K. and Taylor, A.G.** (2001) Time course study of ethanol production by corn and soybean to optimize the use of ANA ethanol index as an accurate seed quality test. *Journal of New Seeds* **3**, 1–17.
- Matthews, S. and Khajeh-Hosseini, M.** (2006) Mean germination time as an indicator of emergence performance in soil of seed lots of maize (*Zea mays*). *Seed Science and Technology* **34**, 339–347.
- Matthews, S. and Khajeh-Hosseini, M.** (2007) Length of the lag period of germination and metabolic repair explain vigour differences in seed lots of maize (*Zea mays*). *Seed Science and Technology* **35**, 200–212.
- McDonald, M.B.** (1980) Assessment of seed quality. *Hortscience* **15**, 784–788.
- McGee, D.C.** (1995) Epidemiological approach to disease management through seed technology. *Annual Review of Phytopathology* **33**, 445–466.
- Perry, D.A.** (1978) Report on the vigour test committee 1974–1977. *Seed Science and Technology* **6**, 159–181.
- Rasmusson, A.G., Soole, K.L. and Elthon, T.E.** (2004) Alternative NAD(P)H dehydrogenase of plant mitochondria. *Annual Review of Plant Biology* **55**, 23–39.
- Rutzke, C.F.J., Taylor, A.G. and Obendorf, R.L.** (2008) Influence of aging, oxygen, and moisture on ethanol production from cabbage seeds. *Journal American Society for Horticultural Science* **133**, 158–164.
- Singer, T.P. and Pensky, J.** (1952) Isolation and properties of the α -carboxylase of wheat germ. *Journal of Biological Chemistry* **196**, 375–388.
- Smith, A.M. and Rees, T.** (1979) Pathways of carbohydrate fermentation in the roots of marsh plants. *Planta* **146**, 327–334.
- Taylor, A.G., Johnson, C.F. and Kataki, P.K.** (1999) Ethanol production by hydrated seeds: A high resolution index of seeds quality. *ISHS Acta Horticulturae* **504**, 153–157.
- Woodstock, L.W. and Taylorson, R.B.** (1981) Ethanol and acetaldehyde in imbibing soybean seeds in relation to deterioration. *Plant Physiology* **67**, 424–428.