

NN 8201

no 505

C

Inactivation of
peroxidase,
pectinesterase and
alkaline phosphatase in
polymers
as a model for
irradiation of
dried foodstuffs

J. P. Roozen

BIBLIOTHEEK
DER
IDBOUWHOGESCHOOL
N. FOULKESWEG 1a
WAGENINGEN

NN08201.505

Inactivation of peroxidase, pectinesterase and alkaline phosphatase in polymers as a model for irradiation of dried foodstuffs

Dit proefschrift met stellingen van Jacobus Petrus Roozen, landbouwkundig ingenieur, geboren te Oud-Gastel op 28 februari 1943, is goedgekeurd door de promotor, dr. W. Pilnik, hoogleraar in de levensmiddelenleer.

De Rector Magnificus van de Landbouwhogeschool
J. M. Polak

Wageningen, 15 januari 1972

J. P. Roozen

Inactivation of peroxidase, pectinesterase and alkaline phosphatase in polymers as a model for irradiation of dried foodstuffs

Proefschrift
ter verkrijging van de graad van
doctor in de landbouwwetenschappen,
op gezag van de Rector Magnificus, Mr. J. M. Polak,
hoogleraar in de rechts- en staatswetenschappen
van de westerse gebieden,
in het openbaar te verdedigen
in de aula van de Landbouwhogeschool te Wageningen
op vrijdag 18 februari 1972 te 16 uur



1972 *Centre for Agricultural Publishing and Documentation*
Wageningen

The material of this thesis is being published in detail in the following papers:

Roozen, J. P. & W. Pilnik,

Über die Stabilität adsorbierter Enzyme in wasserarmen Systemen.

1970. I. Die Stabilität von Peroxydase bei 25°C. Lebensm.-Wiss. u. Technol. 3: 37–40.

1971. II. Der Einfluss von pH-Verschiebungen auf die Stabilität von Peroxydase bei 2°C. Lebensm.-Wiss. u. Technol. 4: 24–26.

1971. III. Der Effekt von Elektronenbestrahlung auf die Stabilität von Peroxydase. Lebensm.-Wiss. u. Technol. 4: 93–96.

1971. IV. Der Effekt von Lagerung und Elektronenbestrahlung auf die Stabilität von Pektinmethylesterase. Lebensm.-Wiss. u. Technol. 4: 158–162.

1971. V. Der Effekt von Lagerung und Elektronenbestrahlung auf die Stabilität von alkalischer Phosphatase. Lebensm.-Wiss. u. Technol. 4: 196–200.

1972. VI. Möglichkeiten zur Inaktivierung von Enzymen in getrockneten Lebensmitteln durch Bestrahlung. Betrachtungen im Lichte der Literatur und eigener Untersuchungen. Lebensm.-Wiss. u. Technol., in press.

ISBN 90 220 0384 1

This thesis will also be published as Agricultural Research Reports 773.

© Centre for Agricultural Publishing and Documentation, Wageningen, 1972.

No part of this book may be reproduced and/or published in any form, by print, photoprint, microfilm or any other means without written permission from the publishers.

Stellingen

I

Wanneer het de bedoeling is op artikelen te promoveren, verdient het aanbeveling, dat tijdens het bewerken van het proefschrift de promotor en de promovendus de resultaten gezamenlijk publiceren.

II

Een mondelinge mededeling is ondeugdelijk als referentie in een wetenschappelijke publikatie.

W. D. Garvey & B. C. Griffith, in: **B. Montgomery**, *The foundations of access to knowledge*, Syracuse University, Syracuse, N.Y., pp. 129–151 (1968).

III

De begrippen radappertisatie, radicidatie en radurisatie zijn gekunsteld en daarom niet ingeburgerd.

H. E. Goresline, M. Ingram, P. Macuch, G. Mocquot, D. A. A. Mossel, C. F. Niven & F. S. Thatcher, *Nature* 204: 237–238 (1964).

IV

Naast een verlenging van de houdbaarheid zou de behandeling van levensmiddelen met ioniserende stralen het doden van pathogene kiemen kunnen beogen.

E. H. Kampelmacher, inaugurele rede, Landbouwhogeschool, Wageningen (1971).

V

De verschillen in gevoeligheid voor bestraling tussen choline-esterase en arylesterase als functie van hun vochtgehalte, zoals dat is beschreven door Augustinsson et al., zijn waarschijnlijk toe te schrijven aan verontreinigingen in de gebruikte enzym-preparaten.

K. B. Augustinsson, G. Jonsson & B. Sparrman, *Acta Chem. Scand.* 15: 11–15 (1961).
Dit proefschrift.

VI

In droge, enzymrijke poeders kan men de hoeveelheid 'gebonden water' bepalen met behulp van ioniserende stralen.

Dit proefschrift.

VII

De gevoeligheid voor bestraling van gezuiverde enzymen in oplossing verschilt dermate van die van enzymen *in vivo*, dat de verwachtingen van Pansolli ten aanzien van de toepassing van ioniserende stralen voor de conservering van fruit ongegrond zijn.

P. Pansolli, *Experientia* 26: 499-500 (1970).

Dit proefschrift.

VIII

Het is af te raden om voor het instellen van het evenwichtsvuchtgehalte van gedroogde systemen deze te plaatsen boven zuur-water mengsels.

L. Acker & E. Lück, *Z. Lebensmittelunters. u.-Forsch.* 108: 256-269 (1958).

IX

Naar het zich laat aanzien zal kiemremming van aardappelen met behulp van ioniserende stralen geen aanleiding zijn tot extra verliezen aan vitamine C.

H. Woggon, *Ernährungsforschung* 8: 387-405 (1963).

X

Voor de bewaring van bereide maaltijden in het 'réguéthermic' systeem blijkt bestraling een weinig zinvolle conserveringsmethode te zijn.

J. C. Botman, Ir-verslag, L.H. afd. Voeding (1970).

XI

Kinderen, die verplicht worden om dagelijks een lepel 'echte' levertraan te slikken, kunnen tegenwoordig letterlijk giftig worden.

T. Sloeker-van der Eijden, E. J. van Luytelaar & J. A. Lukkenaer, *Voeding* 32: 401-402 (1971).

Acknowledgments

I wish to express my sincere gratitude to my promotor, Prof. Dr. W. Pilnik, for giving me the opportunity of undertaking this work and for his suggestions which helped me considerably with my research. The method of publishing the results of the various phases of this research proved to be very valuable as the preparation of each paper stimulated and encouraged me further.

Furthermore, I am most grateful to Dr. Ir. D. de Zeeuw, director of the Institute for Atomic Sciences in Agriculture (ITAL), who suggested the topic and provided research facilities.

My thanks are due to the Davis (California) student Ann E. Vaughan and the Wageningen students F. H. Joosten, H. Hessels and H. Stegeman for their co-operation.

I am very grateful to Miss Helga E. Belling for typing the manuscript, to Mr. J. C. Baars for assisting at the irradiations, to Messrs. C. Rijpma and M. Schimmel for drawing the figures and to Ir. J. Eelkman Rooda for the computer programme. Finally I wish to thank Mrs. E. Brouns-Murray for correcting the English text and Mr. R. J. P. Aalpol for editing.

Abstract

ROOZEN, J. P. (1972) Inactivation of peroxidase, pectinesterase and alkaline phosphatase in polymers as a model for irradiation of dried foodstuffs. Doctoral thesis, Wageningen. ISBN 90 220 0384 1, (xii) + 11 p., 3 tbs, 33 refs, Eng. and Dutch summaries.

Also: Agric. Res. Repts (Versl. landbouwk. Onderz.) 773.

Aqueous solutions containing an enzyme (peroxidase, alkaline phosphatase, pectinesterase) and a polymer (carboxymethylcellulose, soluble starch, polyvinylpyrrolidone, sodium pectate) were freeze-dried. The systems thus obtained were incubated over saturated salt solutions at various levels of water activity. They were then irradiated by an electron generator. The storage and radiation sensitivity of the enzymes depended on the polymer used in the model, the enzyme concentration, the humidity and the pH.

Samenvatting

De inactivering van peroxydase, pektine-esterase en alkalische fosfatase in polymeren als modelstudie voor de bestraling van gedroogde levensmiddelen

Het effect van ioniserende stralen op de inactivering van enzymen als functie van het vochtgehalte werd onderzocht.

De benodigde systemen werden verkregen door de volgende enzym-polymeer oplossingen te vriesdrogen:

peroxydase-carboxymethylcellulose (POD-CMC)
peroxydase-zetmeel (POD-sS)
peroxydase-polyvinylpyrrolidon (POD-PVP)
pektine-esterase-carboxymethylcellulose (PE-CMC)
pektine-esterase-zetmeel (PE-sS)
pektine-esterase-polyvinylpyrrolidon (PE-PVP)
pektine-esterase-natriumpektaat (PE-NaP)
alkalische fosfatase-carboxymethylcellulose (AP-CMC)
alkalische fosfatase-zetmeel (AP-sS) en
alkalische fosfatase-polyvinylpyrrolidon (AP-PVP).

Het gevriesdroogde materiaal werd tot poeder vermalen en daarna verdeeld over 42 polyethyleen zakjes. De verkregen monsters werden één week in een exsiccator met een drooggemiddel bewaard en daarna twee weken in exsiccatoren met verzadigde zoutoplossingen om de monsters op het gewenste vochtgehalte te brengen. Vervolgens werden de monsters bestraald bij een constante temperatuur, in water opgelost en op enzymactiviteit getest.

Tijdens de behandeling, die aan de bestraling voorafging, was de stabilitet van POD afhankelijk van temperatuur, vochtgehalte, pH en de aard van de polymeren. Dezelfde factoren beïnvloeden de bestralingsgevoeligheid van POD. Een na-effect (Eng.: after-effect) van de bestraling werd ontdekt in de oplossingen van de POD-CMC- en de POD-sS-, maar er was geen na-effect in de oplossingen van POD-PVP systemen.

Bij incubatie met silicagel verloren de PE systemen de minste enzymactiviteit. In deze systemen was het enzym het minst bestralingsgevoelig bij 40% relatieve vochtigheid, behalve in de PVP systemen waarin de bestralingsgevoeligheid van het enzym scheen af te nemen met toenemende relatieve vochtigheid.

In alle systemen had de bestralingsgevoeligheid van alkalische fosfatase dezelfde orde van grootte, maar boven 40% relatieve vochtigheid daalde de gevoeligheid in de PVP systemen en steeg in de CMC- en sS systemen. Een toename van de enzymconcentratie deed de bestralingsgevoeligheid van AP dalen. Er was geen na-effect

in de oplossingen van de AP-PVP systemen, terwijl de na-effecten het sterkst tot uitdrukking kwamen in de oplossingen van de 'lage enzymconcentratie' systemen van CMC en sS.

In de CMC- en sS systemen en bij neutrale pH's was de volgorde in intrinsieke bestralingsgevoelighed van de onderzochte enzymen als volgt:

PE < POD < AP.

Deze staan in dezelfde volgorde als de molecuulgewichten van de enzymen en daarom is een regressieanalyse uitgevoerd van de intrinsieke bestralingsgevoelighed als functie van het molecuulgewicht. De intrinsieke bestralingsgevoeligheden van PE in het NaP systeem en van de enzymen in de PVP systemen pasten niet in dit schema. Deze afwijkingen zouden veroorzaakt kunnen worden door de vorming van een enzym-produkt complex (PE-NaP) en door de hoge reactiviteit van gehydrateerde electronen met PVP.

Contents

1	Introduction	1
	Dried foodstuffs	1
	Irradiation of macromolecules	1
	Radiosensitivity of enzymes	1
	Defining purpose of study	2
2	Definitions and abbreviations	3
3	Methods	4
4	Results	5
	Peroxidase systems	5
	Pectinesterase systems	5
	Alkaline phosphatase systems	6
5	Discussion and conclusions	7
	Radioresistance of the enzymes	7
	Intrinsic radiosensitivity	7
	After-effects	8
	Summary	9
	References	10

Curriculum vitae

De auteur begon, na het behalen van het einddiploma HBS-B op het Norbertus-Lyceum te Roosendaal, in 1961 zijn studie in de levensmiddelentechnologie aan de landbouwhogeschool te Wageningen. In januari 1966 behaalde hij het kandidaatsdiploma en in januari 1968 volgde het ingenieurs-diploma met als specialisatie kennis van levensmiddelen en als bijkijken voedingsmiddelenmicrobiologie, fysische chemie en industriële bedrijfskunde. Hierna kwam hij als wetenschappelijk assistent in dienst bij de landbouwhogeschool (laboratorium voor levensmiddelenchemie en -microbiologie) en sinds juli 1971 maakt hij deel uit van de staf als wetenschappelijk medewerker.

1 Introduction and literature

Dried foodstuffs Since ancient times foodstuffs have been dried to preserve them from spoilage which can be microbiological, chemical or enzymatic. The shelf life of dried products depends on their water content. Acker (1962) has written a detailed review on enzymatic spoilage in relation to the water activity of food. Most enzymes do not react at moisture contents below the inflection point of the sorption isotherm (Acker & Lück, 1958). However, lipids may be oxidized at such low moisture contents (Purr, 1970). Even if there is no problem of oxidation it is often difficult to achieve or maintain the low moisture content. In practice therefore it is still necessary to blanch dried and frozen vegetables to prevent enzymic deterioration. Replacement of this heat treatment by irradiation could improve some of the existing products; then even vegetables that wither on heating could be stored in a frozen or dried state.

Irradiation of macromolecules The most important effects of ionizing radiation on macromolecules are degradation and crosslinking. The disruption of secondary structures have so far only been described in proteins and deoxyribonucleic acid (Bacq & Alexander, 1966). Chapiro (1964) discussed some aspects of the radiation chemistry of polymers. In irradiated solid polymers, degradation and crosslinking can largely be ascribed to reactions by free radicals. When polymers are irradiated in solution, these reactions are influenced by free radicals from the solvent. Thus in the solid state, variations in water content have been shown to cause changes in the radiosensitivity of deoxyribonucleic acid (Lett & Alexander, 1961), carboxymethylcellulose (Munzel & Schenkel, 1964), starch (Reuschl & Guilbot, 1966), pectin (Wahba & Massey, 1966), and polyvinylpyrrolidone (Alexander & Charlesby, 1957).

Radiosensitivity of enzymes From a survey of literature, Marples & Glew (1958) concluded that enzymes in food are extremely radio-resistant. A great measure of protection for the enzymes is provided by the extraneous material in food. The irradiation energy taken up by foodstuffs is probably distributed between the enzyme and the other material present.

A decrease in radiosensitivity of enzymes, when enzyme-polymer complexes are formed, has been found for

1. chelated complexes such as carboxypeptidase A-cysteine (Fultz, 1968), catalase-glycerol (Lohmann et al., 1964) and catalase-hydrogen cyanide (Sutton, 1956),

2. enzyme-substrate complexes such as deoxyribonuclease I-deoxyribonucleic acid (Okada, 1957), polyphenol oxidase-tyrosin and actomyosin-adenosine triphosphate (Arnaud & Coelho, 1969),
3. covalent-bound complexes such as papain-poly-p-aminophenylalanine (Giovanazzi-Sermannini & Di Marco, 1965) and
4. unidentified complexes such as trypsin-agar (Holladay et al., 1966).

In biological material all the various complexes are likely, because of the diversity of the material. For intercellular yeast catalase, activity increases by irradiation of cell suspensions (Aronson et al., 1956). This phenomenon is accompanied by a significant lowering of the energy of activation for the catalase-hydrogen peroxide reaction and a considerable increase in the sensitivity of the enzyme to inactivation by heat. The explanation of the authors is, that ionization can lead to the dissociation of a ribonucleic acid-catalase complex and the release of the enzyme from the interface. A combined irradiation and heat treatment might therefore inactivate catalase in yeast cells very effectively.

Defining purpose of study Loss of enzyme activity by storage and irradiation of foodstuffs was simulated in systems of enzymes and polymers. For ease, the polymers used were water soluble. The enzymes were selected from different groups for their importance in food technology, availability in pure form and for easy assay (Table 1).

The purpose of the study was to look for a set of conditions which would increase the radiosensitivity of enzymes in foodstuffs. Augenstein (1962), in his excellent review, said: 'The radiosensitivity of a given enzyme is affected not only by temperature, pH, gases and the assay method employed, but more importantly by the nature of the system in which they are irradiated. Thus, analysis of the mechanisms of enzyme inactivation in heterogeneous systems, which more nearly simulate cellular conditions, is complicated by the effects of moisture content, the presence of other solutes and the possibility that the enzyme is adsorbed at an interface.'

Table 1. Data on the enzymes.

Enzyme	Molecular weight	Prosthetic group	Bound metal	Supplier	Assay
POD	40.000 ¹	+	Fe	Worthington	spectrophotometer
AP	100.000 ¹	—	Zn	Boehringer	spectrophotometer
PE	26.000 ²	—	—	Worthington	titrimeter

1. Dixon & Webb, 1965
2. Delincée & Radola, 1970

2 Definitions and abbreviations

Before irradiation the samples were freeze-dried and stored for three weeks. The residual enzyme activities were determined and plotted as a function of water activity. The curves obtained were a measure for the *storage stability* of the enzyme in question. The enzyme activity immediately before irradiation is called *the blank*.

The enzyme activities of the irradiated samples were expressed as a percentage of the blank and plotted as a function of the dose. The inactivation-dose relationship is exponential for many enzymes (Augenstein, 1962), but linear for papain (Sanner & Pihl, 1963) and neither exponential nor linear for α -amylase (Gorin et al., 1969). In my systems exponential inactivation was observed so that the D_{37} value could be used as an index for the radiosensitivity. The D_{37} value is the dose that leaves 37% of the original enzyme activity.

After storage of the solutions of the irradiated systems for a week in the refrigerator the D_{37} value was determined again and called the D'_{37} value. Then the *after-effect* could be defined as:

$$\frac{D_{37} - D'_{37}}{D_{37}} \times 100\%$$

In a general kinetic analysis of enzyme inactivation in solution by ionizing radiation, Sanner & Pihl (1963) defined the *intrinsic radiosensitivity* as the probability that a radical which becomes inactivated by the enzyme will concurrently inactivate the enzyme. For papain the authors worked out a graphic method which was used for processing the data of this study. The reciprocal of the slope of the curve $D_{37} = f$ (enzyme concentration), is a measure of intrinsic radiosensitivity. At water activities higher than 45% the samples contain 'free water', in which radicals can be generated by ionizing radiations. These samples can therefore be considered as *concentrated solutions*, and an intrinsic radiosensitivity can be calculated. For this purpose the enzyme concentration was taken as the reciprocal of the water content (% H₂O) of the system.

CMC : carboxymethylcellulose
NaP : sodium pectate
PVP : polyvinylpyrrolidone
sS : soluble starch

AP : alkaline phosphatase
PE : pectinesterase
POD : peroxidase

3 Methods

According to Table 2 an enzyme solution was added to a 2% polymer solution, after which the mixture was freeze-dried. Precautions were taken to avoid changes in pH of the enzyme-polymer systems during the handling period (Roozen & Pilnik, 1971a). The freeze-dried material was ground, divided between 42 polyethene bags, incubated for one week in a desiccator and then for a fortnight over saturated salt solutions. The bags were then sealed and the samples were irradiated up to $6 \cdot 10^6$ rad by a vertical beam Van de Graaf electron generator. Blue cellophane was used as a dosimeter.

To determine the enzyme activities, the samples were dissolved in distilled water and then stored overnight in a refrigerator. The assays used were standard methods (Roozen & Pilnik, 1970, 1971c, 1971d) which were adapted to process 50 assays per day.

For the various water activities the D_{37} values of the enzyme-polymer systems were calculated with the help of linear regression analysis ($\ln \%$ residual enzyme activity versus dose). The D_{37} values were plotted against the water activities but results with a correlation coefficient less than 0.9 were omitted. The intrinsic radiosensitivities were then calculated from linear regression analysis of the D_{37} values versus the reciprocals of the water contents.

As control the pH values of the solutions of the irradiated systems were also measured. The solutions of the irradiated systems were stored for a week in the refrigerator after which the D'_{37} values were determined and the after-effects calculated.

Table 2. Enzyme concentrations of the systems
(in mg enzyme per gram dry polymer).

Polymers	Enzymes		
	POD	PE	AP
		I	II
CMC	1	4	0.05
sS	1	4	0.05
PVP	1	4	0.05
NaP	-	4	-

4 Results

In addition to the results given in Table 3 some other aspects of the enzyme-polymer systems were studied.

Peroxidase systems At 25°C and the characteristic pH values of the polymers, big differences in the storage stability of POD were observed (Roozen & Pilnik, 1970). In the CMC systems POD had highest storage stability at the extremes of the water activities used. This enzyme was very unstable in the PVP systems and lost all its activity at high water activities. In the sS systems the activity of POD could be regenerated by water after incubation at the lower water activities.

The D_{37} values of the POD-polymer systems were calculated with the help of linear regression analysis. The extrapolated curves, found for the sS systems did not intersect at the blank point on the activity coordinate, whereas the curves of the CMC and PVP systems cut the activity coordinate in the blank point.

Pectinesterase systems For this enzyme the same methods and polymers were used as for POD. NaP-PE systems were also studied. In these systems one can expect enzyme-product-complexes (Jansen, 1960) that provide information about

Table 3. Results, obtained with enzyme-polymer systems, that illustrate influence of water activity on storage stability and radioresistance of enzymes.

System	Storage stability: residual activity curve ¹	Radioresistance: D_{37} -curve ²	Intrinsic radio-sensitivity ratio	After-effect
POD-CMC	minimum	maximum	3	≈ 40%
POD-sS	horizontal	descending	4	≈ 50%
POD-PVP	minimum	maximum	2	none
PE-CMC	descending	maximum	2	none
PE-sS	descending	maximum	2	none
PE-PVP	horizontal	ascending	—	none
PE-NaP	horizontal	maximum	1	none
AP-CMC	horizontal	maximum	3	≈ 25%
AP-sS	horizontal	maximum	5	≈ 30%
AP-PVP	horizontal	ascending	-4	none

1. % enzyme activity versus increasing water activity (shape of curve)

2. D_{37} -versus increasing water activity (shape of curve)

the radiosensitivity of enzymes in such complexes. The storage stability of PE depended little on the water activity of the systems studied. It is rather the nature of the polymer which determines the stability of the enzyme.

The \ln activity versus dose curves of all the PE systems studied cut the activity axis in the blank point. As mentioned before the forming of enzyme-polymer complexes has been proved in the NaP-PE systems (Jansen, 1960). Both facts indicate that in all systems the active PE molecules are complexed with polymer molecules.

In solutions of the sS systems, the pH decreased as a function of the radiation dose. Over the whole range the change of the pH was 0.2 units per 10^6 rad.

Alkaline phosphatase systems For this enzyme two different AP concentrations were chosen (Table 2) to obtain information about the influence of the concentration on the storage and irradiation behaviour of enzymes. The storage stability of AP was the same in the high and low concentration systems and was not strongly influenced by the water activity. The storage stability depends mainly on the nature of the polymers.

The dose versus \ln activity curves of all the AP systems studied can be extrapolated to the blank point on the activity axis. In AP systems, all the active enzyme molecules seemed to be complexed with the polymers. The calculated D_{37} values were plotted as a function of their water activities. The plots of both enzyme concentrations were similar to the corresponding plots of the PE systems. A 20-fold increase of the enzyme concentration increased the radioresistance (D_{37}) of these systems by a quarter.

PVP systems showed no after-effects. For the sS and CMC systems the after-effect depended on the enzyme concentration: it was about 25% for the low concentration systems and ranged from 5 to 15% for the high concentration systems.

As in the sS-PE systems the pH of the sS-AP systems was decreased by the action of ionizing radiation. Here the decrease of the pH was 0.07 units per 10^6 rad.

5 Discussion and conclusions

Radioresistance of the enzymes The radioresistance of an enzyme in an enzyme-polymer system was calculated from a linear regression analysis of the $\ln \%$ residual enzyme activities versus irradiation doses. The fact that the extrapolated curves of the sS-POD systems did not intersect at the blank point on the activity axis, whereas the curves of all the other enzyme-polymer systems cut the activity axis in the blank point can be explained as follows. In the sS-POD systems there are both bound and free enzyme molecules. The latter are much more radiosensitive than the bound ones, so that there is a discontinuity in the curve $\ln \%$ residual enzyme activity versus irradiation dose. In the other systems this discontinuity does not occur, because the active enzyme molecules are only present in a bound form.

In the systems used in this study it was found that the enzymes are least radioresistant at high water activities ('free water') except for the PVP systems of PE and AP. An explanation could be that water radicals are more reactive to the enzymes than to the polymers, but for the exceptions the water radicals are more reactive to PVP. The radioresistance of AP in its systems was increased in the higher enzyme concentration systems. This protective behaviour could be caused by a higher concentration of impurities including inactivated enzyme molecules. Therefore environmental factors strongly determine the radioresistance of an enzyme.

Intrinsic radiosensitivity As shown in Table 3 the intrinsic radiosensitivities of the enzymes studied in the CMC and sS systems are in the following order:

$$\text{PE} < \text{POD} < \text{AP}$$

The molecular weights of these enzymes have the same sequence. Although there were not sufficient data for establishing a clear relationship, a regression analysis was carried out for intrinsic radiosensitivity as a function of the molecular weight (Roozen & Pilnik, 1972). The intrinsic radiosensitivities of PE in the NaP systems and of the enzymes in the PVP systems are out of this order. These deviations could be caused by the forming of an enzyme-product complex (PE-NaP) and by the high reactivity of hydrated electrons with PVP (Hart, 1964). In the experiments of Okada (1957) the protective effect of proteins seemed to be a function of their respective molecular weights. His results imply that the number of reactive sites on the protein molecule increase with the size of the molecule. In analogy, in aqueous systems the intrinsic radiosensitivity of an enzyme is partly determined by its molecular weight (Roozen & Pilnik, 1972).

After-effects For some time after the actual irradiation of aqueous solutions of deoxyribonucleic acid with X-rays there is a slow decrease of viscosity which is called an after-effect. Daniels et al. (1953) considered this after-effect to be due to a slow hydrolysis of labile substances which were formed during the irradiation process. I found similar after-effects with the enzyme activity of dissolved enzyme-polymer systems. It is therefore possible that unstable enzyme molecules are formed during irradiation and that these molecules will be inactivated more quickly than the original ones.

Summary

The effect of ionizing radiation on the inactivation of enzymes was studied as a function of their moisture contents. For this purpose the following enzyme polymer solutions were freeze-dried: POD-CMC, POD-sS, POD-PVP, PE-CMC, PE-sS, PE-PVP, PE-NaP, AP-CMC, AP-sS and AP-PVP. The freeze-dried material was ground and then divided between 42 polyethene bags. The samples were incubated for one week in a desiccator and then for a fortnight over saturated salt solutions in order to obtain samples with the desired moisture content. These samples were then irradiated, dissolved and assayed for enzyme activity.

During the treatment before irradiation the stability of POD depends on temperature, moisture content, pH and the nature of the polymers. The same factors influence the radiosensitivity of POD. An after-effect of the irradiation was detected in the solutions of the CMC-POD and the sS-POD systems, but no after-effect was found in the solutions of POD-PVP systems.

Incubated over silica gel the PE systems lost minimum enzyme activity. In all systems the enzyme was least radiosensitive at 40% relative humidity except in the PVP system in which the enzyme decreased in radiosensitivity with increasing relative humidity.

In all the systems the radiosensitivity of AP had the same magnitude but over 40% relative humidity the sensitivity decreased in the PVP systems and increased in the CMC and sS systems. An increase in enzyme concentration decreased the radiosensitivity of AP. There was no after-effect in the solutions of the AP-PVP systems, while the after-effects were more pronounced in the solution of the CMC and sS systems with the lower enzyme concentrations.

At neutral pH, the intrinsic radiosensitivities of the studied enzymes in the CMC and sS systems were in the proportion of:

$$\text{PE} < \text{POD} < \text{AP}$$

The molecular weights of these enzymes have the same sequence. Therefore a regression analysis was carried out from the intrinsic radiosensitivity as a function of the molecular weight. The intrinsic radiosensitivities of PE in the NaP systems and of the enzymes in the PVP systems were out of the order. These deviations could be caused by the formation of an enzyme-product complex (PE-NaP) and by the high reactivity of hydrated electrons with PVP.

References

- Acker, L., 1962. Enzymic reactions in foods of low moisture content. *Adv. Food Res.* 11: 263-330.
- Acker, L. & E. Lück, 1958. Über den Einfluss der Feuchtigkeit auf den Ablauf enzymatischer Reaktionen in wasserarmen Lebensmitteln. *Z. Lebensmittel-Unters. u. -Forsch.* 108: 256-269.
- Alexander, P. & A. Charlesby, 1957. Effect of x-rays and gamma rays on synthetic polymers in aqueous solution. *J. Polymer Sci.* 23: 355-375.
- Arnaud, Y. & R. Coelho, 1969. Rôle des substrats dans l'irradiation des enzymes. *Bull. Soc. Chim. Biol.* 51: 275-282.
- Aronson, D. L., M. J. Fraser & C. L. Smith, 1956. Enzyme alteration by ionizing radiation. *Radiat. Res.* 5: 225-237.
- Augenstein, L. G., 1962. The effects of ionizing radiation on enzymes. *Adv. Enzymol.* 24: 359-413.
- Bacq, Z. M. & P. Alexander, 1966. Fundamentals of radiobiology. Pergamon Press, Oxford.
- Chapiro, A., 1964. Radiation chemistry of polymers. *Radiat. Res. suppl.* 4: 179-191.
- Daniels, M., G. Scholes & J. Weiss, 1953. 'After-effect' in aqueous solutions of deoxyribonucleic acid irradiated with x-rays. *Nature* 171: 1153-1154.
- Delincée, H. & B. J. Radola, 1970. Some size and charge properties of tomato pectin methyl-esterase. *Biochim. Biophys. Acta* 214: 178-189.
- Dixon, M. & E. C. Webb, 1965. Enzymes. Longmans, Green and Co. Ltd., London.
- Fultz, E. D., 1968. The role of metal chelation in chemical radiation protection of enzyme systems, with application to carboxypeptidase A. *Radiat. Res.* 34: 544-554.
- Giovannozzi-Sermanni, G. & G. Di Marco, 1965. X-ray inactivation of papain coupled with a synthetic polypeptide. *Radiat. Res.* 26: 63-68.
- Gorin, G., S. F. Wang & T. D. Seth, 1969. Radiolysis of α -amylase in dilute solution, a non-exponential inactivation-dose relationship. *Int. J. Radiat. Biol.* 16: 93-97.
- Hart, E. J., J. K. Thomas & S. Gordon, 1964. A review of the radiation chemistry of single-carbon compounds and some reactions of the hydrated electron in aqueous solution. *Radiat. Res. suppl.* 4: 74-88.
- Holladay, W., E. Augenstein & L. Augenstein, 1966. The effects of added agar on the radiation sensitivity of trypsin in solution. *Radiat. Res.* 28: 132-142.
- Jansen, E. F., R. Wang & J. Bonner, 1960. Orange pectinesterase binding and activity. *J. Food Res.* 25: 64-72.
- Lett, J. T. & P. Alexander, 1961. Crosslinking and degradation of deoxyribonucleic acid gels with varying water contents when irradiated with electrons. *Radiat. Res.* 15: 159-173.
- Lohmann, W., A. J. Moss, W. H. Perkins & C. F. Fowler, 1964. Intermolekulare Wechselwirkung zwischen Glyzerin und Katalase, Einfluss auf die Strahlenempfindlichkeit des Enzyms. *Biophysik* 2: 16-21.
- Marples, A. & G. Glew, 1958. The effects of ionizing radiations on enzymes. Atomic Energy Research Establishment, Harwell. I/R 2726: 2-26.
- Münzel, F. & J. Schenkel, 1964. Bestrahlung von Cellulose und Cellulosederivaten. *Radio-nuklide*, 141-164.

- Okada, S., 1957. Inactivation of deoxyribonuclease I by x-rays. III. Modification of the effect of radiation on the deoxyribonucleic acid - deoxyribonuclease system. Arch. Biochem. Biophys. 67: 113-120.
- Purr, A., 1970. Zum Ablauf chemischer Veränderungen in wasserarmen Lebensmitteln. II. Versuche mit Lipoxygenase sowie enzymatischen Wirkstoffgemischen und deren Auswirkung auf den autoxydativen Fettverderb in Trockenprodukten in Abhängigkeit von der Gleichgewichtsfeuchtigkeit. Fette-Seifen-Anstrichm. 72: 725-732.
- Reuschl, H. & A. Guilbot, 1966. Der Einfluss von Gammastrahlen auf Kartoffelstärke in Abhängigkeit von ihrem Wassergehalt. Stärke 18: 73-77.
- Roozen, J. P. & W. Pilnik, 1970. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. I. Die Stabilität von Peroxydase bei 25°C. Lebensm.-Wiss. u. Technol. 3: 37-40.
- Roozen, J. P. & W. Pilnik, 1971a. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. II. Der Einfluss von pH-Verschiebungen auf die Stabilität von Peroxydase bei 2°C. Lebensm.-Wiss. u. Technol. 4: 24-26.
- Roozen, J. P. & W. Pilnik, 1971b. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. III. Der Effekt von Elektronenbestrahlung auf die Stabilität von Peroxydase. Lebensm.-Wiss. u. Technol. 4: 93-96.
- Roozen, J. P. & W. Pilnik, 1971c. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. IV. Der Effekt von Lagerung und Elektronenbestrahlung auf die Stabilität von Pektinmethylsterase. Lebensm.-Wiss. u. Technol. 4: 158-162.
- Roozen, J. P. & W. Pilnik, 1971d. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. V. Der Effekt von Lagerung und Elektronenbestrahlung auf die Stabilität von alkalischer Phosphatase. Lebensm.-Wiss. u. Technol. 4: 196-200.
- Roozen, J. P. & W. Pilnik, 1972. Über die Stabilität adsorbiertes Enzyme in wasserarmen Systemen. VI. Möglichkeiten zur Inaktivierung von Enzymen in getrockneten Lebensmitteln durch Bestrahlung. Betrachtungen im Lichte der Literatur und eigener Untersuchungen. Lebensm.-Wiss. u. Technol., in press.
- Sanner, T. & A. Pihl, 1963. Kinetics of enzyme inactivation by ionizing radiation. Radiat. Res. 19: 12-26.
- Sutton, H. C., 1956. Effects of radiation on katalase solutions. II. The protection action of complexing agents. Biochem. J. 64: 456-460.
- Wahba, I. J. & L. M. Massey, 1966. Degradation of pectin by gamma-radiation under various moisture conditions. J. Polymer Sci. A 4: 1751-1771.