

## Distribution and transport of Fennoscandinavian indicators

A synthesis of data from the literature leading to a reconstruction of a pattern of flowlines and ice margins of the Scandinavian ice sheets

CENTRALE LANDBOUWCATALOGUS



0000 0086 9822

Dit proefschrift met stellingen van Cornelis Johannes Overweel, doctorandus in de geologie, geboren te Haarlem op 25 april 1921, is goedgekeurd door de promotor, dr. J. D. de Jong, hoogleraar in de geologie aan de Landbouwhogeschool te Wageningen en de tweede promotor, dr. G. C. Maarleveld, hoogleraar in de fysische geografie van periglaciaire en glaciale landschappen aan de Universiteit van Amsterdam.

De Rector Magnificus  
van de Landbouwhogeschool,  
H. C. van der Plas

Wageningen, 16 september 1977

HN 0204

701

701

III

C. J. Overweel

# Distribution and transport of Fennoscandinavian indicators

A synthesis of data from the literature leading to a reconstruction of a  
pattern of flowlines and ice margins of the Scandinavian ice sheets

Proefschrift

ter verkrijging van de graad van  
doctor in de landbouwwetenschappen,  
op gezag van de rector magnificus,  
dr. H. C. van der Plas,  
hoogleraar in de organische scheikunde,  
in het openbaar te verdedigen  
op woensdag 23 november 1977  
des namiddags te vier uur in de aula van de  
Landbouwhogeschool te Wageningen

## STELLINGEN

1

Het genetisch uitgangspunt maakt de classificatie van gesteenten onduidelijk en ondoorzichtig.

2

Het calciumphosphaat van het tandemail van het levende organisme neemt geen fluor op van fluoriden in drinkwater of tandpasta.

Gezondheidsraad 1960. 'Cariës preventie met fluoriden'. Verschl. Med. Volksgez. 7.

3

Zoals onder andere gebleken uit het onderzoek naar fossiele beenderen op de bodem van de Zeeuwsche wateren is het Pretiglien niet in de Westerschelde ontsloten.

Vlerk, I. M. van der & F. Flörschütz, 1953. 'The palaeontological base of the subdivision of the Pleistocene in the Netherlands.' Verh. Kon. Ned. Akad. Wetensch. afd. Nat., 1e r., 20, no 2: 1-58.

4

Wanneer de noodzakelijke registratie van museumcollecties geschiedt volgens conventionele methoden worden meer manuren bespaard dan bij het registreren met de computer.

Germraad, J. H., et al., 1972. 'A computer based registration system for geological collections.' Scripta Geol., 9: 1-12.

5

De Röntgenografische toepassing van de fluordateringsmethode van fossiele beenderen, die op het ionen uitwisselingsmechanisme berust waarbij hydroxylapatiet wordt omgezet in fluorapatiet, is een illusie.

6

De algemene bewegingsrichtingen van het Scandinavische ijs tijdens het Weichselien en het Saalien stemmen met elkaar overeen.

7

De door Torrel in 1865 geïntroduceerde Baltische ijsstroom heeft nooit bestaan.

Charlesworth, J. K., 1957. 'The Quaternary era, Vol. 2.' E. Arnold Ltd, London: 595-1700.

8

Gedurende het Weichselien is het Scandinavische ijs noch tot de Shetland of Orkney eilanden, noch tot Schotland doorgedrongen.

9

Waar het doubleren bij het AVO is afgeschaft wordt het kind dat niet wordt bevorderd thans meedogenlozer aangepakt dan voorheen.

Stellingen, behorende bij: C. J. Overweel, Distribution and transport of Fennoscandinavian indicators; A synthesis of data from the literature leading to a reconstruction of a pattern of flowlines and ice margins of the Scandinavian ice sheets.

Aan de nagedachtenis van mijn vader.  
Aan mijn moeder.

## Samenvatting

Op grond van de voornaamste algemene ijsbewegingsrichtingen van het laatste glaciaal in Fennoscandinavië zijn de volledig gepubliceerde zwerfsteen tellingen uit gebieden ten zuiden en ten oosten van de Oostzee geheel opnieuw bewerkt. Uit deze herbewerking is een distributiepatroon van gidsgesteenten voortgevloeid, dat ten dele samenvalt met het patroon van de voornaamste richtingen van ijsbeweging. Dit geldt zowel voor het Weichselien als het Saalien. Het andere deel van de distributie behelst gesteenten van Baltische oorsprong.

Een verdere analyse en uitwerking van het verband tussen het distributie- en richtingspatroon leidde tot een reconstructie van bewegingsrichtingen en omstrekkervormen van het Fennoscandinavische landijs. Het perifere afwateringssysteem is met deze reconstructie verenigd tot één geheel en de hiermee samenhangende mogelijkheid dat drijvend ijs zwerfstenen van Baltische oorsprong langs de oerstroomdalen en over de glaciale meren zou kunnen hebben vervoerd is nader uitgewerkt.

Op verzoek van het College van Dekanen van de Landbouwhogeschool te Wageningen, volgt hier een beknopt curriculum vitae van de schrijver.

Hij werd op 25 april 1921 te Haarlem geboren. Zijn middelbare schoolopleiding genoot hij aan het Christelijk Lyceum te Bandung, Java, waar hij in mei 1941 met goed gevolg het eindexamen 5-jarige HBS-B aflegde.

In september 1941 werd hij als student ingeschreven aan de Landbouwhogeschool in het toenmalige Batavia (Djakarta) waaraan hij tot begin december 1941 studeerde. Van die datum af tot aan maart 1947, was zijn studie onderbroken door het vervullen van zijn militaire dienstplicht in actieve dienst, tijdens de tweede wereldoorlog.

In oktober 1947 hervatte hij de studie aan de Landbouwhogeschool te Wageningen en werd hij door de colleges en practica in geologie van Prof. Dr. D. J. Doeglas zodanig geboeid, dat hij zich in oktober 1948 liet inschrijven aan de Rijksuniversiteit te Groningen om zich verder in die studierichting te bekwaamen. Hij mocht zijn studie verrichten onder leiding van de hoogleraren Dr. Ph. H. Kuenen en Dr. P. Terpstra en behaalde 28 maart 1950 het candidaatsexamen Wis-, en Natuurkunde (letter i). Hij vervolgde zijn studie in Leiden onder leiding van de hoogleraren: Dr. B. G. Escher, Dr. E. Niggli, Dr. L. U. de Sitter en Dr. I. M. van der Vlerk en verrichtte veldwerk onder leiding van Dr. J. D. de Jong, toen geoloog bij de Rijks Geologische Dienst te Haarlem. Op 2 juni 1953 werd met goed gevolg het doctoraalsexamen geologie afgelegd, met als eerste bijvak palaeontologie en als tweede bijvak petrologie.

Na een korte periode als volontair bij de Geologische Dienst te Haarlem, was hij van maart 1954 tot mei 1957 als mijngeoloog bij de Cerro de Pasco Corporation in Peru werkzaam. Van juli 1957 tot september 1974 was hij verbonden aan het Rijksmuseum van Geologie en Mineralogie, te Leiden, als conservator in de petrologie. Thans is hij als petrologisch en mineralogisch adviseur werkzaam bij het Instituut voor Prehistorie van de Rijksuniversiteit te Leiden.

Alhoewel het zwerfsteen beheer één van de zaken was die onder zijn verantwoording vielen tijdens zijn museum periode, begonnen de plannen voor het onderhavige speciale onderzoek eerst omstreeks 1972 vorm te krijgen en werd, na enkele gesprekken met Prof. Dr. J. D. de Jong, die zich zeer voor dit onderwerp interesseerde, in begin 1973 aan het eigenlijke promotie onderzoek begonnen. Het werd uitgevoerd onder leiding van Prof. Dr. J. D. de Jong aan de Landbouwhogeschool te Wageningen als promotor en Prof. Dr. G. C. Maarleveld, hoogleraar aan de Universiteit van Amsterdam, als tweede promotor.

# Distribution and transport of Fennoscandinavian indicators

A synthesis of data from the literature leading to a reconstruction of a pattern of flowlines and ice margins of the Scandinavian ice sheets

C. J. Overweel

Overweel, C. J., Distribution and transport of Fennoscandinavian indicators: A synthesis of data from the literature leading to a reconstruction of a pattern of flowlines and ice margins of the Scandinavian ice sheets. — Scripta Geol., 43: 1-117, 54 figs., 1 folding map, Leiden, October 1977.

Detailed indicator counts in the literature on areas south and east of the Baltic were reevaluated on the basis of the main ice movement directions of the last glacial in Fennoscandia. This revaluation resulted in a distribution pattern of indicators which partially coincides with the general trends of ice movement. This holds for both the Weichselian and the Saalian. The other part of the indicator distribution is of Baltic origin.

The interrelationship found between flowline and distribution pattern was further analysed and elaborated, which led to a reconstruction of the directions of ice movement and outline forms of the Scandinavian ice sheets. The peripheral glacier-margin drainage was incorporated into the reconstruction, and the resulting possibility that drift ice could have transported indicators of Baltic origin through the Urstromtäler and the glacial lakes is discussed in some detail.

C. J. Overweel, Instituut voor Prehistorie der Rijksuniversiteit Leiden, Breestraat 87, Leiden, The Netherlands.

Introduction	2
Acknowledgements	3
Part 1. Areal distribution of indicators	4
History and complexity of indicator investigation	4
Indicator identification	6
Indicator fans	8
The quantitative approach	9
Determination of generalized mean trends of ice flow	9
Considerations	12
Regrouping of provenances	13
Criteria for the estimation of the directions of ice flow	13
Flowline pattern in Fennoscandia	17
Source areas adapted to ice flow directions	19
Rearrangement of indicator counts	21
The Weichselian	21
The Saalian	31

Resulting distribution pattern of indicators	43
Correspondence between the Weichselian and the Saalian distributions	43
The double-peaked character of the indicator diagrams	46
The Weichselian distribution and flowline pattern in Fennoscandia	47
Corresponding directions of ice movement in the Weichselian and the Saalian	48
<b>Part 2. Interpolation of indicator transport between distribution and source areas</b>	<b>49</b>
The inconsistent distribution unit	49
The force of the consistent and inconsistent ice movements	49
The enigmatic Baltic ice stream	52
Waterborne transport	54
The consistent distribution unit	61
Further analysis and elaboration of the flowline pattern	61
The inferred structure of flowlines in relation to the dispersal of indicators	71
Geometrical aspects	85
Integration of consistent and inconsistent distribution processes	93
The Weichselian recession in Denmark	93
Ice halts in the Baltic water basin	101
Conclusions	110
References	111

## Introduction

In 1957 the study of Fennoscandinavian indicators became the author's concern as part of his assignment as the keeper of rocks at the Rijksmuseum van Geologie en Mineralogie (National Museum of Geology and Mineralogy) in Leiden. He began to acquaint himself with the indicators in the museum and those of some large, well-founded, private collections. With others and alone, he visited field occurrences of indicators in The Netherlands, West Germany, Denmark, and England, and also examined the outcrops in source areas in south Sweden.

During this introductory period it became clear that the investigation of Fennoscandinavian indicators was still in the stage of data collecting, and consisted mainly of the analysis of the indicator association per locality. Attempts had been made to correlate the results regionally on distribution maps, as exemplified by the chart of ter Wee (1962) of the Saalian ice front phases in The Netherlands. The localities of these counts are indicated here by Hesemann numbers. Although a distribution pattern can be distinguished, these numbers do not help us to visualize the relationship with the source areas. The distribution pattern is much more pronounced on, for instance, the porphyry map of Denmark: Table I of K. Milthers' publication (1942). Here, too, the relationship between the distribution

pattern, ice movement, and provenance is difficult to grasp, but the zones of the tripartite indicator distribution in Lithuania given by Tarvydas (1960) trend toward their sources.

Thus, the quantitative approach of indicator investigation resulted in numberless counts, which are generally represented as separate, unrelated data. With the use of different procedures, some solitary regional correlations were made, but the overall picture remained one of separate and independent observations.

To advance indicator investigation, these loose entities needed to be fitted together. To achieve this, the counts based on different methods had first to be reduced to a common denominator. The Hesemann method and Lüttig's TGZ estimations were available, but these procedures did not yield a very clear picture of the indicator distribution pattern, perhaps because these approaches are too statistical since glacial control was not taken into account.

Therefore, I developed a new strategy, reckoning with glacial aspects. Thus, Part 1 of this paper gives a detailed regional analysis of the indicator distribution with reference to the traces of former ice movements in Fennoscandia. This analysis provides data from which a correlation was attempted of the published indicator counts of the circum Baltic. The resulting distribution pattern has two components, one corresponding to the Fennoscandian main ice flow trends, the other lacking this correspondence.

In the light of these findings, the relationships of the distribution pattern with glacial, periglacial, geomorphological, and geometrical features were investigated, as described in Part 2. The total forms a synthesis, a functional representation, of the Fennoscandinavian ice sheets.

#### ACKNOWLEDGEMENTS

Foremost among those to whom I am indebted are Prof. J. D. de Jong and Prof. G. C. Maarleveld, who gave me their experienced guidance, constructive criticism, and valuable advice. Their aid when necessary, their encouragement, and their enlightening discussions contributed greatly to the ultimate result.

In 1957, the late Prof. I. M. van der Vlerk, in his capacity as Director of the Rijksmuseum van Geologie en Mineralogie, asked me to set up a section on northern crystalline erratics in the museum. He gave me full liberty in this commission for which I have always been grateful. He advised me to get in touch with the non-professional prominents in this field and to visit the parent rocks in Fennoscandia.

My contacts with the experts in crystalline erratics in The Netherlands have been very helpful indeed. Their cooperation, their readiness to communicate and to take me over their collections, will be always remembered. They contributed greatly to my present knowledge of indicator identification. Special mention in this respect should be made of the late Mr P. van der Lijn, Mr A. P. Schuddebeurs, and Mr J. G. Zandstra. The instructive excursions of the Nederlandse Geologische Vereniging, conducted by Messrs. W. F. Anderson and A. G. Koenderink, gave me a better understanding of the Quaternary geology of the countries adjacent to The Netherlands. My friend and colleague Mr M. W. ter Wee was so kind as to take me over the exposures in The Netherlands to acquaint me with the Saalian.

I am indebted to the Netherlands Organisation for the Advancement of Pure Research (Z.W.O.) for financial support to visit the southern part of Sweden to

study the parent rocks of the indicators in 1959. Besides this opportunity to study the source rock, I benefited greatly from the help given me by, and the enlightening discussions with the Swedish geologists, especially Dr E. Åhman, Prof. S. Gavelin, and Dr P. H. Lundegårdh, who took me to the field and allowed me to join them on the excursions to the Vestervik archipelago in preparation for the 21 International Geological Congress in 1960.

I am indebted to Prof. E. den Tex and the other members of the Geologisch en Mineralogisch Instituut in Leiden for their hospitality. Prof. P. Hartman was kind enough to read the section on the geometrical aspects and gave me valuable advice. The most essential part of this paper are the figures, maps, and diagrams which elucidate the text, which were prepared by Messrs. J. Bult and F. J. Fritz. I am immensely grateful to them for their cooperation, skill, patience, and helpful advice. The librarians Messrs. C. H. Duvekot, O. J. Groot, and H. B. van der Meer gave me every possible assistance.

I am indebted to Mrs D. de Munter for correcting the English of large parts of the manuscript in its initial stage, and to Mrs I. Seeger for the final correction of the completed manuscript.

A word of special thanks is due to Prof. P. J. R. Modderman and his coworkers of the Instituut voor Prehistorie in Leiden for the way in which they accepted me in their midst, for the many instructive discussions about prehistory in particular and scientific inquiry in general, and for the chance they gave me to bring the present study to a close.

Finally, I cannot adequately express my gratitude for the aid and encouragement given by my wife, who undertook the preparation and typing of the manuscript. Without the support and loyalty of my wife and children, this study would probably never have been completed.

## Part 1. Areal distribution of indicators

### History and complexity of indicator investigation

To find out more about the shape and dynamics of the Quaternary ice sheets it is necessary to appraise the glacial deposits stratigraphically and to unravel the spatial trends of the ice stream movements. On this basis, the results of about two centuries of research on the dispersal of erratics will be discussed.

Each of the two objectives requires its own, specific approach, as shown by a discussion of Fig. 1. Figure 1a gives a schematic representation of a segment of an ice sheet flowing over outcrops of various parent rocks. An imaginary vertical plane has been indicated parallel to the direction of the stream and such that its line intersecting the subglacial surfaces marks an intervening boundary between different types of source rocks. On one side of this boundary line rock types A and B are present, on the other rock types C and D. Since the intervening boundary lies parallel to the direction of the stream, the highest densities of the indicators




Fig. 1. Schematic representations of segments of an ice sheet. Source areas separated by lines parallel (a) and transverse (b) to the flow direction.

deriving from source rocks A and B will be found on one side of the produced part of the intersection line and the indicators deriving from source rocks C and D on the other side. Hence, when the respective areas of provenance are situated between lines running parallel to the direction of the ice movement, the distribution of the indicators in the area of deposition permits the drawing of conclusions concerning the general trend of the ice movement.

If, however, the source areas are separated by the base lines of vertical planes taken at right angles to the direction of flow (Fig. 1b), differences in the quantities of the indicators in areas situated far enough from the source, have a bearing on the duration and the rate of ice flow. The one region may have been covered during a relatively short period, the other region for a much longer period. This is closely related to such conditions as the duration of halts of the ice front and the rate of expansion or recession of the ice sheet. Thus, unless unknown factors played a role,

the greater number of indicators derived from area E as compared with areas F and G (Fig. 1b) points to a longer period of ice erosion in area E.

In sum, the fact that the ice sheets in different glacials change in different ways means that provenances enclosed by lines at right angles to the direction of ice movement can be used for stratigraphic purposes. Contrarily, provenances contained by lines parallel to the direction of ice movement are required to unravel the trend of ice movement in relation to the area of deposition.

With these observations on the objectives of this field of research in mind, we are better equipped to evaluate the present state of affairs, which will be discussed in terms of the historical approach.

#### INDICATOR IDENTIFICATION

The advance of research during the last hundred and forty years can be roughly divided into three phases, i.e., indicator identification, reconstruction of indicator fans, and application of the quantitative approach.

The initial stage, which started around 1830 and culminated in the early part of the twentieth century, is characterized by qualitative work. One of the pioneers in The Netherlands was S. J. Brugmans, who as early as 1781 proved in his doctoral dissertation that rock species occurring in the province of Groningen were related to Fennoscandia (Capadose, 1825). (This pioneer was brought to my attention by Professor L. D. Brongersma). The erratics found in the glacial deposits of northwestern Europe were described, classified, and compared with their possible counterparts from the solid bedrock in Fennoscandia. These investigations resulted in several standard works on the petrography of Fennoscandinavian crystalline indicators, such as those by Korn (1927), Hesemann (1936, 1939b, 1975), and van Calker (1912), as well as publications by Kruizinga (1918), Hucke (1917, 1967), and several articles by Martin published in the last quarter of the nineteenth century (brought to my attention by Professor L. D. Brongersma) concerning the sedimentary rocks. In this phase contributions of lasting importance were made by non-professional geologists who — working alone or as members of associations — collected, described, and compared erratics from the glacial deposits in their own region. It is worth mentioning here some of the periodicals issued by these enthusiastic societies: *Grondboor en Hamer* in The Netherlands and the *Zeitschrift für Geschiebeforschung* and *Der Geschiebe Sammler* in Germany.

Because indicator investigation is based on indicator identification, consideration must be given to the question of the extent to which a given indicator is a true specimen of the source to which it refers. Discussion of this problem will be more to the point if we start by asking ourselves what the main features of an ideal indicator are. By definition, an ideal indicator should be a unique, easily recognizable rock type whose area of origin is clearly defined. Good examples of this among the relatively small number of groups of Fennoscandinavian indicators, are the typical Oslo rocks and the group of rapakivi granites from Finland and the Åland islands. These externally striking specimens immediately catch the eye of a collector searching through the lag concentrates of till. Due to differential weathering of the component parts, the surface of the indicator is usually more vividly marked than that of the original source rock, but in any case the fresh fracture of a broken specimen bears comparison with the rough unweathered faces of the Fennoscandinavian equivalents.

The macroscopic similarity of indicator and possible source rock can be evaluated by microscopical examination. In addition to a macroscopic description, Hesemann (1936, 1975) also gave a microscopic specification of the indicator types he described. Van Calker and his school set a high value on the microscopical approach. Once the identity of an ideal indicator, e.g. as a rhomb-porphyry, is established both macroscopically and microscopically, macroscopic determination of other specimens of this kind suffices.

As mentioned by Korn (1927), another method of further verification of an indicator can be obtained by determining whether members of coexisting rock types in the source area of a certain indicator also occur as components of the indicator association. Thanks to Milthers' counts it is well known that the northern part of Jutland is strewn with rhomb-porphyrries. Nonetheless, in visiting this region in the summer of 1971, the present author was impressed by the huge amount of these porphyries on the western beaches. In addition to the rhomb-porphyrries there are many larvikites, albeit in distinctly smaller amounts. This indicator assemblage of typical Oslo rocks in the erratic association of northern Jutland confirms the identity of the rhomb-porphyrries. On the other hand, the porphyries in their turn confirm the authenticity of the larvikites.

By reason of their rarity as rock types, the Oslo rocks and the rapakivis cause few problems as to their reliability as indicators. The distribution of the bedrock of the former is restricted to the Oslo area and that of the latter to the Åland islands, the sea-bed around these islands, and the southern part of Finland, so that the condition of a limited regional extension of the rock types is also satisfied, although this holds more strongly for the Oslo rocks than for the rapakivis.

Would further subdivision of these categories of indicators be useful? This depends on the problem under study.

1. For proving the authenticity of a given kind of indicator, collation of the divergences of the single specimens with variations of its source rock leads to greater certainty. However, use of this elaborate approach, especially when it requires microscopical investigation, will only make sense in some essential cases of dubious identity.
2. Subdivision into varieties of a kind of indicator whose main dispersal is confined to a restricted area of deposition, might provide more information about the directions and behaviour of movement of the ice streams that traversed the region. Hence, it could be useful to divide the rhomb-porphyrries or the larvikites in northern Jutland into subgroups, but only if the source rocks of these varieties occur in separate, distinctly defined areas.
3. As opposed to local investigations where the main or overall features of the Scandinavian ice sheets are concerned, there is little need to carry out further subdivision of indicators to obtain information about areas outside of their centres of accumulation. For instance, subdivision of rhomb-porphyrries or larvikites, outside their area of distinct concentration in northern Jutland, does not give much additional data about the general features of the ice sheet.

After these general remarks on the reliability of the identification of indicators from the Oslo area and the rapakivis, some other important groups of indicators will be dealt with. For more extensive and detailed information, the reader is referred to the publications of Hesemann (1936, 1975).

The third group of indicators to be considered are those from Dalarna. The Bredvad porphyries in the glacial deposits not only show macroscopic and micro-

scopic correspondence with their source rock but also predominate over the other varieties of Dalarna porphyries and porphyrites, which strongly supports their reliability as indicator. V. Milthers (1909) pointed out that the Bredvad porphyry occurs more frequently than the other Dalarna porphyries in Denmark, which he explained by its greater distribution in the source area. According to Hesemann (1936) the Bredvad porphyry predominates in Germany too, and van der Lijn (1974) found it to be the most common of the Dalarna group in The Netherlands. The association of the Dalarna porphyries in the glacial deposits with the Jotnian sandstones, Grönklitt porphyrite, and Garberg granite porphyry, which also crop out in Dalarna, supports their reliability as well.

The Småland granites and porphyries form a fourth group of indicators. The average Småland granite is medium or coarse grained and has a yellowish to reddish-brown hue, determined by the colour of its conspicuous microcline-perthites. This granite is a characteristic indicator. The source area in the eastern half of the southern part of Sweden is roughly defined in the south by a line running slightly north of Karlshamn to Karlskrona; on the west by a line from Karlskrona to the Vättern lake, on the north from lake Vättern to Västervik, and on the east by the coast south of Västervik. Of the Småland porphyries, the granite-porphyries series from Påskallavik, Sjögelö, and Emarp, with their marked alkali-feldspar phenocrysts, measuring 1-3 cm, predominate as indicator.

A fifth group is formed by the so-called black and white granites from Stockholm and Uppland. The former is a fine-grained grayish-black, biotite-granite, of which there is also a red variety. The Uppland granites are granodiorites containing hornblende. The outcrops of the Bohuslän granite on the west coast of Sweden contain subordinate bodies of a granite variety that could be mistaken for the Stockholm granite. As indicator, the typical Bohuslän granite with its porphyric crystals of microcline is seldom found in The Netherlands or northern Germany. Consequently, the subordinate variety resembling the Stockholm granite should be even rarer in these countries and because of its rarity as erratic, there is little chance of confusion with indicators of the true Stockholm granite.

The easily recognizable, red-spotted granular gneis granites from the island of Bornholm may be mentioned as a sixth group of indicators.

#### INDICATOR FANS

As qualitative stock-taking continued, several investigators began to relate the localities of these indicators with their source areas, which brings us to the second phase of the research, in which the sorting and reconstruction of indicator fans ranked above the qualitative classification of the individual rocks. A now famous compilation of these fans was given by Caldenius and Sandegren in a map in the Atlas of Finland (1910), which has been reproduced in many textbooks on Quaternary or glacial geology (e.g. Magnusson, Lundqvist & Regnell, 1963). The reconstruction of these fans was the first logical attempt to unravel the general trends of ice movements of the Fennoscandinavian ice sheets. In a general sense the compiled directions on this map are radial, but strongly disturbed by fans intersecting the general radial pattern. The apices of most of these interfering fans are situated in the Baltic area. It is hardly surprising that the enigmatic configuration of the line-bundles of the indicator fans gave rise to controversy about the ice stream directions. In addition, the fact that these directions could be attributed not just to one but to several ice ages caused further confusion.

## THE QUANTITATIVE APPROACH

To find a way out, the problem was taken in hand by supporters of the quantitative approach, which characterizes the third phase of investigation. An enormous number of counts were made, but in three different ways, according to Madsen, Hesemann, and Milthers.

For Madsen's method (Madsen, 1928), a 10 kg dry boulder-clay sample is passed through a 6 mm-mesh sieve. The erratics in the residue are counted and subdivided into a number of major systematic groups such as crystalline rocks, limestones, dolomites, and flintstones, after which the proportion of flints to crystalline erratics is determined. This ratio or flint coefficient is used as an index to distinguish glacial deposits of different ages. Since the source areas are left out of consideration, Madsen's method may be considered purely stratigraphic.

In contradistinction to the methodology of Madsen, the source areas are a condition *sine qua non* in the counting procedures of Hesemann (1930) and K. Milthers (1942). Hesemann divided Fennoscandia into four areas, which he designated I to IV. The indicators from a given locality are grouped accordingly, and their percentages in tens are collocated in conformity with the identification numbering of the selected areas in what is called the Hesemann number. Milthers confined himself to porphyry counts from three source areas, namely the rhomb-porphries of the Oslo district, the porphyries from Dalarna, and the Baltic porphyries. These instigators and their adherents performed numerous counts, and as a result we now have at our disposal a dense regional network of counts covering Denmark, West and East Germany, and The Netherlands.

Woldstedt and Duphorn (1974) made a general estimate of the stratigraphic value of different methods of indicator investigation. The counts after Milthers, and particularly those of Hesemann, proved to be useful, inasmuch as different glacials and subglacials are characterized by specific, prevailing Hesemann numbers. On the basis of Woldstedt (1955), east Fennoscandinavian indicators culminate in the Elster, the numbers fluctuating round 6310. Numbers such as 2170, 2260, or 1180, bear upon the Saalian moraines. Similar numbers of the first three source areas such as 4330 or 3340 are characteristic of the Warthe Stage. To a certain extent, the numbers of the Weichselian are the same as those of the Warthe, but considerable deviations occur frequently here.

## DETERMINATION OF GENERALIZED MEAN TRENDS OF ICE FLOW

*Hesemann's method*

We have seen that Hesemann's counts have stratigraphic value, and that provenances enclosed by lines perpendicular to the direction of ice movement are useful for stratigraphic purposes. Conversely, use of the common trend of the boundary lines of Hesemann's source areas should make it possible to arrive at an approximation of the direction of the ice flow toward The Netherlands and the western and eastern parts of Germany, where most of his counts were made. Since three of the four provenances proved to be involved, only the dividing lines between areas I, II, and III should be considered for the reconstruction. Because these lines are curved, straight lines were superimposed. In Fig. 2 the direction in question has been drawn perpendicular to these auxiliary lines, giving a generalized,




Fig. 2. Generalized mean ice stream directions tentatively reconstructed on the basis of the trends shown by the boundary lines of Hesemann's source areas.

mean ice flow direction running roughly from NE to SW. When source areas are separated by a line running parallel to the ice flow, the highest density zones of their indicator distribution are expected on either side of this line or its extension. Since the accumulation loci of the Oslo indicators lie on the west side of the extension of the line dividing Hesemann's areas IV and III and the main distribution of the Swedish indicators occurs on the east side, this dividing line should be parallel to the just estimated flow direction, which, as Fig. 2 shows, is in a general sense the case.

Hence, besides its stratigraphic merits, the Hesemann method yields an

approximation of the general direction of ice flow in the southwestern part of the ice sheet. Nevertheless, his more or less W-E bearing stretches of source areas are too large for a more detailed analysis of the ice stream direction pattern.

### *Lüttig's TGZ estimations*

Lüttig (1958) reduced the size of Hesemann's hypothetic source areas by estimating the theoretical indicator centre (Theoretisch Geschiebezentrum, TGZ) of a large number of these counts. The longitude and latitude of the actual source of each indicator from a count of a given locality, were determined. The arithmetical mean of these longitudes and latitudes gives a geographical point that is the theoretical centre of the count in question. Working along these lines, Lüttig revised almost all of the published Hesemann counts, and some important points emerge from this approximation based on synthesis.

In the first place, Lüttig's TGZ characteristics provide greater stratigraphic




Fig. 3. The mean direction of ice movement toward the circumscribed area inferred from Lüttig's TGZ estimations.

refinement than the Hesemann numbers, because the conversion of the Hesemann counts into the TGZ estimations clarifies the stratigraphic positions and relations of glacial deposits in northern Germany and The Netherlands. The favourable effect of the application of Lüttig's method may have been due to the reduction of the vast Hesemann provenances to a restricted area. The situation of this synthetic mean, or rather an aggregate of its stratigraphical components, was established by Lüttig (1958) and is shown in Fig. 3. Since there is an interconnection between source area, direction of ice movement, and depositional area, the remaining elements have been added to Lüttig's original illustration (1958). For this reason, Figure 3 shows the additional outline, embracing the areas of deposition that Lüttig reconsidered. From the centre of this circumscribed region a line is drawn connecting and bisecting the related synthetic source area. In Fig. 3 this line, shown as an arrow, indicates the mean direction of ice movement looked for.

One of the main points to emerge from this sum total of Lüttig's synthesis is the stringent stratigraphic function of this method. It exposes an elongated mean ice flow with overlapping source areas upstream. Each of these source areas pertains to the predominant composition of indicators from a given glacial or stadial in the region of deposition. The backward and forward shifts along the stream direction of the locations of these provenances may be attributed to variations in the regimen of the diverse icecaps involved. A noteworthy factor here is the permanence of the mean ice stream direction, despite the different glacials and glacial phases.

It is of great importance that the mean direction runs about parallel to those which were roughly estimated on the basis of the locations of Hesemann's source areas. To reconstruct these directions, we began from two opposite starting-points, i.e. the source areas (by putting those of Hesemann to use) and the area of deposition (by taking advantage of Lüttig's TGZ estimations).

#### CONSIDERATIONS

In spite of the inferred and confirmed general mean trend of ice movement, our review of the quantitative methods did not lead to a conception of the actual streamline forms. As we have seen, to determine the trend of ice movement with respect to an area of deposition on the basis of indicators, provenances are required that can be defined by lines running parallel to the ice stream direction, which means that the data we need are at the same time the relations to be deduced. In other words, we find ourselves in a vicious circle, which must be broken through in one way or another if our main objective is to attempt to unravel the ice stream pattern. In this respect it is of great advantage that the advocates of indicator counts and indicator fans have shown us how the land lies and have supplied us with a rich store of valuable information to draw upon.

Thus, although the indicators as modus operandi have put us in a fair way to gain some insight into the directions of ice stream, we have reached a deadlock and it is now time to adopt a different method of approach. When we strike the balance of the rich store of useful information available, our attention is drawn by the immense accumulation of data and facts concerning the direct and indirect glacial effects and features that have been carefully brought together by generations of geologists from Fennoscandia and the countries around the Baltic.

Of special interest, for instance, are the compilations on a regional scale of

the directions of ice movement in Norway, Sweden, and Finland, the general outlines of the limits of the glacial phases of the diverse ice sheets, the inferred recession lines of the last glaciation, and the esker systems in Sweden and Finland. These features, which reflect the morphology of the former ice sheets, may provide additional information that lends itself to investigate the functional relationship between the location, the route or mode of transport, and the source of the indicators.

The first part of this report concerns the attempt to gain more insight into the regional distribution of indicators. It starts with an evaluation of the possible directions of ice flow in the source areas, to permit regrouping of these areas by lines that parallel the most plausible directions of the main ice movement. The indicator counts in the circum-Baltic area were rearranged accordingly, and the resulting distribution pattern is discussed in relation to the inferred design of ice flow in Fennoscandia.

## Regrouping of provenances

### CRITERIA FOR THE ESTIMATION OF THE DIRECTIONS OF ICE FLOW

To regroup the provenances, it is essential to know the former directions of ice motion in the source areas of the indicators. In this section, the methods used to determine these directions in Fennoscandia are commented upon. The flowline pattern proper will be discussed in the next section.

Different types of traces left by former ice movements are outlined in Table 1. The indicated characteristics and their possibilities and limitations for the assessment of ice flow directions are based mainly on data given by Charlesworth (1957), Flint (1971), and Embleton & King (1975). The relation of directional data to the sense of ice movement will be evaluated on the basis of Table 1.

To start with the striae, it must be kept in mind that changing ice flow directions may erase the earlier marks. Concerning this uncertainty, Flint (1971, p. 95) remarked that 'probably most of the striations that have been mapped were made near glacier margins during deglaciation and therefore do not indicate the direction of flow that characterized glacial expansion'. Here, in conformity with his cautious view — even though the striae could have been formed elsewhere under the ice sheet — they are considered to indicate glacial activities that took place toward the periphery of the ice mantle. The main glacial phase may be represented by well-defined striae, like those found by Bergersen & Garnes (1972) in the Gudbrandsdal area in Norway. Less well defined striations are younger, and there are strong differences in their trends. Because of variations from the main direction of the ice flow caused by the unevenness of the bedrock topography as well as the divergent direction of flow at the ice margin, Virkkala (1951, 1960) recommended that a statistical analysis of a large number of striae over a reasonably large area should be made to gain insight into the main trend of ice movement. The value of the results should be weighed against that of the measurements of other directional indicators such as grooves, fluted surfaces, and crescentic marks. Bergersen and Garnes (1972) found that lee-side plucklings and roches moutonnées had the same general orientation as the striae of the main glaciation phase. A

Table 1. Traces of directions of ice flow.

Features	Mode of origin	Glacial distribution	Approach to interpretation of main ice flow directions	Deviations from main ice flow directions	Complications due to nonglacial processes	Scale of application in Fennoscandia
Glacial striæ	Glacial basal abrasion	Toward periphery of the ice sheet	Statistical analysis of large numbers of striae	Unevenness of bedrock topography Events at the ice margin	Striations made by: drift ice tectonic movements mud flows snow avalanches rock avalanches	Regional
Roches moutonnées	Glacial basal abrasion	Throughout the ice sheet	Measurement of directions of the major axes	Slight deviations		Local
Drumlins	Moulded at the base of the ice sheet	Throughout the ice sheet	Measurement of directions of the major axes	Slight deviations		Local
Indicators	Dislodgement from the overridden bedrock As loose fragments picked up by the ice sheet	Throughout the ice sheet	Statistical analysis of large numbers of indicators	Nonglacial events	Waterborne transport before or after their incorporation into the ice	Local
Fabric of till	Alignment in englacial or basal till by ice movement Till flowage	Throughout the ice sheet	Fabric analysis of joints Measurements of orientation of long axes of stones	Alignment depends on genetic till type Till flowage Slumping at ice edge	Disturbances by: crustal rebound drying up of till layers geochemical processes decalcification of till Pryoturbation	Local

possible source of error for which we should be on our guard in dealing with striae are the non-glacial striae (Embleton & King, 1975). In Fennoscandia the measurement, plotting, and analysis of striae has been applied on a regional scale.

Roches moutonnées are rock masses which have been rounded off by glacial basal abrasion. They have been found in the area of the Scandinavian ice divide, on transfluence passes, and at the heads of valleys, but also at low levels or on flat ground, where they are best developed. The rather powerful abrasion action that must have shaped these bodies and the parallelism of the whaleback forms to inferred directions of ice flow, suggest that the overriding ice was active and had a persistent direction of movement. Consequently, the roches moutonnées are ideal criteria for the estimation of main directions of ice flow. Unfortunately, they are shown only locally on the Fennoscandinavian Quaternary maps.

The elongation of drumlins is generally considered to conform to the direction of the basal ice movement (Embleton & King, 1975). Accordingly, Aario et al. (1974) found that the majority of the striations in Koillismaa, Finland, lie parallel to the drumlin orientation. Flint (1971) thinks the drumlins to be more reliable indicators of former ice movement, because they are less influenced than striae by the local topography. Although it has been argued that drumlins are formed submarginally, Embleton and King state that they are also developed more toward the centre of the ice sheet. Therefore, they are considered here to be distributed throughout the ice sheet. In Fennoscandia they are recorded in local patches.

By definition, an indicator refers to its area of origin, or more precisely to the area where the source rock crops out as hard bedrock. However, the ice also removed clasts of the source rock around the source area proper. These disintegrated parent rock fragments were transported by non-glacial agencies. Of the non-glacial terrestrial agents (gravity, rivers, and wind), the rivers carry rudaceous rocks the farthest from their source, and fluvial transport may be called the leading non-glacial agency in the distribution of disintegrated source rock. Hence, pre-glacial rivers or river systems whose course through a source area was not parallel to the later direction of the main ice flow, tended to enlarge this provenance. The greatest effect can be expected where the river flowed at right angles to the ice stream direction.

Indicators in transit arriving at the edge of the ice could have been transported further, in a direction different to that of the ice, by proglacial rivers or river systems. Therefore, it is not inconceivable that the European ice-marginal streams carried indicator boulders from east to west. These boulders had a chance of being re-incorporated by an advancing ice sheet.

Consequently, preglacial and proglacial transport may be worth considering in attempts to explain cases of incongruity of indicator distribution with respect to inferred main ice stream directions.

Till is deposited throughout the ice sheet. There are several types of till. In the review given by Dreimanis (1976) they are subdivided genetically into two groups: superglacial tills and subglacial tills. Englacial till is assigned to the superglacial group. Another two-part classification mentioned by Dreimanis (1976) is based on the environment of deposition. Ortho-tills are deposited on land, para-tills in water. It is difficult to differentiate the genetic type on the basis of the lithologic character of a till.

In para-till, or water-laid till in the terminology proposed by Dreimanis, the alignment of clasts is found to be random. The alignment of boulders in basal till




Fig. 4. Directions of main ice movement in Fennoscandinia in relation to general orientation of drumlins, distribution of erratics in southern Sweden, and boulder trains in Finland.

may be either parallel or transverse to the direction of the ice flow. In englacial till the long axes of stones lie parallel to the direction of glacial movement. Evidently, the alignment depends on the type of the till in question, but the genetic type of till as sediment is hard to distinguish. Even water-laid till is seldom recognized as such (Dreimanis, 1976). Moreover, there are till fabrics which, according to Embleton & King (1975) are not directly related to the basal movement of the ice, but are determined by other processes such as till flowage due to hydrostatical pressure or the squeezing of water-soaked till from under the ice. Fabrics may also have been disturbed by slumping at the ice edge. This makes the interpretation of the results of fabric analyses very complicated.

Post-depositional non-glacial forces add to these complications. Grisak et al. (1976) mention a crustal rebound after glacial loading, the drying up of till layers, and geochemical processes during various periods of ground-water circulation. To these may be added cryoturbation and solution of limestones. Consequently, the younger the drift under investigation, the less chance of post-depositional disturbances. Hence, a fabric study of a Weichsel drift could carry more weight than an analysis of the fabric in a Saale till. Richter (1932, 1933) examined a large number of exposures of the Weichselian in the northwestern and eastern parts of Germany, Poland, and Denmark, and estimated the alignment of boulders and the sense of joint patterns in the till. The general impression given by the results is one of orderliness. The main regional tendency of the longitudinal position of the boulders in the vicinity of Rügen is northeast. The further toward the east, the more long axes trend north-south, resulting in a roughly north-south sense just beyond the meridian of 16° E (Richter, 1936).

The outcome of fabric analyses of Saalian till in The Netherlands is not order but disorder. This led Boekschoten and Veenstra (1967) to point out that this method of estimating the direction of ice movement falls short here. Despite the promising results obtained in the Weichselian in Germany, the limited applications so far and the inadequacy with respect to altered and older strata make the estimation of orientations of indicators inadequate to approximate the regional directions of ice flow for the Scandinavian ice sheets.

Because we are primarily concerned with the main directions of active ice, the *roches moutonnées* and the drumlins are the most reliable direction indicators for the present purpose. Unfortunately, these features have only been recorded locally. The striae were mapped on a regional scale, but their formation towards the periphery of the ice sheet and their susceptibility to local variations of ice flow mean that their reliability is uncertain. The dispersal of indicators gives an over-all view of main ice stream directions, but even so, the non-glacial events may lead us astray here.

In view of all this, we must exercise the greatest care, and weigh one directional feature against another, in judging main directions of active ice flow.

#### FLOWLINE PATTERN IN FENNOSCANDIA

In 1965 Andersen published two glacial maps, one of southern and the other of northern Norway, on which the direction and location of numerous striae of the last glaciation are shown. Lundqvist (1965) compiled the main directions in Sweden, on the basis of the directions of striae and boulder transport, and here and there on rock forms. A map of the more recent movements of the ice in Finland

was given by Donner (1965). In the defining of the general movement of the ice in Finland, striae have been used almost exclusively.

Because the main directions in Sweden were not derived from striae only, but also from boulder transport and rock forms, they have been defined on a sounder basis than those in Norway and Finland. Nevertheless, if the directional data from these three maps are combined, the pieces fit well together in the general sense. The result is shown in Fig. 4.

Although the regional consistency of the streamline features suggests a certain reliability, these features should also be evaluated by more trustworthy direction indicators, say drumlins, which, as we have seen, only occur locally. Compiled from various sources (mentioned in the legend), the locations and general orientations of drumlin occurrences are also shown in Fig. 4. Their directions agree with those of the ice movements.

The same holds for the direction of spread of some Swedish indicators. Magnusson et al. (1963) provided us with a representation of the dispersal of Jotnian sandstones (J). (Capital letters between parentheses refer to Fig. 4). Instances of the glacial distribution of Cambro-Silurian gravel on the east (E) and west (W) sides of Lake Vättern were given by Gillberg (1965). The same author published the results of investigations of the dispersal of Cambro-Silurian gravel in Närke (N) and Uppland (U) in 1967. Figure 4 indicates that the southward trends of the more or less tongue-shaped frequency distributions lie parallel to the pattern of the inferred ice flow.

The directions of boulder trains and trace element concentrations in Finland are given in Table 2 and are also depicted by numbered arrows in Fig. 4.

Table 2. Boulder trains in Finland.

number of arrow	area	lithology	author
1	Laitila	rapakivi	Sauramo (1929)
2	Satakunta	olivine diabase, sandstone	Sauramo (1929)
3	Viipuri	rapakivi	Sauramo (1929)
4	Viipuri	rapakivi	Sauramo (1929)
5	Ladoga	rapakivi	Sauramo (1929)
6	Virtasalmi	ore boulders	Hyvärinen (1969)
7	Seinäjoki	plagioclase porphyrite antimony bearing blocks	Pääkkönen (1966)
8	Korsnas		Hyvärinen (1967)
9	Kolima	trace element concentrations	Kauranne (1967)
10	northern Savo	diabase	Sauramo (1929)
11	Paukkajanvaara		Wennervirta (1967)
12	Ylivieska		Mutanen (1971)
13	Moisionvaara	quartzites	Virkkala (1951)

These directions are in reasonable agreement with the general tendency of the young ice movements. As a result, the flowline pattern is linked up at its north, east, and south ends with swarms of drumlins and the dispersal of certain indicators. This could be taken as confirmation of the traces of ice movements under consideration.

Drumlins are shaped and indicators are transported by ice in motion. Hence,

the flowline pattern must reflect the directions of ice movement. This is obviously the case for advancing ice, but it is not as self-evident for receding ice. Flint (1971) mentions the pronounced shrinkage of most of the present ice bodies in the first half of this century. In spite of the retreat of their termini, however, these glaciers are still in motion, as shown by an illustrative case cited by Bischop, to which Flint (1971) refers. Here, a zone near the terminus of the ice sheet near Thule in north-western Greenland stagnates while the ice flow upstream proceeds, and in this case the stagnant ice is even overridden by its mobile counterpart. Embleton and King (1975, p. 93) remarked that: 'A dynamically active glacier is one which is flowing fast, whether it is retreating or advancing at its snout'.

It is therefore uncertain whether the Fennoscandinavian flowline pattern was formed at the beginning or the end of the last glaciation, but the chances are that, in the main, the present directional remnants are related to mobile ice of the final regional recession of the last ice sheet.

#### SOURCE AREAS ADAPTED TO ICE FLOW DIRECTIONS

As stated at the outset of this chapter, unravelling of the general trends of an ice flow pattern on the basis of the distribution of indicators requires provenances contained by lines running parallel to the directions of ice movement. Since these directions bear roughly north to south in southern Sweden, the source areas should be chosen such that they are consistent with a north-south trend. In the eastern part of Sweden the source areas have to be consistent with the general south-easterly direction.

To fulfil these conditions, twelve compound source areas which could be combined into five rows, were chosen in Fennoscandinavia. These north-south trending arrays are represented in a west to east sequence and are indicated by Roman numerals (see Fig. 5) in ascending progression. In this way, the left side of a paper refers to west, the west to east order is maintained when the numbers are put in writing in regular succession. The component parts of the north-south trending arrays are designated by the same number as the unit to which they pertain. Arranged according to the five major units, the twelve source areas are: I: Oslo, and the Swedish west coast including the provinces of Västergötland, Bohuslän, and Dalsland; II: Scania, Småland, and Dalarna; III: Bornholm and Blekinge; IV: Stockholm and Uppland; V: Ångermanland, forming a single group with the island of Rödön and the Ragunda area, the Baltic and the Gulf of Bothnia, Åland, and Finland.

On the south side of the circum-Baltic area the rapakivis and Baltic quartz porphyries account for the lion's share of the indicators from unit V. As a result this large unit may be narrowed down to a more restricted zone comprising the coastal parts of southern Finland, the Åland islands, and a southwestward extension including the areas where the Baltic porphyries may crop out. The compound source areas and units are shown in Fig. 5. It must be kept in mind that these groups are adjustable and can be rearranged into other units should the need arise.

Although the eastward-trending increase of north Baltic elements in the coastal areas bordering the south Baltic, at the expense of the southern Swedish indicators (Woldstedt, 1955) supports the proposed approach, a more detailed analysis must be carried out to evaluate this new strategy properly.




Fig. 5. Source areas chosen according to directions of ice flow. For geographic names, see folding map.

## Rearrangement of indicator counts

Whether or not the Fennoscandinavian flowline pattern should be ascribed to ice movement of an advancing or receding ice sheet, one thing is certain — it belongs to the last glaciation. Therefore, the indicator distribution of that time should be considered first. Hence, diverging current practice, our investigations will be carried out from young to old, and begin with the Weichselian.

### THE WEICHSELIAN

#### *Denmark (Milthers, K.)*

K. Milthers (1942) made an extensive and systematic investigation into the




Fig. 6. Indicator distribution of the Weichselian in Denmark. Counts of K. Milthers (1942).

distribution of indicators throughout Denmark. To obtain statistically comparable results, he confined himself to rhomb-, Dalarna, and Baltic porphyries. These are tough and easily recognizable indicators, but in addition their resistance to mechanical wear and disintegration is comparable. Schematic representations of their source areas are given in Fig. 6.

The three-component system allows grouping of the results into three categories, i.e. localities with prevailing Oslo, Dalarna, and Baltic indicators. To characterize these categories on his porphyry distribution map, Milthers indicated the locations of his counts by distinctive symbols. On referring to this map (Table I in Milthers' publication) we find that localities with predominantly Oslo indicators are concentrated in and virtually restricted to the northern half of Jutland. A sharp, slightly curved, southern boundary line can be drawn that separates the area in which rhomb-porphyries predominate from an area characterized by Dalarna porphyries. North of this line, the localities of the counts are indicated on Milthers' distribution map by one symbol only, i.e. the one for the prevailing rhomb-porphyries. South of this line, however, next to the symbol for predominance of Dalarna porphyries, which occurs the most frequently, the symbol for counts with predominance of Baltic porphyries occurs, especially on the present coastlines. Toward the south, this mixed zone in which Dalarna porphyries predominate can be delimited by a more or less straight line, which runs parallel to the northern boundary line through Jutland and from the southeasternmost part of Jutland to the northern end of the Öresund.

Counts in which Baltic porphyries predominate abound south of this line. In some places there is, however, an occasional locality in which Dalarna porphyries predominate.

Figure 6 shows the over-all porphyry distribution pattern in Denmark. A north-east-trending belt in which Dalarna porphyries predominate is flanked on its northern side by a major distribution zone of rhomb-porphyries and on its southern side by an area where the Baltic porphyries predominate. North of the dividing line through Jutland, the rhomb-porphyries average 80%. Between the lines through Jutland and the Danish archipelago the average percentages of the Dalarna, Baltic, and rhomb-porphyries are 50, 40, and 10, respectively. South of the line through the Danish archipelago, the average ratio of Baltic to Dalarna porphyries is 75 : 25, and almost no rhomb-porphyries occur here (less than 1%).

#### *An area west of the Oder (Hesemann)*

Hesemann (1932, 1938) made about 70 counts in a tract south of the island of Rügen and west of the Oder. He drew his indicator samples from the following stretches of moraines: Barth-Grimmen (1); Greifswald-Zirchow (2); Gnoien-Rosenthal (3); west Oder end moraine of the Pomeranian Stage (4); Gransee-Rüdnitz (5) (see Fig. 7). For each of these areas I combined the localities and regrouped their indicators according to the twelve newly chosen compound source areas. The resulting percentages per combination of localities are shown in Table 3A.

It is obvious that the indicators from Småland and the Baltic unit are in the majority. This is shown diagrammatically in Fig. 7, where the average shares of these areas in the total indicator association of the 70 counts for the west side of the Oder are shown superimposed on the corresponding source area. The density shading indicates that the indicator distribution is split into two groups, the Småland and Baltic indicators falling in the 41-60% range and those from Dalarna,

Table 3. Rearranged Hessmann counts in the Weichselian glacier area west of the Oder (in percentages)

		Regional combinations of count localities (see Fig. 7)					A.				B.				
		Source areas					Total of unit V		100- total of V		Source areas without V; the remaining percentages brought up to 100				
		I II III IV V					I II		I II III IV		I II III IV				
Localities	Count	Barth-Grimmen	1	—	40	3	3	2	1 21 30	52	48	—	84	6	
Greifswald-Ziechow			2	—	37	3	1	2	10 47	57	43	—	86	7	
Gnoien-Rosenthal			3	—	37	3	1	3	6 50	56	44	—	84	7	
West Oder end moraine			4	—	68	5	8	1	5 13	18	82	—	83	6	
Pomeranian Stage			5	—	58	8	1	1	5 27	32	68	—	85	12	
Gransee-Rüdnitz												1	2	Hesemann, 1933	
		Oslo Västergötland Bohuslän Dalsland					Småland Dalarna					Bornholm Blekinge			
		Stockholm Uppland					Angermanland, Ragunda, Rödön the Baltic + Gulf of Bothnia Åland + Finland					Bothnian-Baltic			
		Bothnian-Baltic					Oslo Swedish W. Coast					Småland Dalarna			
		Småland Dalarna					Bornholm Blekinge					Bornholm Blekinge			
		Stockholm Uppland					Stockholm Uppland					Stockholm Uppland			

Uppland, Stockholm, and Bornholm in the 1-5% range. Indicators from the Oslo area and the Swedish west coast are lacking.

Figure 7 visualizes the relationship of the indicator association to the twelve compound source areas, but is not equally suitable for the evaluation of the distribution of the five major units. To overcome this drawback, the data from Table 3A have been incorporated into Fig. 8 as an indicator diagram in which the twelve source areas are combined in the five major units as indicated. The units are set out geographically from west to east. The indicator diagram shows that the individual line-graphs for the various stretches of moraine are comparable. They emphasize the supremacy of the Småland-Dalarna and the Bothnian-Baltic units with respect to the others, which from a quantitative point of view are rather thin.

Since the west side of the Oder is in a direct line with the flowline pattern in Dalarna and Småland but far from the conceivable extension of the flowlines through the Bothnian-Baltic unit, the values for the latter were omitted from Table 3A and the remaining percentages recalculated to total 100. The results shown in Table 3B are represented schematically in Fig. 8. Now the percentages of the Scania, Småland, and Dalarna unit lie in a narrow range of 89 to 97%. As a result, the lines connecting the percentages of this compound source area with those of the Bornholm-Blekinge unit almost coincide.




Fig. 7. Schematic representation of indicator associations of the Weichselian of an area west of the Oder. For numbered stretches, see also Table 3.




Fig. 8. Indicator diagrams of rearranged Hesemann counts for the Weichselian west of the Oder.

*An area east of the Oder (Hesemann)*

As can be seen from Fig. 9, the Weichselian area east of the Oder is situated between two lines, one of them a NNE-SSW-bearing line that passes through the Lower Oder, and the other the 16°E meridian. Hesemann (1932, 1937, 1938) made 140 counts in this area, which are subdivided in Table 4A into four regional sections: the N-S trending stretch of the east Oder end moraine of the Pomeranian Stage (6), the eastern section of the W-E track of the Pomeranian end moraine (7), the Züllichauer Bogen (8), and an area between Zossen and Unruhstadt (9).

The percentages in Table 4A clearly reflect the deficiency of the Bornholm indicators as opposed to the demonstrable quantities on the west side of the Oder. This could mean that Bornholm lay in the path of ice movement towards the west side of the Oder but outside the course towards the east side of the river. In accordance with this, Bornholm is tentatively placed to the left of the Småland-Dalarna groups in Table 4, in order to maintain the west-east arrangement of the source areas for a given region.

The values for the Bothnian-Baltic and the Småland groups stand out again,




Fig. 9. Distributional features of the Weichselian indicator associations of an area east of the Oder. For numbered stretches, see also Table 4.

Table 4. Rearranged Hesemann counts for the Weichselian glacier area east of the Oder (in percentages)




Fig. 10. Indicator diagrams of rearranged Hesemann counts for the Weichselian east of the Oder.

but those for Dalarna are better represented on the east than on the west side of the Oder. These three groups predominate as can be seen from Fig. 9. The indicator diagram (Figure 10), shows that the Småland-Dalarna unit ranges from 41 to 64% and the subdivisions on either side, i.e. the Bornholm and the Uppland-Stockholm groups, do not exceed 2%. At first sight the Småland-Dalarna unit seems on the wane as compared with the corresponding values in the western Oder area, but when the Bothnian-Baltic unit is subtracted and the remaining values recalculated to 100 (Table 4B), the Småland-Dalarna unit on the east side of the Oder outranges the same association on the west side, due to the higher content of Dalarna indicators.

#### *Lithuania (Korn and Tarvydas)*

No Hesemann counts have been carried out in Lithuania, but Korn (1895)




Fig. 11. Lithuania: the tripartite indicator distribution given by Tarvydas (1960).

investigated the erratics in material collected by drilling in Kaliningrad and surroundings. He found that the Swedish indicators in east Prussia appear to originate from areas lying north of the 59° parallel, because the material lacked Scanian basalts and indicators from Småland, but contained indicators from Stockholm, Uppland, Dalarna, Jämtland, Ångermanland, and Norrland. The same tendency is evident from the percentages published by Tarvydas (1967). In the Kaliningrad region the Swedish indicators total 46%, including south Swedish ones accounting on average for 7%, whereas no south Swedish indicators are encountered in the Lithuanian areas east of this region.

An over-all picture of the erratic distribution of the Pomeranian Substage in Lithuania is given by Tarvydas (1960). In west Lithuania, indicators from Sweden and the bed of the Baltic predominate; in the centre, indicators from the Åland islands and SW Finland, and in NE Lithuania, indicators from SW Finland, Vyborg and neighbouring regions of Karelia. His representation of the three areas of distribution is reproduced in Fig. 11.

Thus, two boundary lines divide Lithuania into three indicator zones. One of these lines delimits the dispersal areas of some central-Swedish indicators, including the Dalarna porphyries, as well as the Uppland and Stockholm granites to the east. The other line separates an area in which Åland indicators predominate from a corner in the northeast of Lithuania, where south Finnish indicators prevail.

#### *Latvia (Eskola)*

Eskola (1933), who identified 961 erratic boulders from the environs of Riga, came to the conclusion that they originated from southwest Finland, including the Åland islands. He drew his conclusion on the basis of the percentages of the different groups of the total association in relation to their distribution in Finland.

Because his enumeration is not divided into the usual indicator categories, it is not possible to compare his values with those of the Hesemann counts. Fortunately, however, the Åland indicators and the Baltic porphyries are exceptions. Since Eskola also took into account the Dalarna porphyries, which were absent, the three groups of indicators necessary for a Milthers count are available. The result is shown in Table 5. These percentages have been recalculated to give a total of 100 in the last column.

The Baltic porphyries and the Åland rapakivis seem to add up to 45.1%, but in reality this is only a minimum value because, according to Eskola, a large

Table 5. A Milthers count derived from Eskola's tabulation of erratic boulders from Riga.

	percentages (from Eskola, 1933)	percentages recalculated to 100%
Dalarna porphyries	—	—
Red Baltic porphyries	0.7	7.5
Brown Baltic porphyries	1.5	16.1
Åland rapakivis	2.0	21.5 }
Rapakivis unspecified	5.1	54.9 }
		100.0

part of the unspecified rapakivis shown in Table 5 may be ascribed to the Åland islands and the sea bottom south of these islands. If this share (in Table 5 in the column on the right) is greater than 5%, which in view of the low verge is likely, the indicators from the Åland islands and the bottom of the Baltic will prevail in Riga. It therefore seems plausible to assign Riga to the central Lithuanian indicator zone defined by Tarvydas.

#### THE SAALIAN

##### *The Netherlands (Hesemann counts)*

In 1972, Zandstra reviewed all of the published Hesemann counts from The




Fig. 12. Regional grouping of the Hesemann counts in The Netherlands.

Netherlands, at that time amounting to 53. To this number we can now add 5 which are counts made by Zandstra (1974), who was also kind enough to place at our disposal the results of two more, which were made in Emmerschans and will be published in 1977.

Thus, there are 60 counts to consider, the localities being placed in nine regional combinations roughly analogous to the subdivision made by Zandstra (1972). The indicators of the lenses of the red boulder clay were assigned to a tenth assemblage. These regions are indicated in Table 6 and Figs. 12 and 13.

In Table 6A (leaving out of account the Hondsrug, the region west of it, and the lenses of red boulder clay, because of their high quantities of Baltic material) the cumulative percentages of the Scania, Småland, and Dalarna groups lie higher than 29%. At the same time, except for the Utrecht ridge where the Bornholm-Blekinge indicators amount to 13%, the shares of this unit do not exceed 10% and those of the Oslo-Swedish west coast 2%.

We should pause here to consider an aspect of the distribution of Oslo indicators that does not find expression in the general mean value just mentioned,




Fig. 13. Distributional features of Saalian indicator associations in The Netherlands. For letters, see Table 6.

Table 6. The Netherlands, percentages of rearranged Hesemann counts.

Regional combinations of count localities (see Fig. 13)		A.					B.				
		Source areas		Total of unit V			100 - total of V		Source areas without V; the remaining percentages brought up to 100		
		I	II	III	IV	V	I	II	III	IV	
Oslo											
Utrecht ridge	UR	1 20	8	6	7	13 14	1 24	6	31	69	1 29 12
E Central Netherlands	EN	1 34	14	8	2	8 1	3	6	23	32	1 50 21
Västergötland Bohuslän Dalsland											
Phase d - ter Wee											
Texel-Wieringen	TW	1	20	10	6	5 3	1	6	38	45	55
SW Friesland	WF	1	28	20	4	7 2	2	5	30	37	63
NE Polder (normal boulder clay)	NP	23	15	3	1	12 6	3	6	31	40	60
Stretch Vollenhove-Coevorden	VC	1	18	18	7	1 5 3	3	5	36	44	56
Scania Småland Dalarna											
Bornholm Blekinge											
Stockholm Uppland											
Angermanland, Ragunda, Rödön the Baltic + Gulf of Bothnia Åland + Finland											
Bothnian-Baltic											
Oslo											
Västergötland Bohuslän Dalsland											
Scania Småland Dalarna											
Bornholm Blekinge											
Stockholm Uppland											
Bothnian-Baltic											
Oslo											
Västergötland Bohuslän Dalsland											
Scania Småland Dalarna											
Bornholm Blekinge											
Stockholm Uppland											
Bothnian-Baltic											
Arithmetic mean of phase d											
Region west of Hondsrug	WH	1	2 14	12	3	1	4	2	2 8	50	60
Hondsrug	H	1	1	12	10	3	1	2	2 5	62	69
E Groningen	EG	1	2 22	19	3	2	4	4 43	47	53	2
Lenses of red boulder clay		9 4	1	13 4	3	3 63	69	31	4 41	35	6
									29	13	4 8
									3	3	42 13




Fig. 14. Distribution of the Oslo indicators in The Netherlands.

i.e. the combined percentage of the Oslo-Swedish west coast unit. The Oslo group itself accounts for one per cent and a half at most (see Table 6A). To show these values more clearly, the grouped frequencies of the percentages of the counts are given by region in Table 7A, from which it can be seen that the regional means range from 0.00 to 1.44%. The percentages above 1.00 are found in the middle of the eastern part of The Netherlands, i.e. in the Hondsrug, the area west of it, and a stretch between Vollenhove and Coevorden. The remaining regional averages are distributed more randomly.

The regional scarcity of the Oslo rocks in The Netherlands is expressed in Table 7B. Since no Oslo indicators have been found in more than 43 of the 60 investigated localities or two-thirds, the mode is 0. The arithmetic mean is 0.83%. There are two exceptions to the homogeneous pattern of indicator associations with hardly any Oslo indicators, but due to the lack of complete counts, these cases could not be included in Table 6. For an association with 20% rhombporphyries near Bergum in Friesland, van der Lijn (1941) mentioned the Hesemann number only. These walnut-sized porphyries were found in a ground moraine, which became accessible during the reclamation of some lakes. In spite of the large amount of Oslo rocks, Schuddebeurs (1959) did not count the indicator association brought up by dredging work near Drachten as it underwent an essential change. The larger pieces of stone were separated out and transported by ship to be dumped elsewhere. The smaller fraction that remained had an appreciable content of Oslo indicators, roughly estimated by Schuddebeurs at some 25%.

These two cases have three things in common: the amounts are so divergent from what would be expected that they may be regarded as isolations, they form part of a smaller fraction, and their location is in the middle of the eastern part of The Netherlands where Oslo indicators occur in percentages above 1%. The isolation is brought out clearly by Fig. 14, a graphical representation of the general distribution of the Oslo indicators in The Netherlands. If these exceptional cases are added to the 60 complete counts, the total number of counts becomes 62 and the arithmetical mean percentage of Oslo indicators 1.53, which means that despite the unusually high amounts of Oslo rock around Bergum and Drachten, the average still falls within the 1 to 5% category.

The density shading in Fig. 13 shows five indicator categories in the over-all distribution pattern in The Netherlands. The mean percentages of the indicators fall in the following ranges: Bothnian-Baltic: 41-60%, Småland: 21-30%, Dalarna: 11-20%, Stockholm: 6-10%, and Oslo, the Swedish west coast, Scania, Blekinge, Bornholm, and Uppland: 1-5%. Four groups stand out: the Bothnian-Baltic, the Småland, the Dalarna, and the Stockholm.

In Fig. 15A, the values from Table 6A are represented as an indicator diagram in which the twelve source areas are combined into the five major units indicated on the base line. From west to east, the distributions of the first three units exhibit symmetry, their lines hardly intersecting, as opposed to the more irregular configuration of the two remaining units. The indicator diagram has two peaks, one reflecting the Scania-Småland-Dalarna unit, the other the Bothnian-Baltic. Apart from these peaks, there is the less pronounced Uppland-Stockholm combination.

For Table 6A the Bothnian-Baltic unit was omitted and the remaining percentages recalculated to 100. The results (see Table 6B) are represented schematically in Fig. 15B. In eight of the Dutch regional categories, the distribu-




Fig. 15. Indicator diagrams of rearranged Hessmann counts in The Netherlands. 1) See Table 6.

Table 7. Percentages of Oslo indicators in The Netherlands.

A. Regional frequency distribution		B. General frequency distribution		
Regional combinations of localities	frequency × percentage	frequency × percentage	totals in %	
			general mean %	
Utrecht ridge	8 × 0% 1 × 7%	0.78	43 × 0% 5 × 1% 5 × 2% 1 × 3% 2 × 4% 2 × 5% 2 × 7%	0 5 10 3 8 10 14
E Central Netherlands	3 × 0%	0.00	subtotal	60
Wieringen-Texel	3 × 0% 1 × 3%	0.75	Bergum	20
SW Friesland	3 × 0%	0.00	Drachten	25
NE Polder, normal boulder clay	6 × 0% 1 × 2%	0.29	total	—
Vollenhove-Coevorden	5 × 0% 2 × 1% 1 × 4% 1 × 7%	1.44	62	95
Region W. of Hondsrug	7 × 0% 1 × 1% 2 × 5%	1.10		1.53
Hondsrug	5 × 0% 1 × 1% 4 × 2% 1 × 4%	1.18		
E Groningen	2 × 0% 1 × 1%	0.33		
Lenses of red boulder clay	1 × 0%	0.00		

tions of the first three Scandinavian units bear a stronger resemblance to each other than is the case in the initial indicator diagram. They comprise more than 60% of the Scania-Småland-Dalarna unit, whereas the Oslo-Swedish west coast and the Bornholm-Blekinge units do not exceed the 10 and 15% limit.

The indicator associations of the two remaining categories, i.e., the Utrecht ridge and the lenses of red boulder clay, also seem to correspond to each other, as suggested by the striking parity of the percentages of the Scania-Småland-Dalarna unit and the higher amount of Uppland-Stockholm granites relative to the other regional categories. This is in accordance with the correspondence of the gravel association of the 3.5 mm fraction of boulder clays in the environs of the Utrecht ridge and Lunteren with that of the decalcified clay of red boulder clay lenses in the Noordoost Polder (Overweel & Zandstra, 1967).

#### *The area between the Rhine and the Elbe (Hesemann)*

In an area adjoining The Netherlands and lying between the Rhine and the Elbe, Hesemann (1939a) performed 71 counts which can be geographically subdivided




Fig. 16. Indicator distribution of the Saalian between the Rhine and the Elbe. For letters, see Table 8.

as follows: a stretch south of Hamburg (SH), and area indicated in Fig. 16 as Münsterland (M), the drainage areas of the Lippe and the Ruhr (LR), and regions encompassing Hannover (H), Braunschweig (B), and Stendal (S). Schuddebeurs (1959, 1967) made 10 counts in the Hümmling and in Ost Friesland (HO in Fig. 16). The Saalian erratic boulder assemblages have been rearranged and averaged in the same way as for the Dutch counts shown in Table 8A and Fig. 17. The counts in the area south of Hamburg and in the Stendal region belong to the Warthe Stage. Figure 16 shows that in the over-all distribution of the counts in this area, the indicators from Småland, the Bothnian-Baltic, and Dalarna again predominate. As usual, the indicator diagram (Fig. 17) is double-peaked, but this time the Småland-Dalarna unit predominates over the Baltic-Bothnian unit. Except for the large amount of Oslo indicators, the Hümmling-Ost Friesland curve is consistent with the other diagrams.

If the Bothnian-Baltic unit is excluded and the remaining percentages recalculated, so they again amount to 100, the sum total of the Scania-Småland-Dalarna unit ranges from 97 to 100% in six of the eight regions. Except for the Hümmling and Ost Friesland, the recalculated percentages of the Småland granites between the Rhine and the Elbe range from 59 to 90, lying above 70 in five of the eight regional categories (see Table 8B). The large amount of Oslo rocks in the Hümmling and Ost Friesland stands out; these rocks even reach 50% at Werpeloh, Fresenburg, and Langefeld. Schuddebeurs (1959, 1967) pointed out that the exceptional abundance of these indicators belongs to part of a fraction (smaller than some 8 to 10 cm) of deposits underlying the boulder clay.

Although Marczinski (1968) did not specify the individual indicators in his counts made in an area between the Lower Elbe and the Lower Weser (EW in Fig. 16), he classed them in some of the same source area groups and units as are used here. He denoted the Bothnian-Baltic unit as rapakivi rocks, the Stockholm-Uppland unit as black-white granites, the Småland group as blue quartz granites and porphyries, the Dalarna group as Dalarna porphyries, and the Oslo group as rhomb-porphyries.

Notwithstanding the fact that the remaining source areas were not taken into consideration, Marczinski's values are comparable to the Hesemann counts in nearby areas. Marczinski subdivided the Saalian of the area he investigated into a series of three strata. His percentages contained in the total for crystalline erratics, have been recalculated to total 100% for each stratum in Table 8. The resulting values with their relatively small share of indicators of the Bothnian-Baltic unit approach those of the Warthe Stage south of Hamburg and of the Saalian of the Hannover area. Compared to the other counts for the area between the Rhine and the Elbe, the percentages of the Småland-Dalarna groups predominate here too, but not to the same degree. The decrease might result from the suppletory co-presence of indicators of the Stockholm-Uppland unit, which are absent in the Hesemann counts.

The inclusion of this unit in the counts made by Marczinski (1968) facilitates comparison with the Dutch association. If the Bothnian-Baltic unit is left out of account, the percentages in both areas are approximately the same except for the Småland group (compare Table 8 with Table 6).

*Poland and White Russia - Belo Russkaya - (Milthers & Milthers)*

Except for those in the eastern Oder area, and those made by Ladwig (1938), there are no Hesemann counts for Poland, but fortunately V. and K. Milthers (1938) performed a systematic study based on porphyry counts of Saalian deposits in an area roughly circumscribed by latitudes 50° and 54° N and meridians 19° and 29° E of Greenwich. The larger, western part of this area lies in Poland, the remainder in White Russia (see Fig. 18). These investigators divided their study area into 6 regions, which I have adjusted slightly. Figure 18 shows not 6 but 5 regions, which are numbered I to V from west to east. The recalculated percentages of the erratic associations of these regions are given in Table 9, where the source areas are shown in the same sequence.

Strictly speaking, there are only two relevant source areas: the Baltic-Bothnian and Dalarna. The Baltic-Bothnian group is represented here by the brown and red Baltic porphyries and the Åland indicators. Since the variance of the Baltic porphyries (i.e. the sum of the brown and red porphyries) is limited to a narrow range of 2% (between 7 and 9%), marked differences in the Baltic-Bothnian must be due to variations of the Åland indicator percentages. Figure 18 shows an




Fig. 18. The eastward trend of the Saalian indicator distribution in Poland and White Russia. For Roman numerals, see Table 9.

Table 9. Milthers counts in Poland and White Russia (in percentages).

Regional combinations of count localities	Indicator type			
	Dalarna porphyries	Brown Baltic porphyries	Red Baltic porphyries	Åland indicators
I Area southwest of Lodz	20	4	3	73
II Area south of Warsaw	13	5	4	78
III Siedlce area	7	3	6	84
IV Area south of Grodno	4	1	6	89
V Area south of Vilna	4	1	7	88

increase of the Baltic indicators and a decrease of the Dalarna indicators, both from west to east, the latter dropping from 20 to 4%. Compared with the 15 to 60% Baltic indicators in the Hesemann counts to the west of this area, the 80% in region I, south of Lodz, is certainly higher. From there to region IV, the percentage of the Baltic indicators rises from 80 to 96%.

In Milthers' method of boulder counting, only the porphyries are used. The percentages in Table 9 would have been much lower, favouring other groups, if the remainder of the indicators had been included as well. The interesting question is, which of the other major groups of indicators besides these porphyries occur in Poland? Kreutz and Glowinska (1932) offer an answer in their statement that the indicators from the Stockholm and Uppland areas are second in importance, after those from Finland.

### Resulting distribution pattern of indicators

#### CORRESPONDENCE BETWEEN THE WEICHSELIAN AND THE SAALIAN DISTRIBUTIONS

Of the twelve basic source areas from which we started, four stand out in the indicator distribution on the south side of the circum-Baltic. This is especially clear from the Hesemann counts where, in contrast with the method of Milthers, there is no restriction on the choice of the species of crystalline indicators.

Thus our new approach reveals four main groups of indicators, i.e. the groups of Oslo, Dalarna, Småland, and the Baltic. In the regional distribution of the Weichselian, which is shown in Fig. 19, the predominance of the Oslo group




Fig. 19. Distribution of predominant indicator groups of the Saalian and the Weichselian.

is restricted to the northwestern half of Jutland. On its east side lies an area where Dalarna and Baltic indicators prevail. The main dispersal of the Dalarna indicators extends to the dividing line in Lithuania separating the zone of central Swedish indicators from the zone of the Åland indicators. The distribution of the Baltic indicators, however, continues further eastward, although those from the Åland islands predominate in the central part of Lithuania and those from southern Finland in the northeastern part.

Although in almost every location in north Jutland where a Milthers' count has been made, the rhomb-porphyry content amounts to 70 to 80% or even more, there are some instances where Baltic or Dalarna porphyries predominate. The transition between the zone of the outranking rhomb-porphries and the zone characterized by the Dalarna-Baltic groups is abrupt. The Oslo rocks lie within the 0 to 25% range comprising most of the cases of indicator associations in a 100 km belt on the southern side of the boundary line. With a few exceptions, the Oslo indicators do not exceed 50% here, which means that in almost every locality in this belt the combined Dalarna-Baltic components are in the majority. Farther to the southeast, the rhomb-porphyry percentages decrease to 0. In short, the area with mainly Oslo rocks in north Jutland is dotted with some localities where Dalarna or Baltic porphyries predominate, particularly close to the boundary line. Conversely, the homogeneous area where the Dalarna-Baltic combination pre-

dominates shows some specks with predominantly Oslo indicators.

The zone of the Weichselian distribution characterized by Dalarna and Baltic indicators shows a gap between the western border of Lithuania and 16° longitude where no counts have been performed. Further westward, on both the east and west sides of the Oder, Småland indicators show up in addition to the indicators from the Baltic and Dalarna.

The same three main groups stand out in the Saalian distribution of The Netherlands and in the area between the Rhine and the Elbe. No counts have been made in the Saalian deposits between the Elbe and the Warthe, but east of the Warthe the association of Dalarna and Baltic porphyries is encountered again in the area investigated by Milthers & Milthers (1938). As can be seen from Table 8, the Dalarna porphyries decrease here from west to east.

To compare the Saalian with the Weichselian we return to the delimitation of the Dalarna indicators of the Weichselian to the east. This lateral limit is the western boundary of the central Lithuanian zone. Where this line is extended southwards, it intersects the Saalian deposits south of Grodno, roughly between the regions III and IV (see Fig. 18). From region III to region IV the Dalarna porphyries decrease from 7 to 4%. The latter percentage, which is in fact a Dalarna-Baltic porphyry quotient, would have been lower if it had included other kinds of indicators as well, say the Stockholm and Uppland granites. The reduction to less than 4% suggests that here we have reached the margin of the main Saalian distribution of the Dalarna porphyries. If this is really the case, the horizontal extension to the east of the Dalarna indicators of the Saalian would coincide with that of the Weichselian.

To the west, the dispersal of the Dalarna porphyries of the Weichselian is distinctly bounded by the area of predominance of rhomb-porphyries in the northwestern part of Jutland. Figure 19 shows that the southward extension of this boundary line of the Weichselian distribution pattern is located in the North Sea, west of The Netherlands. This location is not only consistent with the predominance of the Dalarna-Baltic combination in The Netherlands and the western part of Germany but also with the absence of a coherent group of localities with more than 70% rhomb-porphyries. Just as in the Weichselian of the Danish archipelago, the area characterized by the Dalarna-Baltic combination of the Saalian is dotted by some localities in which Oslo indicators predominate, i.e. near Drachten in The Netherlands and near Werpeloh, Fresenburg, and Langefeld in the northwestern part of Germany.

In general, therefore, the distribution patterns of the Weichselian and the Saalian are similar. In the first place, the width of the dispersal area of the Dalarna porphyries of the Weichselian, although not yet completely investigated, is distinctly delimited to the west and the east. The southward extensions of the boundary lines seem to comprise the Saalian distribution of the Dalarna indicators as well. This holds more clearly for the east than for the west side, where the extended boundary line is in the North Sea.

Secondly, in both the Weichselian and the Saalian, the Dalarna indicators are accompanied as far as their eastern limits, by the Baltic suit, which in both stages extends beyond the Dalarna porphyries farther to the east.

Thirdly, where Hesemann counts have been made the Småland indicators stand out in the Weichselian and the Saalian alike. Topographically, the Saalian distribution of the Småland, Dalarna, and Baltic groups in The Netherlands and in the area between the Rhine and the Elbe seems to link up with the Weichselian

west and east of the Oder. There seems to be a continuous distribution of these groups roughly from the North Sea as far as 16° longitude, but since no Hesemann counts have been performed either in Denmark or in Poland and Russia, we are still in the dark as to the actual dispersal of these indicators.

#### THE DOUBLE-PEAKED CHARACTER OF THE INDICATOR DIAGRAMS

Unlike the counts of Milthers, which comprise only a limited number of indicator categories, the Hesemann method includes a wide selection of the crystalline indicators and therefore lends itself to the compilation of indicator diagrams.

These diagrams are distinctly double-peaked. Irrespective of whether they are from the Saalian or Weichselian, one peak reflects the Scania-Småland-Dalarna unit and the other the Bothnian-Baltic. Although the indicator diagrams are on the whole similar, there are some differences in detail which deserve further attention.

The Weichselian distribution of Oslo indicators in Denmark drops from




Fig. 20. Indicator distribution in relation to flowline pattern.

80% in Jutland to less than 1% in the eastern islands of the Danish archipelago. This tendency seems to persist, because hardly any Oslo indicators have been found on the west or the east side of the Oder. In the Saalian the distribution of Oslo indicators in The Netherlands is comparable with that in the area between the Rhine and the Elbe.

When the percentages of the Bothnian-Baltic unit are omitted and the remaining values recalculated to total 100, the percentage of the Scania-Småland-Dalarna unit ranges from 42 to 84 in The Netherlands, from 97 to 100 in the area between the Rhine and the Elbe, from 89 to 97 on the west side of the Oder, and from 92 to 100 on the east side. The wide range of the recalculated values for The Netherlands contrasts with the narrow range of the percentages in the other areas. Since the area between the Rhine and the Elbe refers, as in The Netherlands, to the Saalian, the nature of the contrast cannot be stratigraphic.

#### THE WEICHSELIAN DISTRIBUTION AND FLOWLINE PATTERN IN FENNOSCANDIA

When source areas are delimited by a line running parallel to the ice movement, the highest densities of their indicators will be found at the corresponding sides of the extension of that line. Figure 20 shows the boundary line in Jutland separating the area with predominantly rhomb-porphyrries on its west side from an area on its east side characterized by Dalarna porphyries. When the northeastward extension of this dividing line is adjusted to the course of the striae, the provenance of the Oslo rocks will lie to the west and the line will pass between this source and that of the Dalarna indicators. Similarly, another boundary, i.e. delimiting the areas of the central-Swedish and Åland indicators in Lithuania, can be extended across the Baltic. On reaching the Swedish coast, the extension is drawn parallel to the general trend of the striae. The Åland islands then lie on one side of the line and the provenance of the Dalarna indicators on the other. The area with predominantly Åland indicators in central Lithuania is separated from the area in which south-Finnish indicators predominate. When the parting is extended northeastwards, it will pass between the Åland islands and the southwest coast of Finland.

Thus, the dividing lines of the Weichselian indicator distribution pattern link up with equivalent features in the flowline pattern. The over-all structure shows a central more or less fan-shaped part encompassing the provenance and the main distribution range of the Dalarna indicators. This fan-like area is flanked on the west by the source area and main distribution area of Oslo clasts. On the east, it is true, it is flanked by a broad strip covering the Åland islands and the central Lithuanian zone in which Åland indicators predominate, but these erratics, together with others from the Baltic, are also in evidence in the major distribution zone of the Dalarna indicators.

In addition, the major distribution zone of Dalarna indicators includes representatives of the Småland group. The lateral extension of the spread of this group is not yet known, but the distribution of the Småland indicators, like their provenance, seems to fall within the fan-shaped range of the Dalarna clan. The western margin of the Dalarna suit approaches both of the source areas, but the eastern margin, although still close to the provenance of the Dalarna porphyries, lies far from that of the Småland indicators. Therefore, to approximate the lateral margin of the distribution of the Småland group on the east, an additional line has

to be drawn parallel to the trend of the striae and close to the bedrock source. Passing through the Baltic, the extension of this line crosses the coast in the Gulf of Gdansk, west of Kaliningrad, and then lies just in the gap in the Weichselian distribution pattern where no counts have been performed. If the extension of this line really denotes the roughly drawn easternmost limit of the major distribution zone of the Småland indicators, these specimens should be scarce farther to the east. This is consistent with the counts done by Korn (1895) because he found no indicators from Småland in the surroundings of Kaliningrad. Moreover, Tarvydas (1967) found only 7% south-Swedish indicators in the Kaliningrad region and no indicators from southern Sweden in the Lithuanian areas east of that region.

So far, the double peak of the indicator diagrams seems to recur in the over-all distribution pattern, because the latter falls into two parts. One part conforms to the Fennoscandinavian ice flow pattern. The other, which does not, is a unit of Baltic origin. At the circum-Baltic south and southwest sides the divergence of the Baltic constituent from the flowline pattern is distinct, but toward the southeast the component parts of the Baltic indicator association of the Weichselian conform to the directions of ice flow in central Sweden and southern Finland.

#### CORRESPONDING DIRECTIONS OF ICE MOVEMENT IN THE WEICHSELIAN AND THE SAAlian

In the preceding section the relationship between the Weichselian distribution and the flowline pattern emerged. But earlier in this chapter we saw that the distribution of indicators of the Weichselian and the Saalian correspond. The same groups stand out in the Saalian distribution as in the Weichselian. The indicator association of the Saalian is also bipartite, one half being characterized by Baltic and the other by Småland and Dalarna indicators. Furthermore on p. 45 we saw that the southward extensions of the boundary lines of the distribution zone of the Dalarna porphyries of the Weichselian seem, in spite of the gap where no counts have been performed, to include this equivalent of the Saalian as well. Since the source areas of the Dalarna and Småland indicators lie in the same sector of the flowline pattern, the erratics from Småland are also included in the distribution zone of the Dalarna porphyries, in both the Weichselian and the Saalian (see Fig. 20). These source areas can, however, be separated by a line running perpendicular to the former direction of ice flow. Because, as shown above, provenances enclosed by lines lying at right angles to the direction of the ice flow may be stratigraphically useful, the ratio between the Småland and Dalarna indicators could have stratigraphic value.

Here, at the end of the first part of this paper, some conclusions may be drawn. Firstly, because the distributions of Weichselian and Saalian indicators are not contradictory, there is no reason to assume that in a general sense the directions of ice movement differed in these stages. This is consistent with the permanence of the mean directions of ice movement for the Saalian and the Weichselian inferred from Lüttig's TGZ estimations (see p. 12), and also with the results of the investigation of Veenstra (1963), who studied the Bryozoa enclosed in various boulder clays from The Netherlands and other regions, and came to the conclusion that in both glacials the main direction of the ice in NW Germany was the same, 'as it must have passed the Danish archipelago to pick up similar microfaunal assemblages'.

Secondly, reference to the Fennoscandinavian flowline pattern shows the importance of the Småland indicators for a comprehensive picture of the distribution pattern. Furthermore, the general approach shows where no counts have been performed.

Thirdly, the indicator distribution on the south side of the circum-Baltic area has two main components. One of these, whose regional distribution harmonizes with the flowline pattern, will be referred to as consistent, and the other, which holds the Baltic indicators, will be called inconsistent. The wide-spread inconsistent Baltic distribution being the larger (see Fig. 19), it will be dealt with first in the following discussion.

## Part 2. Interpolation of indicator transport between distribution and source areas

Agents of the transport of indicators and the pattern of ice movements stand central in the second part of this paper. They are viewed in the light of the two classes into which the indicator distribution of the southern circum-Baltic region falls: the consistent unit, with its distribution showing correspondence with the flowline pattern, and the inconsistent, with indicators from the Baltic predominating. After considering the origin of the inconsistent distribution, I shall give a further elaboration of the streamline pattern on the basis of the consistent unit, which will result in a geometric approximation of the outlines and lines of flow of the ice sheet. Further adjustment and completion of the theoretical picture will lead to an integration of the hypothetical reconstructions of the consistent and inconsistent distribution processes.

### The inconsistent distribution unit

#### THE FORCE OF THE CONSISTENT AND INCONSISTENT ICE MOVEMENTS

During the revision of the indicator counts, little attention was paid to the processes and agents of transport which underlie the dispersion of indicators. Because the consistent distribution agrees with the pattern of ice flow in Fennoscandia, the ice movement tendency belonging to it may be denoted as consistent too. Because the Baltic distribution unit is intersected by the trend of the Fennoscandinavian flowline pattern, the transport of the Baltic indicators is defined inconsistent.

Since the second half of the nineteenth century it has been generally assumed that the Baltic indicators were dispersed by an ice stream flowing through the Baltic basin. Since this basin is now submerged, most of the traces of the Baltic ice stream are covered by water and pretty well inaccessible. Denmark and Scania

form an exception due to their location in the outlet of the Baltic. There, the central axis of flow of the huge ice stream through the Baltic depression must have lain straight across the Danish archipelago; but Denmark was also crossed by the consistent ice. Consequently, it is possible to compare here the consistent and inconsistent ice streams of the last glaciation.

#### *Indicator distribution in Denmark*

The Danish geologists not only accumulated ample information but also represented the data in a way making them highly accessible. In this connection the porphyry distribution map (table I) given by K. Milthers in 1942 is of special interest. On first inspection, the introduction of Baltic elements from east to west is obvious from the significant percentages of Baltic porphyries throughout the Danish




Fig. 21. Indicator distribution in Denmark, modified after K. Milthers (1942).

archipelago as far as an area northwest of Aarhus in Jutland, for which many porphyry counts show a composition with 50% (or slightly more) Baltic elements.

But, as we have seen, some essential dividing lines can be drawn on this distribution map between areas in which rhomb-, Dalarna, or Baltic porphyries predominate (Fig. 21).

The ratios for porphyry distribution in Denmark are fairly consistent in the areas separated by the dividing lines. In other words, there is a consistency in the porphyry distribution in a NE-SW direction over a distance of roughly 400 to 500 km. In a SE-NW direction, within a distance of 250 km, the distribution undergoes a distinct change from an area with an absolute majority of Baltic porphyries, via an area with Dalarna porphyries predominating, to an area with an absolute majority of rhomb-porphyries.

From the consistency of the porphyry distribution in the NE-SW direction in all three areas it follows that the main Weichselian ice movement in Denmark was also from NE to SW. This direction is consistent with the NE-SW trend of the boulder train of the Cyrena and Katholm erratics and the sense of the dispersal of Kinnediabaas (Baartman & Christensen, 1975). The consistency in the NE-SW direction, together with the inconsistency of the porphyry distribution in the SE-NW direction, indicates that the effect of any Baltic ice stream was very small during the Weichselian.

The moderate force of the Young Baltic ice stream as compared with the vigour of the NE flow is confirmed by the comprehensive reports by Hansen (1965) and Lundqvist (1965). Hansen's publication shows that whereas the islands of the Danish archipelago were hardly real obstacles for the NE ice stream (which must have ridden roughshod over them on its way to Jutland), those same islands were virtually insurmountable obstacles for the Young Baltic ice stream, which only succeeded in passing through the channels the Little Belt, the Great Belt, and the Öresund. The impotence of the Young Baltic ice stream is also evident from the 1965 paper by the Swedish geologist J. Lundqvist, who saw the Young Baltic ice stream, which was surrounded by melt-water, as the result of a shortliving effort of an advance of the inland ice during the deglaciation of Scania.

#### *Striae in Denmark and the surrounding region*

According to Charlesworth (1957), Torrel's recognition in 1865 of a Younger Baltic ice stream was based partially on the double systems of striae of Scania, Bornholm, Gotland, and northern Germany. There are NE striae which are consonant with the consistent ice stream and others which are attributed to the inconsistent stream.

As far back as the second half of the nineteenth century the striae deviating from the NE direction in southern Scania and Bornholm, and from the N to NW trends in Öland and Gotland, intrigued the Swedish geologists. In 1885, G. de Geer made a comprehensive study and pointed out that the divergent striae post-date the retreat of the land ice and occur only below the 70 m mark above present sea level. Detailed striae analyses made by Mattsson (1960) confirmed that the aberrant striae in Scania and Bornholm were made during or after the retreat of the NE ice. According to Hillefors (1967), the same holds for the divergent striae on Rön and Malö, two islands lying a few miles offshore halfway between Göteborg and Varberg (see Fig. 21).

Both Mattsson and Hillefors point to the force of the NE ice, which can be

moraines, were deposited in the sea, the Skagerrak must be abandoned as a possible region of alimentation of the Kattegat ice stream. A much more northerly open sea is consistent with the attribution of these aberrant striations to the action of floating ice. In 1974, Hillefors reconsidered the SSE striae. He questioned their meaning and gave preference to main NE-SW ice movements from the Göta river valley into the Kattegat.

#### *The Young Baltic moraine in southern Scania*

The main direction of ice movement in Scania was from NE to SW. On top of the moraine of this so-called NE ice, but often separated by intermorainic glaciofluvial sediments, lies the Young Baltic moraine. What is its horizontal extent? Fig. 22 shows that the Young Baltic deposits are found in a 15 km wide strip running roughly parallel to the coast. Topographically, the Young Baltic moraine is restricted to the coastal plains, or, as stated by G. de Geer in 1885, the Baltic ice was only able to transgress the lowest and flattest parts of Scania.

The thickness of the upper Baltic moraine ranges from 2 to 6 m. In contrast to the moraine of the NE ice, which is generally deficient in clay, the Young Baltic is clayey. According to Vortisch (1972) the Baltic clay bears a resemblance to water-laid sediments. Carbonates are virtually absent in the moraines of the NE ice, but the lime content of the Baltic deposits is high. The Baltic moraines are characterized by indicators from the Baltic, in the form of boulders from the Åland islands and red and brown Baltic porphyries, which is in agreement with the porphyry counts made by Grey (1932), who found that these Baltic indicators predominate in a narrow coastal strip (15-25 km wide) in western and southern Scania.

Unlike the NE ice stream, the Young Baltic in Scania must have been weak.




Fig. 22. Horizontal extension of the Young Baltic moraine in Scania, after Vortisch (1972).

On this point the investigations of Gustafsson & Stjernkvist (1966), who studied a moraine section occurring in the northern part of the province of Lund, are of interest. In their opinion, the intermorainic sands of this section were deposited on beaches of lakes, and when the water became deeper clay covered the sand. Later, the Baltic ice entered the area but the intermorainic sediments were not affected by ice pressure. These authors thought it quite possible that the ice advanced into a lake.

The age of the Young Baltic is still uncertain. The clay-varve method of G. de Geer, pollen-analysis correlations done by Nilsson, and radiocarbon datings have yielded an absolute chronology reaching back to 12 000 years B.P. (Håkansson, 1971). On this basis, the oldest reliably dated recession line in Sweden lies roughly on the northern boundaries of Scania and Blekinge. Owing to absence of varved Late Glacial sediments in most of the area south of this line, it has not been possible to perform a continuous line of varve measurements further southward.

In central Denmark the absolute time scale covers the Holocene. No  $^{14}\text{C}$  dates are yet available for the main stationary line in Jutland or the ice-border stages (C-H) in eastern Denmark, so that here too the absolute age of the Young Baltic is unknown. We know from the review by Hansen (1965) that the Young Baltic penetrated the Little Belt, the Great Belt, and the Øresund. The Øresund glacier, which also covered Scania, is thought to have been the final phase of the Young Baltic ice stream.

Relatively, the Young Baltic in Scania is younger than the NE ice stream, because the Baltic Upper moraine lies on top of the NE moraine. The intermorainic sediments and the Baltic moraine covering the osar of the NE ice stream, as observed by G. de Geer (1885), means that there must have been an ice-free region before the final advance of the Baltic.

The assumption of the existence of a Young Baltic ice stream in the interpretation of the field data leads to the following unharmonious picture. One ice stream retreating to the north during a general improvement of the climate, was trailed by another ice stream — now coming from the south — despite the same improved climatic condition. The advancing ice stream even invaded the partially submerged deglaciated area from which the receding ice stream had withdrawn. This contradictory picture lies at the root of the differences of opinion about the age relations between the NE and the Baltic ice defined by Lundqvist (1965), Vortisch (1972), and Mohrén (1973).

But another approach, based on the gradually contracting ring of marginal glacier lakes and the receding Scandinavian ice sheet as integral parts of a whole, leads to a more regular development of retreat. In this view, the circum-Baltic zone of the Urstromtäler swelled to a cohesive body of water in the Baltic depression and tailed off into a valance of separate lakes and lacustrines against the upgrading grounds of Sweden. Thus, during the climatic amelioration the margin of the ice sheet in the Baltic depression bordered on a ice lake under arctic conditions. In this light the restriction of the Baltic moraine to the coastal plains of Scania, its clay resembling water-laid sediments, and the weakness of the presumed ice stream, seem to point to a para-till, as Harland, Herod, and Krinsley (1966) defined deposits formed by ice-rafting. In this context it is worth noting that Spjeldnoes (1973) suggested the possibility that 'a number of tills which have been described as bottom moraines — especially in the marginal part of a glaciated area — were really ice-rafted'.

*The glacial morphology of the Weichselian in Denmark*

The features of the Quaternary landscape in Denmark are well illustrated by Hansen and Milthers. Their survey map, published by Hansen in 1965, is repro-




Fig. 23. The Quaternary landscape in Denmark, after Hansen (1965).

duced in Fig. 23. From this map it is evident that the glacial morphology of the Weichselian in Jutland, i.e. the greater part of Denmark, is clearly discernible. On the Danish archipelago, however, this structure is less pronounced. The glacial pattern is even obscure, and it is hardly surprising that the interpretation of these incoherent field data (in contrast with those in Jutland) led to the many question marks that Hansen and Nielsen (1960) had to put on their map of the glacial morphology in southern Denmark.

There is no consensus among the Danish geologists on the 'significance of the different ice borders and recession lines in southern and eastern Denmark and the stratigraphic models accordingly', as Berthelsen (1973) stated. More recently, Binzer (1974) questioned whether the glacial advances from different directions really took place in Denmark. The difficulty underlying the conflicting views is that the islands do not form a related whole. Glacio-morphologically, the Danish archipelago looks like a jig-saw puzzle with some of the parts missing. This complicates attempts at correlation, the more so because of the lack of palaeontological evidence.

Another reason for the indistinct glacio-morphological pattern of southern Denmark are the glacial features that intersect the trend of the NE ice stream. These features are ascribed to the 'Younger Baltic' or the 'south-east ice' which is thought to have followed the NNW-SSE-running depressions between Jutland, Fyn, Sjaelland, and Scania. The Little and the Great Belt glaciers only touched the coastal parts of W, S, and E Fyn (Hansen & Nielsen, 1960), but crossed the coastal plains of southern Jutland and of western Sjaelland. Even so, the intersecting features may be seen in another way, and we will return to that point in the section on the Weichselian recession in Denmark.

### The consistent distribution unit

#### FURTHER ANALYSIS AND ELABORATION OF THE FLOWLINE PATTERN

##### *Flowline pattern, eskers, moraines, and recession lines*

After taking stock of the indicator distribution in the marginal belts of the Weichselian and Saalian ice sheets, we found that the flowline pattern in Fennoscandia satisfies a consistent distribution. Conformity between the distribution and the flowline pattern means that the glacial directions refer to the advance.

The trends of most eskers, as indicated by Flint (1971), parallel in a general sense the directions of ice flow during deglaciation. These features of retreat have been mapped on a regional scale in Fennoscandia, and shown in combination with the directions of movement in Fig. 24, from which it is obvious how well the patterns of flowlines and the system of Fennoscandinavian eskers blend. Consequently, the pattern of ice flow is consistent with the retreat as well.

Just as a set of stained rings on the inside of an emptied bowl bears witness to the changing form of the liquid it contained, the recession lines at intervals of a thousand years — shown schematically by E. H. de Geer (1954) — throw light on the transformation of the last Scandinavian ice sheet during its retreat. These lines, which are shown here in Fig. 25, are intersected perpendicularly by the




Fig. 24. Pattern of eskers in relation to ice movement.

Swedish and Finnish eskers. Because eskers parallel the directions of flow during deglaciation and the main directions of ice movement must in general have been transverse to the ice margin, the esker pattern agrees with that of the streamlines of the Weichselian ice sheet during its retreat. The recession lines together with the

limits of the Weichsel (We) and Saale (S) glaciations, the Saalian ice front phases a, b, c, d, and e, after ter Wee (1962), and the terminal moraine of the Warthe (Wa) Substage, all seem to form part of the same concentric pattern. If this is actually the case and since the maximum extents indicate the extremes of the growth of the ice sheets, an advance of the north European ice mantle along the same course as the retreat seems likely here too.

This may also have been the case in Canada. Flint (1971) discusses the close association of the streamline features of a large part of this country with end moraines and esker systems in particular, both of which he takes as features of deglaciation. This flowline pattern also conforms, however, to that of the maximal ice sheet obtained by analogy simulation with a field plotter. As a result, Flint was uncertain as to whether these glacial directions were related to the advance or to the retreat of the ice sheet. This uncertainty itself might be an indicator that in the broad sense the general pattern of movement was similar in the advance and the retreat.

The re-expansion effect occurring during the general shrinkage of a glacier suggests the same thing. As reported by Flint (1971), radiocarbon investigations of end




Fig. 25. Recession lines and maximum extension of ice sheets in relation to esker pattern. For further explanation, see text.

moraines from the last 10 000 years give the impression that on the whole these forms are the result not of pauses in the course of general shrinkage of a glacier, but of culminations of re-expansion during the period of retreat. Small advances occurring while the ice was retreating indicate that the ice was still flowing at least by fits and starts. Thus, on the whole, the moraines left by the retreating ice may after all promote information about the advance as well.

The Fennoscandian recession lines are shown in Fig. 25 together with some limits of various glaciations and the flowline pattern. The two systems intersect roughly perpendicularly, which suggests that they contribute to a whole. The flowline pattern is in agreement with the esker system but also with the consistent distribution of indicators; the recession lines are features of shrinkage but the lines of maximum extent belong to a growing ice sheet. Thus, there are features which belong to both advance and retreat. Thus far, in a tentative way we obtained a general picture of a possible flow pattern and outlines of the last Scandinavian ice sheet. The further investigation, completion, and elaboration of this integrated structure will be discussed in the following subsections.

*Approximation of the outlines and flow pattern of the ice sheets by a set of confocal ellipses and hyperbolas*

As shown in Fig. 26, the recession lines may be regarded as rough ellipses lying concentrically around common foci. Particularly in the southeastern direction they waver around means of nearly true elliptical contour, but south of Oslo, in southwest Sweden, in the Danish archipelago, and in Jutland, a clear indentation in these series of concurrent ovals attracts the attention. Deviations from the true elliptical contour should of course be expected. Ice of moderate thickness tends to follow the depressions, but with greater thickness the ice movement is more independent of the subglacial relief. It is interesting to note that elliptical outlines reappear in the reconstruction of the Scandinavian ice sheets made by Aseev (1968), by Aseev et al. (1973), and in the isobases of the domelike upwarping in Fennoscandia.

If the last ice sheet in northwest Europe maintained a roughly elliptical shape while it shrank, the theoretical direction of ice movement would be perpendicular to the circumference of each ellipse, and this raises the question of the extent to which the trends of ice movement found in Fennoscandinavia coincide with the theoretical directions pertaining to the concentrical ellipses.

The normal curves being perpendicular to the greatest diameter of the central ellipses, this diameter should intersect the equivalent sections of the flow pattern at right angles too. The strong curvation in Dalarna, as well as the perpendicularity of the directional trends north of Maan Selkä to those in central Sweden, offer *prima facie* an indication of the rough lie of the greatest median. To approximate the true position, the assumed median should be adjusted to intersect the trends of movement in central Sweden at right angles. Fig. 27 shows the result to be a remarkable fit, the normal curves of the concentrical ellipses running parallel to the Fennoscandinavian flow pattern. There is a deviation in the extreme southwestern part of Sweden, but a glance at Fig. 26 shows that for south Sweden the deviation concerns the divergence of the recession lines from true ellipses.




Fig. 26. Recession lines after E. H. de Geer (1954) in relation to confocal ellipses. The approximate dates are given in years B.P. Heavy lines: ellipses; dotted and dashed lines: recession lines.

#### *Location and form of the main ice divide*

Curves normal to confocal ellipses are hyperbolas, and the major axis of the ellipses, which is also the median of the set of confocal hyperbolas in Fig. 27, may be considered a theoretical ice divide.

East of this median, the directions of arrows and lines match fairly well. On




Fig. 27. The trends of curves normal to the confocal ellipses in relation to directions of ice movement.

the west side, however, there is some disagreement in the axial zone. This incongruity may have been brought about by the westward movement of the ice divide at the end of the last glacial, the shifting ice mass disturbing and obscuring




Fig. 28. The Weichselian ice sheds (Andersen, 1956; Lundqvist, 1956) and the median F<sub>1</sub>F<sub>2</sub> of the confocal hyperbolas. The ruled parallelogram in Finnish Lapland is a zone of the ice divide, and the arrows show trends of the ice movement of the Weichselian main glaciation (Kujansuu, 1967). The trends of ice movement can be compared with the directions of some hyperbolas in this area.

the features of the main glaciation. At some distance from the median, east of the Swedish-Norwegian border and in Norway, the trend of ice movements and hyperbolas run more or less parallel again, except for the southern part of Norway, to which we will return below.

Although the divide of the ice sheet presented by Aseev et al. (1973) lies further to the east, since their construction is also based on the isostatic subsidence, the difference in position might be explained by the structure of earth crust and substratum in this area. The easternmost ice divide proposed by Lundqvist (1974) approximates the Swedish component of our median. To the north, however, the divide of Lundqvist continues eastward across northern Finland and follows the longitudinal axis of the Kola peninsula. The directions of ice flow consistent with

this deviation to the east, should be approximately northward and southward. But the observed directions, which are from northwest to southeast, do not support Lundqvist's hypothesis.

Our view of the topographical location of the median of the confocal hyperbolas is supported by the general agreement between the directions of ice movement and those of the hyperbolas. The principal axis separates the concordant trends of ice movement to the east from the deviating trends to the west. Furthermore, the ice divide zone of the Weichselian main glaciation in the western part of Finnish Lapland, which Kujansuu (1967) defined after careful glacio-geological investigations, is consistent with the hypothetical zone, the trends of the main glaciation lying parallel to the northern hyperbolic directions of ice movement (see Fig. 28). Apart from the conformity between the hyperbolic trends and both the directions of movement and the distribution pattern of the indicators in northwest Europe, the criteria mentioned above imply that the main ice shed would roughly coincide with the median in question. On this basis, the westward deviation of the northern end of the main ice divide, as represented in Fig. 28, seems to be part of one of the younger ice sheds.

The southern end of the Weichselian ice shed also diverges from the theoretical median. The aberrant component, running roughly parallel to the Norwegian coast (like the greater part of the median), and the opposite ice movement directions on both sides of the deviation, confirm the impression that this southern feature represents a bend of the main ice divide. If this is the case, the bend should manifest itself in the streamline pattern.

#### *The zigzag bend in the south end of the divide; resulting flowline pattern and morainic contours*

Figure 29 depicts the re-entrant and salient angle of a bend with straight streamlines perpendicular to the sides. These lines diverge in the larger angle and converge in the smaller angle. Since the theoretical streamlines are not straight but hyperbolic, the straight lines are replaced by hyperbolas in Fig. 30. Divergent streamlines give rise to a convex ice front, convergent streamlines to a concave ice front; in other words, re-entrant and salient angles are reflected by the shape of their respective ice fronts.

Armed with this knowledge, we return to the flowline pattern in Fig. 31. The divergent streamline system in southeast Sweden, with its apex adjoining the re-entrant angle of the ice divide, confirms the postulated bend in the main ice shed. Since the convex south-Swedish recession lines and moraines represent the re-entrant angle of the bend in the ice divide, the concave Ra moraines of the Oslo district could reflect a salient angle in the divide. The Ra-Gothiglacial Substage moraines form a kind of Z bend that also shows up in the main stationary line in Denmark and matches the crooked Norwegian ice divide as well (see Fig. 31).

It seems as though the shape of the bend in the ice divide is expressed in the outline of the southwestern moraines and as though a rotation of the ice front took place. Figure 32 shows that a common centre of circles describing the rotation can be determined.

Because these circles run parallel to the esker system in southwest Sweden and to the dividing lines between the area with prevailing rhomb-, Dalarna, and Baltic porphyries in Denmark, they are represented in Fig. 33 next to the diverging bundle of hyperbolic lines at the southern end of the divide to permit comparison




Fig. 29. Straight streamlines in re-entrant and salient angles.




Fig. 30. Hyperbolic streamlines in same angles as in Fig. 29. Dashed lines: resulting ice fronts.




Fig. 31. The zigzag bend in the ice divide, in the Ra Central Swedish moraines (RCS), and in the main stationary line (M) in Denmark.

with the main directions of ice movement. The constructed directions match those of the ice movement fairly well. This supports the idea of an ice flow across the Skagerrak (Holtedahl & Bjerkli, 1975). Nevertheless, there are traces of ice movement in southwest Sweden that deviate. Because in general the esker and stream patterns correspond rather well with the circular lineature (see Fig. 32 and 33), the deviations might be explained by topographic control during deglaciation.

The general outline of this flow pattern related to the zigzag bend in the ice divide is correlated with the dispersal of some major groups of indicators (see Fig. 33). On its east side, the Oslo district is encircled by circular lines of ice flow. The westernmost circle runs west of the Oslo fjord and touches the northwest coast of Denmark. This configuration confines the glacial dispersal of the Oslo indicators to a relatively narrow circular zone that corresponds with the presence of these indicators in north Jutland, which lies in this zone, and the absence of these indicators in The Netherlands, which lies well outside this zone.

The immense dispersal area of the Dalarna indicators, reaching from Denmark and The Netherlands up to Lithuania, follows from the position and shape of the divergent system, whose apex lies at the Swedish-Norwegian border in Dalarna and whose westernmost streamline follows the easternmost circular line.




Fig. 32. The circular pattern.

#### THE INFERRED STRUCTURE OF FLOWLINES IN RELATION TO THE DISPERSAL OF INDICATORS

The analysis of the linear trend of the flowline pattern resulted in a generalized descriptive picture with three geometric components: a set of confocal hyperbolas, a divergent system, and a zone of concentric circles. This picture not only visualizes the outline and interrelationships of the flowline pattern, but also permits interpolation or extension of the initial delineation. As indicated above, the dividing lines of the indicator distribution were lengthened toward the flowline pattern in Fig. 20; now, in Fig. 33, the extension starts from the flowline pattern.

This new approach again shows the close relationship between distribution

and flowline pattern. It also enables us to fill in the interspaces between the few connection lines of our first attempt, which means that a fresh and a more detailed appraisal of the interrelationships between the inferred lines of ice flow and consistent distribution is feasible.

Thus, the rest of this section concerns the dispersal of several species of indicators, with emphasis on the corresponding tracts of the flowline pattern. Most of these indicators have already been taken into consideration, but there are others that require attention.

#### *The tract of flowlines toward The Netherlands*

The key position of The Netherlands with respect to Pleistocene stratigraphy — according to Brouwer (1963) and elaborated by de Jong (1967) — also holds glaciologically, because the southwesternmost continental traces of the Saalian ice sheet lie within the Dutch borders. In addition, since the mean ice movement in the southwestern sector of the ice mantle that covered The Netherlands, Denmark, and southern Sweden was roughly NE-SW (see Fig. 33), the corresponding actual paths of ice movement are accessible in this area, because at present they lie almost entirely overland. This amounts to an ideal situation for a reconnaissance of the route of consistent transport in the southwestern sector of the Scandinavian ice sheets.

There is one drawback, however. The flowline pattern is indistinct here and not yet understood. On the other hand, thanks to the investigations of Maarleveld (1953), ter Wee (1962), and Zagwijn (1974), the locations of various ice front phases of the Saalian are known. Starting at the southernmost location, flowline directions are drawn normal to the general regional trend of these phases. This procedure is permissible because of the correspondence between the directions of ice movement in the Saalian and Weichselian and the similarity between the general patterns of ice flow of the expanding and the shrinking ice sheets.

Figure 34 shows this retraced tract, which with its southern end in north-eastward direction turns to the east. It is interesting to note that this course parallels the valleys of the Saalian ground moraine in the northern part of The Netherlands and northwestern Germany (Edelman & Maarleveld, 1958). Maarleveld thought it likely that the prevailing direction of these valleys reflects the direction of movement of the Saalian ice sheet. Further to the north, the route perpendicularly intersecting the moraine of the Warthe Substage proves to be normal to the general trend of the end moraines of the Last Glacial as well, and, then turning through east northeast in northwest Germany and the Danish archipelago, the tract goes straight north in southwest Sweden, and even takes a northwest direction beyond Lake Vänern. The excellent maps published by G. de Geer in 1910 make it possible to compare the south-Swedish part of this path of retreat with the directions of åsar and thrust striae; the results show a close resemblance in general trend.

If the transport of erratics, with The Netherlands as ultimate destination, took place along the roughly Z shaped route starting in the Dalarna area, an appreciable percentage of Dalarna indicators would be expected in the Dutch boulder clay. This is confirmed by the findings of van Calker (1912) and de Waard (1949). The latter made a special study of the erratic associations in the boulder clay of the Noord Oost Polder, which was once part of the Zuiderzee. Other investigators (van der Kley & de Vries, 1941) have also reported that Dalarna




Fig. 33. Inferred flowline pattern (heavy lines) of the ice sheet in relation to directions of ice movements (arrows) and the distribution of indicators (See also Fig. 4, p. 16).

porphyries, Dala sandstones, and diabases from Dalarna occur generally in The Netherlands.

Although several authors ascribed the provenance of the numerous flints found in most of the Dutch glacial sediments to the Cretaceous outcrops in Jutland,

the Danish archipelago, southern Sweden, or northern Germany, Veenstra (1963) traced these flints back to the Danish archipelago and southern Sweden. His studies concerned, for instance, the Bryozoa enclosed in the boulder clay. The locality of this provenance argues strongly for a transport of the flints along the southern part of the Z shaped course. This would explain the exceedingly high percentages of flint of the coarse fraction of erratics in the Weichselian glacial deposits on the




Fig. 34. Reconstructed tract of flowline pattern through The Netherlands. I: topographically incorporated indicator diagrams (Fig. 15). The key also applies to Figs. 35-37, 50-53.

east side of Schleswig Holstein, where there are no Cretaceous outcrops. During an excursion conducted by A. C. Koenderink and P. Schuddebeurs in July, 1968, the present author made the following rough estimates (see Table 10) in a number of exposures situated in the hatched area in Fig. 34.

Table 10. Approximate flint percentages of the coarse fraction of erratics on the east side of Schleswig Holstein.

Locality	Kind of deposit	Percentage of flint
Stendorf am Bungsberg	fluvioglacial	70 to 80
Kasseedorf south	fluvioglacial	10 to 20
Kasseedorf north	fluvioglacial	30 to 40
Kassau	fluvioglacial	25
Kreuzfeld 1	fluvioglacial	20 to 30
Kreuzfeld 2	fluvioglacial	30 to 40
Luschendorf	fluvioglacial	30
Malente south	fluvioglacial	30
Sieversdorf 1	fluvioglacial	70
Sieversdorf 2	fluvioglacial	70
Weissenhaus	fluvioglacial	30
Övelgönne (Süsel)	fluvioglacial	70
Retin (Neustadt)	boulder clay (coastal cliff)	50 to 60

Finally, the puzzling scarcity of Oslo indicators and Bohuslän granites in The Netherlands tallies with a route of transport bypassing their provenances.

Are the ratios of the indicators in The Netherlands also consistent with this transport route? Returning to Table 6, we learn that — excluding the Hondsrug and the lenses of red boulder clay on account of their large quantities of Baltic material — the added percentages of the Scania, Småland, and Dalarna groups amount to more than 30%. At the same time, in cases other than the Utrecht ridge, where the Bornholm indicators amount to 11%, the shares of the Oslo-Swedish west coast and the Bornholm-Blekinge unit do not exceed 4%.

Thus, the predominance of the combined percentages of indicators from Dalarna, Småland, and Scania over those from the adjacent areas also supports a Netherlands-bound transport route starting from Dalarna and proceeding roughly through the median part of southern Sweden, bypassing the Oslo area and the Swedish west coast to the west, and to the east, the province of Blekinge and the island of Bornholm. If the flints from Scania and the Danish archipelago had been included in the Hesemann counts the results would have been much more pronounced, since the flint content of the recognizable indicator association in the normal grey boulder clay in The Netherlands amounts to at least 28%. Thus, for a better understanding of the directions of ice flow, there is much to be said for not restricting counts to the crystalline or sedimentary erratics only, as has been done because of the two widely divergent specialities. Nevertheless, for the present study the Dutch Hesemann counts were not altered in this respect, leaving them suitable for comparison with crystalline counts made in other areas.

In concluding the discussion of the route of indicator transport toward The Netherlands, reference should be made to the danger involved in inferring spatial relations of ice movements from the Hesemann numbers with the reserve that Hesemann numbers lent themselves above all to stratigraphic ends. Ter Wee (1962)

directed attention to an interesting relationship between the Hesemann numbers and the Saalian ice front phases in The Netherlands. From the maximum extension of the ice northward, the indicators from southern Sweden (Hesemann area III) decrease and the north Baltic elements (Hesemann area I) increase, which ter Wee attributed to a west-to-east shifting of the ice stream directions upstream. Despite the rational conclusion ter Wee drew from the Hesemann numbers, the frequency curves in Fig. 15 do not account for a west-to-east shift of the source areas in the successive recessional Saalian ice front phases in The Netherlands. On the contrary, as illustrated by the incorporated indicator diagrams, the curves of Fig. 34 pertaining to the three westerly oblong source units show an almost perfect symmetry, which supports a fixed path of the Netherlands-bound inland ice.

The lines of motion of the Saalian ice movement represented by Münnich (1936) are also based on Hesemann numbers. His flowlines from Dalarna toward The Netherlands — showing a convex curve to the northwest — pass through Bohuslän, the west coast of Sweden, and Jutland, instead of curving to the southeast and transversing the Danish archipelago and Schleswig Holstein as indicated by the dispersal of indicators in The Netherlands.




Fig. 35. The consistent distribution of the areas between the Rhine and the Elbe and on both sides of the Oder. For key, see Fig. 34. The capital letters refer to Table 8 (p. 41), the numbers to Tables 3 (p. 23) and 4 (p. 27).

### *The consistent distribution east of The Netherlands to the Oder and beyond*

Compared with the route of indicator transport toward The Netherlands, a section of the reconstructed flowline pattern covering most of the area between the Rhine and the Elbe, includes more of the source area of the Småland granites and lies further from Oslo (Fig. 35). This conforms to a higher amount of Småland granites and a greater scarcity of Oslo indicators in the Saalian indicator association of Western Germany.

The route of the indicator transport toward the region west of the Oder is still further away from the Oslo district and the Swedish west coast, which might account for the absence of indicators from these sources on the west side of the Oder. In accordance with the transport route running through Dalarna, Småland, Scania, and Bornholm, indicators from these provenances are here abundantly represented. Except for the Baltic association, their combined percentages range from 93 to 99, as is apparent from Table 3A.

When the percentages in Table 4 for the Weichselian on the east side of the Oder are considered, the deficiency of the Bornholm indicators is manifest, as could be expected with a course of ice flow that bypassed this island.

### *The gap in the distribution pattern*

Although the gap in the distribution pattern is due to the lack of counts, Ladwig (1938) did some 80 Hesemann counts for the Weichselian of eastern Pomerania in an area west of Gdansk denoted by EP in Fig. 36. Unfortunately, he gave only the Hesemann numbers, which means that his results cannot be regrouped. Besides making these counts, however, he investigated the frequency of occurrence of a number of sedimentary erratics and found that the Neksö sandstones cropping out on the island of Bornholm were scarce in eastern Pomerania, as distinct from the indicators of *Paradoxides tessini* Sandstone, whose source is in the southern part of Öland. The highest density of the distribution pattern of this sandstone in the central part of the area investigated by Ladwig declines to the east and west, which tallies with the flow pattern, since the reconstructed streamlines cutting across the southern part of Öland run virtually through the centre of this region. The streamlines traversing Bornholm, which pass well to the west of the site in question, agree with the observed scarcity of the Neksö sandstone.

In analogy with this, the major dispersal of the 'Beyrichien Kalk' in Poland (Reuter, 1885) and its main source area southeast of Gotland (Hucke, 1967), both of which lie west of the flowline through Kaliningrad (see Fig. 36), is consistent with the inferred directions of ice flow.

### *Comparison of the Saalian in Poland and White Russia with the Weichselian in Lithuania*

As shown in Fig. 36, the segment of the inferred flow pattern covering the regions investigated by Milthers & Milthers in Poland and White Russia is made up of two parts. One contains the main source areas of the Dalarna porphyries together with regions I, II and III, and the other the Åland islands with regions IV and V. Thus, the inferred tracks toward regions I, II and III have their seat in the provenance of the Dalarna indicators, which is consistent with the fair amount of Dalarna porphyries in the Saalian indicator associations of these regions.




Fig. 36. The Weichselian in Lithuania in relation to the Saalian in Poland and White Russia. EP: area west of Gdansk (in the gap in the distribution pattern). Source areas as in Fig. 34, I-V, regions where Milthers & Milthers performed their counts. W: west, C: central and NE: northeast Lithuania after Tarvydas (1960). Solid-line segments: dividing lines between areas with predominant porphyries in Denmark. Dashed lines: inferred flowline pattern.

Apparently, the Dalarna porphyries were transported along a corresponding route in the Weichselian, because G. de Geer (1885) concluded from the presence of Alvdal porphyries on Gotland and their absence in the Uppsala region, that they passed west of Uppsala and reached Gotland from the NW. Königsson and Svantesson (1975), however, questioned the genuineness of the Alvdal porphyries. They refer to Hedström, who pointed out in 1894 that the porphyries on Gotland originated from the bottom of the Baltic north of the island. This may be partly true, but the high percentages of Jotnian sandstones in the gravel deposits imply that, likewise, porphyries from Dalarna are bound to occur on Gotland. The decline of the Dalarna porphyries from west to east in the investigated area in Poland also agrees with the position of the tracks under study, since the distal parts of those toward regions IV and V lie outside the source areas, and such that the farther they lie to the east, the greater the distance to the outcrops. Although the main provenance of the Dalarna indicators is not covered by the routes to regions IV and V, low percentages of Dalarna porphyries are still found in these parts of White Russia. If, however, the distal sides of the routes to IV and V are

extended northwestward, they come to lie over outcrops of Dalarna, or more precisely Bredvad, porphyries along the Swedish-Norwegian boundary in Jämtland. The locations of these porphyries, it is true, lay on the wrong side of the main ice shed for their boulders to be carried in glacial transport toward the southeast. Even so, since the outcrops are situated on the east side of the main watershed, boulders of the Bredvad porphyries must have been carried to the east by river transport in the pre- and interglacials in order to have been picked up in the glacials by the southeastward-moving ice stream. This might explain the presence of part of the Bredvad porphyries in regions IV and V. On the other hand, in view of the opinion of Eskola (1933) it is conceivable that outcrops of porphyries similar to the Dalarna types occur on the sea-bed of the Baltic or the Gulf of Bothnia.

The areas of the theoretical flow pattern that connect the Åland islands with regions IV and V in White Russia cover the central part of Lithuania, where, as already mentioned, indicators from Åland predominate. Thus, in analogy with the conveyance of the Dalarna porphyries, the route by which the Åland indicators were conducted to regions IV and V in the Saalian seems to coincide with the transport route of these indicators in the Weichselian as well, but with central Lithuania as the destination. Viewed in this light, the predominance of the Åland indicators in central Lithuania as compared with the 89 and 88% of such indicators in the White Russian regions is explained. A noteworthy point is the parallelism of the dividing lines separating Tarvydas' meridional zone with the theoretical lines of ice movement.

As can be seen from Fig. 36, the hypothetical route of ice movement close to the southwestern tip of the Finnish mainland and over the adjoining archipelago connects up with the area of NE Lithuania where indicators from southwest and south Finland predominate. It is true that Viborg and the neighbouring regions of Karelia lie outside this track, but Eskola (1933) pointed out that rock varieties from these areas also crop out on the sea-bottom and the archipelago southwest of Finland. This is the case, for instance, for the Viipuri or Viborg rapakivi. Therefore, there is little or no point in distinguishing between indicators from southeast and southwest Finland. On the other hand, east Finnish indicators could have been shifted westward by floes and icebergs before the advance of the ice. According to Hyypä (Donner, 1965), such transport took place in the Finnish Gulf during the last retreat of the inland ice.

The introduction of south Finnish indicators into the lower moraine horizons of the Eopleistocene and Mesopleistocene in SE Lithuania could just equally well have been promoted by partially waterborne transport. However, Gaigales (1963, 1965), who called attention to these lower horizons, suggested an ice movement from north to south to explain the presence of the south-Finnish material, and held a NW-SE ice stream responsible for the indicators from northern Sweden and the Baltic, which are encountered in the upper horizons of the moraines of the Eopleistocene and Mesopleistocene and the moraines of the Neopleistocene in southeastern Lithuania.

Gaigales and Gudelis (1965) specified percentages of the various groups of indicators in the Brandenburgian, Frankfurtian, and Pomeranian substages in SE Lithuania, a region which ranges under central Lithuania, as designated by Tarvydas (1960) in his tripartite regional distribution of the Pomeranian Substage. The indicators of the Åland islands, which fluctuate between 51 and 54% predominate not only in the Pomeranian but in the two older substages as well. Besides the Åland group, there are those of the Baltic sea-floor and of Sweden,

with average percentages of 16 to 20 for the three substages. The indicators from southwestern Finland are in the minority, their percentages ranging between 5.4 and 13. The distribution of the association's percentages in each substage agrees with a NNW-SSE drift over the Åland islands toward SE Lithuania. This also holds for the Pomeranian Substage, despite the N-S ice stream proposed by Gaigales & Gudelis in 1965, because the 5.2% difference between the south Finnish associations of the Brandenburgian and Pomeranian substages is not sufficient to make an exceptional case of the direction of the Pomeranian glacier movement. On closer analysis of the erratic association of the last glacial, Tarvydas (1967) too arrived at the conclusion that in both the Brandenburgian and the Pomeranian the direction of the ice flow toward Lithuania was more or less the same, i.e. roughly NNW-SSE.




Fig. 37. The indicator fans of Kraus, Tschirwinsky and Hausen in relation to directions of the confocal hyperbolas. See also key to Fig. 34.

Further inland, southeast of Lithuania and Latvia, lie White Russia and Wolhynia. To what extent do the indicator associations of these regions fit into the dispersal pattern of the Baltic coastal areas? Tschirwinsky (1926) found that the indicators from Aland predominate in Grodno, a region lying between central Lithuania and Milthers' region IV in White Russia (see Fig. 36 and enclosure map). These indicators also predominate in the western part of Wolhynia, where, according to Gagel & Korn (1918) there is also a small amount of Dalarna porphyries.

In the eastern part of Wolhynia (the Kiev and Tschernigov districts), and the eastern part of White Russia (southeast of Minsk and the Mogilev districts) Tschirwinsky (1926) found neither Dalarna nor Baltic porphyries and only a few Aland indicators, but indicators from southern and southwestern Finland were in evidence. Since these areas lie north of the line of ice flow (see Fig. 36 and folding map), which all but coincides with the dividing line separating the central part of Lithuania from the northeast corner of that country, where indicators from Finland predominate, the indicator dispersal pattern of the eastern coastal areas of the Baltic continues further inland to the southeast, in accordance with the inferred lines of ice movement.

#### *Indicator fans in the eastern circum-Baltic*

Kraus (1934) is one of the investigators who worked on the reconstruction of indicator fans. Some of his fans, whose eastern limits are shown in Fig. 37, were subjected to further investigation in the present study. Because, as Eskola (1933) pointed out, the provenance of the Viipuri rapakivi is not restricted to its type locality, the western lateral limit of this indicator fan as shown by Kraus cannot be maintained. Since rapakivi granites related to the Viipuri type also occur in the rapakivi areas on the west coast of Finland, the initial point of the western lateral limit of this fan would have been west of the Finnish west coast and not on the west side of the Viipuri massif. Accordingly, the shifted limit trends roughly NW to SE instead of N by E to S by W, as in Kraus' representations.

The easterly lateral limits of the indicator fans represented by Kraus (1934) run more or less parallel from NW by W to SE by E and intersect the hyperbolas at an angle of 20-30°. It should be pointed out that the eastern limits of the indicator fans, whose apices lie north of the Baltic and the Finnish Gulf, are more reliable for determination of the consistent directions of ice movement than the western lateral limits in this area. There are two main reasons for this. Firstly, the westward extension of the provenances of some of the important indicator types cannot be directly observed, e.g. outcrops of the Viipuri rapakivi and the Suursaari (Hogland) porphyries are theoretically present under the Baltic. Secondly, for the western limits lying mainly in the Baltic, i.e. the territory of the Baltic ice stream, these limits are more closely related to the inconsistent than to the consistent ice movement. This means that the eastern limits of the indicator fans are indicated to learn us more about the directions of movement of the inland ice in this area.

As represented by Kraus (1934), these limits intersect the inferred lines of ice movement at an angle of 20 to 30°. Figure 37 shows that the eastern limits defined by Tschirwinsky (1926) and Hausen (1912) correspond much better with the trends of the hyperbolas especially those of the Viipuri granites, the Suursaari (Hogland) porphyries, and the Tammela uralite-porphyrates. The eastern limit of the Aland indicator fan deviates distinctly from the direction of the hyperbolas,




Fig. 38. Dispersal of indicators in northwestern Russia as described by Ramsay & Hackman.

which might be due to the lack of information about the exact submarine extension of the Åland rapakivi massif.

These three investigators are in complete agreement about the shape and position of the eastern limit of the Dalarna porphyries, the latter is indicated in Fig. 37 by the symbol used for Hausen's view. The Swedish continental part of this limit coincides rather well with the inferred direction, but on the Baltic coast both the limit and the direction diverge widely, the inferred trend continuing toward Grodno, the eastern limit crossing the northern tip of Gotland and zigzagging through Estonia. As we have seen (p. 79), this deviation could be due to the outcrops of Bredvad porphyries in Jämtland and/or possible submarine occurrences of Dalarna porphyries in the Baltic, but there might be another reason.

As the dates of the quoted publications show, the fans under consideration were defined at a time when the quantitative method was rarely or not yet used for the study of indicators. It can be inferred from these publications that in the mapping of indicator-fans, localities with only a few or even only one indicator were used. Therefore, deviations from the main ice movements during glacial and non-glacial transport are apt to manifest themselves in such representations, especially along the lateral limits. Thus, the main ice movements may be obscured by a deceptive background effect. Conceivably, this effect could be suppressed by estimating the pro rata contributions of every recognizable indicator group in each locality, and excluding those groups whose percentages lie below a certain minimum threshold. The height of the threshold for each separate indicator category could be determined by trial and error. This might yield lateral limits of indicator fans giving a better approximation of the main flow directions of the inland ice than those of the initial fans.

Although appropriate handling, including the making and analysis of supplementary counts, is required to carry the proposed method into effect, a possible case in point is of interest here. The afore mentioned eastern limits of the Åland indicator fans were based on occurrences of Åland indicators, which were in the minority. In contrast, they predominate in the central zone of Tarvydas's tripartite division (Fig. 37) of the Pomeranian Stage in Lithuania. His investigations show that the indicators of the Åland indicator fan are not evenly distributed and are distinctly predominant in the central Lithuanian zone. This zone with its parallel boundaries trending NNW to SSE is directed toward the Åland islands. It need hardly be said that this density zone is more likely to indicate the direction of the main ice flow over the Åland islands than is the eastern lateral limit of this indicator fan.

*The distribution of indicators observed by Ramsay and Hackman  
in northwestern Russia*

Ramsay and Hackman (1894) were of the opinion that during the main extension of the ice sheet the glacial flow over the Kola peninsula was from west to east, the ice stream having transported boulders from north Finland toward this area. Because the glacial drift in Fennoscandia formed during the last glaciation (drift older than the Weichselian is exceptional here), the glacial deposits of the main extension in Kola are likely to be considered Weichselian too, the more so because according to Ramsay subsequent the nunatak phase developed, around the time of ice halt at the Salpausselkä. In 1912, however, Ramsay reconsidered his view in that the ice stream direction from west to east followed the main phase, in which

the glacial flow direction over the Kola peninsula was to the SE. He did so because he had found that the extension of the dispersal area of the nepheline-syenite from Umptek and Lujavr-Urt was much larger than he had initially thought. While sailing down the rivers Suchona and Dwina (Fig. 38) he saw the first nepheline-syenite in the pavements of Totma. Further downstream, syenite indicators occurred among the river boulders, but the first embodied in moraine were found in Weliky Ust-Jug and Kotlas. The syenite finds along the Suchona and the Dwina form part of a small fraction representing only one per cent of the whole body of crystalline erratics. The syenite indicator density is greater on Kanin peninsula, where it amounts to about one per cent of the total number of crystalline erratics, which seems to suggest that the main ice flow over the Kola peninsula was after all from west to east.

This direction is supported by additional information obtained by Ramsay (1912) about the dispersal of some other indicator groups. Typical Karelian rocks that crop out north and northwest of Lake Onega are found in the Wologda area




Fig. 39. The circular zone round  $F_3$  further to the south.  $F_1$ ,  $F_2$ : foci confocal ellipses.  $F_3D$ : base of circular pattern.  $NF_3$  is normal to  $F_3D$ .  $M$ : main stationary line,  $W$ : moraines of the Warthe glaciation. a, b, c, d, e: Saalian ice front phases.  $F_n$ : centre of circular configuration in The Netherlands.

and in the Wjatka district. As shown in Fig. 38, this dispersal points to a consistent transport direction. The same holds for some indicators belonging to the Ladoga rapakivi that were found in the rivers Unscha and Wertuga. White quartzites and uralite-diabases, whose provenances lie in a zone running from Lake Ladoga and Lake Onega into Lappland, occur generally along the rivers Suchona and Dwina.

Since the dispersal of the additional indicators fits reasonably well into the framework of hyperbolas and the higher density of the nepheline-syenites on Kanin also fits into the theoretical streamline pattern, there is after all good support for a west-to-east ice flow over the Kola peninsula during the main extension of the ice sheet. The main ice movement directions indicated by Rudovitz over Kola peninsula are also shown in Fig. 38. They hardly deviate from the inferred streamlines. This is not the case for the Saalian main ice flow directions toward Moscow represented by Jakowlev on a map of the Russian Quaternary, reproduced in Woldstedt (1958).

The evaluation in this chapter of the consistent indicator distribution on the basis of the inferred flowline pattern, has led to the following results. It not only confirmed the conclusions arrived at thus far, but also contributed to a better understanding of the relationships between the source, the route of transport, and the consistent distributions in The Netherlands and in Germany as far as the east side of the Oder. It partially filled the gap in the distribution pattern in which no Hesemann or Milthers counts had been performed. It furnished detailed evidence confirming the corresponding stream directions in the Weichselian and in the Saalian, in this case toward Lithuania and toward Poland and White Russia. The eastern limits of indicator fans and the dispersal of a number of species of indicators in Russia proved to coincide passably with the inferred pattern of ice flow.

Crude as the results may be, they justify a further elaboration of the flowline pattern, which might be useful in considering the changing shapes of the Fennoscandinavian ice sheets. Accordingly, an attempt to work out the inferred flow structure will be discussed in next section.

#### GEOMETRICAL ASPECTS

##### *The circular zone further southward*

We have shown above that the pattern of ice flow and successive outlines of the Fennoscandinavian ice sheets may be compared to a set of confocal hyperbolas and ellipses adapted in the southwestern sector to a zigzag bend in the ice divide. As expounded on p. 39, the flowlines between the ice divide and the main stationary line in Denmark resemble arcs of a set of concentric circles. In Fig. 39,  $F_1$  and  $F_2$ , the coequal foci of the ellipses and hyperbolas, are indicated by circlets.  $F_3$  is the focus of the circular pattern. This pattern returns in the general tendency of the river courses in this particular zone of former ice movement.

We have followed the circular feature as far as the main stationary line in Denmark, but how will this pattern continue in the southward direction? In central Jutland the stationary line strikes north, but farther to the south the moraines strike east-to-west. The direction of ice flow being normal to the moraines, a shift in the orientation of the moraines means a shift in the trends of flowlines. Accordingly, the encircling structure flattens out toward The Netherlands and even turns to the southwest.

For a better understanding of this zigzag tract we may turn to The Netherlands, where diverse glaciation phases of the Saalian have been studied by, among others, Brouwer (1948, 1950), Maarleveld (1958, 1960), and ter Wee (1962). A detailed review has been given by de Jong (1967) and Jelgersma & Breeuwer




Fig. 40. The pattern of the Saalian morphology in The Netherlands in relation to circular tracts about  $F_n$ . Saalian after retreat of inland ice from Zagwijn (1974). a, b, c, d, e: Saalian ice front phases according to Maarleveld (1958, 1960) and ter Wee (1962). EM: part of the confocal ellipse to which the maximum of the Saalian in The Netherlands approaches.

(1975). Fig. 39 shows the phases of the Saalian compiled by ter Wee (1962). It would appear from the diverging trends of ice front phases that a wheeling movement of the ice margin took place, but there is other evidence to be considered.

Zonneveld (1975) pointed out that many of the ice-pushed ridges show 'drumlinized' forms. Those of phase d, being parallel to each other and striking NE to SW, argue – according to Zonneveld – for an inherent ice stream from the NE. The drumlins of phase e, on the other hand, strike N60°E to S60°W. Thus, as compared with phase d, a change of directions as one should expect with a wheeling movement. In both cases the directions of ice flow agree with the trends of the glacial ridges they belong to.

Since not only the main stationary line and the moraines of the Warthe glacial but also phase e in Groningen are collateral, the ice flow directions pertaining to these phases must have been much alike, which would virtually restrict the change in direction to The Netherlands. Therefore, the centre of rotation of the wheeling movement should be sought nearby, and a centre of revolution was indeed found, by trial and error, just east of the Ems. Figure 40 presents a set of confocal circles drawn by means of this centre ( $F_n$ ), and shows how the circular structure and the pattern of the glacial landscape interlock. This circular pattern of movement also answers the question as to why, despite the different directions of ice flow in different parts of The Netherlands, the consistent indicator distribution of the diverse glacial phases remained the same.

How is  $F_n$  related to the flowline pattern geometrically? In Fig. 39 the distance between the centre,  $F_n$ , and the focus,  $F_3$ , equals the distance between  $F_3$  and D. Since  $F_3D$  coincides with the ice divide in the circular zone, it may be looked upon as a base or a line of symmetry of the circular pattern.

Although the measured distances are equal, the neglect of the curvature of the earth surface with respect to the scale and projection of the map in use might lead to inaccuracies that are not negligible.

To evaluate this point, I applied the formula  $\cos X = \sin M \sin M_1 + \cos M \cos M_1 \cos (L-L_1)$  from spherical trigonometry (Donnay, 1945) to the coordinates M, L ( $68^{\circ}36'N$ ,  $25^{\circ}50'E$ ) of  $F_2$  and  $M_1$ ,  $L_1$  ( $52^{\circ}18'N$ ,  $7^{\circ}36'E$ ) of  $F_n$ , and obtained a spherical distance x of  $18^{\circ}15'$ . If 6377.36 km is taken as the radius of the mean curvature of the earth's surface, the actual distance is 2049.875 km, which on a scale of 1 : 12 070 000 (the original scale of Fig. 39) amounts to 16.98 cm. Measured on this scale, thanks to the type of map projection used,  $F_2$  and  $F_n$  are 16.99 cm apart. Thus, the difference between the spherical and measured distance is within the magnitude of the error of measurement, which permits direct measurements from maps of the same size or smaller. No greater accuracy will be achieved by the application of spherical trigonometry.

In Figure 39, angle  $F_3DF_1$  measures  $20^{\circ}$ .  $NF_3$  is perpendicular to  $F_3D$ , and angle  $NF_3F_n$  measures  $20^{\circ}$  too. The recurrence of the angle of  $20^{\circ}$  and the equality of  $F_3F_n$  and  $F_3D$  may indicate that  $F_n$  is a natural component of the geometrical framework. This could mean, in effect, that  $F_n$  is an inevitable result of the unhampered development of the ice flow pattern.

On the other hand,  $F_n$  may have been induced not by internal but by external forces, i.e. a trigger effect from the resistance offered by the Harz, the Teutoburger Wald, and the interjacent range of mountains to the advancing ice. This 300-400 m high massif which rises 200 to 300 m from the north-German plains, was the largest obstacle the ice encountered on its way southwestward. A spur of this obstacle, the Teutoburger Wald, must have been overridden, while the rest of the Harz massif created a reentrant ice margin in the maximum extent of the Saalian glaciation.

Such an obstacle slackens an ice flow, and in this case must have slowed down the advance on the east flank of the ice stream to the southwest, a stream whose width may be put at 300 km. The impediment to movement on the east side of the ice in combination with a free passage to the west may account for the wheeling effect of the ice front in The Netherlands. An obstruction interfering with the natural development of the growing ice sheet, as in this case, can lead to stresses which may build up until they trigger off glacial surges. This fits into the view of Zagwijn (1974), who proposed sudden, torrential ice flows as the active agents in the formation of the ice-thrust ridges or lobes at the southernmost extension of the ice sheet in the central part of The Netherlands.

In sum, it appears that the circular pattern round  $F_3$  of the Weichselian continues from the main stationary line in Denmark as far as the Saalian phase e in The Netherlands, where it seemingly changes over into another circular unit with its centre in  $F_n$ .




Fig. 41. Morainic contours on the inside of the circular zone, constructed by symmetry operations. For explanation, see text.

*Provisional flowline and ice margin pattern in the North Sea*

In the foregoing, a flowline pattern was analysed and developed as far as the maximum extension of the Saalian ice sheet. Traced on land, this alignment is very clear in Fennoscandia. It disappears into the Gulf of Bothnia and the Baltic, to reappear, partially blurred, in the circumjacent area south of the Baltic. The perceptible pattern has several elements of symmetry, such as the median of the set of confocal ellipses and hyperbolas,  $F_1F_2$ , the diameter of the concentric circles which coincides with  $F_3D$ , and the centres of rotation  $F_3$  and  $F_n$ . Symmetry elements mean symmetry operations, and this led to the question of whether, on the basis of our flowline pattern and with the help of these elements, a flowline structure could be obtained in the North Sea, and Atlantic. The following discussion attempts to answer this question.

The broad sinuous zone toward The Netherlands gives some clues about the position of the former ice borders. It provides a correlation of end moraines and recession lines in The Netherlands, Denmark, southern Sweden and Norway, with plausible elliptical counterparts to the east. Before this feature can be used as a kind of gauge, however, further elaboration of the pattern is necessary.

In the circular zone with  $F_3$  as centre, the morainic contours hardly underwent any change, as appears from the persistence of shape of the zigzag bends (Fig. 39). To simplify matters, the Z bends are represented as straight segments of lines through  $F_3$  in Fig. 41. The diameter coinciding with  $F_3D$  is the most plausible axis of symmetry for the circular pattern. Equivalent straight-line segments of the circular zone on either side of  $F_3D$  are connected by thin lines to the focus  $F_3$ , and are coupled together by symmetrical arched curves. The arches, still narrow at the beginning of the movement, become wider and wider the more the ice sheet grows, and even become rectilinear when the straight segments of the circular zone are in line with each other. This right-line meets  $DF_3$  perpendicular at  $F_3$ , the centre of rotation of the circular pattern. That is to say, at an angle of rotation of  $90^\circ$  a straight stretch appears in the contour of the corresponding moraine. In this tentative approach, the moraines at the inside of the circular zone are initially convex. The moraine through  $F_3$  is rectilinear, and between  $F_3$  and  $E$  the moraines are concave.

The angle between the straight line through  $F_3$  and the line denoted by  $F_nF_3$  is  $20^\circ$ . Figure 41 shows the concave contour that coincides with  $F_nF_3$ . A concave margin which, at least on its southern branch, forms the limit of the movement that can be reconstructed by means of  $F_3$ .

This descriptive appraisal of possible relationships between several elements of the movement pattern provides the following features with which a provisional flowline and outline pattern of the Scandinavian ice sheets can be constructed (Fig. 41):

- 1) The most essential element is the sinuous zone, the combination of two circular units. The major, northern part round  $F_3$  trails off in a smaller, southern part round  $F_n$ . As a result, a rough correlation of assumed or subsistent ice margins in this zone with counterparts on either side is possible. Since the foci and the location of the transition of the circular zones are known, the circular lines of flow are easy to reconstruct. The result is a winding course which may be seen as a dial to correlate ice fronts on both sides.
- 2) Any stretch of end moraine in the sinuous zone can be linked up with an appropriate member of the set of confocal ellipses. This gives a rough idea of the

extension of the ice sheet in question on the east side of our dial.

3) On the west side of the sinuous zone, the assumed configuration of ice front phases on the inside of the circular unit around  $F_3$  offers a useful basis for an attempt at a theoretical reconstruction of Fennoscandinavian ice sheet margins.

Let us try, with these tools, to determine a maximum Weichselian extension of the Scandinavian ice mantle in the North Sea and Atlantic. In Fig. 42 the sinuous-shaped zone, a combination of the circular units about  $F_3$  and  $F_n$ , contains three Z bends that refer to the ice divide, the Ra central Swedish moraines (RCS), and the main stationary line in Denmark (M). According to the method specified in Fig. 41, a convex ice margin that links up with the west side of the Ra moraine has been drawn in as a dotted line in the inner circle. The actual terrestrial equivalent (solid line), however, curves further outward. It appears from the direction of the glacial striae represented by Andersen (1965) that the north-to-south-striking mountain spur in southwestern Norway held a local ice divide. Consequently, the terrestrial moraines in the southernmost part of Norway extend further southward with respect to their theoretical position. North of  $F_3$ D the theoretical ice border skirts the west coast of southern Norway and continues into




Fig. 42. A reconstruction of the maximum extension of the Weichselian ice sheet in the North Sea.

the Atlantic. The southern end being normal to the adjoining fjords, it may approximate the corresponding former ice margin.

According to Fig. 41, the dotted line indicating the theoretical continuation of the main stationary line in the North Sea, as shown in Fig. 42, is only slightly convex owing to its nearness to  $F_3$ . But because of the difference between the theoretical and actual ice margins in southern Norway, this part of the hypothetical maximum Weichselian extension should protrude also. The dashed line in Fig. 42 gives an instance of this situation.

The development of the ice sheet, as expounded here, tends to have an elliptical outline, which means a symmetrical shape. Flint (1971) proposed, however, an asymmetric shape for the Scandinavian ice sheet, the southeastern expansion going beyond the extension on the northwestern side of the ice divide because the deep Atlantic water impeded expansion to the west. An asymmetric ice sheet determined by an underdeveloped northwest flank involves a possible smaller extension in the northern part of the North Sea and in the Atlantic than Fig. 42 indicates. Hence, apart from the bulging lobe between  $F_3$  and the main stationary line, which could protrude differently, as suggested the Weichselian maximum north of  $F_3$  as represented in Fig. 42 indicates an outer limit. Thus, according to these theoretical deductions, the Scandinavian ice sheet did not reach either the Faeroes and the Shetland islands or Scotland in the Weichselian.

Some authors are of the opinion, however, that in the Weichselian the Scandinavian ice sheet crossed the North Sea and linked up with its British counterpart. Figure 42 shows that one of the most recent views (Jansen, 1976) differs north of  $F_3D$  from our reconstruction, but that south of this line the two are generally comparable. Likewise, the location of the Weichselian non-Scandinavian gravels between the Doggersbank and the English coast (Veenstra, 1970, 1971), conforms to a remote Scandinavian ice margin toward the north.

What are the implications of the preceding theoretical line of approach for the Saalian ice fronts in the North Sea area?

In Fig. 43 the Z bend is encircled farther to the south and joins with the Warthe Substage. Because the rotation passed through the  $90^\circ$  angle, the theoretical ice margin bridging the area of the inner circle of the circular zone must be concave (see Fig. 41). This ice front (dotted line) may bend outward again (dashed line) because of the disturbing effect of the local ice divide in southern Norway.

The ice front phase e of ter Wee (1962) lies in the neighbourhood of the transition zone of the encircling movement about  $F_3$  to that around  $F_n$ . Its continuation in the North Sea is shown in Fig. 43.

South of ter Wee's phase e, the ice fronts wheel about the pivot  $F_n$  to adjust themselves to the elliptical contours. Since  $DF_3$  and its produced part form an axis of symmetry, a coexisting counterpart of this motion would have had its centre west of the Faeroes. No distinct traces of this movement have been found, however, in the Faeroes, the Shetlands, the Orkneys, or in Scotland, possibly because at the time of the Scandinavian ice approach these islands were covered by their own local glaciers, which were probably even merged with the Scottish Highland ice. This armour of ice may have protected the rock surface against the charge of the Scandinavian ice sheet. Another point to be considered is that the location of the focus west of the Faeroes coincides with a part of the Atlantic which at present is over 500 m deep. It is a question whether this focus lying on a buoyant member of the ice sheet would have had behaved in the same way as its terrestrial counter-




Fig. 44. An array of concave curves and concentric circle fragments set against the glacial landscape in northern Jutland and the ice-marginal channels and lakes of the circum-Baltic.

part. On the other hand, since the wheeling movement about  $F_n$  might also have been due to the resistance of the Harz massif, it could have been a solitary case sought in vain at the other side of  $DF_3$ .

Returning to The Netherlands, we note with interest that the wheeling movement bypassed the ellipse  $EF_n$  through  $F_n$ . South of this ellipse, within a range of 150 km to the west of focus  $F_n$ , the ground is more heavily ploughed than on the north side. This zone between  $EF_n$  and the ellipse  $EM$ , through the maximum extension of the Saalian ice sheet, is a zone where the hampering of the natural

belt to be roughly 60 km. On this basis, the provenance of the brown porphyries would be limited to the area NW of Gotland below the 120 m contour. This trough, which is rather broad west of Gotska Sandön, tapers toward the southwest.

If the outcrops of brown porphyries are evenly distributed over this trough, the wide stretch will contain more porphyries than the narrow stretch. Therefore, a higher percentage of brown porphyry indicators is to be expected on Gotska Sandön and the northern part of Gotland and smaller percentages to the south, which agrees with the actual distribution of brown porphyries in these islands indicated by the counts made by Milthers in 1933.

According to these arguments, there is much to be said for a location of the provenance of the brown porphyries between the mainland of Sweden and the islands of Gotland and Gotska Sandön, which is southwest of the area proposed by Milthers and shown by Ludwig on his 1967 map.

We may now return to the fresh-water basin assumed to have arisen when the southern Baltic became ice free (Fig. 50). At that time conditions for calving were favourable in the rather deep water north of the Gulf of Gdansk. Since this calving ice forming part of the segment of ice sheet that crossed the source area of the brown porphyries, some of these indicators could have become waterborne. For the red porphyries, however, the conditions for waterborne transport were not




Fig. 51. Sketch of an ice halt in the broad trans-Baltic area where Silurian and Ordovician rocks crop out. For explanation, see Fig. 50 and the key to Fig. 34.

yet fulfilled at this stage. Consequently, the brown porphyries had the lead in the westward dispersion of the Baltic porphyries.

The drift to the west corresponded with the counterclockwise circulation and the easterly and northeasterly winds prevailing in the summer. A possible outlet for the rising waters of the marginal lake must be sought in Denmark. Even though the natural westward discharge of this lake was dammed by the ice sheet or lobes from the ice sheet in this area, the receding zigzag-shaped ice front, the eustatic and isostatic movements, and the continuous rise of the proglacial lake that was fed not only by meltwater but also by the Oder and the Weichsel, were weak points with respect to a hermetic sealing off of the lake water. Consequently, intermittent subglacial outflow and/or desultory overflow in areas of lowest relief are conceivable. On this basis, the waterborne brown porphyries — unlike the red ones — would have had the marginal lake themselves for some time in the course of the retreat or advance of the ice mantle. Therefore, higher percentages of brown porphyries than of red porphyries would be expected in associated indicator counts made in the southern circum-Baltic. This would explain why the counts of K. Milthers (1942) with twice as many browns as reds are restricted mainly to the coastal areas of the southeastern part of Denmark.

The described course of events offers an explanation of the predominance of brown porphyries in central Swedish indicator associations and the predominance of red porphyries in the assemblage of Åland and Finnish indicators in The Netherlands (Schuddebeurs, 1955, 1956) and northern Germany (Hesemann, 1936). We will go into this point further in the following.

#### *Ice halts and indicator associations*

Figure 50 shows that the segment of the ice sheet overlying the source rock of the brown porphyries partially coincides with Uppland, Södermanland, Västmanland, and Dalarna. This explains the affinity of the brown porphyries to middle Swedish indicators in connatural glacial deposits.

In the same way, the source area of the red porphyries is related to the Åland islands and part of the south-Finnish archipelago by a common segment of the ice mantle. However, in the situation sketched in Fig. 50 waterborne transport of this association would be suspended until a further retreat of the ice front.

To turn back to the brown porphyries, it is evident that under the conditions shown in Fig. 50 the Småland granites, like the middle-Swedish indicators, had a chance of being transported by floating ice. Since the ice edge lay on the island of Bornholm, the indicators from this island could also have entered this association. With respect to the question as to whether equivalent glacial deposits are known, a publication by Schuddebeurs (1956) deserves attention. He evaluated the high percentages of brown porphyries as against the low percentages or even absence of red porphyries in the glacial deposits of the region of het Gooi, which forms part of the Utrecht ridge, as shown in Table 6 and Fig. 13. Zandstra (in Overweel & Zandstra, 1967) demonstrated that the Saalian till debris in this area contains a gravel association comparable to the rock compositions of decalcified lenses of red boulder clay. This resemblance to the red boulder clay may be an indication of waterborne transport, which is supported not only by the high percentages of Stockholm and Uppsala granites in these deposits of het Gooi, but also by the exceptionally large quantity of Bornholm indicators (see Table 6).

Thus, the area of the Utrecht ridge shows Saalian deposits with an indicator

clay stand out. Those of the Saalian in The Netherlands contain an indicator association which agrees to some extent with that of an ice front situated in the area of Gotland and Öland. However, the Baltic porphyries are poorly represented in the lenses, in contrast with the Småland indicators. There are high percentages of Aland indicators, not only in the lenses and the red boulder clay of the Hondsrug but also in the average Saalian deposits in The Netherlands, which as a rule are intermixed with elements from waterborne transport. These high percentages may account for a prolonged halt of the ice front close to the Aland islands during its advance in the Saalian. But there is no denying that we should also take into consideration the glacial prior to the Saalian. For if another glacial should occur in the far future, the Fennoscandinavian and circum-Baltic deposits, upon which the Weichselian set its seal, will be redistributed. Consequently, the events deducible from the new sediments will refer not only to the added glacial but also to the Weichselian. In the same way, components of the Elsterian may have been recycled in the Saalian boulder clays.

Unlike the red lenses of boulder clay, the aforementioned Saalian deposits of the Utrecht ridge seem to offer an ideal example, due to their indicator association, which is completely consistent with that of an ice halt in Latvia, as set out in Fig. 54.

Notwithstanding the complications, it might be possible, along the lines expounded here, to remove a little of the veil that covers the standstill phases of the Scandinavian ice sheets in the Baltic, of substages that are even older than the Upper Pleniglacial, as defined by van der Hammen et al. in 1967.

In the last three of our hypothetical standstill phases of the ice edge across the Baltic the Kattegat was open, and this implies — depending on the height of the relative sea level — that ice-rafted Oslo indicators had a chance of slipping through to the Baltic proper via a possible tide-water glacier in the Skagerrak. Sauramo (1923), who compared the Salpausselkä ridges in Finland with the Raend moraines in the Oslo area, pointed out that both slope away from the inland ice, which would have promoted calving.

In this connection two publications by Schuddebeurs (1959, 1967) are relevant. In The Netherlands, where Oslo indicators are scarce, he found a remarkable high percentage of Oslo pebbles (about 25) near Drachten in the province of Friesland. He related this occurrence to a count made by van der Lijn and Bos at Bergum, where Oslo indicators accounted for 20%. Furthermore, he came across such accumulations near Langefeld, Werpeloh, and Fresenburg in NW Germany, an area which may be compared with the northeastern part of The Netherlands. Although Schuddebeurs (1959) saw this situation as the result of a direct ice stream, waterborne transport is suggested by the occurrence of clusters of Oslo indicators in regions otherwise virtually without these erratics, their confinement to deposits underlying the boulder clay and the scattering of single Oslo boulders in a direction opposite to their natural southwest trend of distribution. Occasional finds of these Oslo indicators in the coastal belt of the southern Baltic as far east as Thorn in Poland (Casper, 1931; Charlesworth, 1957) are shown on the distribution maps by Schulz (1973).

In conclusion, it may be mentioned that as early as 1896, G. de Geer took drift ice into account (Charlesworth, 1957) as one of the possible mechanisms, to explain the presence of Norwegian erratics on the east coast of England.

APPROXIMATE POSITIONS OF ICE-FRONT	INDICATORS						
	Åland	Uppland Stockholm	Dalarna	Red Baltic porphyries	Brown Baltic porphyries	Småland	Bornholm
Åland islands							
Stockholm							
Gotland-Öland							
Latvia							

Fig. 54. Tentative outline of indicator associations according to different parts of the ice front in the Baltic.

## Conclusions

The approach to the re-evaluation of indicator counts on the basis of the general trend of the traces of ice movement in Fennoscandinavia has shown that the trends of these traces link up with the consistent distribution pattern of the circum-Baltic indicators. Because this holds not only for the Weichselian but also for the Saalian, the general directions of ice flow must have been similar in these two stages.

The interrelationship between the flowline and the consistent distribution pattern allowed an integration of these units, permitting a reconstruction of the Fennoscandinavian ice sheet. Other reconstructions have been made, for instance those of Aseev (1968) and Aseev et al. (1973). There are differences between those and our reconstruction such as variations of a structural or morphologic nature, for instance the location of the ice divide, but above all there is a difference of fundamental nature. The ice sheets defined by Aseev (1968) and Aseev et al. (1973) are static, representing standstill phases of the maximum extension of the Fennoscandinavian ice mantles in the Weichselian and the Saalian. The ice sheet proposed here, however, is dynamic. Dynamic in a functional way, because it is susceptible to expansion and reduction. With respect to indicator transport, the ice mantle should not be analysed independent of the peripheral water system. Whether receding or advancing, the ice sheet and its inseparable, closely following outer margin of water constituted a whole. It follows that without the alternating play of ice and water, which was taken into account in the chapter on the Integration of consistent and inconsistent distribution processes, the reconstruction would have appeared incomplete.

Since Lyell's drift theory was replaced by the idea of continental ice sheets, ice rafting has been neglected as a means of indicator transport. As a result, the share of ice rafting in the distribution of the Fennoscandinavian indicators has not been given the attention it deserves. To remedy this neglect, the role of the peripheral water system as a possible transporting medium for indicators has been stressed in this paper, and it was included in the discussion on the nature of the Baltic ice stream. Were the indicators of the inconsistent distribution conveyed by such an ice stream or by drift-ice in the peripheral water channels? In our reconstruction, the evidence favours waterborne transport.

In sum, the pattern inferred in this study revealed a relationship between indicator distribution and glacial and periglacial features. The flexibility of this pattern enabled us to simulate advance, retreat and standstill phases of the Fennoscandinavian ice sheets. Its application provided answers to some questions and again raised other questions. Whatever this reconstruction may be worth, it offers a new way of looking at old problems.

## References

Aario, R., L. Forsström & P. Lahermo, 1974. Glacial landforms with special reference to drumlins and fluting in Koillismaa, Finland. — *Geol. Survey Finland, Bull.* 273: 1-30.

Agrell, H., 1976. The highest coastline in south-eastern Sweden. — *Boreas*, 5: 143-154.

Andersen, B. G., 1965. The Quaternary of Norway. In: Rankama, K. (ed.). *The geologic systems. The Quaternary 1.* — J. Wiley & Sons Inc., New York: 91-138.

Aseev, A. A., 1968. Dynamik und geomorphologische Wirkung der europäischen Eisschilde. — *Petermanns Geogr. Mitt.*, 112: 112-115.

Aseev, A. A., V. V. Bronguleev & A. N. Makkaveev, 1973. Reconstruction of the last (Upper Pleistocene) European ice sheet. Reconstruction of the maximum (Middle Pleistocene) European ice sheet. In: Gerasimov, J. P. (ed.). *The paleography of Europe during the Late Pleistocene.* — IX I.N.Q.U.A. Congress New Zealand, 1973, Inst. Geogr. USSR Acad. Sci.

Baartman, J. C. & O. B. Christensen, 1975. Contributions to the interpretation of the Fennoscandinavian Border zone. — *Danmarks Geol. Undersøg.*, 2, 102: 1-47.

Bahnson, H., 1973. Lithological investigations in some Danish boulder-clay profiles. — *Bull. Geol. Inst. Univ. Uppsala, N.S.*, 5: 93-109.

Belderson, W. H. & J. B. Wilson, 1973. Iceberg plough marks in the vicinity of the Norwegian Trough. — *Norsk Geol. Tidsskr.*, 53: 323-328.

Bergersen, O. F., 1970. Undersøkelser av steinfraksjonens rundingsgrad i glacigene jordarter. — *Norges Geol. Undersøk.*, 266 (Arbok 1969): 252-262.

Bergersen, O. F. & K. Garne, 1972. Ice movements and till stratigraphy in the Gubrandsdal area. Preliminary results. — *Norsk Geol. Tidsskr.*, 26: 1-16.

Berthelsen, A., 1973. Weichselian ice advances and drift successions in Denmark. — *Bull. Geol. Inst. Univ. Uppsala, N.S.*, 5: 21-29.

Binzer, K., 1974. Sedimentological and geochemical features of Weichselian tills and pre-Quaternary sediments in Denmark. — *Danmarks Geol. Undersøg. Årbog* 1973: 111-131.

Boekschoten, G. J. & H. J. Veenstra, 1967. Over stenen-oriëntatie in het Nederlandse keileem. — *Geologie Mijnbouw*, 46: 195-205.

Brouwer, A., 1948. Pollenanalytisch en geologisch onderzoek van het Onder- en Midden-Pleistoceen van Noord-Nederland. — *Leidse Geol. Meded.*, 14 B: 259-346.

—, 1950. De glacigene landschappen in Nederland. — *Tijdschr. Kon. Ned. Aardrijksk. Genoot.*, 2, 67: 20-32.

—, 1963. Cainozoic history of the Netherlands. — *Verh. Kon. Ned. Geol. Mijnb. Genoot., Geol. Ser.*, 21-1: 117-148.

Calker, F. J. P. van, 1912. Die kristallinischen Geschiebe der Moränen-Ablagerungen in der Stadt und Umgebung von Groningen. — *Mitt. Miner.-Geol. Inst. Reichs-Univ. Groningen*, 2, 3: 175-390.

Capadose, A., 1825. Lofrede op Sebaldus Justinus Brugmans. — *Werken Hollandsche Maatsch. Fraaije Kunsten Wet.*, 7: 401-403, 551-554 (refers to the doctoral dissertation of S. J. Brugmans, 1781. *Lithologia Groningana, juxta ordinem Walleri digesta, cum synonymis aliorum, imprimis Linnaei et Cronstedii, cum figuris aeneis*).

Casper, C., 1931. Ein Rhombenporphyr-Geschiebe bei Thorn. — *Zeitschr. Geschiebeforsch.*, 7: 1-93.

Cepek, A. G., 1969. Zur Bestimmung und stratigraphischen Bedeutung der Dolomitgeschiebe in den Grundmoränen im Nordteil der D.D.R. — *Geologie*, 18: 657-673.

Charlesworth, J. K., 1957. The Quaternary era, Vol. 1 & 2. — E. Arnold Ltd., London: 1-1700.

Donnay, J. D. H., 1945. Spherical trigonometry after the Cesàro method. — Intersci. publ. Inc., New York: 1-83.

Donner, J. J., 1965. The Quaternary of Finland. In: Rankama, K. (ed.). *The Geologic Systems. The Quaternary 1.* — J. Wiley & Sons Inc., New York: 199-272.

Dreimanis, A., 1976. Tills: their origin and properties. In: *Glacial Till.* — Roy. Soc. Canada, Sp. Publ., 12: 11-49.

Edelman, C. H. & G. C. Maarleveld, 1958. Pleistozän-geologische Ergebnisse der Bodenkartierung in den Niederlanden. — *Geol. Jahrb.*, 73: 639-684.

Embleton, C. & A. M. King, 1975. Glacial geomorphology. — E. Arnold Ltd., London: 1-573.

Eskola, P., 1933. Tausend Geschiebe aus Lettland. — *An. Acad. Sci. Fennicae*, A, 39: 5-41.

Faber, F. J., 1960. Aanvullende hoofdstukken over de Geologie van Nederland. Deel IV. — Noorduyn & Zoon N.V., Gorinchem: 1-607.

Feyling-Hanssen, R. W., 1971. Weichselian interstadial foraminifera from the Sandnes-Jæren area. — Bull. Geol. Soc. Denmark, 21: 72-116.

—, 1972. The Pleistocene/Holocene boundary in marine deposits from the Oslofjord area. — Boreas, 1: 241-246.

Flinn, D., 1967. Ice fronts in the North Sea. — Nature, 215: 1151-1154.

Flint, R. F., 1957. Glacial and Pleistocene Geology. — J. Wiley & Sons Inc., New York: 1-553.

—, 1971. Glacial and Quaternary Geology. — Ibidem: 1-892.

Frödin, G., 1916. Über einige spätglaziale Kalbungsbuchten und fluvoglaziale Estuarien im Mittleren Schweden. — Bull. Geol. Inst. Univ. Uppsala, 15: 149-174.

Gagel, C. & J. Korn, 1918. Der Geschiebeinhalt des wohlynischen Diluviums. — Zeitschr. Deutschen Geol. Ges., 70: 83-94.

Gaigales, A., 1963. Reconstruction of the glacier movements of Pleistocene on the territory of Lithuania according to the data of petrographical investigations of tills. — Lietuvos Geologijos Klausimai, Geol. Inst. Vilnius: 527-544 (in Russian).

—, 1965. Eigentümlichkeiten grosser Gesteinsbrocken pleistozäner Moränen verschiedenem Alters im Südostlitauen und ihr Wert für die Stratigraphie. — Strat. Quartärer Ablagerungen Südostlitauens Antropogäne Paläogeographie, Geol. Inst. Vilnius, 2: 104-156.

Gaigales, A. & V. Gudelis, 1965. Erratic boulders of the main stages of the last glaciation in the S.E. Baltic Area and the dynamics of ice sheet. — Baltica 2: 213-232 (in Russian with English and German summaries).

Geer, E. H. de, 1954. Skandinaviens Geokronologi. — Geol. Fören. Förh., 76: 299-329.

—, 1957. Old and new datings of Swedish ice lakes and the thermals of Bölling and Alleröd. — Ibidem, 79: 93-100.

—, 1961. G. de Geer's Chronology. — Geokronolog. Inst. Stockholm Univ.: 1-28.

—, 1963. G. de Geer's part in exploring the history of the Baltic sea. — Baltica, 1: 15-33.

Geer, G. de, 1885. Ueber die zweite Ausbreitung des skandinavischen Landeises. — Zeitschr. Deutsch. Geol. Ges., 37: 177-206.

—, 1910. Quaternary sea-bottoms in Western Sweden. — Geol. Fören. Förh., 32: 1139-1195.

—, 1919. Om moränmaterial i sekundärt läge. — Ibidem, 41: 193-201.

—, 1932. Stockholmstraktens kvartärgeologi. — Sveriges Geol. Undersökn., Ser. Ba., 12: 3-89.

—, 1938. Varve datings contra suppositions. — Geol. Fören. Förh., 60: 236-242.

Gillberg, G., 1965. Till distribution and ice movements on the northern slopes of the South Swedish Highlands. — Ibidem, 86: 433-484.

—, 1967. Further discussion of the lithological homogeneity of till. — Ibidem, 89: 29-49.

Glückert, G., 1971a. Drumlinlandschaft auf der Wasserscheide zwischen Pieksämäki und Haukipuori im Mittelfinnland. — Bull. Geol. Soc. Finnland, 43: 141-161.

—, 1971b. Bewegungen des Inlandeises im Lohja-Seebecken, Südfinnland. — Ibidem, 43: 173-202.

Grey, H., 1932. Undersøgelser over Ledeblokke i Skaane. — Meddel. Dansk Geol. Fören., 8: 143-166.

Gripp, K., 1964. Erdgeschichte von Schleswig-Holstein. — K. Wachholtz Verlag, Neumünster: 1-411.

—, 1975. Eisrandstudien ausgehend von Sermeq, SW-Grönland. — Meddel. Grönland, 195: 1-19.

Grisak, E. G., J. A. Cherry, J. A. Vonhof & J. P. Blumele, 1976. Hydrogeologic and hydrochemical properties of fractured till in the Interior Plains Region. In: Glacial Till. — Roy. Soc. Canada, Sp. Publ., 12: 304-333.

Gustafsson, O. & S. Stjernkvist, 1966. Några iakttagelser i en moränskärning i Lund. — Geol. Fören. Förh., 87: 431-440.

Håkansson, S., 1971. Absolute chronology of the Late-Quaternary for southern Sweden — a brief review. — Geol. Fören. Förh., 93: 477-493.

Hammen, T. van der, G. C. Maarleveld, J. C. Vogel & W. H. Zagwijn, 1967. Stratigraphy, climatic succession and radiocarbon dating of the Last Glacial in the Netherlands. — Geologie Mijnbouw, 46: 79-95.

Hansen, S., 1965. The Quaternary of Denmark. In: Rankama, K. (ed.). The Geologic systems. The Quaternary 1. — J. Wiley & Sons Inc., New York: 1-90.

Hansen, S. & A. V. Nielsen, 1960. Glacial geology of southern Denmark. — *Internat. Geol. Cong.*, 21 Sess., Norden, Guide excurs. A44 & C39: 1-56.

Harland, W. B., K. N. Herod & D. H. Krinsley, 1966. The definition and identification of tills and tillites. — *Earth-sci. Rev.*, 2: 225-256.

Hausen, H., 1912. Studier öfver de Sydfinska ledblockens spridning i Ryssland. — *Fennia*, 32, 3: 3-32.

Hellström, R. H., 1913. Bidrag till Kändedomen av isförhållandena i Bottenvägen. — *Ibidem*, 33: 1-364.

—, 1922. Talviliukenne Dohjanlahdella IV kokemuksia talvena 1916. — *Ibidem* 43, 6: 1-24.

Hesemann, J., 1930. Wie sammelt und verwertet man kristalline Geschiebe? — *Preusz. Geol. Landesanst. Sitz. Ber.*, 5: 188-196.

—, 1932. Zur Geschiebeführung und Geologie des Odergletschers. — *Jahrb. Preusz. Geol. Landesanst.*, 53: 70-84.

—, 1936. Zur Petrographie einiger nordischer kristalliner Leitgeschiebe. — *Abh. Preusz. Geol. Landesanst.*, N.F., 173: 5-168.

—, 1937. Zur Geschiebeführung und geologie des Odergletschers. II Der Züllichauer Bogen. — *Jahrb. Preusz. Geol. Landesanst.*, 58: 455-471.

—, 1938. Geschiebeuntersuchungen zwischen Pommerschem und Warthe-Stadium in der mittleren Mark Brandenburg. — *Ibidem*, 59: 45-54.

—, 1939a. Diluvialstratigraphische Geschiebeuntersuchungen zwischen Elbe und Rhein. — *Abh. Naturw. Ver. Bremen*, 31: 247-285.

—, 1939b. Zur Petrographie weiterer, hauptsächlich ostfennoskandischer Leitgeschiebe. — *Zeitschr. Geschiebeforsch.*, 15: 68-98.

—, 1975. Kristalline Geschiebe der nordischen Vereisungen. — *Geol. Landesamt Nordrhein-Westfalen*, Krefeld: 1-267.

Hillefors, Å., 1967. Die Glazialschrammen auf Hallands Svartskän und der Verlauf des Würmzeitlichen Eisrandes längs der Küste Nordhallands Westschweden. — *Medd. Dansk Geol. Fören.*, 17: 76-89.

—, 1968. Die Würmzeitlichen Eisbewegungen unter der Verlauf der Deglaziation im Kattegat und im südöstlichen Skagerrak. — *Ibidem*, 18: 315-343.

—, 1974. The stratigraphy and genesis of the Dösebacka and Ellesbo drumlins. A contribution to the knowledge of the Weichsel glacial history in Western Sweden. — *Geol. Fören. Förh.*, 96: 355-374.

Holtedahl, H. & K. Bjerkli, 1975. Pleistocene and Recent sediments of the Norwegian continental shelf (62°N - 71°N), and the Norwegian Channel area. — *Norges Geol. Undersök.*, 316: 241-252.

Hoppe, G., 1951. Drumlins i Nordöstra Norrbotten. — *Geogr. An.*, 33: 157-165.

Hucke, K., 1917. Die Sedimentärgeschiebe des norddeutschen Flachlandes. — *Verlag von Quelle & Meyer*, Leipzig: 1-195.

—, 1967. Einführung in die Geschiebeforschung. — Herausgegeben und erweitert von Ehrhard Voight, Oldenzaal, Verlag Ned. Geol. Ver: 1-132.

Hurtig, T., 1971. Jahreszeitliche Beobachtungen an den mecklenburgischen Boddengewässern zwischen dem Festland und der Insel Rügen. — *Natur Museum*, 101: 481-492.

Hyvärinen, L., 1967. Geochemical prospecting for lead ore at Korsnäs. In: Kvalheim, A. (ed.). *Geochemical prospecting in Fennoscandia*. — J. Wiley & Sons, Intersci. publ., New York: 171-179.

—, 1969. On the geology of the copper ore field in the Virtasalmi area, Eastern Finland. — *Bull. Commis. Géol. Finlande*, 240: 1-82.

Hyppä, E., 1963. On the Late-Quaternary history of the Baltic sea. — *Fennia*, 89, 1: 37-49.

Jansen, J. H. F., 1976. Late Pleistocene and Holocene history of the northern North Sea, based on acoustic reflection records. — *Netherlands Jour. Sea Research*, 10: 1-43.

Järnefors, B. & E. Fromm, 1960. Chronology of the ice recession through middle Sweden. — *Internat. Geol. Cong. Rept. 21 Sess. Norden*, pt 4, proc. sec. 4: 93-97.

Jelgersma, S. & J. B. Breeuwer, 1975. Toelichting bij de kaart glaciale verschijnselen gedurende het Saalien. In: Zagwijn, W. H. & C. J. van Staalduin (eds.). *Toelichting bij geologische overzichtskaarten van Nederland* (1 : 600.000). — Rijks Geologische Dienst, Haarlem: 93-103.

Jong, J. D. de, 1967. The Quaternary of the Netherlands. In: Rankama, K. (ed.). *The Geologic systems. The Quaternary 2*. — J. Wiley & Sons Inc., New York: 301-426.

Kauranne, L. K., 1967. Trace element concentrations in layers of glacial drift at Kolima, central Finland. In: Kvalheim, A. (ed.). *Geochemical prospecting in Fennoscandia*. — J. Wiley & Sons, Intersci. publ., New York: 181-191.

Kley, K. van der & W. de Vries, 1941. *Gidsgesteenen van het Noordelijk Diluvium*. — J. A. Boom & Zn, Meppel: 1-191.

Knudsen, K. L., 1971. Late Quaternary foraminifera from the Løkken area. — Bull. Geol. Soc. Denmark, 21: 130-158.

König, D., 1976. *Waddengebied van Schleswig-Holstein*. In: *Waddenzee*. — Landel. Ver. Behoud Waddenze, Harlingen; Ver. Behoud Natuurmonumenten Nederland, 's-Gravenland: 77-89.

Königsson, L. K. & S. I. Svantesson, 1975. The petrography of gravel deposits on the island of Gotland in the Baltic. — Geol. Fören. Förh., 97: 312-320.

Korn, J., 1895. *Ueber diluviale Geschiebe der Königsberger Tiefbohrungen*. — D. Sc. Diss., L. Schade, Berlin: 1-66.

—, 1927. Die wichtigsten Leitgeschiebe der nordischen Kristallinen Gesteine im Norddeutschen Flachlande. — Preusz. Geol. Landesanst., Berlin: 1-64.

Kraus, E., 1934. Über die Geschiebe in Lettland. — Zeitschr. Geschiebeforsch., 10: 67-80.

Kreutz, St. & A. Głowinska, 1932. Die polnischen Geschiebe. — Ann. Soc. Géol. Pologne, 8: 219-221.

Krog, H. & H. Tauber, 1974. C-14 chronology of Late- and Post-glacial marine deposits in North Jutland. — Danmarks Geol. Undersøg., Årbog 1974: 93-105.

Kruizinga, P., 1918. *Bijdrage tot de kennis der sedimentaire zwerfstenen in Nederland*. — Mouton & Co., 's-Gravenhage: 1-271.

Kujansuu, R., 1967. On the deglaciation of western Finnish Lapland. — Bull. Commis. Géol. Finlande, 232: 1-98.

Kummerow, E., 1930. Fortschritte der Geschiebeforschung. (Bewegungsrichtung des Inlandeises. Drift- und Schollentransport). — Centralbl. Min. Geol. Paläont., B, 1930: 136-145.

—, 1935. Neuere Forschungen über Herkunft, Transportwege und Verteilung der Geschiebe in Nordeuropa. — Ibidem, 1935: 129-140.

—, 1950/51. Fortschritte in der Diluvialgeologie. (Drift- und Inlandeistheorie. Herkunft und Verbreitung der Geschiebe). — Die Erde, 2: 16-24.

—, 1954. Grundfragen der Geschiebeforschung. — Geologie, 3: 42-54.

Kvasov, D. D., I. P. Bakanova & N. N. Davidova, 1970. Main questions of the Late-Glacial history of the East Baltic. — Baltica, 4: 90-91.

Ladwig, H., 1938. Geschiebezählungen als Forschungsmittel zur Diluvial-stratigraphie und Tektonik Ostpommerns. — Abh. Geol.-Palaeont. Inst. Greifswald, 19: 1-45.

Lijn, P. van der, 1941. Twee ijsbedekkingen in Nederland op grond van zwerfsteen tellingen. — Natura, 40: 70-90.

—, 1974. Het keienboek. — B.V. W. J. Thieme & Cie., Zutphen: 1-361.

Ludwig, A. O., 1967. Der präquartäre Untergrund der Ostsee. Teil I: Nordliche und Mittlere Ostsee. — Wiss. Zeitschr. Univ. Rostock, Math.-Naturw. R., 16: 1105-1136.

Lundqvist, J., 1965. The Quaternary of Sweden. In: Rankama, K. (ed.). *The Geologic Systems. The Quaternary 1*. — J. Wiley & Sons Inc., New York: 139-198.

—, 1974. Outlines of the Weichsel Glacial in Sweden. — Geol. Fören. Förh., 96: 327-339.

Lüttig, G., 1958. Methodische Fragen der Geschiebeforschung. — Geol. Jahrb., 75: 361-418.

Maarleveld, G. C., 1953. Standen van het landijs in Nederland. — Boor Spade, 4: 95-105.

—, 1956. Grindhoudende Midden-Pleistocene sedimenten. — Meded. Geol. Sticht., Ser. C-VI-6: 1-105.

—, 1960. Glacial and periglacial landscape forms in the central and northern Netherlands. — Tijdschr. Kon. Ned. Aardrijksk. Genoot., 2, 77: 298-304.

Madsen, V., 1928. Übersicht über die Geologie von Dänemark. — Danmarks Geol. Undersøg., 5, 4: 1-225.

Madsen, V., V. Nordmann & N. Hartz, 1908. Studier over Cyprinaleret og andre Eem-Aflejninger i Danmark, Nord-Tysland og Holland. — Ibidem, 2, 17: 1-302 (+ Atlas: pls. 1-12).

Magnusson, N. H., G. Lundqvist & G. Regnell, 1963. *Sveriges geologi*. — Svenska Bokförlaget, Norstedts: 1-698.

Marczinski, R., 1968. Zur Geschiebekunde und Stratigraphie des Saaleglazials (Pleistozän) im nördlichen Niedersachsen zwischen Unterweser und Unterelbe. — Rotenburger Schr., Sonderh. 11: 1-132.

Martin, K., 1875. Die Geschiebe von Jever im Groszherzogthum Oldenburg. — Abh. Naturw. Ver. Bremen, 4: 385-391.

—, 1877a. Silur-, Devon-, Trias-, Kreide- und Tertiär-Geschiebe aus Oldenburg. — Ibidem, 5: 289-298.

—, 1877b. Untersuchungen über die Organisation von *Astylospongia*, Ferd. Roem. und Bemerkungen über die Natur der Wallsteine, Meyn. — Arch. Ver. Freunde Naturgesch. Mecklenburg, 31: 1-32.

—, 1882. Ueber das Vorkommen eines gemengten Diluviums und anstehenden Tertiärgebirges in den Dammer Bergen, im Süden Oldenburgs. — Abh. Naturw. Ver. Bremen, 7: 311-334.

—, 1883. Aanteekeningen over Erratische gesteenten van Overijssel. — Overijsselsche Ver. Ontwikkeling Prov. Welvaart, W. E. J. Tjeenk Willink, Zwolle: 1-14, 1pl.

—, 1888. Ein neues Untersilurisches Geschiebe aus Holland. — Versl. Meded. Kon. Akad. Wet., Afd. Natuurk., 3, 4: 293-296.

—, 1889. Het eiland Urk, benevens eenige algemeene beschouwingen over de geologie van Nederland. — Tijdschr. Kon. Ned. Aardrijksk. Genoot., Versl. Aardr. Med., 6, 1889: 1-37.

Martinsson, A., 1960. The submarine morphology of the Baltic-Cambrosilurian Area. — Bull. Geol. Inst. Univ. Uppsala, 38: 11-35.

Mattsson, A., 1960. Die glaziale Felsskulptur und die jüngsten Eisströme über Südschonen-Nordbornholm. — Zeitschr. Geomorph., N.F., 4: 129-140.

McLellan, A. G., 1971. Ambiguous 'glacial' striae formed near water bodies. — Canadian Jour. Earth Sci., 8: 477-479.

Milthers, K., 1942. Ledeblokke og landskabsformer i Danmark. — Danmarks Geol. Undersøg., 2, 69: 9-137.

Milthers, V., 1909. Scandinavian indicator-boulders in the Quaternary deposits. Extension and distribution. — Danmarks Geol. Undersøg., 2, 23: 1-153.

—, 1911. Preliminary report on boulders of Swedish and Baltic rocks in the south-west of Norway. — Meddel. Dansk Geol. Fören., 17: 511-512.

—, 1933. Leitgeschiebe auf Gotland und Gotska Sandön sowie die Heimat der Ostseeporphyre. — Geol. Fören. Förh., 55: 19-28.

Milthers, V. & K. Milthers, 1938. Die Verteilung einiger wichtiger skandinavischer Leitgeschiebe in einem Teile Polens. — Bull. Serv. Géol. Pologne, 5: 1-27.

Mohrén, E., 1973. The driftbeds of western Skåne in southern Sweden. — Bull. Geol. Inst. Univ. Uppsala, N.S., 5: 209-215.

Mörner, N. A., 1969. The Late Quaternary History of the Kattegat sea and the Swedish West Coast: deglaciation, shorelevel displacement, chronology, isostasy and eustasy. — Sveriges Geol. Undersök., C, 640 (Årsbok 63, 3): 1-487.

Münnich, G., 1936. Quantitative Geschiebeprofile aus Dänemark und Nordostdeutschland mit besonderer Berücksichtigung Vorpommerns. — Abh. Geol. Palaeont. Inst. Greifswald, 15: 1-52.

Mutanen, T., 1971. An example of the use of boulder counting in lithologic mapping. — Bull. Geol. Soc. Finland, 43: 131-140.

Nilsson, E., 1960. Södra Sverige i senglacial tid. — Geol. Fören. Förh., 82: 134-149.

Oele, E., 1969. The Quaternary geology of the Dutch part of the North Sea, north of the Frisian Isles. — Geologie Mijnbouw, 48: 467-479.

—, 1971. The Quaternary geology of the southern area of the Dutch part of the North Sea. — Ibidem, 50: 461-473.

Overweel, C. J. & J. G. Zandstra, 1967. Een zwerfsteen van Uppsala-graniet in Lage Vuursche en enkele opmerkingen over keileem in Midden-Nederland. — Grondboor Hamer, 1967: 175-187.

Pääkkönen, V., 1966. On the geology and mineralogy of the occurrence of native antimony at Seinäjoki, Finland. — Bull. Commis. Géol. Finlande, 225: 2-70.

Price, R. J., 1973. Glacial and fluvoglacial landforms. — Oliver & Boyd, Edinburgh: 1-242.

Ramsay, W., 1912. Über die Verbreitung von Nephelinsyenit-geschieben und die Ausbreitung des Nordeuropäischen Inlandeises im Nördlichen Russland. — Fennia, 33, 1: 1-17.

Ramsay, W. & V. Hackman, 1894. Das Nephelinsyenitgebiet auf der Halbinsel Kola. — Ibidem, 11, 2: 1-225.

Reinhard, H., 1967. Geschiebeverfrachtung durch Eisschollen. — Wiss. Zeitschr. Ernst-Moritz-Arndt-Univ. Greifswald, Math. Naturw. R., 16: 209-220.

Reite, A. J., 1967. Lokalglaciasjon på Sunnmøre. — Norges Geol. Undersøg., 247 (Årbok 1966): 262-287.

Repo, R. & R. Tynni, 1971. Observations on the Quaternary geology of an area between the 2nd Salpausselkä and the ice-marginal formation of central Finland. — Bull. Geol. Soc. Finland, 43: 185-202.

Reuter, G., 1885. Die Beyrichien der Obersilurischen Diluvialgeschiebe Ostpreussens. — Zeitschr. Deutsch. Geol. Ges., 37: 621-679.

Richter, K., 1932. Die Bewegungsrichtung des Inlandeises, rekonstruiert aus den Kritzen und Längachsen der Geschiebe. — Zeitschr. Geschiebeforsch., 8: 62-66.

—, 1933. Gefüge und Zusammensetzung des Norddeutschen Jungmoränengebietes. — Abh. Geol. Paläont. Inst. Greifswald, 11: 1-63.

—, 1936. Ergebnisse und Aussichten der Gefügeforschung im Pommerschen Diluvium. — Geol. Rundschau, 27: 196-206.

—, 1958. Geschiebegrenzen und Eisrandlagen im Niedersachsen. — Geol. Jahrb., 76: 223-234.

Ringberg, B., 1971. Glacialgeologi och isavsmältning i Östra Blekinge. — Sveriges Geol. Undersök., C, 661: 1-174.

Roemer, F., 1885. Lethaea erratica. — Palaeont. Abh., 2: 250-420.

Rosa, B., 1970. Einige Probleme der Geomorphologie, Paläographie und Neotektonik des Südbaltischen Küstenraumes. — Baltica, 4: 197-210.

Rudowski, S., 1972. Influence of freeze on active processes in shore zone and on beach structure under moderate climatic conditions. — Bull. Acad. Polonaise Sci., Sér. Sci. Terre, 22: 139-144.

Sauramo, M., 1923. Studies on the Quaternary varve sediments in Southern Finland. — Fennia, 44, 1: 3-164.

—, 1929. The Quaternary geology of Finland. — Bull. Commis. Géol. Finlande, 13, 86: 1-110.

Schell, I. I., 1966. Icebergs. In: Fairbridge, R. W. (ed.), The encyclopedia of oceanography. — Reinhold Publ. Corp., New York: 367-369.

Schuddebeurs, A. P., 1955. Mededelingen over drie gesteentetellingen en enige opmerkingen betreffende de Oostzee-porfieren en hun verspreiding. — Grondboor Hamer, 2: 60-62.

—, 1956. Drie gesteentetellingen, Oostzee-porfieren en hun verspreiding (II en slot). — Ibidem, 3: 76-83.

—, 1959. De verspreiding van de zwerfstenen uit het Oslogebied. — Ibidem, 13: 316-333.

—, 1967. Opnieuw over de verspreiding van de zwerfstenen uit het Oslogebied. — Ibidem, 1967, 3/4: 41-49 (= 121-129).

Schulz, W., 1973. Rhombenporphyr-Geschiebe und deren östliche Verbreitungsgrenze im nordeuropäischen Vereisungsgebiet. — Zeitschr. Geol. Wiss. Berlin, 1, 9: 1141-1154.

Serebryanny, L., A. Raukas & J. M. Punning, 1970. Fragments of the natural history of the Russian Plain during the Late Pleistocene with special reference to radiocarbon datings of fossil organic matter from the Baltic region. — Baltica, 4: 351-366.

Spjeldnoes, N., 1973. Moraine stratigraphy, with examples from the basal Cambrian ('Eocambrian') and Ordovician glaciations. — Bull. Inst. Univ. Uppsala, N.S., 5: 165-171.

Strübing, K., 1970. Satellitenbild und Meereiskundung. — Deutschen Hydrol. Zeitschr., 23: 193-213.

Tarvydas, R., 1960. Crystalline boulders of the last and penultimate glaciations in Lithuania. — Collectanea Act. Geol. Lithuania, Coll. Pap. 21 Sess. Internat. Geol. Cong.: 215-220.

—, 1967. Dynamics of Neopleistocene glacier in the territory of Lithuania and the Kaliningrad region. — Lietuvos Kvartero Geolog. Paleograf. Klausimai, Geol. Inst. Vilnius, 5: 161-179 (in Russian with an English summary).

Termier, H. & G. Termier, 1963. Erosion and Sedimentation. — D. van Nostrand Co., London: 1-433.

Tschirwinsky, P., 1926. Petrographische Untersuchung der Diluvialgeschiebe des Dongebietes. — Zeitschr. Geschiebeforsch., 2: 128-140, 145-152.

Vaitiekūnas, P. & J. M. Punning, 1970. Some results of investigations into the paleogeography and absolute geochronology of the last glacial period in the Baltic region. — Baltica, 4: 346-347.

Veenstra, H. J., 1963. Microscopic studies of boulder clays. — Doctoral thesis, 'Stabo' N.V., Groningen: 1-211.

—, 1970. Quaternary North Sea Coasts. — Quaternaria, 12: 169-184.

—, 1971. Sediments of the southern North Sea. In: Delany, F. M. (ed.). The Geology of the East Atlantic Continental Margin. I.C.S.U./S.C.O.R. Working party, 31 Symposium. — Inst. Geol. Sci. Rept., 70/15: 9-23.

Virkkala, K., 1951. Glacial geology of the Suomussalmi area, East Finland. — Bull. Commis. Géol. Finlande, 155: 1-66.

—, 1960. On the striations and glacier movements in the Tampere region, southern Finland. — Ibidem, 188: 159-176.

Vortisch, W., 1972. Untersuchungen im Pleistozän SO-Schonens. — Geol. Fören. Förh., 94: 35-68.

Waard, D. de, 1949. Glacigen Pleistoceen, een geologisch detailonderzoek in Urkerland (Noordoostpolder). — Verh. Ned. Geol. Mijnb. Genoot., Geol. Ser., 15: 70-246.

Wee, M. W. ter, 1962. The Saalian Glaciation in the Netherlands. — Meded. Geol. Sticht., N.S., 15: 57-76.

Wennervirta, H., 1967. Geochemical methods in uranium prospecting in Finland. In: Kvalheim, A. (ed.). Geochemical prospecting in Fennoscandia. — J. Wiley & Sons, Intersci. publ., New York: 155-169.

Winterhalter, B., 1967. The Sylen and Solovjeva shoals as observed by a diving geologist. — Geol. Fören. Förh., 89: 205-217.

—, 1972. On the geology of the Bothnian Sea, an epeiric sea that has undergone Pleistocene glaciation. — Geol. Survey Finland, Bull. 258: 1-66.

Woldstedt, P., 1955. Norddeutschland und angrenzende Gebiete im Eiszeitalter. — K. F. Koehler, Stuttgart: 1-467.

—, 1958. Das Eiszeitalter. Grundlinien einer Geologie des Quartärs. Zweiter Band. — F. Enke Verlag, Stuttgart: 1-438.

Woldstedt, P. et al., 1971. International Quaternary map of Europe. Blatt 7 Moskva, Blatt 3 Nordkapp. — Bundesanstalt Bodenforsch., Hannover; UNESCO, Paris.

Woldstedt, P. & K. Duphorn, 1974. Norddeutschland und angrenzende Gebiete im Eiszeitalter. — K. F. Koehler Verlag, Stuttgart: 1-500.

Zagwijn, W. H., 1974. The palaeogeographic evolution of the Netherlands during the Quaternary. — Geologie Mijnbouw 53: 369-385.

Zandstra, J. G., 1967. Een zwerfsteentelling bij Wijldemerk (Gaasterland). — Grondboor Hamer, 1967, 1: 12-19.

—, 1972. Notities over heidegrond en hun ontgining en twee nieuwe zwerfsteentellingen in Drente. — Ibidem, 1972, 1: 7-18.

—, 1974. Over de uitkomsten van nieuwe zwerfsteentellingen en een keileemtypen indeling in Nederland. — Ibidem, 28, 5: 95-108.

Zeuner, F. E., 1959. The Pleistocene Period. — Hutchinson, London: 1-447.

Zonneveld, J. I. S., 1975. Zijn de noordnederlandse stuwwallen overreden of niet? — Ber. Fys. Geogr. Afd. Rijks Univ. Utrecht, 9: 1-13.

Manuscript received 24 June 1977.

