

**Intestinally Secreted C-Type Lectin Reg3b
Attenuates Salmonellosis but Not
Listeriosis in Mice**

Marleen T. J. van Ampting, Linda M. P. Loonen, Arjan J. Schonewille, Irene Konings, Carolien Vink, Juan Iovanna, Mathias Chamillard, Jan Dekker, Roelof van der Meer, Jerry M. Wells and Ingeborg M. J. Bovee-Oudenhoven
Infect. Immun. 2012, 80(3):1115. DOI: 10.1128/IAI.06165-11.
Published Ahead of Print 17 January 2012.

Updated information and services can be found at:
<http://iai.asm.org/content/80/3/1115>

	<i>These include:</i>
REFERENCES	This article cites 28 articles, 13 of which can be accessed free at: http://iai.asm.org/content/80/3/1115#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Intestinally Secreted C-Type Lectin Reg3b Attenuates Salmonellosis but Not Listeriosis in Mice

Marleen T. J. van Ampting,^{a,b} Linda M. P. Loonen,^{a,c} Arjan J. Schonewille,^{a,b} Irene Konings,^{a,c} Carolien Vink,^{a,b} Juan Iovanna,^d Mathias Chamailard,^e Jan Dekker,^a Roelof van der Meer,^a Jerry M. Wells,^{a,c} and Ingeborg M. J. Bovee-Oudenhoven^{a,b}

TI Food and Nutrition, Wageningen, The Netherlands^a; Department of Health, NIZO food research, Ede, The Netherlands^b; Host Microbe Interactomics Group, Animal Sciences Department, Wageningen University, Wageningen, The Netherlands^c; INSERM U624, Marseille, France^d; and INSERM U801, Lille, France^e

The Reg3 protein family, including the human member designated pancreatitis-associated protein (PAP), consists of secreted proteins that contain a C-type lectin domain involved in carbohydrate binding. They are expressed by intestinal epithelial cells. Colonization of germ-free mice and intestinal infection with pathogens increase the expression of Reg3g and Reg3b in the murine ileum. Reg3g is directly bactericidal for Gram-positive bacteria, but the exact role of Reg3b in bacterial infections is unknown. To investigate the possible protective role of Reg3b in intestinal infection, Reg3b knockout (Reg3b^{-/-}) mice and wild-type (WT) mice were orally infected with Gram-negative *Salmonella enteritidis* or Gram-positive *Listeria monocytogenes*. At day 2 after oral *Listeria* infection and at day 4 after oral *Salmonella* infection, mice were sacrificed to collect intestinal and other tissues for pathogen quantification. Protein expression of Reg3b and Reg3g was determined in intestinal mucosal scrapings of infected and noninfected mice. In addition, *ex vivo* binding of ileal mucosal Reg3b to *Listeria* and *Salmonella* was investigated. Whereas recovery of *Salmonella* or *Listeria* from feces of Reg3b^{-/-} mice did not differ from that from feces of WT mice, significantly higher numbers of viable *Salmonella*, but not *Listeria*, bacteria were recovered from the colon, mesenteric lymph nodes, spleen, and liver of the Reg3b^{-/-} mice than from those of WT mice. Mucosal Reg3b binds to both bacterial pathogens and may interfere with their mode of action. Reg3b plays a protective role against intestinal translocation of the Gram-negative bacterium *S. enteritidis* in mice but not against the Gram-positive bacterium *L. monocytogenes*.

In many countries in the industrialized world, food-borne intestinal infections are continuing to increase (11). For example, outbreaks of salmonellosis and listeriosis have been reported for decades, but within the past 25 years, their incidence has increased on many continents (29). To inhibit the colonization (adhesion to the intestinal epithelium) and translocation (invasion of host tissues) of food-borne pathogens and commensals, the intestinal mucosal surfaces are armed with an array of physical and chemical defense mechanisms. The low pH of the gastric compartment and bile acids secreted in the proximal intestine reduce the number of bacteria that will survive within the gut (7, 8). Other important defense mechanisms against pathogens include competition with commensal bacteria for nutrients and adhesion sites, a thick mucus layer on the luminal side of the epithelium, and the rapid innate response of the intestinal epithelium (6, 23).

Studies in rodents indicate that innate recognition of bacteria or bacterial components triggers epithelial expression of secreted C-type lectins Reg3g and Reg3b (3, 12, 22, 25). In rodents, mucosal and fecal levels of the Reg3 protein are associated with the severity of infection (25). The human homologue of this protein (i.e., they are, on the amino acid level, the most identical proteins between the two species [bidirectional best hit]) is pancreatitis-associated protein (PAP). PAP is also detectable in feces, and its use as a biomarker to monitor or discriminate human intestinal disease is the subject of ongoing studies.

Currently, the role and function of the Reg3 family members are of much interest. The murine Reg3g and human PAP have been shown to have antimicrobial activity against Gram-positive bacteria but not Gram-negative *Escherichia coli* (3). PAP and Reg3g were shown to bind peptidoglycan carbohydrate, which is critical for bacterial killing (14). Expression of Reg3g is greatly reduced in mice deficient in MyD88, an intracellular adaptor pro-

tein involved in most Toll-like receptor (TLR)-mediated signaling. MyD88-deficient mice have been shown to be more susceptible to *Listeria monocytogenes* infection than wild-type (WT) mice (2). Injection of recombinant Reg3g into the lumen of MyD88-deficient ligated ileal loops before inoculation with *L. monocytogenes* was shown to reduce *L. monocytogenes* survival in the gut. Other *in vivo* studies showed that genetic ablation of Reg3b, which is another Reg3 murine isoform, resulted in impaired clearance of the bacterial load in ileal Peyer's patches during *Yersinia pseudotuberculosis* infection without affecting luminal bacterial levels (5). It remains unclear, however, how Reg3b protects against bacterial translocation and whether this is specific for Gram-negative pathogens.

In this study, we used Reg3b knockout (Reg3b^{-/-}) mice and their WT littermates to investigate the protective role of Reg3b following oral challenge with Gram-negative *Salmonella enteritidis* or Gram-positive *L. monocytogenes*. Additional *ex vivo* experiments were performed to gain insights into the mechanisms involved in the protective function of Reg3b.

Received 16 November 2011 Returned for modification 12 December 2011

Accepted 3 January 2012

Published ahead of print 17 January 2012

Editor: J. L. Flynn

Address correspondence to Ingeborg M. J. Bovee-Oudenhoven, Ingeborg.Bovee@nizo.nl.

M. T. J. van Ampting and L. M. P. Loonen contributed equally to this study.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/IAI.06165-11

MATERIALS AND METHODS

Animals and diet. The experimental protocol was approved by the animal welfare committee of Wageningen University (Wageningen, The Netherlands). Reg3b^{-/-} mice with a C57BL/6 × 129O1a background were generated as described previously (15). Knockout mice were obtained by breeding heterozygote mice. Genotyping was performed as described previously (15), and Reg3b^{-/-} mice were used as the knockout mice and Reg3b^{+/+} littermates were used as the WT control. The mice were selected based on genotype and age and were fed a semipurified AIN93-G (21) diet (Abdiets, Woerden, The Netherlands). At 7 to 9 weeks of age, mice with an average body weight of 22 g (range, 16 to 31 g) were housed individually. They received a purified diet containing (per kg of feed) 200 g acid casein, 326 g cornstarch, 174 g glucose, 160 g palm oil, 40 g corn oil, 50 g cellulose, and a vitamin and mineral mix (without calcium) according to AIN93 (21). Diets were supplemented with CaHPO₄ · 2H₂O (Merck, Darmstadt, Germany) to a final concentration of 30 mmol/kg. To mimic the composition of a Western human diet, the prepared diets were relatively low in calcium and high in fat compared to standard rodent diets (21). Food and demineralized drinking water were supplied *ad libitum*. Mice were separated into groups, and age and gender were randomized over all groups as much as possible. The power analysis used to calculate the minimum number of animals needed per group was based on infection studies previously performed in this model (our unpublished data). In total, 3 Reg3b^{-/-} and 3 WT groups were formed. From each mouse strain, one group was orally infected with *Salmonella* (Reg3b^{-/-}, *n* = 10; WT, *n* = 12), a second group of each strain was orally infected with *Listeria* (Reg3b^{-/-}, *n* = 7; WT, *n* = 8), and a final group was sham treated (Reg3b^{-/-}, *n* = 5; WT, *n* = 6). Body weight was measured every 2 days before infection and daily after infection.

Nramp1 genotype of *Salmonella*-infected mice. To monitor susceptibility to *Salmonella* infection, the natural resistance-associated macrophage protein 1 (Nramp1) genotype was determined in the *Salmonella*-infected Reg3b^{+/+} and Reg3b^{-/-} mice. A PCR product was created from the genomic DNA by using primers 5'-GGAATGAATGTCAAGCAGCCAG-3' and 5'-ATCCCACCTCATAGCCGAAG-3' in order to sequence the area of the Nramp1 gene that is known to vary between different mouse strains (G or A at position 596) (9). The PCR cycle was 3 min at 95°C, 35 cycles of 30 s at 95°C, 30 s at 58°C, and 1 min at 72°C, and 5 min at 72°C. The PCR product was subsequently sequenced by Baseclear (Leiden, The Netherlands) using the primers 5'-CACCCATTCCAGTAGGG-3' and 5'-CCTGTGACCTGGATGTTTC-3' to determine the Nramp1 genotype of the *Salmonella*-infected mice.

Infection. After adaptation to individual housing and the diet for 8 days, mice were orally infected by gavage with 0.2 ml saline containing 5 × 10⁸ CFU of *S. enteritidis* (clinical isolate, phage type 4, strain NIZO1241; NIZO food research, Ede, The Netherlands) or *L. monocytogenes* (animal isolate, strain EGD-e, serotype 1/2a; NIZO food research). The virulence of each strain was sustained by routine oral passage in Reg3b^{+/+} mice and subsequent isolation of the pathogen from extraintestinal organs. *S. enteritidis* was grown on brilliant green agar modified (BGAM) plus sulfamandolate supplement (Oxoid, Basingstoke, United Kingdom) and quantified as described previously (18). *L. monocytogenes* was grown and enumerated by similar culturing methods using Palcam agar (28) plus Palcam selective supplement (Oxoid).

Collection of biological samples and bacterial quantification. A pilot study indicated that the determination of the number of *L. monocytogenes* and *S. enteritidis* bacteria in organs of mice was optimal, which means that the numbers of *Listeria* and *Salmonella* CFU could be detected in organs at 2 and 4 days after oral infection, respectively (our unpublished data). Noninfected mice were sacrificed 3 days after oral sham treatment. Mice were anesthetized by isoflurane, and blood was collected via an orbital puncture to isolate heparin plasma. After cervical dislocation, the colon and the distal third of the small intestine, representing the ileum, were excised. Approximately 1 cm was cut from the middle of each intestinal segment to quantify the number of viable *Salmonella* or *Listeria* cells pres-

ent in the colonic and ileal tissue. This piece was cut open longitudinally, briefly flushed in sterile saline, and homogenized in 250 μl of saline, and 10-fold dilutions were plated for bacterial growth on selective agar for *Salmonella* or *Listeria* (as described above). The remaining parts of the ileum and colon were cut open longitudinally and flushed with saline, and the mucosa was isolated by scraping with a spatula. Mucosal samples were immediately frozen in liquid nitrogen for protein analysis. Furthermore, the mesenteric lymph nodes (MLN), spleen, and liver were removed and, after homogenization in sterile saline, directly used for *Salmonella* or *Listeria* quantification as described above. Bacterial counts were expressed as the total log₁₀ CFU per gram of tissue.

Reg3b and Reg3g protein analysis in the ileal mucosa. Reg3g and Reg3b expression was determined in the ileum because this is the area where expression is upregulated during infection of rats (22, 25) and upon microbial colonization of germ-free mice (3). Frozen mucosal scrapings of the ileum were pulverized under liquid nitrogen. Approximately two-thirds of the pulverized tissue was suspended in a 0.2-mol/liter sucrose buffer of pH 7.4 containing 20 mmol/liter Tris and a protease inhibitor cocktail (Complete, Roche Diagnostics). To homogenize the samples, they were sonicated on ice for 30 s at level 4 (Sonicator XL2020; Heat Systems, Farmingdale, NY), and protein concentrations were determined using the bicinchoninic acid (BC) assay (Omnilab, Breda, The Netherlands) according to the manufacturer's protocol. Samples from each group were pooled, and 100 μg protein was denatured at 95°C for 10 min in sample buffer (4× buffer containing 160 mM Tris [pH 6.8], 8% SDS, 40% glycerol, 0.05% [wt/vol] bromophenol blue), subjected to SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (4.8% stacking gel [pH 6.8], 12.5% separation gel [pH 8.8]), and transferred to a polyvinylidene difluoride (PVDF) membrane (Immobilon-P; Millipore, Billerica, MA). After blocking, the membranes were incubated with polyclonal anti-Reg3b (1:50,000) or anti-Reg3g (1:10,000) antibodies (custom made by Eurogentec, Seraing, Belgium). These antibodies were generated in rabbits against synthetically produced peptides by using the peptide sequences GEDSLKNIPSARISC (Reg3b) and EVAKKDAPSSRSSC (Reg3g). The chosen peptide sequences correspond to unique sequences within the Reg3 protein and allow differentiation between the Reg3b and Reg3g proteins. Serum from immunized rabbits was affinity purified using the same peptides. The signal of the secondary horseradish peroxidase (HRP)-conjugated antibody (goat anti-rabbit HRP, 1:100,000; Jackson ImmunoResearch, Suffolk, United Kingdom) was detected by using the ECL Plus chemiluminescent detection kit (GE Healthcare, Den Bosch, The Netherlands).

Myeloperoxidase analysis in the ileal mucosa and serum amyloid A detection in plasma. A mouse myeloperoxidase (MPO) enzyme-linked immunosorbent assay (ELISA) kit (Hycult Biotechnology, Uden, The Netherlands) was used according to the manufacturer's guidelines to determine the concentration of MPO in ileal mucosal scrapings.

Levels of serum amyloid A (SAA) 2, an acute-phase apolipoprotein present in plasma (24), were determined in order to investigate systemic inflammation. A mouse SAA 2 ELISA kit (Life Diagnostics, Inc., West Chester, Pennsylvania) was used according to the guidelines of the manufacturer.

Pulldown assay. To investigate the possible mode of action, it was evaluated in a pulldown assay whether Reg3b present in the ileal mucosa binds to *S. enteritidis* or *L. monocytogenes*. *S. enteritidis* and *L. monocytogenes* strains identical to those used in the infection study were grown overnight (37°C in a horizontal shaker at 250 rpm) in LB and brain heart infusion (BHI) broth, respectively.

Bacteria grown overnight were spun down (3 min at 2,350 × *g*), and 1 × 10⁸ bacteria were resuspended in 20 μl ileal mucosal scraping extracts from *Salmonella*-infected Reg3b^{-/-} or WT mice which were diluted 1:5 in the above-described sucrose buffer. The bacteria were incubated for 30 min at 37°C. After the bacteria were spun down (5 min at 1,500 × *g*), the presence of Reg3b was investigated in the supernatant (15 μl) by SDS-PAGE as described above. Proteins were transferred to an Immobilon-FL




FIG 1 Western blot analysis of Reg3b and Reg3g in pooled ileal mucosa samples of Reg3b^{-/-} mice (KO) or WT mice (Wt). Reg3b^{-/-} mice ($n = 10$) and WT mice ($n = 12$) were orally infected with *S. enteritidis* (S), or Reg3b^{-/-} mice ($n = 7$) and WT mice ($n = 8$) were orally infected with *L. monocytogenes* (L). Noninfected mice ($n = 5$) received saline as a control treatment.

membrane (Millipore) and, after blocking, exposed to rat anti-mouse Reg3b antibodies (MAB5110, 1:2,000; R&D Systems, Minneapolis). The secondary antibody used was goat anti-rat IRDye 680CW (1:20,000; Li-Cor, Nebraska), and the blots were scanned using the Odyssey scanner (Li-Cor). Experiments were performed in triplicate.

Statistics. All data are the means and the standard errors of the means (SEM), and statistical analysis was performed by using Prism 5.0 software (GraphPad, San Diego, California). Our aim was to investigate the role of Reg3b in infection. Therefore, the predefined comparisons were between Reg3b^{-/-} and the WT except for *in vitro* investigations to evaluate the binding of Reg3b to *Salmonella* and *Listeria* when binding to these pathogens was investigated. Data were tested for normality by the Kolmogorov-Smirnov normality test and the Shapiro-Wilk normality test. If the data were normally distributed, the differences were tested for significance using Student's *t* test (two-tailed). Data with unequal variances were tested by using the Mann-Whitney U test. *Salmonella* output in feces was determined at multiple time points, and therefore, these data were analyzed by a repeated-measures two-way analysis of variance (ANOVA) (mixed model). Differences were considered statistically significant when the *P* value was < 0.05 .

RESULTS

Body weight. Body weight was determined during the experiment to monitor the general condition of the mice. At the start of the experiment, the average body weight was 22 ± 0.7 g. During the study, there was no significant difference between the body weight of Reg3b^{-/-} mice and that of WT mice (data not shown). The average body weights before infection (average weight during 2 weeks prior to infection) were 22.8 ± 1.4 g and 23.1 ± 1.4 g for the WT and Reg3b^{-/-} groups, respectively. The postinfection body weights of the WT and Reg3b^{-/-} groups were 23.2 ± 1.4 g and 23.2 ± 1.5 g (average of the postinfection period).

Bacterial infection increases Reg3b and Reg3g in the ileum mucosa. To evaluate the presence of the Reg3b and Reg3g proteins in the ileal mucosae of WT and Reg3b^{-/-} mice and the subsequent levels upon oral infection, mucosal scrapings were evaluated by immunoblotting using antibodies specific for Reg3b and Reg3g. As expected, Reg3b was not detected in the ileal mucosa of Reg3b^{-/-} animals (Fig. 1). After *Salmonella* and *Listeria* infection, the protein level was higher in the mucosal samples of WT mice than it was before infection (Fig. 1). The isoform Reg3g was




FIG 2 *Salmonella* (triangles) and *Listeria* (circles) excretion in feces. Individually housed mice were orally infected at day 0. Reg3b^{-/-} mice (filled symbols) were infected with 5×10^8 CFU of *S. enteritidis* ($n = 10$) or *L. monocytogenes* ($n = 7$), and WT littermates (open symbols) were infected with identical numbers of CFU of *S. enteritidis* ($n = 12$) or *L. monocytogenes* ($n = 8$). Values are the means and SEM.

detectable in mucosal scrapings of noninfected Reg3b^{-/-} mice (Fig. 1). In contrast, the WT mice had detectable levels of this Reg3 isoform and levels increased during *Salmonella* and *Listeria* infection in WT and Reg3b^{-/-} mice.

Reg3b^{-/-} mice are more susceptible to salmonellosis, but not listeriosis, than WT mice. In order to ensure similar Nramp1 genotypes in all *Salmonella*-infected animals, these mice were sequenced for a previously reported point mutation in this gene which could affect infection susceptibility (9). All mice were shown to be homozygous for the Nramp1 susceptibility allele (data not shown). To investigate the role of Reg3b in intestinal colonization, the numbers of viable *Salmonella* cells were determined in fresh fecal samples of infected WT and Reg3b^{-/-} mice. Colonization of *Salmonella* in Reg3b^{-/-} mice was not different from that in WT mice (Fig. 2). In the colonic tissue, which was first flushed with saline to remove intestinal content, higher numbers of *Salmonella* cells were recovered in Reg3b^{-/-} mice than in WT mice (Fig. 3A) ($P < 0.05$). In the ileal tissue, *Salmonella* levels were identical in Reg3b^{-/-} and WT mice. Furthermore, in MLN, spleen, and liver, these levels were higher in Reg3b^{-/-} mice than in their WT counterparts (Fig. 3B) ($P < 0.05$).

Intestinal colonization of *Listeria*, as found for *Salmonella*, was identical in Reg3b^{-/-} and WT mice (Fig. 2). To further investigate the effects of Reg3b on *Listeria* translocation, the numbers of CFU in ileal and colonic tissue and extraintestinal organs were determined. The number of *Listeria* cells recovered from the WT did not differ from that recovered from Reg3b^{-/-} mice in ileal and colonic tissues (Fig. 3C) and in MLN, spleen, and liver (Fig. 3D).

Mucosal and systemic inflammation. The role of Reg3b in inflammation was investigated by measuring the levels of the inflammation marker myeloperoxidase (MPO) in the ileum mucosa at 2 and 4 days after oral infection of *Listeria* and *Salmonella* groups, respectively. MPO levels of *Salmonella*-infected (0.49 ± 0.05 ng/mg protein) and *Listeria*-infected (0.39 ± 0.06 ng/mg protein) groups did not differ significantly from that of the noninfected groups (0.35 ± 0.05 ng/mg protein) (data not shown). To study systemic inflammation, SAA levels in plasma were determined at the same time points. The levels were elevated from 0.1 ± 0.001 μ g/ml (noninfected mice) to $1,164 \pm 186$ and 502 ± 174 μ g/ml upon *Salmonella* and *Listeria* infection, respectively ($P < 0.05$) (data not shown). However, there were no significant differences between WT and Reg3b^{-/-} mice.




FIG 3 Levels of *Salmonella* (A and B) and *Listeria* (C and D) in intestinal and extraintestinal tissues at 4 and 2 days after oral infection, respectively. Numbers of CFU of pathogens in the ileal and colonic mucosas (A and C) and in mesenteric lymph nodes (MLN), spleen, and liver (B and D) of Reg3b^{-/-} mice ($n = 10$; black bars) or WT mice ($n = 12$; white bars). Values are the means and SEM. An asterisk indicates that the results are significantly different from those of the WT ($P < 0.05$).

Reg3b binds to *Salmonella* and *Listeria*. To investigate the possible mode of action, it was evaluated in a pull-down assay whether Reg3b present in the ileal mucosa binds to *S. enteritidis* and *L. monocytogenes*. Upon *ex vivo* incubation of mucosal samples from infected mice with these bacteria, it was shown that Reg3b protein levels decreased in the supernatant (Fig. 4). It was mainly the lower-molecular-weight form of the proteolytically processed protein that decreased after incubation with bacteria. This indicated that it is the cleaved form of Reg3b that binds to the pathogens, not the full-length polypeptide. We have shown previously that the Reg3 protein is not precipitated even when centrifuged at $15,000 \times g$ (25). Therefore, the results presented here indicate that ileal mucosal Reg3b is able to directly bind both *Salmonella* and *Listeria*.

DISCUSSION

Both microbial infections of conventional mice and intestinal colonization of germ-free mice by commensals increase ileal expression of the secreted C-type lectins Reg3g and Reg3b (3, 12, 22, 25). While Reg3g has been shown to have a bactericidal activity against Gram-positive bacteria (3), the physiological function of Reg3b

remains unknown. This study demonstrates a role for Reg3b in protection against infection with Gram-negative *S. enteritidis*. Despite the fact that similar levels of viable *Salmonella* cells were recovered from feces, higher numbers were present in MLN, spleen, and liver of infected Reg3b^{-/-} mice than in WT mice, indicating that the translocation and dissemination of *Salmonella* in host tissues are increased in the absence of Reg3b. As the mice were bred on a mixed background (C57BL/6 and 129O1a) and these strains differ in their susceptibility to *Salmonella* infection due to a mutation (G to A) in the Nrampl1 gene, all mice were genotyped individually (9). All the *Salmonella*-infected animals were shown to be homozygous for the Nrampl1 susceptibility allele, ruling out the possibility that the Nrampl allele status would affect the results. In colonic tissue, *Salmonella* levels were also increased in Reg3b^{-/-} mice compared to those of WT mice. The MLN drain the small intestine, which is a main site of *Salmonella* invasion (19). Therefore, reduced levels of *Salmonella* in the MLN of WT mice suggest protective effects of Reg3b on *Salmonella* translocation in the ileum. Although we did not observe differences in *Salmonella* counts in the ileal tissues of WT and Reg3b^{-/-} mice, this may be due to a lack of statistical power in this assay, as SEM levels are higher in ileal tissue than in colonic tissue. Moreover, the investigated ileal tissue did not contain Peyer’s patches and M cells, which, in addition to absorptive enterocytes, are considered to be important sites of *Salmonella* translocation (10, 27). All together, the experimental infection data strongly suggest that Reg3b inhibits *Salmonella* translocation from the gut lumen into intestinal tissues and further extraintestinal tissues but does not kill *Salmonella* in the gut lumen. These results are supported by the recent findings of Dessein et al., who showed that genetic ablation of Reg3b did not affect the bacterial load of the Gram-negative pathogen *Yersinia pseudotuberculosis* in the intestinal lumen (5) but significantly increased the bacterial burden in the Peyer’s




FIG 4 *Listeria* and *Salmonella* pull-down assays. Presence of Reg3b, detected by Western blot analysis, in supernatant fractions of ileal mucosa samples from *S. enteritidis*-infected WT (Wt) or Reg3b^{-/-} (KO) mice incubated with *Salmonella* (S) or *Listeria* (L). Mucosal samples incubated without bacteria served as the negative control (-). The prestained protein marker is indicated by M.

patches, which are the main route of entry for this pathogen. In contrast, Reg3b did not influence resistance to infection with Gram-positive *Listeria monocytogenes*, as identical numbers of this pathogen were recovered from intestinal and extraintestinal tissues in Reg3b^{-/-} and WT mice, as found in our study.

As murine Reg3g and human PAP are directly bactericidal for Gram-positive bacteria *in vitro* (3, 17), it seemed possible that Reg3b would have similar antimicrobial activity. However, in this study, there was no effect of Reg3b on the colonization and translocation of *L. monocytogenes*. The strain of *L. monocytogenes* used in our infection study was identical to that used previously for *in vitro* antibacterial assays with Reg3g (3). Previous investigations with rat PAP1, which is the homologue of the murine Reg3b, also showed no direct bactericidal effect against *L. monocytogenes in vitro* (25). Others also report a lack of bactericidal effect of this protein (5). The coexistence of the two closely related Reg3 proteins in the mouse strongly suggests a different function for each of these proteins. Here we provide evidence that Reg3b has a protective role against Gram-negative *Salmonella* translocation but not *Listeria* infection. The exact mechanism by which Reg3b inhibits *Salmonella* translocation into host tissues is unclear.

The observed difference in Reg3b functionality between listeriosis and salmonellosis might be due to differences in physical appearance and host invasion mechanisms of the two pathogens but might also be related to differences in host responses induced by these pathogens. We can only speculate about the latter, as we did not monitor the exact (immune) responses of the mice. Previous studies by Brandl et al. (2) indicated that the innate immune defense against *L. monocytogenes*, which requires Toll-like receptor (TLR)-mediated signals, plays a crucial role in Reg3g-related protection against this pathogen. As *S. enteritidis* and *L. monocytogenes* may trigger different (TLR-mediated) innate immune signals (1), this might be a cause of differential regulation of Reg3b expression and, therefore, the subsequent protective effects of this protein. On the other hand, it might be that Reg3b modulates the immune response directed against *Salmonella* but not the response against *Listeria*. Although measured at a single time point in the current study, Reg3b did not affect the systemic inflammatory response as measured by SAA levels in serum. The local inflammatory response in our infection models was relatively low, as MPO levels did not increase upon infection in the ileal mucosa. This was likely due to the relatively short period that mice were exposed to the orally administered pathogens because our focus was on pathogen colonization and translocation, which normally precede inflammation. Thus, based on our study design, we cannot fully exclude the absence of effects of Reg3b on the inflammatory response. However, previous investigations in our lab with a similar *Salmonella* infection model in rats indicated that mucosal inflammation in this infection model is relatively low at day 4 postinfection (26). Further analysis of the collected intestinal tissues from the current study might give insight on the role of the innate (immune) response in Reg3b-related protective effects. For example, microarray analysis of intestinal tissues may identify whether and which host defense pathways are associated with protective effects of Reg3b. Performing additional *Listeria* and *Salmonella* infection experiments in mice deficient in specific TLRs might reveal the involvement of specific innate immune signaling in Reg3b-mediated protection against intestinal infection.

Besides the proposed association of the host response with Reg3b-related protection of intestinal infection, this lectin might

also directly and differentially affect *Listeria* and *Salmonella*. The outer membranes of these pathogens are different; for example, *Salmonella* contains immunoreactive lipopolysaccharides (LPS) which are absent in Gram-positive bacteria, such as *Listeria*. This influences host signaling pathways, e.g., TLR recognition. Our *ex vivo* analysis revealed that ileal mucosal Reg3b showed binding affinity to both *Salmonella* and *Listeria*. Bactericidal activities of Reg3g depend on binding to cell wall peptidoglycan, a molecule exposed on the Gram-positive bacterial surface (8). This peptidoglycan recognition is determined by the so-called Loop1 EPN tripeptide motif, which is also present in Reg3b (14). This motif is involved in peptidoglycan binding, and binding affinity for carbohydrate ligands depends on carbohydrate chain length (14). Moreover, Reg3b binds to the Gram-positive *Bacillus subtilis* peptidoglycan, mannose polysaccharides, and chitin, which is a long-chain polymer of *N*-acetylglucosamine (14). Our results suggest that Reg3b possibly recognizes carbohydrate structures on the bacterial surface of microorganisms which enter the intestinal mucosa. Currently we do not know what the specific target of Reg3b is on the outer membrane of bacteria. It may be related to LPS, which are large molecules consisting of a lipid and a polymeric carbohydrate structure on most Gram-negative bacteria (20). This may be a binding target for the EPN motif in Reg3b. This motif is likely, as shown for Reg3g, also involved in the binding of Reg3b to Gram-positive bacteria. The molecular mechanism by which Reg3b may recognize its binding target remains to be determined. Studies with purified Reg3b are needed to determine what ligands specifically bind to this protein. To address this question, it is necessary to have purified, biologically active Reg3b protein, and efforts are under way to produce and purify this protein in an active form. Although binding may be crucial for the function of Reg3b, it does not necessarily result in a protective effect *in vivo*, as possible binding of Reg3b to *Listeria* in the intestinal mucosa does not result in reduced *Listeria* levels in extraintestinal organs. This might be caused by another important difference between the pathogens, which is that *Listeria* and *Salmonella* use complex and very different mechanisms of cellular attachment and invasion (4, 19). *Listeria* can invade the human intestinal epithelium via the epithelial receptor E-cadherin. A single amino acid change in the murine E-cadherin, however, makes mice relatively resistant to intestinal infection (13). As also shown in the present study, relatively high doses of *Listeria* do achieve a significant invasive infection in rodents (13). This probably involves a second but less-well-characterized surface protein of *L. monocytogenes*, internalin B, which enables *Listeria* to enter and survive within epithelial cells via interaction with three different epithelial membrane receptors (4, 19). In the case of *Salmonella*, the bacteria are not highly adherent, but their invasion machinery is particularly efficient (19). *Salmonella* has another mechanism to invade cells which involves binding to epithelial cells by its type III secretory system (TTSS) (10, 16). The TTSS allows direct activation of components of the host-cell cytoskeleton by intracellular delivery of dedicated bacterial effectors, resulting in membrane ruffling and endocytosis. By this event, *Salmonella* species invade and reside in an atypical acidic compartment called the *Salmonella*-containing vacuole (SCV) and survive inside the cell (4). It is possible that Reg3b interferes with the expression or function of *Salmonella* pathogenicity island 1 (SPI-1) TTSS invasion machinery.

In conclusion, Reg3b inhibits intestinal bacterial translocation

upon oral infection with Gram-negative *Salmonella*. In contrast, this protein does not have protective effects against intestinal infection with Gram-positive *Listeria*. Inhibitory effects of Reg3b on bacterial infection may be linked to its observed binding to pathogens. The protective mechanism of Reg3b in salmonellosis is not associated with direct bactericidal effects but may be related to interference with *Salmonella* virulence mechanisms or host responses to this pathogen.

ACKNOWLEDGMENTS

We thank the biotechnicians at the Small Animal Centre of Wageningen University (Wageningen, The Netherlands) for excellent assistance. We also thank our colleagues at the Department of Health of NIZO food research and members of the Host-Microbe Interactomics group at Wageningen University for fruitful discussions.

REFERENCES

1. Abreu MT, Fukata M, Arditi M. 2005. TLR signaling in the gut in health and disease. *J. Immunol.* 174:4453–4460.
2. Brandl K, Plitas G, Schnabl B, DeMatteo RP, Pamer EG. 2007. MyD88-mediated signals induce the bactericidal lectin RegIII gamma and protect mice against intestinal *Listeria monocytogenes* infection. *J. Exp. Med.* 204:1891–1900.
3. Cash HL, Whitham CV, Behrendt CL, Hooper LV. 2006. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science* 313:1126–1130.
4. Cossart P, Sansonetti PJ. 2004. Bacterial invasion: the paradigms of enteroinvasive pathogens. *Science* 304:242–248.
5. Dessein R, et al. 2009. TLR2 is critical for induction of REG3{beta} expression and intestinal clearance of *Yersinia pseudotuberculosis*. *Gut* 58:771–776.
6. Duncan HE, Edberg SC. 1995. Host-microbe interaction in the gastro-intestinal tract. *Crit. Rev. Microbiol.* 21:85–100.
7. Giannella RA, Broitman SA, Zamcheck N. 1973. Influence of gastric acidity on bacterial and parasitic enteric infections. A perspective. *Ann. Intern. Med.* 78:271–276.
8. Gorden J, Small PL. 1993. Acid resistance in enteric bacteria. *Infect. Immun.* 61:364–367.
9. Govoni G, et al. 1995. Genomic structure, promoter sequence, and induction of expression of the mouse *Nramp1* gene in macrophages. *Genomics* 27:9–19.
10. Grassl GA, Finlay BB. 2008. Pathogenesis of enteric *Salmonella* infections. *Curr. Opin. Gastroenterol.* 24:22–26.
11. Kaur T, Ganguly NK. 2003. Modulation of gut physiology through enteric toxins. *Mol. Cell. Biochem.* 253:15–19.
12. Keilbaugh SA, et al. 2005. Activation of RegIIIbeta/gamma and interferon gamma expression in the intestinal tract of SCID mice: an innate response to bacterial colonisation of the gut. *Gut* 54:623–629.
13. Lecuit M. 2007. Human listeriosis and animal models. *Microbes Infect.* 9:1216–1225.
14. Lehotzky RE, et al. 2010. Molecular basis for peptidoglycan recognition by a bactericidal lectin. *Proc. Natl. Acad. Sci. U. S. A.* 107:7722–7727.
15. Lieu HT, et al. 2006. Reg2 inactivation increases sensitivity to Fas hepatotoxicity and delays liver regeneration post-hepatectomy in mice. *Hepatology* 44:1452–1464.
16. Ly KT, Casanova JE. 2007. Mechanisms of *Salmonella* entry into host cells. *Cell. Microbiol.* 9:2103–2111.
17. Mukherjee S, et al. 2009. Regulation of C-type lectin antimicrobial activity by a flexible N-terminal prosegment. *J. Biol. Chem.* 284:4881–4888.
18. Oudenhoven IM, Klaasen HL, Lapre JA, Weerkamp AH, Van der Meer R. 1994. Nitric oxide-derived urinary nitrate as a marker of intestinal bacterial translocation in rats. *Gastroenterology* 107:47–53.
19. Pizarro-Cerda J, Cossart P. 2006. Bacterial adhesion and entry into host cells. *Cell* 124:715–727.
20. Raetz CR, et al. 1991. Gram-negative endotoxin: an extraordinary lipid with profound effects on eukaryotic signal transduction. *FASEB J.* 5:2652–2660.
21. Reeves PG, Nielsen FH, Fahey GC, Jr. 1993. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J. Nutr.* 123:1939–1951.
22. Rodenburg W, Bovee-Oudenhoven IM, Kramer E, van der Meer R, Keijzer J. 2007. Gene expression response of the rat small intestine following oral *Salmonella* infection. *Physiol. Genomics* 30:123–133.
23. Sarker SA, Gyr K. 1992. Non-immunological defence mechanisms of the gut. *Gut* 33:987–993.
24. Uhlar CM, Whitehead AS. 1999. Serum amyloid A, the major vertebrate acute-phase reactant. *Eur. J. Biochem.* 265:501–523.
25. van Ampting MT, et al. 2009. Ileal mucosal and fecal pancreatitis associated protein levels reflect severity of *Salmonella* infection in rats. *Dig. Dis. Sci.* 54:2588–2597.
26. van Ampting MT, et al. 2009. Intestinal barrier function in response to abundant or depleted mucosal glutathione in *Salmonella*-infected rats. *BMC Physiol.* 9:6.
27. van Asten AJAM, Koninkx JFJG, van Dijk JE. 2005. *Salmonella* entry: M cells versus absorptive enterocytes. *Vet. Microbiol.* 108:149–152.
28. van Netten P, Perales I, van de Moosdijk A, Curtis GD, Mossel DA. 1989. Liquid and solid selective differential media for the detection and enumeration of *L. monocytogenes* and other *Listeria* spp. *Int. J. Food Microbiol.* 8:299–316.
29. WHO. 2007. Food safety and foodborne illness. <http://www.who.int/mediacentre/factsheets/fs237/en/index.html>.