

Young mangrove stands produce a large and high quality litter input to aquatic systems

B.T. Nga^{1,*}, H.Q. Tinh¹, D.T. Tam¹, M. Scheffer² and R. Roijackers²

¹Department of Environmental and Natural Resources Management, College of Agriculture, Cantho University, Vietnam; ²Department of Environmental Sciences, Aquatic Ecology and Water Quality Management Group, Wageningen University, Wageningen, The Netherlands; *Author for correspondence (e-mail: btnga@ctu.edu.vn; phone: +84-71-830635; fax: +84-71-830814)

Received 30 March 2004; accepted 10 November 2004

Key words: Biomass, Mangrove, Nitrogen, Phosphorus, Vietnam

Abstract

Mangrove swamps are key ecosystems along the Vietnam coast. Although mangrove litter is thought to represent an important input of organic matter and nutrients to the coastal aquatic systems, the factors determining the quality and size of this litter flux have not been studied so far. We monitored leaf, stipule, twig, and reproductive litter monthly in monocultures of *Rhizophora apiculata* mangrove forests of 7, 11, 17 and 24 years old in the Camau province, Mekong Delta, Vietnam. Litter traps were used to measure litter fall production from June 2001 till May 2002. Total litter fall was in the range of 8.86–14.16 t DW ha⁻¹ year⁻¹. Leaves were the main component, and represented 70% of litter fall production in all stands. Total litter fall was lower in the older stands but the amount of reproductive litter was significantly higher in these stands (17 and 24 years). Biomass of leaf litter was highest between the end of the wet season and the beginning of the dry season. Phosphorus and nitrogen levels in leaf litter were significantly higher in younger than in older stands. Overall, our study indicated that young stands produced the highest input of litter and particularly of nitrogen and phosphorus to the surrounding aquatic system. Consequently, these stands contribute significantly to the fisheries.

Introduction

Mangrove swamps are extremely productive ecosystems, which export organic matter to support a variety of organisms (Odum and Heald 1975; Lee 1989). The export of these large amounts of organic material can have substantial effects on food webs in coastal waters (Alongi 1990). In the past decade, mangrove forests have been under severe pressure by a rapidly increasing human population, large scale deforestation practices and conversion of forests into aquaculture farms especially in Southeast Asia. The fast developments in

aquaculture have resulted in non-sustainable farming systems (Graaf and Xuan 1998). Preservation of the mangrove forests is an important issue to protect the coastal ecosystem and to improve the water quality in the coastal areas. Therefore, the need for protection of the coastal forest belt has been an important issue (Loi et al. 2002). In recent years, studies on species classification, tree growth, succession, silviculture, forest utilization, and litter fall of mangroves have been carried out in Vietnam (e.g. Nam and Thuy 1997; Clough et al. 2000). However, information on litter fall from mangroves in Vietnam is lacking.

The mangrove forest in the coastal areas of Vietnam suffered severe damage over the past 50 years; it decreased from 400,000 to 290,000 ha due to overexploitation of forest timber, firewood and charcoal (Hong 1996). Since then, the deforestation continued by changing forests into agricultural land and primitive extensive and intensive shrimp farming i.e. about 53,969 ha of forest was left on the Ca Mau peninsula in 1993 (Department of Agriculture, Forestry and Fishery 1996). The Camau province (Camau Peninsula), situated in the south western part of the Mekong Delta, has a long shoreline and is the province with the most severe mangrove destruction. Following the cessation of hostilities in Vietnam, mangrove forests in the Camau peninsula initially recovered as a result of both natural regeneration and manual planting of the preferred forestry species, *Rhizophora apiculata* Blume. Although extension of shrimp farming in mangrove areas occurs at a rapid rate, world shrimp production has leveled off in recent years, as many aquaculture farms have either collapsed or experienced declining yields. For instance, production figures varied dramatically between the various shrimp culture systems with semi-intensive farms producing between 1000 and 2000 kg ha⁻¹ year⁻¹, whereas extensive farms produced 100–400 kg ha⁻¹ year⁻¹ in 1996 (Johnston et al. 2000), and 80–250 kg ha⁻¹ year⁻¹ in 2000 (Ministry of Fisheries 2001). Fish production is believed to be dependent on mangrove areas, and the dependence of many penaeid shrimp species on mangroves has also been shown (Christensen 1978; Barbier 2000). In fact, the decline in mangrove areas in the Mekong delta certainly had an impact on the decrease in coastal fisheries production over the last decades (Graaf and Xuan 1998). The study described here was carried out as part of a larger study for Integrated Management of Coastal Resources (MHO8-project) in the Mekong Delta, Vietnam, where a number of provincial governments have established shrimp farming-forestry enterprises. In these mixed aquaculture and mangrove forestry systems, shrimp and other cultured species are dependent mostly on the natural food chains for their food supply. It is widely recognized that mangrove derived detritus might be a major source of carbon for estuarine food webs along tropical and subtropical coastlines (e.g. Odum and Heald

1972; Malley 1978; Robertson and Daniel 1989; Daniel and Robertson 1990), but data for the systems we studied in Vietnam were lacking. The present research focused on the question how the dynamics and quality of mangrove litter input to aquatic systems differ between tree stands of different age.

Materials and methods

Description of the research area

Sampling was carried out from June 2001 till May 2002 in monocultures of the planted *Rhizophora apiculata* of respectively 7, 11, 17, and 24 years old. The mangrove stands of 7 and 11 years old were selected from the mixed shrimp farming-mangrove forestry systems Tam Giang 3 enterprise. The *Rhizophora apiculata* stands of 17 and 24 years were selected in the 184, and Kien Vang enterprises respectively in Camau province, Mekong Delta, Vietnam (Figure 1). The average tree density varied from 7000–10,000 trees per hectare. The mixed shrimp-mangrove system is characterized by mangrove stands, surrounded by ditches. The area of each of the mangrove stands is 7–12 ha of which approximately 30% are ditches. Average width and depth of these ditches are 7 and 1 m respectively. The research area is located at a latitude of about 8°50' N. The climate is humid tropical and dominated by monsoons. Mean annual rainfall was over 2400–3460 mm in 1996–2002. Annually, the temperature is relatively uniform with an average of 27–28 °C; the annual sunshine amounts 1918–2390 h and the humidity ranges from 80–84%. During the research period, average rainfall and humidity were the lowest, while temperature and sunshine were the highest recorded over the last 5 years. The rainy season lasts from May till November, and the rest of the year is the dry season with very little rainfall. During the research period the average rainfall, humidity, temperature, and sunshine were 1756 mm, 83.3%, 27.6 °C, and 891 h respectively in the wet season and 336 mm, 75%, 28.8 °C, and 1403 h in the dry season. There was a large difference in rainfall between the wet and dry seasons. The rainfall was very low in March and April and in February there was no rain at all.




Figure 1. Map of the Camau province, showing the sampling locations.

Sample collection and analyses

Litter traps of 1 m² were constructed from nylon nets (1 mm mesh size) and positioned in each *Rhizophora apiculata* stand above the highest tidal water level. In each stand the content of four traps was collected at monthly intervals for 1 year. Total 16 traps were used in the whole. The samples were sorted into leaves, stipules, twigs, and reproductive litter. Dry weight (DW) and ash-free dry weight (AFDW) were analyzed for litter samples. From the leaf litter, total nitrogen (TN), total phosphorus (TP) were analyzed directly after sampling. The dry weight was determined by the drying the samples at 105 °C for 24 h, AFDW was determined by changing of weight before and after burning at 550 °C during 3 h. The Kjeldahl method was used for determining TN, colorimetry for determining TP.

Statistics

The difference of litter fall production between the different mangrove stands during one year were

statistically compared using Repeated Measure in the General Linear Model (GLM – SPSS version 10.0 for windows). Significant differences (at 95% confidence level) were distinguished using Tukey's post-hoc comparison test.

Results

Litter fall

Total litter fall varied from 2.43–3.88 g DW m⁻² day⁻¹ and was significantly highest at the 11 year old stand (Figure 2). A seasonal pattern in the total litter fall was not clear (Figure 3). The oldest stands had a significantly higher proportion of reproductive parts in the litter compared to the younger stands (Figure 2). Twig litter fall was relatively low throughout the year, and was significantly highest in the 7 and 11 year old stands. About 70% of the total litter was leaf litter (decreasing with increasing age) and stipule, twig and reproductive litter covered respectively 8, 12, and 10%. Leaf and stipule litter were significantly




Figure 2. Average biomass (g DW m⁻² year⁻¹) of litter fall components from June 2001 till may 2002 in the Camau province, Vietnam. Different letters indicate significant differences (*p* < 0.05) between the age classes for the total leaf litter (top) and for the different leaf litter components (next to the bars).




Figure 3. Total litter fall (mg DW m⁻²) from June 2001 till May 2002 in Camau province, Vietnam. The dry period has been indicated.

highest in the 11 year old stands, and the lowest leaf litter fall was recorded in the oldest stands. The average ash-free dry weight accounted for 90% of the dry weight of the total litter fall. The highest AFDW was recorded in the 7 and 11 year old stands. Litter fall in the Camau forest is high compared to other mangrove systems (Table 1)

Nutrients in the leaf litter

The nitrogen and phosphorus contents in the leaf litter as well as the total yearly input of N and P via leaf litter were significantly higher in the younger stands (Figure 4). The input of nutrients via the leaf litter showed a clear seasonal pattern in both young and old stands (Figure 5a and b). As




Figure 4. Nitrogen and phosphorus in leaf litter from June till May 2002 in the Camau province, Vietnam (a) nutrients in the leaves (mg g⁻¹ DW), (b) total nutrient input through leaf litter (g m⁻² year⁻¹).

Table 1. Literature data on total litter fall of *Rhizophora* spp. mangrove forests around the world.

Species (type of mangal)	Latitude of study area	Total litter (g DW m ⁻² day ⁻¹)	References
Australia			
<i>Rhizophora apiculata</i>		3.10	Bunt 1982
<i>Rhizophora stylosa</i>		2.55	Duke et al. 1981
Brazil			
<i>Rhizophora</i> spp.	23° S	2.38	Silva et al. 1998
Ecuador			
<i>Rhizophora</i> spp		1.75–2.90	Twilley et al. 1986, 1997
Florida			
<i>Rhizophora</i> spp.	27°41' N	2.21–3.50	Pool et al. 1975
<i>Rhizophora</i> spp.	27°41' N	3.10	Dawes et al. 1999
<i>Rhizophora mangle</i>		1.69–2.10	McKee and Faulkner 2000
Fr. Guyana			
<i>Rhizophora</i> spp		2.38–3.45	Betoulle et al. 2001
Hawai			
<i>Rhizophora mangle</i>		6.90	Cox and Allen 1999
India			
<i>Rhizophora apiculata</i>	6–14° N	1.95	Mall et al. 1991
<i>Rhizophora apiculata</i>		3.21–3.23	Wafar et al. 1997
Kenya			
<i>Rhizophora mucronata</i>		2.70	Slim et al. 1996
<i>Rhizophora mucronata</i>		3.22–4.66	Woitchik et al. 1997
Malaysia			
<i>Rhizophora apiculata</i>	4°50' N	2.70	Gong et al. 1984
<i>Rhizophora</i> spp	3°15' N	4.32	Sasekumar and Loi 1983
Mexico			
<i>Rhizophora</i> spp.		3.40	Day et al. 1987
Papua New Guinea			
<i>Rhizophora</i> spp.	9°31' S	3.91	Leach and Burgin 1985
Tanzania			
<i>Rhizophora mucronata</i>		3.84	Shunula and Whittick 1999
Thailand			
<i>Rhizophora apiculata</i>	8°03' N	2.70	Christensen 1978
<i>Rhizophora apiculata</i>	9°97' N	2.43–3.54	Aksornkoae 1993
<i>Rhizophora apiculata</i>	8°03' N	3.20	Nielsen and Anderson 2003
Tuvalu			
<i>Rhizophora stylosa</i>	7°28' S	2.12	Woodroffe and Moss 1984
Vietnam			
<i>Rhizophora apiculata</i>	8°50' N	2.58–5.15	Clough et al. 2000
<i>Rhizophora apiculata</i>	8°50' N	2.43–3.88	present study

the period between two sampling visits was rather long (1 month), particularly during the wet season leaching could play an important role. Therefore, the wet period data presented here are most probably an under-estimation.

Discussion

Litter fall production

Loi et al. (2002) showed that the soils at the sites in the Tam Giang 3 and Kien Vang enterprises are

fairly homogeneous with respect to pH, salinity, total-N, total-P, and available-P. Therefore, differences in litter fall of *Rhizophora apiculata* stands, as we found, are directly correlated to the age of the trees. This is in line with findings of Clough et al. (2000) who published similar results for leaves in 6–10 year old stands of *Rhizophora apiculata*, and found a further drop in 36 year old stands. This may be due to the more densely leaved canopy in the young stands compared to the more open old stands. Also biological characteristics of the old trees play a role, as these trees produce much woody materials, such as stems, branches




Figure 5. Monthly nutrient input *via* leaf litter (g m^{-2} ; June 2001–May 2002) compared with shrimp yields (kg ha^{-1}) in the Camau province, Vietnam. (a) phosphorus, (b) nitrogen, (c) shrimp yield (kg ha^{-1} ; from Johnston et al. 2000).

and reproductive parts, while the young trees produce more leaves (Hong and San 1993). Further, Clough et al. (2000) concluded that the leaf area index (LAI) declined as stands age in Camau forest. These certainly affect the litter fall. In conclusion, the decline of litter fall with age seems likely to be causally related to the forest ageing indeed.

Nutrient input through leaf litter

Much of the litter from mangroves can be exported *via* creeks to adjacent waters (Robertson et al. 1992; Bunt 1995), and the role of this matter in supporting a rich aquatic and benthic food supply for fisheries is well documented (Mackey and Smail 1996). Nutrients (N, P)

derived from mangrove litter may also boost aquatic primary productivity. Nitrogen and phosphorus levels were considerably higher in the leaf litter of younger stands (7 and 11 years). Mangrove plants in the younger stands are able to take up more nitrogen and phosphorus than those in the older stands. Overall, our results suggest that young stands produce a large quantity of higher quality litter as input to the aquatic system. This may directly or indirectly enhance food availability to the shrimps in younger stands. Indeed, Morrissey et al. (2003) showed that numbers of faunal taxa were higher in younger stands, and numbers of individuals of several taxa were also higher at these sites.

The input of phosphorus and nitrogen peaked during the second part of the wet period and again during the first part of the dry period. This input of nutrients (and organic matter) to the aquatic systems surrounding the mangrove forests is a potential boost to the shrimp production. This can be either directly through the high amount of prime quality food, or indirectly *via* algae, fungi and bacteria. In fact, Johnston et al. (2000) found that shrimp yields peaked between July and October (1996/97) in the Ca Mau province (Figure 5c). We suggest that there may be a direct relationship between the input of nutrients *via* leaf litter and dynamics of shrimp production. The shrimp peak followed the N and P input peaks in the wet period when litter derived material may directly be washed into the ditches and become available to the foodwebs. In the subsequent dry period, the material will tend to stay on the mangrove soil for longer and only gradually becomes available to the ditches.

Our results suggest that litter of younger mangrove stands serves as a significant food source for aquaculture and fisheries in coastal areas. As the Mekong Delta is the most important commercial fishery area in Vietnam, local government and managers should focus on preservation and reconstitution of pristine mangrove system in the region.

Acknowledgements

This research was a part of the MHO-8 project for Environmental Management of Coastal Resources in the Mekong Delta, a co-operative project be-

tween Wageningen University, The Netherlands and Can Tho University, Vietnam. This study was financially supported by NUFFIC.

References

- Aksornkoae S. 1993. Ecology and Management of Mangroves. IUCN. Bangkok, Thailand, pp. 175.
- Alongi D.M. 1990. Abundances of benthic microfauna in relation to outwelling of mangrove detritus in a tropical coastal region. *Marine Ecology Progress Series* 63: 53–63.
- Alongi D.M., Boto K.G. and Tirendi F. 1989. Effect of exported mangrove litter on bacterial productivity and dissolved organic carbon fluxes in adjacent tropical nearshore sediments. *Marine Ecology Progress Series* 56: 133–144.
- Barbier E.B. 2000. Valuing the environment as input: review of applications to mangrove-fishery linkages. *Ecol. Econ.* 35: 47–61.
- Betoulle J.L., Fromard F., Fabre A. and Puig H. 2001. Characterization of litter and its contributions to soil nutrient in a mangrove of French Guiana [French]. *Can. J. Bot.* 79(2): 238–249.
- Bunt J.S. 1982. Studies of Mangrove Litter Fall in Tropical Australia. *Mangrove Ecosystem in Australia: Structure, Function and Management*. Australian National University Press, Canberra, pp. 193–210.
- Bunt J.S. 1995. Continental scale patterns in mangrove litter fall. *Hydrobiologia* 295: 135–140.
- Christensen B. 1978. Biomass and primary production of *Rhizophora apiculata* in a mangrove in southern Thailand. *Aquat. Bot.* 4: 43–52.
- Clough B., Tan D.T., Phuong D.X and Buu D.C. 2000. Canopy leaf area index and litter fall in stands of the Mangrove *Rhizophora apiculata* of different age in the Mekong Delta, Vietnam. *Aquat. Bot.* 66: 311–320.
- Cox E.F., Allen J.A. 1999. Stand structure and productivity of the introduced *Rhizophora mangle* in Hawaii. *Estuaries* 22: 276–284.
- Daniel P.A. and Robertson A.I. 1990. Epibenthos of mangrove waterways and open embayments: community structure and the relationship between exported mangrove detritus and epifaunal standing stocks. *Estuarine Coastal and Shelf Sci.* 31: 599–619.
- Dawes C., Siar K. and Marlett D. 1999. Mangrove structure, litter and macroalgal productivity in a northern-most forest of Florida. *Mangroves and Salt Marshes* 3: 259–267.
- Day J.W.Jr., Conner W.H., Ley-Lou F., day R.H. and Navarro A.M. 1987. The productivity and composition of mangrove forests, Languna de Terminos, Mexico. *Aquat. Bot.* 27: 267–284.
- Department of Agriculture, Forestry, and Fishery 1996. Statistical Data of Agriculture, Forestry, and Fishery 1990–1995, pp. 199.
- Duke N.C., Bunt J.S. and Williams W.T. 1981. Mangrove litterfall in northeastern Australia. I. Annual totals by component in selected species. *Austr. J. Bot.* 29: 547–553.
- Gong W.K., Ong J.E., Wong C.H and Dhanarajan G. 1984. Productivity of mangrove trees and its significance in managed mangrove ecosystems in Malaysia. *Proceedings of the*

- Asian Symposium on Mangrove Environment: research and management. University of Malaysia and UNESCO, pp. 216–225.
- Graaf G.J. and Xuan T.T. 1998. Extensive shrimp farming mangrove clearance and marine fisheries in the southern provinces of Vietnam. *Mangroves and Salt Marshes* 2: 159–166.
- Hong P.N. 1996. Mangrove destruction for shrimp rearing in Minh Hai, Vietnam: Its damage to natural resources and the environment. *SEAFDEC Asian Aquacult.* 18(3): 6–11.
- Hong P.N. 1997. Function of Mangroves forest in Vietnam. Agriculture Publishing house, pp. 217.
- Hong P.N. and San T.H. 1993. Mangroves of Vietnam. IUCN, pp.173.
- Johnston D., Trong N.V, Tien D.V and Xuan T.T. 2000. Shrimp yields and harvest characteristics of mixed shrimp-mangrove forestry farms in southern Vietnam: factors affecting production. *Aquaculture* 188: 263–284.
- Leach G.J. and Burgin S. 1985. Litter production and seasonality of mangroves in Papua New Guinea. *Aquat. Bot.* 23: 215–224.
- Lee S.Y. 1989. Litter production and turnover of the mangrove *Kandelia Candel* (L.) Druce in a Hongkong tidal shrimp pond. *Estuarine Coastal Shelf Sci.* 29: 75–87.
- Loi L.T., Tri L.Q. and Tee J. 2002. Biomass of *Rhizophora apiculata* and soil characteristics in the coastal area of Camau province, Mekong Delta, Vietnam. Selected papers of the workshop on Integrated Management of Coastal Resources in the Mekong Delta, Vietnam, pp. 65–70.
- Mackey A.P. and Smail G. 1996. The decomposition of mangrove litter in a subtropical mangrove forest. *Hydrobiologia* 332: 93–98.
- Mall L.P., Singh V.P. and Garge A. 1991. Study of biomass, litter fall, litter decomposition and soil respiration in monogeneric mangrove and mixed mangrove forest of Andaman Island. *Trop. Ecol.* 32: 144–152.
- Malley D.F. 1978. Degradation of mangrove leaf litter by the tropical sesarimid crab *Chiromantes onychophorum*. *Mar. Biol.* 49: 377–386.
- McKee K.L. and Faulkner P.L. 2000. Restoration of biogeochemical function in mangrove forests. *Restoration Ecol.* 8: 247–259.
- Ministry of Fisheries 2001. Report on the Fisheries Development in the Mekong Delta and its Development Plan for the Period from 2001–2005, pp. 49.
- Morrisey D.J, Skilleter G.A., Ellis J.I., Burns B.R., Kemp C.E. and Burt K. 2003. Differences in benthic fauna and sediment among mangrove (*avicennia marina* var. *australasica*) stands of different ages in New Zealand. *Estuarine, Coastal and Shelf Sci.* 56(3–4): 581–592.
- Nam V.N. and Thuy N.S. 1997. The Biomass of a *Rhizophora apiculata* Plantation in Can Gio district, Ho Chi Minh city. Proceeding of National workshop on the relationship between mangrove reforestation and coastal aquaculture in Vietnam. CRES-ACTMANG, Hanoi.
- Nielsen T. and Andersen F. 2003. Phosphorus dynamics during decomposition of mangrove (*Rhizophora apiculata*) leaves in sediments. *J. Exp. Mar. Biol. Ecol.* 293: 73–88.
- Odom W.E. and Heald E.J. 1972. Trophic analysis of an estuarine mangrove community. *Bull. Mar. Sci.* 22: 671–738.
- Odom W.E. and Heald E.J. 1975. The detritus-based food web of an estuarine mangrove community. In: Cronin L.E. (ed.), *Estuarine Research*. Academic Press, New York, pp. 265–286.
- Pool D.J., Lugo A.E. and Snedaker S.C. 1975. Litter production in mangrove forests of Southern Florida and Puerto Rico. In: Barrett G.W. and Rosenberg R. (ed.), *Stress Effects on Natural Ecosystems*. Wiley, New York, pp. 213–237.
- Robertson A.I., Alongi D.M. and Boto K.G. 1992. Food chains and carbon fluxes. In Roberson A.I. and Alongi D.M. (eds.), *Tropical Mangrove Ecosystems*. American Geophysical Union, Washington, DC, pp. 293–326.
- Robertson A.I. and Daniel P.A. 1989. The influence of crabs on litter processing in high intertidal mangrove forest in tropical Australia. *Oecologia* 78: 191–198.
- Sasekumar A. and Loi J.J. 1983. Litter production in three mangrove forests zones in the Malay Peninsular. *Aquat. Bot.* 17: 283–290.
- Shunula J.P. and Whittick A. 1999. Aspects of litter production in mangroves from Unguja Island, Zanzibar, Tanzania. *Estuarine Coastal Shelf Sci.* 49: 51–54.
- Silva C.A.R., Lacerda L.D., Ovalle A.R. and Rezende C.E. 1998. The dynamics of heavy metals through litter fall and decomposition in a read mangrove forest. *Mangroves and Salt Marshes* 2: 149–157.
- Slim F.J., Gwada P.M., Kodjo M. and Hemminga M.A. 1996. Biomass and litterfall of *Ceriops tagal* and *Rhizophora mucronata* in the mangrove forest of Gazi bay, Kenya. *Mar. Freshwater Res.* 73: 25–38.
- Twiley R.R., Lugo A.E. and Patterson-Zucca C. 1986. Litter production and turnover in basin Mangrove forests in south west Florida. *Ecology* 67: 670–683.
- Twiley R.R., Poro M., Garcia, V.H., Rivera-Monroy V.H., Zambrano R. and Bodero A. 1997. Litter dynamics in riverine mangrove forests in the Guayas River estuary, Ecuador. *Oecologia* 111: 109–122.
- Wafar S., Untawale A.G. and Wafer M. 1997. Litter fall and energy flux in a mangrove ecosystem. *Estuarine Coastal Shelf Sci.* 44: 111–124.
- Woitchik A.F., Ohawa B., Kazungu J.M., Rao R.G., Goeyens L. and Dehairs F. 1997. Nitrogen enrichment during decomposition of mangrove litter in an east African coastal lagoon (Kenya): relative importance of biological nitrogen fixation. *Biogeochemistry* 39: 15–35.
- Woodroffe C.D. and Moss T.J. 1984. Litter fall beneath *Rhizophora stylosa* Griff., Vaitupu, Tuvalu, South Pacific. *Aquat. Bot.* 18: 249–255.