

**Original article****Biodiversity****Habitat and diversity of ectomycorrhizal fungi in forests of South Cameroon**

Nérée A. ONGUENE\* &amp; Thomas W. KUYPER

\*Department of Soils, Water & Atmosphere, Institute of Agricultural Research for Development, P.O. Box. 2123, Yaoundé, Cameroon; email: [nereeo@yahoo.fr](mailto:nereeo@yahoo.fr)

Sub department of Soil Quality, Wageningen University, P.O. Box. 8005, 6700 EC, Wageningen, The Netherlands; email: [thom.kuyper@wur.nl](mailto:thom.kuyper@wur.nl)

**Abstract**

Information is lacking on habitat and diversity of ectomycorrhizal (ECM) fungi of African humid forests. For three years, mushroom excursions were carried out in four sites with contrasted soil and altitude characteristics of South Cameroon, during wet seasons. Collected fungi were described in fresh state and dried exsiccates examined for microscopic description before morpho-anatomical identification. ECM fungi abundantly fruited exclusively in mixed caesalp, monodominant *Gilbertiodendron* and *Uapaca* forest clumps, independently of elevation, rainfall, topography and soil texture. 21 ECM tree species in 11 genera belonging to two families, Ceasalpiniaceae and Phyllanthaceae, dominated ECM forest clumps. More than 100 putative ECM fungal species in 27 genera were identified and assigned to seven families and one super family group. Members of Russulaceae and Amanitaceae recruited the highest number of ECM taxa, followed by species of Boletoids and Cantharellaceae. *Paxillus*, *Clavulina*, *Coltricia*, *Scleroderma*, *Cortinarius* and *Inocybe* genera contained limited species. 12 ECM fungal species were locally edible, including *Lactarius gymnocarpus* and all chanterelles. Though, species richness was very large in a mountainous area, all four sites shared a substantial large number of ECM mushroom species. Low plant diversity but high fungal richness recommends ECM forest clumps for biodiversity sanctuaries.

**Key words.** Ceasalpiniaceae – Clumps – Ectomycorrhiza – Fungi – *Gilbertiodendron* – *Uapaca*

**INTRODUCTION**

Little information exists on the habitat and diversity of ectomycorrhizal (ECM) fungi of African humid forests compared with other biomes. ECM fungi are important below-ground components of rain forests. Where they occurred, they form obligatory symbioses with specific plant taxa. Both symbionts are threatened by various scales of deforestation. Lack of data on the distribution of habitat and diversity of ECM tree and fungal species is likely to jeopardize the regeneration of African ECM forests.

Until recently, it has been thought that ectomycorrhizal fungi were uncommon in tropical humid forests [1]. However, recent mycoinventories revealed that native ECM fungal genera in tropical Africa belong, to a large extent, to the same Basidiomycete taxa as those that provide the fungal partners of temperate ECM plants [2, 3, 4]. Common ECM taxa in the tropics belong to the Amanitaceae, Boletaceae, Cantharellaceae, Russulaceae and Sclerodermataceae [5, 6, 7, 8, 9]. However, on

species level, hardly any ECM fungal species (except when introduced) is shared between tropical and temperate regions [2, 4]. Moreover, many ECM fungal species are poorly described or have their microscopic details poorly illustrated: a reliable key to all African ECM fungal species is lacking and many species remained unnamed. Various species of Amanitaceae, boletes, Cantharellaceae and Russulaceae have been described from West African rain forests [10], with virtually no members of the Cortinariaceae [1]. Mycoinventories in forests of Korup National Park, South-West, Cameroon yielded close to 40 species of suspected mycorrhizal fungi [10]. In forests of South-West Burkina Faso, 27 ECM fungal species were described beneath indigenous *Azelia* and *Uapaca* trees [11]. Using a fragment of the mitochondrial large subunit rRNA gene, more than 100 ECM sporocarps were typed from Ceasalpiniaceae in a Guinean tropical rain forest [4]. It was also concluded that in the same Guinean forest, multi-host ECM fungi predominate and are shared between canopy trees and

seedlings, thereby suggesting the formation of common mycorrhizal networks between trees [12] as found in the rain forest of south Cameroon [13]. Hitherto, the ecology of ECM fungi of African rain forests remains poorly known, with only a small fraction of many thousands ECM formally described. The aim of this study was to provide new records on the habitat and diversity of ECM fungi in forests of South Cameroon.

## MATERIALS AND METHODS

### Location, climate and soils

The study was carried out in portions of humid forests of South Cameroon (Fig. 1), located between Bipindi, Lolodorf and Akom II (2°47' – 3°14'N and 10°24' – 10°51'E), approximately 80 km east of the coastal town of Kribi (2°56'N; 9°54'E) and 40 km west of Ebolowa (2°51'N; 11°11'E). The landscape varied considerably. Altitude ranged from 40 m a.s.l. in the northwestern lowland, to more than 1000 m a.s.l. in the mountainous southwest region (Fig. 1). The original vegetation consisted of dense humid evergreen forest [14]. The climate was hot and humid, with two wet seasons from mid-March to mid-June and mid-August to mid-November. Mean annual rainfall ranged from 1500 to 2500 mm [15]. Mean monthly temperatures varied from 23° to 25°C in Ebolowa (638 m a.s.l.) and 25° to 27.5°C in Kribi (10 m a.s.l.). Relative humidity was generally above 80 % year-round [16]. Geologically, the area consisted of Precambrian metamorphic rocks and old volcanic intrusions [17]. Soil characteristics changed considerably with elevation (Table 1).

Within the study area, four sites were selected in Ebimimbang (low elevation; sandy and near neutral soils), Ebom (mid-elevation; high clayey and acid soils), Nyangong and Bityili (high elevation; very high clayey and strongly acid soils). In each site, one 1-ha permanent sampling

plot was selected in near primary forest. All tree species at all growth stages were enumerated and diameter at breast height measured. As trekking towards these plots, observations were also made in other ecosystems including small forest clumps, secondary forests, cocoa (*Theobroma cacao*) plantations, and food crop fields, bush fallow of *Chromolaena odorata*, farmers' trails and homegardens.

### Mushroom collection, description and identification

Mushroom excursions were undertaken during the wet seasons, one week per month, with the help of local assistants. Collected specimens were macroscopically described in fresh state before twilight. After description, mushrooms were dried for 2 – 3 days at about 40°C using a locally made drier. Dried mushrooms were temporarily preserved in a local herbarium in Kribi before being sent to Wageningen, the Netherlands for microscopic examination and identification.

Several provisional keys were elaborated based on morphological and anatomical characters of sporocarps [18]. Voucher specimens were kept at the National herbarium of the Netherlands (Leiden Branch) and some duplicates were preserved at the agricultural research station in Kribi, Cameroon.

Several manuals were used for the identification of most species such as “La Flore Iconographique des Champignons du Congo”, “La Flore illustrée des Champignons d’Afrique Centrale” (Illustrated manual edited by the Belgian National botanical garden), the genus *Lactarius* [19, 20, 21, 22], *Russula* by Buyck B [23, 24, 25] and the genus *Amanita* [26, 27, 28, 29]. When some taxa were new, they were attributed a new name. Their description will be provided in subsequent publications.

Table 1: Localization, elevation, rainfall and soil physico-chemical characteristics of the four research sites

Research sites	Ebimimbang	Ebom	Nyangong	Bityili
Localization	3°02.67'N; 10°28.25'E	3°04.73'N; 10°41.24'E	2°58.11'N; 10°45.18'E	2°56.07'N; 10°49.55'E
Altitude (m.a.s.l)	0 – 350	350 – 500	500 – 800	>800
Rainfall (mm)	1556	1987	1677	1800
Soil texture	Sandy	Highly clayey	Highly clayey	Very highly clayey
pH	5 – 6	4 – 5	3 – 4	3 - 4
Carbon (%)	1.69	2.26	2.21	5.7
Nitrogen (%)	0.15	0.18	0.19	0.36
Phosphorus(µm/ml soil)	0.01	0.005	0.002	0.000




Figure 1: Localization of the villages in south Cameroon where fungal samples were collected for three years

## RESULTS

All species of putative ectomycorrhizal mushrooms were found mainly in three types of forests: mixed oligodominant ceasalp, monodominant *Gilbertiodendron* and *Uapaca* forest clumps. ECM forest clumps varying in physiognomy and aspect, recruited at least 10 to more than 100 ECM tree specimens of the same genus or species, simultaneously occurring on a single spot, at different growth stages. ECM forest clumps had low cover values of shrub species and herb without moss. They slightly varied in canopy closure. In general, ECM forest clumps formed closed-canopy dominated by ceasalps and/or *Uapaca* trees. But the commonest ECM tree species differed with site: mixed ceasalp species in Ebimimbang, *Gilbertiodendron* species in Ebom, *Uapaca* species in Nyangong and a mixture of all ECM tree species in Bityili.

Twenty-one ECM tree species belonging to 11 genera and two families, Ceasalpiniaceae and Phyllanthaceae, were recorded. Most ceasalp species in tribe Amherstieae of the

Ceasalpiniaceae were collectively called “Ekop” by forest prospectors and local populations (Table 2).

Elevation, aspect and landform of ECM forest clumps largely varied with site. Soil humidity was continuously wet during both wet and dry seasons in all sites but Ebimimbang where sandy soils rapidly dried up during the long dry season (mid-November to mid-march). Soil nutrient regimes differed in a relatively narrow range but fell almost exclusively within the range of poor soils (Table 1). Floors of ECM forest clumps were thin, with low litter and little organic matter accumulation at all sites. The most common natural disturbance was wind fall, to a very low extent, though. Ectomycorrhizal habitats were unevenly distributed at the four sites. They were abundant in high (Bityili) and low elevation (Ebimimbang), intermediate in Nyangong and lowest in Ebom. Monodominant clumps of *Gilbertiodendron* tree species were observed only at the Ebom site, in swampy areas, close to small and large rivers, where they stood long periods of inundation.

**Table 2:** The ectomycorrhizal species ranged by type of forest habitat

Habitat types	Family	Species	Pilot name
Mixed ceasalp clumps	Caesalpiniaceae (Tribe Amherstieae)	<i>Anthonotha fragans</i>	Enak
		<i>A.macrophylla</i>	Enak
		<i>Berlinia bracteosa</i>	Ebiara
		<i>Berlinia confusa</i>	Ebiara
		<i>Brachystegia cynometrioides</i>	Ekop naga
		<i>Brachystegia euricoma</i>	Ekop evene
		<i>Brachystegia zenkeri</i>	Ekop gombe
		<i>Didelotia africana</i>	Ekop rouge
		<i>Didelotia letouzeyi</i>	Ekop
		<i>Julbernadia seretii</i>	Ekop blanc
		<i>Monopetalanthus letestui</i>	Ekop mayo
		<i>Monopetalanthus microphyllus</i>	Ekop mayo ngang
		<i>Paraberlinia bifoliolata</i>	
		<i>Tetraberlinia bifoliolata</i>	Ekop beli
		<i>Touabouate brevipaniculata</i>	Ekop ribi Ekop zing
<i>Gilbertiodendron</i> clumps	Caesalpiniaceae	<i>Gilbertiodendron brachystegioides</i>	Abem
		<i>Gilbertiodendron dewevrei</i>	Abem
<i>Uapaca</i> clumps	Phyllanthaceae	<i>Uapaca acuminata</i>	Rikio
		<i>Uapaca buchholizianum</i>	Rikio
		<i>Uapaca guineensis</i>	Rikio
		<i>Uapaca vanhoutei</i>	Rikio

Notes: Pilot names of “Ekop” were taken in Letouzey and Mouranche (1952).

Most ectomycorrhizal habitats were of small to medium size, with one dominating ECM tree species or one genus. But on the Bityili hill, ECM tree species depicted a distinctive distribution pattern: the bottom abrupt hill was mostly overgrown by *Uapaca* tree species, followed in the middle by *Gilbertiodendron* trees and a mixture of caesalp tree species towards hill top.

Abundance of ectomycorrhizal tree species in 1-ha plot varied in the range of 20 to 80 % with site. It was lowest in the youngest stands of Ebom (19% basal area) and the highest in a nearly pristine rain forest of Bityili (79% basal area). However, it reached almost 100% in monodominant *Gilbertiodendron* clumps. ECM habitats occurred throughout the landscape: along roadsides, on hills, on flat terrain, in swamps, in riverine spots, far in the forests or close to villages, even within agricultural farm holdings. They were not cleared for shifting cultivation, though. Rather, local people claimed that soils under ECM forest patches were very poor and used them only as a collecting ground for ECM mushrooming during wet periods.

In the study area, more than 1000 presumed ectomycorrhizal sporocarps were sampled during three years of mushroom excursions, exclusively in ECM forest clumps during both wet seasons. ECM mushrooms were also observed during the dry seasons, to a much lower extent, though. In the three ECM forest types, ECM fungal taxa were very diverse. More than 100 putative ECM fungal species in 27 genera were identified and assigned to seven families: Amanitaceae, Cantharellaceae, Clavulinaceae, Coltriciaceae, Cortinariaceae, Russulaceae, Sclerodematocae and one super family group, Boletoids (Table 3).

Among ectomycorrhizal fungal species identified, *Amanita* and *Russula* species were dominant, with 34 and 32 species, respectively, followed by 28 species in 15 genera of the Boletoids, 15 species in three genera of the Cantharellaceae and 12 *Lactarius* species. Both *Russula* and *Lactarius* recruited the highest number of ECM taxa. *Paxillus* (two species), *Clavulina vanderstii*, *Coltricia* (two species), *Scleroderma* (two species), *Cortinarius* (three species) and *Inocybe* (six species) genera contained only limited number of species (Table 3).

**Table 3:** List of putative ectomycorrhizal mushroom species by family, genus and species in humid forests of South Cameroon

<p><b>Amanitaceae</b>  <i>Amanita afzeliae</i> nov.  <i>A. albidodisca</i> nov.  <i>A. albopulverulenta</i> nov.  <i>A. afrobispora</i> nov.  <i>A. afroflavescens</i> nov.  <i>A. afro rubescens</i> nov.  <i>A. afrovaginata</i> nov.  <i>A. annulatovaginata</i> Beeli  <i>A. aureofloccosa</i> Bas  <i>A. bingensis</i> Beeli  <i>A. brachystegiae</i> nov.  <i>A. calopus</i> Beeli  <i>A. crassiconus</i> Bas  <i>A. elegans</i> Beeli  <i>A. flavovirens</i> nov.  <i>A. fulvosquamulosa</i> Beeli  <i>A. gigavolvata</i> nov.  <i>A. griseofarinosa</i> Hongo  <i>A. lanosa</i> Beeli  <i>A. leucoagaricioides</i> nov.  <i>A. luteoflava</i> Beeli  <i>A. monopetalanthi</i> nov.  <i>A. nigropyramis</i> nov.  <i>A. pseudoafroalba</i> nov.  <i>A. pseudolanosa</i> nov.  <i>A. pustulata</i> nov.  <i>A. roseocinnamomea</i> nov.  <i>A. rubescens</i> (Pers.:Fr) G  <i>A. strobilaceovolvata</i> Beeli  <i>A. subviscosa</i> Beeli  <i>A. sulphurea</i> nov.  <i>A. virella</i> Beeli</p>	<p><b>Boletaceae</b>  <i>Afroboletus luteolus</i> (Heinem) P. &amp; Y.  <i>Boletus boletiformis</i> nov.  <i>B. macrocystis</i> nov.  <i>B. nyangongensis</i> nov.  <i>B. pustulatus</i> Beeli  <i>B. suspinulosus</i> nov.  <i>Boletellus sulcatipes</i> Hein.&amp; Goo.  <i>Chalciporus clypeatus</i> nov.  <i>Gyrodon aberrans</i> nov.  <i>Gyroporus microsporus</i> (Sing.&amp; Grinl.)  Heinem &amp; Rammeloo  <i>Leccinum excedens</i> nov.  <i>Paxillus brunneotomentosus</i> nov.  <i>Paxillus camerunensis</i> nov.  <i>Phlebopus braunii</i> (Bres.) Heinem  <i>P. silvaticus</i> Heinem  <i>Phlebopus</i> sp. nov  <i>Phylloporus depressus</i> Heinem  <i>Pulveroboletus aberrans</i> Heinem  <i>P. viridis</i> Heinem  <i>Rubinoletus luteopurpureus</i> (Beeli)  <i>Strobilomyces echinatus</i> Beeli  <i>S. luteolus</i> Heinem  <i>S. strobilaceus</i> (Berk.)  <i>S. velutipes</i> Corner  <i>Tubosaeta alveolata</i> Heinem  <i>T. brunneosetosa</i> (Singer) E. Horak  <i>T. goosseniae</i> nov.  <i>Tylopus violaceus</i> nov.  <b>Clavulinaceae</b>  <i>Clavulina vanderstii</i> nov.  <b>Sclerodermataceae</b>  <i>Scleroderma sinnamariense</i> Mont  <i>S. roseacarneum</i>  <b>Coltriciaceae</b>  <i>Coltricia spathulata</i> (Hooker) Murill  <i>C. pyrophila</i> (Wakef) Ryvarden</p>	<p><b>Russulaceae</b>  <i>Russula afronigrigans</i> Buyck  <i>R. albospissa</i> Buyck  <i>R. annulata</i> Heim  <i>R. annulatobadia</i> Beeli  <i>R. apсила</i> Buyck  <i>R. intricata</i> Lizoii  <i>R. areolata</i> Buyck  <i>R. aurantiofloccosa</i> Buyck  <i>R. camerunensis</i> nov.  <i>R. cellulata</i> Buyck  <i>R. chrysotricha</i> nov.  <i>R. declinata</i> Buyck  <i>R. diffusa</i> var. <i>fissurans</i>  <i>R. discopus</i> Heim  <i>R. echnosperma</i> R.Heim &amp; Gilles  <i>R. fulvoochrascens</i> Buyck  <i>R. heliochroma</i> Heim  <i>R. intricate</i> Buyck  <i>R. kivuensis</i> Buyck  <i>R. lamprocystidia</i> (Nakasone)  <i>R. liberiensis</i> Buyck  <i>R. macrocystis</i> nov.  <i>R. mimetic</i> nov.  <i>R. pausiaca</i> Buyck  <i>R. pseudocarmesina</i> Buyck  <i>R. pseudopurpurea</i> Buyck  <i>R. pseudostriatoviridis</i> Buyck  <i>R. striatoviridis</i> Buyck  <i>R. testaceoaurantiaca</i> Beeli  <i>R. velutina</i> (DC per Pers.:Fr)P  <i>R. yaeneroensis</i> Buyck</p>
<p><b>Inocybe</b>  <i>Inocybe afronodulosa</i> nov.  <i>I. afrostellata</i> nov.  <i>I. korupensis</i> nov.  <i>I. bipindensis</i> nov.  <i>I. perpusilla</i> Velen  <i>I. zingii</i> nov.  <b>Cortinarius</b>  <i>Cortinarius afroconicus</i> nov.  <i>C. diobensis</i> nov.  <i>C. ionopygmaeus</i> nov.</p>	<p><b>Cantharellaceae</b>  <i>Gomphus brunneus</i> (Heinem)  <i>G. clavatus</i> (Pers.) Gray  <i>Craterullus crispus</i> Fr.  <i>C. cornucopioides</i> Persoon  <i>Cantharellus camerunensis</i> nov.  <i>C. cibarius</i> var. <i>roseocanus</i> Arora &amp; Dunham  <i>C. congolensis</i> Beeli  <i>C. dichrous</i> (Fr.) Bres  <i>C. densifolius</i> Heinem  <i>C. isabellinus</i> Heinem</p>	<p><b>Lactarius</b>  <i>Lactarius acutus</i> Heim  <i>L. annalatoangustifolius</i> nov.  <i>L. claricolor</i> nov.  <i>L. denigrigans</i> Ver.&amp; Kar.  <i>L. densifolius</i> Ver.&amp; Kar.  <i>L. gymnocarpus</i> Heim  <i>L. kivuensis</i> deWitte  <i>L. medusa</i> Verbeken  <i>L. pumilus</i> Verbeken  <i>L. pulcrispermus</i> nov.  <i>L. sesemotani</i> Beeli</p>

	<i>C. luteopunctatus</i> (Beeli) Heinem <i>C. miniatescens</i> Heinem <i>C. microcibarius</i> Heinem <i>C. pseudocibarius</i> Hennings <i>C. rufopunctatus</i> (Beeli) Heinem	<i>L. undulates</i> Verbeke
--	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------

All four sites shared a substantial number of similar ECM fungal species. But, species richness was the highest at the Bityili Mountain.

Twelve species of ECM fungi were edible by native populations, including *Lactarius gymnocarpus*, locally called 'Nyumelane' and most *Cantherellus* species but *Craterellus cornucopioides*. Several folk names of Bantu and Bagyeli pygmy ethnic groups were used for chanterelles, notably, "Bingöbiñdoñg, Mebeum, Nyavem, Ötoye, Otyetya" and "Ambini, Ampembo, Nyanbembo", respectively.

## DISCUSSION

In the Bipindi-Lolodorf-Akom II area, cameroonian ectomycorrhizal mushrooms abundantly fruited exclusively in three types of habitats, viz. mixed oligodominant caesalp, monodominant *Gilbertiodendron* and *Uapaca* forest clumps. Tree-forming ECM clumps belonged only to two families, Ceasalpinaceae and Phyllanthaceae. Ten genera were ceasalp tree-forming monodominant clumps of *Gilbertiodendron* species or mixed caesalp oligo-dominant stands. *Uapaca* tree species chiefly formed small monodominant stands on riversides. Hence, the presence of such tree species and stands should be made known to all forest stakeholders for their ecological specificity and relevance in the management of African humid forests.

Similar to other sites in the Congo basin, clumping of ceasalp and *Uapaca* tree species with associated ECM fungi is commonplace in humid forests of Cameroon. In Korup National Park (SW Cameroon), ECM fungi were recorded in large groves dominated by *Microberlinia* and *Tetraberlinia* species, on sandy soils with two distinct seasons [30, 31]. In the Dja Faunal World Heritage, in South-East, Cameroun, large forest patches dominated by *Gilbertiodendron dewevrei* developed on swampy areas (14; N.A. Onguene and A. M. Bâ, pers.obs.). In our research area in South Cameroon, the distribution of small and medium-size ECM forest clumps was independent of elevation, rainfall, topography and soil texture: they occurred on both sandy and clayey soils. Our results do not agree with earlier observations of

edaphic specialization of ECM trees owing to the competitive ability of ECM fungi on poor soils [30, 32, 33]. The question remains on how ECM clumpiness originated in tropical humid forests and their locally uneven distribution. It was assumed that the gregarious habit of ECM trees could be due to limited seed dispersal of both symbiotic partners [34]. Since both ECM tree and ECM fungus do not possess the capacity to reproduce independently of the other symbiotic partner, establishment of the ECM symbiosis might be a rare chance event [42], which then serve as focal points for subsequent establishment of successive ECM trees, confirming the theory of regeneration niche to explain co-dominance of ECM trees and fungi.

If this hypothesis holds, then it implies that ectomycorrhizal recovery after large scale disturbance (be it natural or manmade) could be very low, requiring special management of native ECM stands of African humid forests. Consequently, ECM forest clumps need a special conservation statute; so far, little information exists on the ecological factors that maintain the current functioning of ECM forest clumps and in general of tropical rain forests.

Ectomycorrhizal forest habitats were the only sites where more than a hundred ECM mushrooms species in 27 genera and eight families were collected and identified. It is the first time in Africa that such a highly diverse and abundant putative ECM mushroom collection is reported from the same area. In the Pakaraima Mountains of Guyana, 75 morpho-species of putatively ECM fungi were identified from discrete groves formed by *Dycimbe* and *Aldina* species [35]. In the Miombo savanna and humid forest of West Africa, only five families of ECM mushrooms were recorded [4, 11, 24]. The same high number of members of the Russulaceae (44 species) from this study supports existing data [1]. Species-rich genera included *Amanita*, *Russula*, members of Boletales and Cantheralles. Less frequent ECM fungal species, *Clavulina*, *Coltricia* and *Scleroderma* were also collected for the first time as well as rare ECM mushroom species, viz. two, three and six species of *Paxillus*, *Cortinarius* and

*Inocybe*, respectively (Table 3), confirming earlier observations elsewhere in West Africa, the Neotropics and the Laojun Mountain region, southwestern China [4, 35, 37]. Lack of frequent inventories and experienced mycologists in Africa could explain the poor accounts for tropical ECM mushroom diversity and their habitat. Only 546 records of wild macrofungi have been made from South Saharan countries [2]. In Benin, one species of *Craterellus*, *Hebeloma*, *Inocybe*, of *Russula* and eight *Lactarius* species were recorded [6]. Five out of six macrofungi in Burkina Faso were identified as *Amanita hemibapha*, *Cantharellus pseudofriessi*, *Lactarius gymnocarpus*, *Phaeogyroporus sudanicus* and *Tubosaeta brunneosestosa* [11, 37] as well as an unidentified *Calvatia* [38]. No less than 40 ECM fungi have been recorded from the republic of Congo including 15 *Cantharellus* species and dozens of *Russula* and *Lactarius*. The highest number of *Lactarius* species (19 species) has been thus recorded in Tanzania. Other African countries are less prolific in ECM fungi [2].

Ectomycorrhizal forest communities could contribute not only to the preservation of forest refuges of South Cameroon with endemic species, but also to the protection of the biodiversity hot spots of Africa's rain forests. Maintenance of such ECM forest communities with their associated edible mushrooms such as chanterelles also helps to preserve valuable source of alternative protein-rich diet for local communities who depend on non-timber forest products [39, 40], in addition to creating new ecological jobs of mushroom collectors. Thus, ECM forest clumps should be protected as biodiversity sanctuaries, owing to lack of knowledge on their regeneration requirements.

Thirteen ectomycorrhizal genera are worldwide edible including, *Amanita*, *Boletus*, *Cantharellus*, *Cortinarius*, *Laccaria*, *Lactarius*, *Leccinum*, *Morchella*, *Russula*, *Suillus*, *Terfezia*, *Tricholoma*, *Tuber* [2]. In Africa, more than 20 edible *Cantharellus* species were described in dozen countries [2, 5] with various tastes, though [37]. 1998). In our research sites, 12 *Cantharellus* species were edible by Bantou and Bagyeli people but *Craterellus cornucopioides*. Local tribes often seek them avidly, had a very high esteem and gave them no less than eight folk names). Yet, chanterelles can not be cultivated but grow vividly only in the vicinity of ECM tree species in ECM forest clumps. Consequently, growing trees that

“produce” chanterelles could serve as a way forward for implicating local populations in sustainable management of remnants of African rain forests. Rejuvenation of degraded forests could be done by mycorrhizal inoculation of tree seedlings with chanterelles.

Cameroonian ectomycorrhizal fungi abundantly fruited exclusively in three types of habitats: mixed oligodominant caesalip, monodominant *Gilbertiodendron* and *Uapaca* forest clumps, independently of elevation, rainfall, topography and soil texture. Though poor in plant diversity, these habitats recruited abundant and various ECM fungal species on which native ECM tree species depend for survival and some others such as chanterelles and *Lactarius gymnocarpus* serve as a ready source of proteins for local people during harsh periods. Therefore, forest clumps should be conserved as biodiversity sanctuaries and the presence of ECM fungi could serve as indicators for sustainable management of African humid forests.

#### Acknowledgements

Financial support by the NWO (Priority Programme – Biodiversity in Disturbed Ecosystems) is acknowledged as well as the logistics provided by the Tropenbos Cameroon Programme and the Institute of Agricultural Research for Development (IRAD). Field assistance was provided by Jean Baptiste Mva, Edouard Nsomoto, Serge Aba'a Aba'a, Roger Eyene and Thomas Mba. The technical assistance of Véronique Anaba spouse Onguene during the course of the project was very much appreciated. Lijbert Brussaard and Ngome E. Francis commented on an earlier version of the manuscript. Thanks to unknown reviewers.

#### References

1. Buyck B., Thoen D. and Watling R. 1996. Ectomycorrhizal fungi of the Guineo-Congo Region. Proc R Soc Eding Sect B 104 (3): 313-333
2. Boa E. 2006. Champignons comestibles sauvages – Vue d'ensemble sur leurs utilisations et leur importance pour les populations. FAO, Rome
3. Brundrett M.C., Bougher N., Dell B., Grove T. and Malajczuk N. 1996. Working with mycorrhizas in forestry and agriculture. ACIAR Monograph 32, Canberra

4. Riviere T., Diedhiou A.G., Diabate M. et al. 2007. Genetic diversity of ectomycorrhizal Basidiomycetes from African and Indian tropical rain forests. *Mycorrhiza*. 17:415-428
5. Corner E.J.H. 1966. A monograph of Cantharelloid fungi. *Annals of Botany Memoirs* 2 : 1-155
6. De Kessel A., Codjia J.T.C, and Yorou N.S. 2002. Guide des champignons comestibles du Bénin. Cotonou, République du Bénin, Jardin Botanique National de Belgique et Centre International d'Ecodéveloppement Intégré (CECODI). Impr.Coco-Mutimedia. 275 pp
7. Natarajan K., Senthilarasu G., Kumaresan V., and Riviere T. 2005. Diversity in ectomycorrhizal fungi of a dipterocarp forest in Western Ghats. *Curr Sci* 88 (12): 1893-1895
8. Smits W.T.M. 1992. Mycorrhizal studies in dipterocarp forests in Indonesia. In : Read DJ, Lewis D.H., Fitter A.H., Alexander I.J. (eds) *Mycorrhizas in ecosystems*, Cambridge, pp283-292
9. Watling R., and Lee S.S. 1995. Ectomycorrhizal fungi associated with members of the Dipterocarpaceae in Peninsular Malaysia. *J Trop For Sci*. 7:657-669
10. Watling R. 1993. Similarities and differences in the Macromycetes floras and biological strategies in selected tropical areas. In Isaac S, Frankland JC, Watling R, Whallen AJJ (eds) *Aspects of tropical mycology*. Cambridge University Press, Cambridge
11. Sanon K.B., Bâ A.M., and Dexheimer J. 1997. Mycorrhizal status of some fungi beneath indigenous trees in Burkina Faso. *For. Ecol. Manag.* 98:61-69
12. Diédhiou A.G., Selosse M.A., Galiana A. et al. 2010. Multi-host ectomycorrhizal fungi are predominant in a Guinean tropical rainforests and shared between canopy and seedlings. *Environ. Microbiol.* 12(8): 2219-2232
13. Onguene N.A. and Kuyper Th.W. 2002. Importance of the ectomycorrhizal network for seedling survival and ectomycorrhiza formation in rain forests of south Cameroon. *Mycorrhiza*. 12: 13-17
14. Letouzey R. 1968. Etude phytogéographique du Cameroun. Ed. P. Le Chevalier, Paris
15. Waterloo M.J., Ntonga J.C., Dolman A.J., and Ayangma A.B. 2000. Impact of shifting cultivation and selective logging on the hydrology and erosion of rain forest land in South Cameroon. 2<sup>nd</sup> revised edition. *Tropenbos Cameroon Documents* 1: 1-84
16. Olivry J.C. 1986. Fleuves et rivières du Cameroun. Collection Monographes hydrologiques d'ORSTOM 9, Paris, France
17. Franqueville A. 1973. Atlas régional sud-ouest. République Unie du Cameroun
18. Onguene N.A. 2000. Diversity and dynamics of mycorrhizal associations in tropical rain forests with different disturbance regimes in south Cameroon. *Tropenbos Cameroon Ser.3*:1-167
19. Verbeken A. 1995. Studies in tropical African *Lactarius* species 1. *Lactarius gymnocarpus* R. Heim ex Singer and allied species. *Mycotaxon* 55:515-542
20. Verbeken A. 1996. New taxa of *Lactarius* (Russulaceae) in tropical Africa. *Bull. Jard. Bot. Nat. Belg* 65:197-213
21. Verbeken A., Walley R., Sharp C., and Buyck B. 2000. Studies in tropical African *Lactarius* species 9. Records from Zimbabwe. *Syst. Geogr. PI* 70: 181-215
22. Verbeken A., and Walley R. 1999. Studies in tropical African *Lactarius* species 7. A synopsis of the section and review on the edible species. *Belg. Journal Bot.* 132 (2): 175-184 (2000)
23. Buyck B. 1993. *Russula* I (Russulaceae). In: Rammeloo J. 1 Heinemann P (eds.). *Fl. III. Champignons d'Afrique Centrale* 15: 337-407
24. Buyck B. 1994. Ubwoba: les champignons comestibles de l'Ouest du Burundi. *Adm. Gén. Coop. Dév. Publi. Agric. Bruxelles* 34, 123p
25. Buyck B. 1997. *Russula* III (Russulaceae). In: Rammeloo J. 1 Heinemann P (eds.). *Fl. III. Champignons d'Afrique Centrale* 17: 545-598
26. Bas C. 1969. Morphology and subdivision of *Amanita* and a monograph of its section *Lepidella*. *Persoonia* 5 (4): 285-579
27. Beeli M. 1935. *Amanita, Volvaria*. *Fl. Iconogr. Champignons Congo* 1 :1-28, pl 1-4
28. Härkönen M., Saarimäki T and Mwasumbi L. 1994. Tanzania mushrooms and their uses 4. Some reddish edible and poisonous *Amanita* species. *Karstenia* 34: 47-60
29. Härkönen M., Saarimäki T., and Mwasumbi L. 1995. Edible mushrooms of Tanzania. *Karstenia* 35 (supplement): 1-92
30. Newbery D.M., Alexander D.W., Thomas D.W., and Gartlan J.S. 1988. Ectomycorrhizal rain forest legumes and soil phosphorus in Korup National Park, Cameroon. *New Phytol.* 109:433-550



31. Newbery D.M., Alexander D.W., and Rother J.A. 1997. Phosphorus dynamics in lowland rain forest: the influence of ectomycorrhizal trees. *Ecol. Mon.* 67:367-409
32. Peh K.S.H., Sonke B., Lloyd J, Quesada C.A., and Lewis S.L. 2011. Soil does not explain monodominance in a central African tropical forest. *PLoS one.* 6(2): 1-9
33. Singer R., and Araujo I. 1986. Litter decomposition and ectomycorrhizal Basidiomycetes in an igapo forest. *Pl. Syst. Evol.* 153: 107-117.
34. Richards P.W. 1966. The tropical rain forest: an ecological study. 2<sup>nd</sup> edition. Cambridge University Press, Cambridge
35. Henkel T.W., Terborgh J., and Vilgalys R.J. 2002. Ectomycorrhizal fungi and their leguminous hosts in the Pakaraima Mountains of Guyana. *Mycological Research* 106, 515-531
36. Ying Zhang., De Qun Zhou., Qi Zhao., Tong X.Z., and Hyde K.D. 2010. Diversity and ecological distribution of macrofungi in the Laojun Mountain region, southwestern China. *Biodivers Conserv.* 19:3545-3563
37. Rammeloo J., and Walley R. 1993. The edible fungi of Africa south of Biafra: a literature survey. *Scripta Bot. Belg.* 5:1-62
38. Walley R., and Rammeloo J. 1994. The poisonous and useful fungi of Africa south of the Sahara. *Scripta Botanica Belgica* 10:1-56
39. Malaisse F. 1967. Se nourrir en forêt claire africaine – Approche écologique et nutritionnelle. Les Presses Agronomiques de Gembloux, Gembloux, Belgique
40. Van Dijk H., Onguene N.A and Kuyper T.H 2003. Knowledge and utilization of edible mushrooms by local populations of the rain forest of South Cameroon. *Ambio.* 32 (1): 19-23
41. Letouzey R., and Mouranche R. 1952. Ekop du Cameroun. Publication 4. Centre Technique Forestier Tropical. NogentS/Marne, France
42. Janos D.P. 1996. Mycorrhizas, succession and rehabilitation of deforested lands in the humid tropics. In: Frankland JC, Magan N, Gadd GM (eds), *Fungin in environmental change*, pp129-161, Cambridge University Press, Cambridge