

---

# Reducing greenhouse gas emissions from dairy farming via feeding & breeding

---


**Corina van Middelaar**

Paul Berentsen, Jan Dijkstra & Imke de Boer



# GHG emissions from dairy farming


Dairy sector: 30% of global GHG emissions by livestock



Based on Van Middelaar et al. (2011) and Gerber et al. (2013).

# How to assess net benefit of a strategy?

## Integrated modelling at chain level



$$CO_2\text{-e: } 1 \times CO_2 + 25 \times CH_4 + 298 \times N_2O$$

---

# Reducing greenhouse gas emissions via feeding?

---

Which strategy is most cost-effective?

## Aim 1

evaluate cost-effectiveness of three feeding strategies to reduce enteric CH<sub>4</sub> in dairy COWS

using integrated modelling

---

# Feeding strategies explored

---

## Nitrate supplementation

- 1% of DM intake; 75% nitrate

## Extruded linseed supplementation


- 1 summer; 2 winter (kg/cow/d); 56% linseed

## Less mature grass (silage)

- grazing: 1400 - 1700 kg DM/ha
- harvesting: 3000 - 3500 kg DM/ha

# Method - feeding

*Average farm: maximize labour income*



- 45 ha
- 603 tonnes milk
- 76 cows; 49 young stock
- milk yield cow: 7968 kg/yr

---

## Method - feeding

---

*Average farm*



*Introduction feeding strategy*




*Optimize farm plan: maximize labour income*



*Difference income : difference GHGs*

# Results feeding strategies



1. Nitrate
2. Linseed
3. Younger grass (silage)



---

# Reducing greenhouse gas emissions via breeding?

---

## Increasing annual milk yield per cow

- Fewer animals to produce same amount of milk
- Dilution of GHGs from maintenance

## Improving longevity

- Fewer female replacements needed

---

# Reducing greenhouse gas emissions via breeding?

---

Which trait offers most potential?


## Aim 2

determine impact of increase of one  $\sigma_g$  in  
milk yield and longevity

using integrated modelling

# Method - breeding

## *Farm 2020: maximize labour income*



- 85 ha; all manure used on farm
- 168 cows; 100 young stock
- milk yield cow: 8758 kg/yr
- Replacement rate: 27%

---

## Method - breeding

---

*Future farm*



*Increase  $\sigma_g$  of trait*



*Optimize farm plan: maximize labour income*



*Impact on GHG emissions*

*$\sigma_g$  milk = 687 kg/y &  $\sigma_g$  longevity = 270 d*

---

# Results breeding strategies

---

## **GHG emissions** (kg CO<sub>2</sub>-eq/ton FPCM)

Reference	882
Milk yield	-36
Longevity	-32

## **Economic value** (EUR per cow/year)

Milk yield	122
Longevity	82

Van Middelaar et al. 2014



---

# Conclusions

---

- ✓ **Feeding & Breeding** offer potential to reduce GHG emissions at chain level
- ✓ **Feeding:** Nitrate largest reduction in emissions  
Reducing grass maturity most cost-effective
- ✓ **Breeding:** Milk yield more important than longevity  
Importance longevity increases with focus on GHG emissions

---

# PhD course

---

## **Environmental impact assessment of livestock systems**

9-13 February 2015

Animal Production Systems, Wageningen University, the Netherlands

Imke de Boer, Wageningen University

Piere Gerber, FAO

Christel Cederberg, SIK

Oene Oenema, Wageningen University

# GHG emissions method-1

kg CO<sub>2</sub>-e/t FPCM\*

	Ref	
Animal emissions		
Enteric CH <sub>4</sub>	445	50% enteric CH <sub>4</sub>
Manure	118	
On-farm feed		
Grass	67	
Maize	37	
Farm inputs		
Maize silage	24	
Concentrates	118	
Synthetic fertilizer	51	
Other	23	
Total	882	Lower than literature

\* FPCM = Fat-and-protein corrected milk



# GHG emissions method-1

kg CO<sub>2</sub>-e/t FPCM\*

	Ref	Milk Yield	
Animal emissions			
Enteric CH <sub>4</sub>	445	-10	Dilution
Manure	118	-5	
On-farm feed			
Grass	67	+6	P application rates
Maize	37	-14	
Farm inputs			
Maize silage	24	+18	Maize cheaper concentrates
Concentrates	118	-28	
Synthetic fertilizer	51	-2	
Other	23	-1	
Total	882	-36	

\* FPCM = Fat-and-protein corrected milk