

## Castration of piglets under CO<sub>2</sub>-gas anaesthesia

M. A. Gerritzen<sup>†</sup>, M. Kluivers-Poodt, H. G. M. Reimert, V. Hindle and E. Lambooij

Wageningen University and Research Centre, Animal Sciences Group, P.O. Box 65, 8200, AB Lelystad, The Netherlands

(Received 26 February 2008; Accepted 19 June 2008; First published online 8 July 2008)

*It has become common practice in pig fattening production systems to castrate young boar piglets without the use of anaesthesia. In this study, we examined whether or not CO<sub>2</sub> gas is capable of inducing an acceptable anaesthetic state during which castration can be performed. The first step was to identify the most promising CO<sub>2</sub>/O<sub>2</sub> mixture. Based on the results from this first experiment, a mixture of 70% CO<sub>2</sub> + 30% O<sub>2</sub> was chosen for further investigation as a potential anaesthetic during the castration of young piglets. Thereby, it was established whether the duration and depth of anaesthesia were acceptable for castration where the animal has to be insensible and unconscious. Physiological effects were assessed based on electroencephalogram (EEG) and electrocardiogram (ECG) measurements, blood gas values and behavioural responses. During the induction phase, the only typical behaviour the piglets exhibited when exposed to the 70/30 gas mixture was heavy breathing. All piglets (n = 25) lost consciousness after approximately 30 s according to the EEG. Heart rate decreased slowly during the induction phase, a serious drop occurred when piglets lost their posture. Immediately after this drop, the heart rate neared zero or showed a very irregular pattern. Shortly after loss of posture, most animals showed a few convulsions. None of the animals showed any reaction to castration in behaviour and/or on the EEG and ECG. On average, the piglets recovered within 59 s, i.e. EEG returned to its pre-induction pattern and piglets were able to regain a standing position. After 120 s, heart rate returned to pre-induction levels. In order to explore the usage range of CO<sub>2</sub> concentration, 24 piglets were exposed to 60% CO<sub>2</sub> + 20% O<sub>2</sub> + 20% N<sub>2</sub> for up to 30 s after loss of consciousness (as registered on EEG), and castrated after removal from the chamber. Sixteen of the 24 animals showed a reaction to the castration on the EEG. To establish the maximum time piglets survive in 70% CO<sub>2</sub> + 30% O<sub>2</sub>, five piglets were placed in this mixture for 3 min. Two of them died. After that, four piglets were placed in this mixture for 2 min after unconsciousness, one died after 2 min. It was concluded from this study that it is possible to anaesthetise piglets with a mixture of 70% CO<sub>2</sub> + 30% O<sub>2</sub>, but that there are limits to its safety in terms of CO<sub>2</sub> concentration and duration of exposure. Before implementation for practical use, further research is essential to assess the limits of gas concentration and exposure times.*

**Keywords:** anaesthesia, carbon dioxide, castration piglets

### Introduction

In most pig-producing countries in the EU, male piglets are castrated at a young age to prevent boar taint in meat. The surgical castration is commonly performed without the use of anaesthesia or analgesia.

However, it has been abundantly substantiated that piglets experience pain during castration (Wemelsfelder and van Putten, 1985; McGlone and Hellman, 1988; McGlone *et al.*, 1993; Weary *et al.*, 1998; Taylor and Weary, 2000; Taylor *et al.*, 2001). Castration induces acute activation of the hypothalamic–pituitary–adrenal axis (HPA) and of the sympathetic nervous system (SNS) (Prunier *et al.*, 2005). Based on this information, there is growing criticism from

consumers, opposition groups and politicians concerning the unanaesthetised castration. Therefore, a lot of attention and effort is being given in different EU countries for the development of acceptable anaesthesia methods for piglets (Kohler *et al.*, 1998; Jäggin *et al.*, 2001; Walker *et al.*, 2004; Svendsen, 2006).

Anaesthesia is generally applied to prevent pain sensation during surgery. Pain is an aversive sensory and emotional experience representing awareness by the animal, of damage or threat to the integrity of its tissues. It changes the animal's physiology and behaviour to reduce or avoid damage (Molony and Kent, 1997). Anaesthesia or reduction of pain during surgery can be performed by local anaesthesia, induced by injection, or by general anaesthesia, induced by injection or inhalation. Anaesthetic gasses can be inhaled, because they evaporate at room temperature.

<sup>†</sup> E-mail: marien.gerritzen@wur.nl

Examples of anaesthetic gasses include ether, chloroform, halothane and methoxy-flurane. Anaesthetic gasses cause depression of the central neural system with induction of unconsciousness and muscle flaccidity. An overdose leads to paralysis of the breathing muscles and the animal is killed by apnea. Induction by an anaesthetic gas can result in excitation, depending on the gas used.

Under practical circumstances, the problem with most anaesthetic gasses is that their usage is restricted to veterinarians. Furthermore, the use of most of these anaesthetic gasses is subject to strict safety regulations and registration. Applying CO<sub>2</sub> as anaesthetic agents could presumably be a useful alternative. It is well recognised that CO<sub>2</sub> is an anaesthetic gas that produces rapid unconsciousness when inhaled at high concentrations (Martoft, 2001). However, increasing stress hormone levels (Kohler *et al.*, 1998), signs of asphyxia and behavioural excitation (Iwarsson and Rehinder, 1993; Raj and Gregory, 1995; Kohler *et al.*, 1998; Leach *et al.*, 2002), were observed due to the occurrence of both hypercapnia and hypoxia. Moreover, CO<sub>2</sub> is an acidic gas and has been found to be painful, causing unpleasant sensations to the nasal mucosa, lips and forehead in humans, when administered as gas-puffed stimuli in concentrations above 65% (Danneman *et al.*, 1997; Hari *et al.*, 1997). The right hemisphere of the human SII cortex is predominantly involved in this response, which may suggest emotional/motivational aspects of trigeminal pain, and is in agreement with the role of the trigeminal pathways as a general warning system. It has been shown that broilers can detect CO<sub>2</sub> in air at concentrations below 10% (Gerritzen *et al.*, 2007). Increases in head shaking and the elicitation of withdrawal from feeding at 55% CO<sub>2</sub> and above suggest that such concentrations of CO<sub>2</sub> in air may be aversive. In rats, a low concentration of CO<sub>2</sub> and addition of O<sub>2</sub> and humidification of the gasses could ameliorate these negative effects (Coenen *et al.*, 1995). In the latter case, almost no signs of asphyxia and excitation were observed. The main affect of CO<sub>2</sub> is not suffocation, but an anaesthetic activity. A problem with O<sub>2</sub>-replacing gasses is their lower efficacy in very young animals (AVMA, 2004).

An electroencephalogram (EEG) is widely used to record brain activity under different circumstances to determine the state of consciousness and brain disorders in humans and animals. The electrical activity recorded on the EEG can be classified into delta (<4 Hz), theta (4 to 8 Hz), alpha (8 to 13 Hz) and beta (>13 Hz) frequency bands. In alpha and beta rhythms, the animal is conscious (Kooi *et al.*, 1978). Increased dosage of most anaesthetic agents results in spontaneous progressive EEG responses with higher amplitude and lower frequencies (theta and delta waves) (Woodburry and Karler, 1960; Mattson *et al.*, 1972). It is also known that a quiescent or near-iso-electric EEG occurs in deep anaesthesia as well as brain death (Eger, 1981). The electrocardiogram (ECG) is used to determine heart function. Changes in frequency (beats/min) and other characteristics of the ECG such as amplitude, wave form and frequency patterns indicate the efficacy of heart function.



Figure 1 Induction test box, recording monitor and gas tank.

The objectives of these experiments were to determine, which concentrations of gasses in a mixture of CO<sub>2</sub> and O<sub>2</sub> are suitable to anaesthetise very young piglets to enable insensible castration. Behavioural, neural and physiological measurements were used to assess animal welfare aspects.

### Material and methods

The study presented here was executed in three successive phases. The first phase – identification of concentrations – was conducted to determine the most promising CO<sub>2</sub>/O<sub>2</sub> mixture to induce an adequate state of anaesthesia and analgesia. In the second phase – castration experiment – piglets were exposed to the most promising mixture from the first phase to provide insensible castration under full anaesthesia. During the final phase – limiting circumstances – an assessment was made as to whether piglets could remain in this mixture for a longer period without dying and whether a lower concentration of CO<sub>2</sub> would induce a sufficient anaesthetic effect. Ethical aspects of the experiment were judged and approved by the Animal Ethical Committees of the Animal Sciences Group and of the University of Utrecht.

#### *Animals and experimental design*

Piglets used during this experiment were from the herd at the experimental pig farm 'De Tolakker', Veterinary Faculty, University of Utrecht. All piglets were commercial cross-breeds ('Topigs') intended for fattening. After the experiment, piglets returned to their farrowing pen and were reared under standard commercial conditions.

Anaesthesia experiments were carried out with individual animals in a Perspex box i.d. 0.8 × 0.8 × 0.8 m (see Figure 1). The test box was fitted with a 1-cm Ø CO<sub>2</sub> inlet located 5 cm above the base of the box and an overflow valve situated at the top of the box. The different CO<sub>2</sub>/O<sub>2</sub> mixtures were injected from a pre-filled compression tank after mixing with a three-phase gas mixer. A gas-measuring tube was placed in the centre of the box at approximately head height (when piglets were standing), and CO<sub>2</sub> and

**Table 1** Experimental conditions (BW and gas concentrations) and behavioural events indicating loss of consciousness and recovery

Treatment	Weight (kg)	Gas concentrations		Behaviour		Recovery	
		O <sub>2</sub> %	CO <sub>2</sub> %	Sit (s)	Lop (s)	Sit (s)	Stand (s)
70% CO <sub>2</sub> + 30% O <sub>2</sub>	3.9 ± 0.4	28 ± 0	72 ± 0	13 ± 4	32 ± 7	67 ± 13	103 ± 32
50% CO <sub>2</sub> + 10% O <sub>2</sub>	3.2 ± 0.2	11 ± 0	51 ± 1	24 ± 5	41 ± 18	38 ± 8	57 ± 17
60% CO <sub>2</sub> + 20% O <sub>2</sub>	3.1 ± 0.7	19 ± 0	60 ± 2	16 ± 5	30 ± 5	45 ± 10	70 ± 14
40% CO <sub>2</sub> + 10% O <sub>2</sub>	3.5 ± 0.3	12 ± 0	43 ± 2	29 ± 6	77 ± 24	57 ± 27	61 ± 26
30% CO <sub>2</sub> + 15% O <sub>2</sub>	3.4 ± 0.4	14 ± 1	31 ± 2	69 ± 52	86 ± 83	6 ± 8	10 ± 11
30% CO <sub>2</sub> + 30% O <sub>2</sub>	3.5 ± 0.4	26 ± 0	30 ± 1	52 ± 9	314 ± 141	27 ± 33	29 ± 38

All values are mean ± s.e.d. of five piglets.

Sit = time taken to sit down (behaviour) or to sit up (recovery); Lop = time taken to loss of posture; Stand = time taken to recover posture (stand up unaided).

O<sub>2</sub> concentrations were measured continuously using an electronic gas analyser<sup>1</sup>. Piglets were placed in the test box after the required concentration had been established.

#### Identify concentrations

In order to identify the most promising CO<sub>2</sub>/O<sub>2</sub> mixture for a painless castration under complete anaesthesia, piglets were exposed to six different gas mixtures (Table 1). Behavioural aspects were judged with five piglets per gas mixture to assess whether they lost posture (indicative for loss of consciousness). After loss of posture, the piglets remained for a further 1 min in the gas mixture before removal from the box and were placed in a recovery crate under normal room conditions. Anaesthetic state was judged based on behavioural aspects, i.e. loss of posture, tonic and clonic seizures, and on clinical reflexes of cornea and to pain stimuli provided by ear and interdigital pinching. The piglets were allowed to recover without assistance. Recovery time was judged to be the moment animals could remain standing without support.

#### Castration experiment

Based on the results of the previous identification phase, 25 piglets were exposed to the most promising gas combination (i.e. 70% CO<sub>2</sub> with 30% O<sub>2</sub>). In order to facilitate identification of loss of consciousness and to control physiological conditions during anaesthesia, animals were equipped with electrodes to measure brain and heart activity. For the registration of the EEG, two needle electrodes<sup>2</sup> (55% silver, 21% copper, 24% zinc) of 10 mm length and 1 mm diameter were positioned subcutaneous on the skull on the line ear to ear, 0.5 cm left and right of the sagittal line. The electrodes were fixated with leucoplast tape. To register heartbeat and rhythmic disorders, two needle electrodes of 30 mm in length were placed subcutaneously at the left and right side of the chest approximately 1 cm behind the front legs. In order to minimise signal distortion, an earth-connecting electrode was placed subcutaneously at the back of the piglets. The electrodes were connected to a registration and recording device using isolated and coaxial shielded wires. The micro-voltage

signals of the brain and heart were amplified using a biomedical-amplifier<sup>3</sup> and continuously recorded using Windaq computer software<sup>4</sup>. Two blood samples were taken from the jugular vein of each animal shortly before introducing the piglets into the gas mixture and directly after castration. One of the blood samples was analysed for pH, pCO<sub>2</sub>, pO<sub>2</sub>, acid–base equilibrium (ABE) and O<sub>2</sub> saturation using a mobile acid–base laboratory<sup>5</sup>; the second sample was analysed for glucose and lactate (LX20 analyzer<sup>6</sup>). After measuring EEG and ECG for 30 s, piglets were individually placed into the pre-filled gas box. Behaviour of the animals was observed and analysed for gasping, head shaking, sitting, loss of posture (falling on its side) and convulsions. At 30 s after loss of posture, animals were taken out of the gas mixture and immediately castrated. Castration was performed by a single incision of approximately 2 cm from the left to the right through the scrotum. Both testicles were taken out and removed by cutting both funiculi. Immediately after castration, the second blood samples were taken from the jugular vein as mentioned previously. Both EEG and ECG measurements were continued for 3 min after removal from the gas mixture.

Blood values as well as EEG and ECG recordings were analysed afterwards to assess whether animals were unconscious at the time of castration and to classify the impact of the anaesthesia and castration on the piglets' welfare. EEG signals were analysed for changes in frequency and amplitude, more specifically, for the suppression of alpha (8 to 13 Hz) and beta (>13 Hz) waves and the occurrence of theta (4 to 8 Hz) and delta (<4 Hz) waves. The ECG signals were analysed for heart rate at the start of the gas flow, at loss of posture and during the last minute of the monitoring period.

#### Limiting circumstances

In order to evaluate the reliability of the method, it is essential to determine the limitations of the method. Decreasing CO<sub>2</sub> and O<sub>2</sub> concentrations are assumed to

<sup>3</sup> MODEL BMA-931, CWE, Inc., Ardmore, USA

<sup>4</sup> DATAQ instruments, Akron, Ohio, USA

<sup>5</sup> ABL 80 Flex, Radiometer, Copenhagen, Denmark

<sup>6</sup> LX20, Beckman Coulter, USA; www.Beckmancoulter.com

<sup>1</sup> PBI Dansensor, Ringsted, Denmark

<sup>2</sup> Engelhard-CLAL, New Jersey, USA

impact the anaesthetic properties of the gas mixture. On the other hand, increasing the duration that animals remain in the gas mixture will increase the duration of anaesthesia, and also the risk of mortality.

To gain an impression of the effect of decreasing CO<sub>2</sub> levels in practice, 25 piglets were exposed to a gas mixture of 60% CO<sub>2</sub> + 20% O<sub>2</sub> in nitrogen. Judgement of anaesthetic as well as of animal welfare aspects was based on the same parameters as described in the castration experiment. Furthermore, all piglets were treated similarly and measurements were taken as described in the castration experiment.

To assess maximal exposure time, five piglets were exposed to the gas mixture 3 min after loss of posture and four piglets were exposed for 2 min after loss of posture. Physiological and animal welfare aspects were assessed comparable to those described for the previous groups.

### Statistics

For assessment of an adequate anaesthesia, each piglet represents an experimental unit for  $n$  piglets, which are treated independently, the number  $x$ , which remain conscious, follows a binomial distribution with total  $n$  and probability  $P$ . A confidence interval can be calculated for probability  $P$  based on a relationship between the binomial and beta distribution. The number of animals that are effectively anaesthetised follows a binomial distribution. A 95% confidence limit on the probability for an effective stun can be obtained by means of a well-known relationship with the beta distribution (Johnson and Kotz, 1969).

Differences in blood values before exposure and directly after castration were analysed as a two-sided paired  $t$ -test (GenStat Committee, 2000).


## Results

### Pilot experiment

Based on behavioural observations, the different gas mixtures were examined on their capability to induce acceptable anaesthesia in young piglets. The results are presented in Table 1. It was observed that all piglets showed intensified breathing during induction with the different gas mixtures. Gasping was observed at a low CO<sub>2</sub> concentration. All animals showed some weak or heavy convulsions after loss of posture. The higher the O<sub>2</sub> concentration, the weaker the convulsions. The intensity of convulsions was not affected by the CO<sub>2</sub> concentration but the duration till onset of convulsion was delayed by lower CO<sub>2</sub> concentrations. The time taken to induce loss of posture was lengthened by a lower CO<sub>2</sub> concentration. More important was the dramatic decrease in duration till recovery after lowering the CO<sub>2</sub> concentration. At an O<sub>2</sub> concentration lower than 14%, all piglets appeared blue (hypoxic). Based on behavioural parameters (loss of posture), it remained doubtful whether piglets became unconscious during immersion in CO<sub>2</sub> concentrations below 31%.

Based on these results, castration experiments were performed using a mixture of 70% CO<sub>2</sub> + 30% O<sub>2</sub>.

**Table 2** Average ( $n = 25$ ) start and duration (s) of behavioural events during exposure to a gas mixture of 70% CO<sub>2</sub> + 30% O<sub>2</sub>



### Castration experiment

A typical behaviour (Table 2) during induction with the gas mixture 70% CO<sub>2</sub> + 30% O<sub>2</sub> was heavy breathing, which started 11 ± 1 s after immersion. Heavy breathing was observed until 6 s after loss of posture. At 16 ± 0.8 s after immersion in the gas mixture the animals sat down until eventual loss of posture at approximately 24 ± 1 s. Immediately after loss of posture all piglets showed a few mild to strong convulsions (at 26 ± 1.7 s).

A change to lower EEG frequencies was observed after approximately 19 ± 1.1 s. Minimal brain activity was observed 33 ± 2 s after immersion. The castration took place on average 19 ± 5 s after the animals were removed from the gas mixture. None of the piglets displayed a behavioural, EEG or ECG response to castration (Figure 2). Recovery started 56 ± 11 s after removal from the gas box.

Based on these observations it was calculated that within a confidence limit of 95% and taking into account the number of animals with a reliable EEG ( $n = 25$ ), the chance that piglets aged 3 to 5 days are effectively anaesthetised is greater than 0.89.


During induction with the gas mixture, the heart rate decreased slowly. After loss of posture the heart rate dropped dramatically to almost zero (Figure 3). Heart activity increased approximately 1 min after the piglets were removed from the gas box. Heart rate returned to the same rhythm as prior to exposure to the gas mixture in approximately 120 s.

Blood pH decreased from 7.4 to 7.1, the pCO<sub>2</sub> increased from 6 to 9.7 kPa and the pO<sub>2</sub> from 3.8 to 4.5 kPa (Table 3). The O<sub>2</sub> saturation increased from 38% to 42%. Due to the increase of pCO<sub>2</sub> and decrease of pH, the ABE shifted from 0 to -10.


Both glucose and lactate levels increased from 7.2 to 9.4 mmol/l and from 7.6 to 13.8 mmol/l, respectively.

### Limits in gas concentration and duration

After immersion in a gas mixture of 60% CO<sub>2</sub> + 20% O<sub>2</sub>, 16 of 24 piglets displayed behavioural and EEG responses during castration. When immersed in a gas mixture of 70% CO<sub>2</sub> + 30% O<sub>2</sub> for 3 min after loss of posture, two of five piglets died. When they remained in this mixture for 2 min,



**Figure 2** Characteristic examples of the electroencephalogram (EEG) and electrocardiogram (ECG) readings: A = prior to anaesthesia, B = loss of consciousness (displaying strong signal disturbance at loss of posture), C = deep anaesthesia (three vertical lines above EEG signal trace indicate the moment of castration), D = recovery of consciousness after anaesthesia.



**Figure 3** Graphical example of the development of the heartbeat during exposure to the 70% CO<sub>2</sub> + 30% O<sub>2</sub> gas mixture and the recovery period.

**Table 3** Average blood gas values (jugular vein) before exposure to 70% CO<sub>2</sub> + 30% O<sub>2</sub> and directly after castration

Parameter	Before exposure	After castration	P values
pH	7.4 ± 0.02	7.1 ± 0.02	<0.001
pCO <sub>2</sub> (kPa)	5.7 ± 0.27	8.2 ± 0.46	<0.001
pO <sub>2</sub> (kPa)	4.2 ± 0.41	7.3 ± 0.91	0.004
O <sub>2</sub> saturation (%)	38.7 ± 2.9	45.6 ± 5.5	0.276
Acid-base equilibrium (mmol/l)	-0.2 ± 0.5	-9.4 ± 0.8	<0.001
Glucose (mmol/l)	7.2 ± 0.16	9.9 ± 0.38	<0.001
Lactate (mmol/l)	7.5 ± 0.5	13.9 ± 0.8	<0.001

Values are mean ± s.e. of the mean based on n = 25. Differences between pairs are significant when P < 0.005 based on two-sample two-sided t-test.

one of four piglets died. Due to this longer immersion period the pH decreased from 7.4 to 7.0, the pCO<sub>2</sub> increased from 6.5 to 15.6 kPa and the pO<sub>2</sub> increased from 2.6 to 9.2 kPa. The O<sub>2</sub> saturation increased from 27% to 73% and the ABE shifted from 0 to -10. The glucose and lactate values increased from 6.5 to 10.3 mmol/l and from

7.2 to 12.5 mmol/l, respectively. No differences were found in the values of these parameters between animals that survived or died. It was observed that the animals that died did so approximately 3 to 4 min after they were removed from the gas box. Conspicuously, the animals apparently regained consciousness (based on behaviour and EEG) but

their heart rhythm did not recover to pre-anaesthesia levels. Moreover, heart rhythm remained irregular in these animals, possibly resulting in death.

## Discussion

Castration of male piglets without the use of any anaesthesia is meeting with increasing resistance from society and from the pig-producing sector itself. Aim of this research was to examine the suitability of CO<sub>2</sub> anaesthesia for castration under practical on-farm conditions. Main reason to focus on CO<sub>2</sub> was because this could be a method that can be used by farmers without intervention of a veterinarian. Moreover, the use of anaesthetic gasses as halothane and isoflurane is in many countries strictly prohibited to veterinarians.

Exposure to CO<sub>2</sub> induces depression of brain activity, which extends progressively from the telencephalon to the diencephalon and then to the mesencephalon. It is well recognised that CO<sub>2</sub> is an anaesthetic gas, which produces rapid unconsciousness and complete analgesia when inhaled at different concentrations (Lauer, 1994; Kohler *et al.*, 1998; Martoft, 2001). However, side-effects like signs of breathlessness and behavioural excitation are observed (van den Bogaard *et al.*, 1985; Raj *et al.*, 1992; Kohler *et al.*, 1998). In rats, a low concentration of CO<sub>2</sub> and the addition of O<sub>2</sub> and humidification of the gasses could help ameliorate the negative effects. In which case almost no signs of asphyxia and excitation were observed (Coenen *et al.*, 1995). The main action of CO<sub>2</sub> is not to suffocate but to anaesthetise (Coenen *et al.*, 1995). In the present experiment, O<sub>2</sub> saturation did not change during exposure to the gas mixture, which could be the reason for the relative mild induction of anaesthesia. However, it must be mentioned that O<sub>2</sub> saturation was low, which is presumed to be due to the stress of handling. After exposure of fattening pigs to high concentrations (80% to 90%) of CO<sub>2</sub>, the pigs remained quiet for the first 10 to 20 s and displayed movement during the subsequent 10 s (Forslid, 1987). It is difficult to perceive what pigs experience during this induction period. At best, it may be mildly unpleasant. At worst, it could be very unpleasant. Following immersion in high CO<sub>2</sub> concentrations, pigs are well stunned and remain so provided they are bled within 1 min of stunning. For stunning of slaughter pigs, a concentration of 80 to 90% CO<sub>2</sub> is recommended for a rapid induction of anaesthesia (Lambooj, 1990). Activity, i.e. wing flapping in broilers, and the number of birds displaying this behaviour were low using the gas mixture 40% CO<sub>2</sub>/30% O<sub>2</sub>/30% N<sub>2</sub>, suggesting a positive effect of O<sub>2</sub> (Lambooj *et al.*, 1999; McKeegan and Smith, 2005). When young piglets were placed in the gas mixture, they usually remained calm, breathing increased and then lost their posture. Most animals showed a few convulsions after they were considered to be unconscious according to the EEG. This observation is supported by results in ducks and turkeys, where these

birds showed low levels of consciousness (according to theta and delta waves on the EEG) during loss of posture (Gerritzen *et al.*, 2006). The piglets recovered on average within 1 min after removal from the mixture, which is rather fast. To be able to guarantee that no piglets are castrated while regaining consciousness and allowing for the biological variation between animals, it is recommended to castrate within 30 s.

Breathing is regulated in the brain by the medulla oblongata. This area of the brain has sensors that regulate the pH (acidity) of the blood. Blood pH is determined by the concentration of CO<sub>2</sub> and O<sub>2</sub> in the blood. When blood pCO<sub>2</sub> increases and subsequently pH decreases, the brain regulates an increase of the ventilation of the lung. Therefore, the animal in the gas box increases its rate of breathing, which under these circumstances results in the inhalation of an increased amount of CO<sub>2</sub>. It appeared that when blood pH decreases, the pH of the cerebrospinal fluid follows (Martoft *et al.*, 2003), resulting in neuronal inhibition and anaesthesia (Woodburry and Karler, 1960). The normal pH of brain fluid is 7.4 and induction of unconsciousness occurs at a pH of 7.1 (Eisele *et al.*, 1967). During our experiment, the blood pH of the piglets decreased to the same value, caused by a shift in the ABE. According to the EEG, the piglets were unconscious in the 70% CO<sub>2</sub> + 30% O<sub>2</sub> mixture as with the 60% CO<sub>2</sub> + 20% O<sub>2</sub> one. However, piglets placed in the 60% CO<sub>2</sub> + 20% O<sub>2</sub> mixture were not anaesthetised long enough and displayed a behavioural and EEG response upon incision and cutting of the funiculus cords during castration, while the decrease in blood pH was the same as for those piglets placed in 70% CO<sub>2</sub> + 30% O<sub>2</sub>. In experiments with fattening pigs (Martoft, 2001) it appeared that pigs have been shown to recover rapidly even when they were exposed to concentrations of 80% CO<sub>2</sub>. It is questionable whether 60% CO<sub>2</sub> in the gas mixture is too low for the anaesthetised castration of young piglets. However, it is possible that a longer exposure time of piglets to this concentration can induce an anaesthetic state that is of sufficient duration to perform a painless castration.

An extraordinary increase of the pCO<sub>2</sub> in the blood causes initially a stimulation and later a depression of the breathing rate, which ends in death (Guyton and Hall, 2000). At slaughter, when the pH in the blood in pigs decreases to a value of 6.6, death is a possibility (Martoft, 2001). The same phenomenon was observed in ducks and turkeys (Gerritzen *et al.*, 2007). Changes in blood pH affect enzyme reactions related to energy production, causing changes in membrane permeability and management of electrolytes. It is hypothesised that this pH point is already too low for the regeneration of the physiological balance. There is an increased risk of death in piglets if they remain in 70% CO<sub>2</sub> + 30% O<sub>2</sub> for longer than 4 min (Svendson, 2006). During our research, piglets died when they remained in this gas mixture for a period longer than 2 min. These differences between the results of this study and Svendson's study cannot be easily explained. Differences

can be introduced due to many factors like stability of the gas concentration in the test box, gas-measuring location, the size of the test box and the breed. The similarities and differences between different studies indicate the variation that can be expected under practical use.

Since blood pH did not decrease below 7.0, recovery should have been possible. However, restoration of the heart rate did not occur. It can thus be concluded that CO<sub>2</sub> concentration in the gas mixture and exposure time to the gas mixture are critical factors for the successful induction of unconsciousness on the one hand or the risk of death in young piglets on the other hand.

## Conclusions

Induction and maintenance of insensibility and unconsciousness during castration of young piglets is possible using 70% CO<sub>2</sub> and 30% O<sub>2</sub>. However, it cannot be neglected that the inhalation and induction of anaesthesia with CO<sub>2</sub> is causing discomfort or stress. However, un-anaesthetised castration or local inter-testicular anaesthesia was assumed to cause more stress and pain than the induction of general CO<sub>2</sub> anaesthesia. It is clear that CO<sub>2</sub> anaesthesia induces an analgetic state for only a short period and therefore the use of additional analgetic drugs is recommended.

For use in practice, the conditions under which insensibility and unconsciousness are induced and the prevention of death are critical. The limits of these conditions should be further assessed before the method can be applied under commercial practice. Moreover, close co-operation with farmers, designers and manufacturers is essential for the development of a practical system for on-farm usage.

## Acknowledgement

The authors gratefully acknowledge the funding of the present study by the Dutch ministry of Agriculture, Nature Management and Food.

## References

AVMA Panel 2004. Report on the AVMA panel on euthanasia. [www.avma.org/issues/animal\\_welfare/euthanasia.pdf](http://www.avma.org/issues/animal_welfare/euthanasia.pdf)

van den Bogaard AJM, van Dam E and Weekers FH 1985. Het gebruik van een koolzuurgas euthanasie apparaat voor ratten. *Biotechniek* 3, 34–38.

Coenen AML, Drinkenburg WHIM, Hoenderken R and van Luitelaar ELJM 1995. Carbon dioxide euthanasia in rats: oxygen supplementation minimizes signs of agitation and asphyxia. *Laboratory Animals* 29, 262–268.

Danneman PJ, Stein S and Walshaw SO 1997. Humane and practical implications of using carbon dioxide mixed with oxygen for anesthesia or euthanasia of rats. *Laboratory Animal Sciences* 47, 376–385.

Eger EIN 1981. Isoflurane: a review. *Anesthesiology* 55, 559–576.

Eisele JH, Eger EI and Muallem M 1967. Narcotic properties of carbon dioxide in the dog. *Anesthesiology* 28, 856–865.

Forslid A 1987. Transient neocortical, hippocampal and amygdaloid EEG silence induced by one minute inhalation of high concentrations CO<sub>2</sub> in swine. *Acta Physiologica Scandinavica* 130, 1–10.

GenStat Committee 2000. The guide to GenStat (ed. RW Payne). VSN Int. Ltd., Oxford.

Gerritzen MA, Lambooi E, Reimert HGM, Spruijt BM and Stegeman JA 2006. Susceptibility of ducks and turkey to severe hypercapnic hypoxia. *Poultry Science* 85, 1055–1061.

Gerritzen M, Lambooi B, Reimert HGM, Stegeman A and Spruijt B 2007. A note on behaviour of poultry exposed to increasing carbon dioxide concentrations. *Applied Animal Behaviour Science* 108, 179–185.

Guyton AC and Hall JE 2000. Chapters 5 and 7 in textbook of medical physiology, 10th edition. W.B. Saunders Co, Philadelphia, Pennsylvania, USA.

Hari P, Portin K, Kettenmann B, Jousmäki V and Kobal G 1997. Right-hemisphere preponderance of responses to painful CO<sub>2</sub> stimulation of the human nasal mucosa. *Pain* 72, 145–151.

Iwarsson K and Reh binder C 1993. A study of different euthanasia techniques in guinea pigs, rats and mice. Animal response and post-mortem findings. *Scandinavian Journal of Laboratory Animal Science* 4, 191–205.

Jäggin N, Kohler I, Blum U and Schatzmann U 2001. Die kastration von neugeborenen ferkeln unter halothananästhesie. *Der Praktische Tierarzt* 82, 1054–1061.

Johnson NL and Kotz S 1969. Discrete distributions. John Wiley, New York.

Kohler I, Moens Y, Busato A, Blum J and Schatzman U 1998. Inhalation anaesthesia for the castration of piglets: CO<sub>2</sub> compared to halothane. *Journal of Veterinary medicine* 45, 625–633.

Kooi AK, Tucker PT and Marshall RE 1978. Fundamentals of electroencephalography, 2nd edition. Harper & Row, Publishers, Maryland.

Lambooi E 1990. The use of CO<sub>2</sub> for stunning of slaughter pigs. Report of a meeting of experts, Heeze, 26–27 January 1990 [Anwendung von CO<sub>2</sub> zur Betäubung von Schlachtschweinen. Bericht über eine von 26. bis 27.1.1990 in Heeze abgehaltene Expertentagung]. IVO B-Rapport 354, 1990, *Zeits. Fleischwirtschaft* 70 1990 10, pp. 1161–1166. (Eng.: pp. 1173–1177).

Lambooi E, Gerritzen MA, Engel B, Hillebrand SJW, Lankhaar J and Pieterse C 1999. Behavioural responses during exposure of broilers to different gas mixtures. *Applied Animal Behaviour science* 62, 255–265.

Lauer S 1994. Die CO<sub>2</sub>/O<sub>2</sub>- Anästhesie zur routinemässigen kastration von Ferkeln. Beurteilung von praxiseignung und Tierschutzrelevanz anhand von Verhaltensbeobachtungen. Diss med vet München.

Leach MC, Bowell VA, Allan TF and Morton DB 2002. Aversion to gaseous euthanasia agents in rats and mice. *Comparative Medicine* 52, 249–257.

Martoft L 2001. Neurophysiological effects of high concentration CO<sub>2</sub> inhalation in swine. PhD, Department of Anatomy and Physiology, Royal Veterinary and Agricultural University, Frederiksberg, Denmark.

Martoft L, Jørgenson HS, Foslid A, Pedersen HD and Jørgenson PF 2003. CO<sub>2</sub> induced acute respiratory acidoses and brain tissue intracellular pH: a <sup>31</sup>P NMR study in swine. *Laboratory Animals* 37, 241–248.

Mattson JL, Stinson JM and Clark CS 1972. Electroencephalographic power-spectral changes coincident with the onset of carbon dioxide narcosis in rhesus monkey. *American Journal of Veterinary Research* 33, 2043–2049.

McGlone JJ and Hellman JM 1988. Local and general anesthetic effects on behavior and performance of two and seven-week-old castrated and uncastrated piglets. *Journal of Animal Science* 66, 3049–3058.

McGlone JJ, Nicholson RI, Hellman M and Herzog DN 1993. The development of pain in young pigs associated with castration and attempts to prevent castration-induced behavioral changes. *Journal Animal Science* 71, 1441–1446.

McKeegan DEF and Smith FS 2005. Behavioural correlation of olfactory and trigeminal gaseous stimulation in chickens, *Gallus domesticus*. *Physiology and Behaviour* 84, 761–768.

Molony V and Kent JE 1997. Assessment of acute pain in farm animals using behavioral and physiological measurements. *Journal of Animal Science* 75, 266–272.

Prunier A, Mounier AM and Hay M 2005. Effects of castration, tooth resection, or tail docking on plasma metabolites and stress hormones in young pigs. *Journal of Animal Science* 83, 216–222.

Raj ABM and Gregory NG 1995. Welfare implications of the gas stunning of pigs 1. Determination of aversion to the initial inhalation of carbon dioxide or argon. *Animal Welfare* 4, 273–280.

Raj ABM, Wotton SB and Gregory NG 1992. Changes in somatosensory evoked potentials and spontaneous electrocardiogram of hens during stunning with carbon dioxide and argon mixture. *British Veterinary Journal* 148, 147–156.

Svensden O 2006. Castration of piglets under carbon dioxide (CO<sub>2</sub>) anaesthesia. *Journal of Veterinary Pharmacology and Therapeutics* 29, 47–59.

## Castration of piglets under CO<sub>2</sub>-gas anaesthesia

Taylor AA and Weary DM 2000. Vocal responses of piglets to castration: identifying procedural sources of pain. *Applied Animal Behaviour Science* 70, 17–26.

Taylor AA, Weary DM, Lessard M and Braithwaite LA 2001. Behavioural responses of piglets to castration: the effect of piglet age. *Applied Animal Behaviour Science* 73, 35–43.

Walker B, Jäggin N, Doherr M and Schatsmann U 2004. Inhalation anaesthesia for castration of newborn piglets: experiences with isoflurane and isoflurane/N<sub>2</sub>O. *Journal of Veterinary Medicine* 51, 150–154.

Weary DM, Braithwaite LA and Fraser D 1998. Vocal response to pain in piglets. *Applied Animal Behaviour Science* 56, 161–172.

Wemelsfelder F and Putten G van 1985. Behavior as a possible indicator for pain in piglets. Zeist, The Netherlands, Instituut voor Veeteltkundig Onderzoek "Schoonoord".

Woodburry DM and Karler R 1960. The role of carbon dioxide in the nervous system. *Journal of American Society of Anaesthesiologists* 21, 686–703.