

SHORT COMMUNICATION

A simple staining method for observation of germinated *Striga* seeds

Shusheng Long^{1,2}, Venasius Lenzemo³, Thomas W. Kuyper⁴, Zhengsheng Kang², Horst Vierheilig^{1,5} and Siegrid Steinkellner^{1*}

¹Department of Applied Plant Sciences and Plant Biotechnology, Institute of Plant Protection, University of Natural Resources and Applied Life Sciences, Peter Jordan-Strasse 82, Vienna A-1190, Austria; ²College of Plant Protection and Shaanxi Key Laboratory of Molecular Biology for Agriculture, Northwest A&F University, Yangling, Shaanxi 712100, China; ³Institute of Agricultural Research for Development, Maroua, Cameroon; ⁴Department of Soil Quality, Wageningen University, Droevendaalsesteeg 4, Wageningen NL-6708 PB, The Netherlands; ⁵Departamento de Microbiología, Estación Experimental de Zaidín, CSIC, Granada E-18008, Spain

Abstract

In vitro techniques are essential for *Striga* research and the development of appropriate control methods. In the laboratory, pre-screening of non-host or false-host plants of *Striga* for trap cropping or the screening of hosts for resistance involves visual evaluation of *Striga* seed germination that may be stimulated by plant parts or root exudates. This technique is presently laborious because the small *Striga* radicles are colourless. A number of solutions were evaluated to visualize the radicles of *Striga hermonthica* to obtain a reliable, simple and fast staining method yielding good contrast for visual observation, with readily available, inexpensive and minimally toxic dyes and staining solutions. The inks Brilliant Blue (Pelikan), Blue (Geha) and Brilliant Green (Pelikan), in either vinegar or lactic acid, produced radicles with excellent contrast, whereas radicles stained with Brilliant Red (Pelikan), Black (Sheaffer), Brilliant Black (Pelikan) gave good contrast. *Striga* radicles stained with Aniline Blue in vinegar or lactic acid, or Coomassie Brilliant Blue R250 in lactic acid, showed excellent contrast. Radicles stained with Ink Black (Sheaffer), Cotton Brown, or Rubin S in either vinegar or lactic acid also showed good contrast. With water as the diluent, only Lactophenol Blue showed excellent contrast. For health, safety and environmental concerns, availability and staining time, Blue ink in household vinegar (5% acetic acid) appears to

be an excellent dye for *Striga* radicles and could be used in routine *Striga* germination assays.

Keywords: germination assay, radicle, staining solutions, *Striga*

Introduction

Witchweed (*Striga hermonthica*) is a root parasitic weed of cereals in tropical agro-ecosystems, most notably in the savannah region of Africa. Frequently, crop yield losses due to this parasite can be as high as 100% under heavy infestation. *Striga* is considered to be one of the most important limiting factors on the yields of cereals (Yoder and Musselman, 2006).

Laboratory investigations are essential for the study of *Striga* for development of appropriate control measurements. Screening of potential trap crops (crops that stimulate suicidal germination of *Striga* seeds) or host plant resistance *in vitro* is vital, as it enables much more material to be evaluated than would be possible with limited field space. Several standard techniques are available for conditioning *Striga* seeds and germination bioassays. Prior to germination the seeds need an after-ripening period of several months, and no germination occurs without appropriate germination stimulants, moisture and temperature (Parker and Riches, 1993). In germination bioassays, preconditioned *Striga* seeds are usually placed on a glass fibre filter disc (Berner *et al.*, 1997; Matusova *et al.*, 2004, 2005) or in a well of a microtitre plate (Gurney *et al.*, 2002) and exposed to a germination stimulant. The final evaluation of

*Correspondence

Email: siegrid.steinkellner@boku.ac.at

germinated seeds is performed by visual examination of the protruded radicle under a dissecting microscope (Berner *et al.*, 1997; Gurney *et al.*, 2002; Matusova *et al.*, 2005). However, the tiny seeds (0.25–0.50 mm in size) (Parker and Riches, 1993), and the rather hyaline and colourless radicles, make observations and counting a laborious task. To our knowledge, staining of *Striga* seeds has been reported only for viability testing using tetrazolium red dye (Berner *et al.*, 1997), but staining protocols to ease evaluation of germinated *Striga* seeds have not been reported. We searched for a reliable, simple, fast and inexpensive staining method to allow investigation of germinated *Striga* seeds. Moreover, we were particularly interested in a method that avoids the use of highly toxic compounds in the staining solution.

Materials and methods

Preconditioning of *Striga* seeds

Seeds of *Striga hermonthica* (Del.) Benth. were harvested from infested sorghum plants in Maroua, Cameroon in October 2005. According to Matusova *et al.* (2004) and Lenzemo (2004), the seeds were surface-sterilized with 2.5% sodium hypochlorite solution (Merck, Darmstadt, Germany) containing two drops of Tween 20 for 5 min, transferred to a funnel lined with a folded filter paper (MN 615 $\frac{1}{4}$, Machery-Nagel, Germany) and thoroughly rinsed with sterile, deionized water. The filter paper with the seeds was removed and placed on a layer of tissue paper to dry. Thereafter, lids of a Petri dish (90 mm diameter) were lined with two layers of filter paper (Whatman No. 2, Dassel, Germany) and moistened with 5 ml deionized sterile water. Thirty glass fibre filter discs (1 cm diameter, Whatman, GF/A, Dassel, Germany) were evenly distributed on the wetted filter paper in each Petri dish, and the *Striga* seeds were spread (c. 80–100 seeds) on each wetted glass fibre filter disc. The Petri dishes were sealed with Parafilm™, wrapped in aluminium foil and incubated at 28°C in the dark for 15–21 d.

Germination bioassay

After the preconditioning period of 15–21 d, the 1 cm glass fibre filter paper discs containing the *Striga* seeds were removed and dried on tissue paper for 1 h to remove excess moisture. The discs were transferred to a new lid of a Petri dish (5 discs per Petri dish) lined with a filter paper ring (outer diameter 90 mm, inner diameter 80 mm) moistened with 1 ml of deionized water. Subsequently, 50 μ l of the synthetic

Table 1. Comparison of different stains diluted in water for staining of germinated *Striga* seeds

Name of dye	Concentration (%)	Staining time	Staining results	Suitability	Indication of danger	Risk phrases ^a
Lactophenol Blue solution	0.78	< 5 min	Radicle blue, excellent contrast	Excellent	Toxic	R23/24/25-34-68-48/20/21/22
Toluidine Blue O	0.05	2 h	Radicle dark purple, medium contrast	Suitable	Not hazardous	
Cotton Brown	0.10	2 h	Radicle light brown, medium contrast	Suitable	Not known	
Methylene Blue	0.01	2 h	Radicle bluish, poor contrast	Not suitable	Harmful	R22
Methyl Red	0.03	2 h	Radicle light brown, poor contrast	Not suitable	Harmful	R40
Trypan Blue	0.03	> 5 h	Radicle partially light blue, poor contrast	Not suitable	Toxic	R45,62,68,63
Evans Blue	0.01	> 5 h	Radicle partially bluish, poor contrast	Not suitable	Toxic	R45
Coomassie Brilliant Blue	0.03	> 5 h	Radicle partially bluish, poor contrast	Not suitable	Not hazardous	
Bengal Rose B	0.01	> 5 h	Radicle partially reddish, poor contrast	Not suitable	Not hazardous	
Cochenille Red	0.05	> 5 h	Radicle partially reddish, poor contrast	Not suitable	Not hazardous	
Black ink (Sheaffer)	0.78	> 5 h	Radicle poorly bluish	Not suitable	Not hazardous	
Bromophenol Blue	0.03	> 5 h	Radicle not stained	Not suitable	Not applicable	
Rubin S	0.03	> 5 h	Radicle not stained	Not suitable	Not hazardous	
Aniline Blue	0.10	> 5 h	Radicle not stained	Not suitable	Not hazardous	

^aStated in the manufacturers' material safety data sheets according to EU regulation (<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32001L0059;EN:HTML>). R23/24/25, Toxic by inhalation, in contact with skin and if swallowed; R48/20/21/22, harmful: danger of serious damage to health by prolonged exposure through inhalation, in contact with skin and if swallowed; R34, causes burns; R68, possible risks of irreversible effects; R22, harmful if swallowed; R40, possible risk of cancer; R45, may cause cancer; R62, possible risk of impaired fertility; R63, possible risk of harm to the unborn child.

Table 2. Comparison of different stains, diluted in vinegar or lactic acid, for staining of germinated *Striga* seeds

Name of dye	Conc. (%)	Diluent	Staining time	Staining results	Suitability
Aniline Blue	0.03	Vinegar	< 5 min	Radicle blue, excellent contrast	Excellent
		Lactic acid ^a	< 5 min	Radicle blue, excellent contrast	Excellent
Coomassie Brilliant Blue R 250 ^b	0.03	Lactic acid	< 5 min	Radicle blue, excellent contrast	Excellent
Ink black (Sheaffer)	0.78	Vinegar	< 5 min	Radicle brownish, good contrast	Good
		Lactic acid	< 5 min	Radicle brownish, good contrast	Good
Cotton Brown ^b	0.25	Lactic acid	< 5 min	Radicle light brown, good contrast	Good
Rubin S	0.03	Vinegar	< 5 min	Radicle reddish, good contrast	Good
		Lactic acid	< 5 min	Radicle reddish, good contrast	Good
Bromophenol Blue ^b	0.03	Lactic acid	< 5 min	Radicle light brown, poor contrast	Not suitable
Methylene Blue	0.03	Vinegar	> 5 h	Radicle not stained	Not suitable
		Lactic acid	> 5 h	Radicle not stained	Not suitable

^a Irritant according to material safety data sheet (Fluka, Switzerland).

^b Hardly soluble in vinegar, therefore not tested.

germination stimulant GR24 (1 mg l⁻¹) was added to each disc, the Petri dishes were sealed with Parafilm™ and incubated at 28°C in darkness for 2 d (Lendzemo, 2004; Matusova *et al.*, 2004).

Staining solutions

The following stains, frequently used in microscopy and plant pathological laboratories (Dhingra and Sinclair, 1995) were tested: Lactophenol Blue solution (Fluka, Buchs, Switzerland), Cotton Brown (Bayer, Leverkusen, Germany), Bengal Rose B (Bayer, Leverkusen, Germany), Coomassie Brilliant Blue R250 (Fluka, Buchs, Switzerland), Bromophenol Blue (Riedel-de Haen, Seelze, Germany), Cochenille Red (Bayer, Leverkusen, Germany), Aniline Blue (Fluka, Buchs, Switzerland), Evans Blue (Sigma-Aldrich, Steinheim, Germany), Methylene Blue (Merck, Darmstadt, Germany), Methyl Red (Merck, Darmstadt, Germany), Toluidine Blue O (Sigma-Aldrich, Steinheim, Germany), Rubin S (= Fuchsin acid) (Merck, Darmstadt, Germany) and Trypan Blue (Fluka, Buchs, Switzerland).

Ink is used in routine staining of the pathogenic fungus *Pseudocercospora herpotrichoides* (Mauler-Machnik and Nass, 1990) and mycorrhizal fungi (Vierheilig *et al.*, 1998). We tested Black ink (Sheaffer, Ft. Madison, USA), Brilliant Black ink (Pelikan, Hannover, Germany), Brilliant Green ink (Pelikan, Hannover, Germany), Brilliant Red ink (Pelikan, Hannover, Germany), Brilliant Blue ink (Pelikan, Hannover, Germany) and Blue ink (Geha, Brilon, Germany). Stock solutions (1%, w/v) were prepared by dissolving the powdery stains in the diluent, storing for 24 h under ambient light and temperature conditions, and subsequent filtering through filter paper (MN 615 1/4, Macherey-Nagel, Germany). The liquid stains (Lactophenol Blue solution and the inks) were used without filtration. Based on preliminary results, the

final concentrations of the staining solutions were adjusted with the respective diluent. Apart from deionized sterile water, we used lactic acid (12.1 M) (Fluka, Buchs, Switzerland), which is known to macerate and clarify plant tissue (Wittmann, 1994), and household vinegar (clear, brownish-yellow, 5% acetic acid) (Mautner-Markhof, Vienna, Austria) as diluents. Fresh working solutions of each dye and ink were prepared for each experiment.

Staining procedure

Thirty microlitres of staining solution were added to each glass fibre disc containing pregerminated *Striga* seeds. The seeds were incubated at room temperature under natural light and observed under a stereo microscope for germination and staining of the radicle after 5 min, 2 h and 5 h. Each staining solution was tested on three individual filter discs.

Results

Out of 14 stains diluted in water, only Lactophenol Blue solution, a substance classified as toxic, resulted in excellent staining of *Striga* radicles within 5 min (Table 1). Compared to the unstained hyaline radicles, Lactophenol Blue solution gave uniformly blue-stained radicles and a pronounced contrast to the brown seeds. Concurrently, the glass fibre filter background was hardly stained. Toluidine Blue O and Cotton Brown showed a slightly reduced staining result. Due to the staining time of 2 h, these stains were classified only as 'suitable'. Bromophenol Blue, Rubin S and Aniline Blue did not stain *Striga* radicles. All other water-based stains were classified as not suitable, due to the poor contrast of the stained radicles and the filter paper background.

The protocols with several non-toxic stains, frequently used in microbiological laboratories, were

Table 3. Comparison of different inks (0.78% in lactic acid or vinegar) for staining of germinated *Striga* seeds

Name of ink (company)	Diluent	Staining time	Comments	Suitability
Brilliant Blue (Pelikan)	Vinegar	<5 min	Radicle blue, excellent contrast	Excellent
	Lactic acid	<5 min	Radicle blue, excellent contrast	Excellent
Blue (Geha)	Vinegar	<5 min	Radicle blue, excellent contrast	Excellent
	Lactic acid	<5 min	Radicle blue, excellent contrast	Excellent
Brilliant Red (Pelikan)	Vinegar	<1 min	Radicle reddish, good contrast	Good
	Lactic acid	<5 min	Radicle reddish, good contrast	Good
Black (Sheaffer)	Vinegar	<5 min	Radicle brownish, good contrast	Good
	Lactic acid	<5 min	Radicle brownish, good contrast	Good
Brilliant Black (Pelikan)	Vinegar	<5 min	Radicle brownish, good contrast	Good
	Lactic acid	<5 min	Radicle brownish, good contrast	Good
Brilliant Green (Pelikan)	Vinegar	<5 min	Radicle light green, excellent contrast	Good
	Lactic acid	<5 min	Radicle light green, excellent contrast	Good

more successful when diluted in lactic acid and vinegar (Table 2). In particular, Aniline Blue (in both diluents) and Coomassie Brilliant Blue R250 resulted in excellent staining results. Black ink (Sheaffer), Cotton Brown and Rubin S showed good suitability. Bromophenol Blue and Methylene Blue were not suitable due to insufficient or absent staining of the *Striga* radicles.

Due to safety concerns, availability and the convenience of handling, other ink products were tested (Table 3). Blue inks (Geha and Pelikan) exhibited better contrast than black inks and could be considered as excellent. All other inks tested showed good staining results. Both diluents, lactic

acid and household vinegar, yielded similar staining results. Representative results are illustrated in Fig. 1.

Discussion

Our results show that *Striga* radicles can be easily stained with minimal effort. Several dyes, frequently used for microscopic examination of fungi (Dhingra and Sinclair, 1995), effectively stained *Striga* radicles. While aqueous Lactophenol Blue gave an excellent radicle stain, it is a potential hazard with known toxicity. The excellent to good contrast of *Striga* radicles




Figure 1. Representative staining results of germinated *Striga* seeds. Staining time is reported in parentheses. (A) Unstained radicles; (B) Blue ink (Geha) diluted in vinegar (<5 min); (C) Lactophenol Blue solution in vinegar (<5 min); (D) Aniline Blue in vinegar (<5 min); (E) Rubin S in lactic acid (<5 min); (F) Black ink (Sheaffer) in vinegar (<5 min); (G) Toluidine Blue O in water (2 h); (H) Black ink (Sheaffer) in water (>5 h); (I) Aniline Blue in water (>5 h).

stained with Aniline Blue, Coomassie Brilliant Blue R250, Rubin S, Cotton Brown or ink is an indication that toxic compounds such as Lactophenol Blue solution can be avoided by using lactic acid or vinegar as diluents. Of the stains tested, Blue ink diluted in vinegar is highly recommended. This solution is easy to handle, readily available, gives excellent staining results and is free of known toxic ingredients. Thus, staining germinated *Striga* seeds can be stained with Blue ink/vinegar without special precautions and can be proposed for routine use in the scoring of *Striga* germination assays.

Acknowledgements

We thank Roswitha Mammerler for technical assistance. We acknowledge funding from the Eurasia-Pacific Uninet in cooperation with the Austrian Council for Research and Technology Development (Technology Scholarship to S.L.).

References

- Berner, D.K., Winslow, M.D., Awad, A.E., Cardwell, K.F., Mohan Raj, D.R. and Kim, S.K. (1997) *Striga research methods* (2nd edition). Ibadan, Nigeria, International Institute of Tropical Agriculture.
- Dhingra, O.D. and Sinclair, J.B. (1995) *Basic plant pathology methods* (2nd edition). Boca Raton, CRC Lewis Publishers.
- Gurney, A.L., Press, M.C. and Scholes, J.D. (2002) Can wild relatives of sorghum provide new sources of resistance or tolerance against *Striga* species? *Weed Research* **42**, 317–324.
- Lenzemo, V.W. (2004) *The tripartite interaction between sorghum, Striga hermonthica, and arbuscular mycorrhizal fungi*. Tropical Resource Management Papers No. 55. Wageningen, Wageningen University and Research Centre.
- Matusova, R., van Mourik, T. and Bouwmeester, H.J. (2004) Changes in the sensitivity of parasitic weed seeds to germination stimulants. *Seed Science Research* **14**, 335–344.
- Matusova, R., Rani, K., Verstappen, F.W.A., Franssen, M.C.R., Beale, M.H. and Bouwmeester, H.J. (2005) The strigolactone germination stimulants of the plant-parasitic *Striga* and *Orobanch* spp. are derived from the carotenoid pathway. *Plant Physiology* **139**, 920–934.
- Mauler-Machnik, A. and Nass, P. (1990) Einfache Methode zur Frühdiagnose von *Pseudocercospora herpotrichoides* mit dem Bayer Getreide-Diagnose System nach Verreet/Hoffmann. *Gesunde Pflanze* **42**, 130–132.
- Parker, C. and Riches, C.R. (1993) *Parasitic weeds of the world: biology and control*. Wallingford, CAB International.
- Vierheilig, H., Coughlan, A.P., Wyss, U. and Piché, Y. (1998) Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* **64**, 5004–5007.
- Wittmann, W. (1994) *Atlas der Zierpflanzenkrankheiten*. Berlin, Blackwell Wissenschafts-Verlag.
- Yoder, J.I. and Musselman, L.J. (2006) *Striga*: A subterranean parasitic angiosperm (witchweed). In *Encyclopedia of plant and crop science*. New York, Taylor & Francis.

Received 25 September 2007
 accepted after revision 25 February 2008
 © 2008 Cambridge University Press