

Evaluation of the performance of dual-purpose cows in European pasture-based systems

Name student: Gerdine Kaptijn

Period: January – July 2016

Farming Systems Ecology Group

Droevendaalsesteeg 1 – 6708 PB Wageningen - The Netherlands



WAGENINGEN UNIVERSITY

WAGENINGEN UR

Evaluation of the performance of dual-purpose cows in European pasture-based systems

Name student: Gerdine Kaptijn
Registration number student: 911227422040
Credits: 36
Code number/
name course: FSE-80436
Period: July 2016
Supervisor: Egbert Lantinga
Professor/Examiner: Jeroen Groot

Abstract

The demand for organic products in Europe is growing, especially organic products of animal origin. Organic dairy milk production requires grazing of cows, which results in a feed ration that is pasture-based. The breed that is kept by far the most for dairy production in Europe is the Holstein Friesian cow, a high-yielding dairy cow. This cattle breed has been developed in North America and was selected for its high milk yield. The disadvantages of this one-sided selection for production manifest in the health, fertility and longevity of the cow. More important, the high milk yield of the Holstein Friesian is only achieved when the cows are fed a diet of high quality and especially concentrates, which is not always possible in a pasture-based system.

In this study the alternative for the Holstein Friesian breed is considered, namely the dual-purpose cow. Most countries feature indigenous cow breeds which were traditionally kept for both milk and meat. Because of the popularity of the Holstein Friesian breed these local breeds are less used in dairy production systems. An extensive literature study was carried out to compare the popular Holstein Friesian breed to the local dual-purpose cattle breeds. This showed that local dual-purpose breeds display in many traits such as health, fertility, meat quality and longevity an advantage over the Holstein Friesian. Additionally, dual-purpose cows are better able to cope with a harsh environment and a diet of low quality.

Furthermore, the grazing behaviour of one of the Dutch dual-purpose breeds, the Dutch Friesian, was compared in the current study to the Holstein Friesian. For this purpose the monitoring system SensOor (AGIS) was used. SensOor registers the four main conducts of the cows, namely eating, ruminating, resting and other activity (such as walking, standing, etc.). The system expresses the time the cows spend on these conducts in percentages per hour or per day. In this study the SensOor system was validated under grazing conditions for the first time. Before, the SensOor system was only validated indoors or in a dirt pen. The validation under grazing conditions was done by observing the behaviour of the cows on the pasture, and comparing these observations with the SensOor output. The observation of the activity 'grazing' corresponded very well with the detection of the activity 'eating' ($R^2 = 0.97$). As the validation of SensOor under grazing conditions was successful, the output of the system was used to compare the behaviour of the two cattle breeds. In the current study no significant differences in behaviour were found, which can be explained by the differences in size of the cows (HF: 650 kg, DF: 575 kg) which have therefore different energy requirements. Another explanation could be that the high quality of the grassland on the studied farm concealed the ability of dual-purpose cows to cope with low-quality roughages.

Besides the study of the behaviour of the cows, the herbage production on two farms was estimated. This was done by the placement of grass cages on the pastures. The herbage growth under the cages was mown, weighed and analysed.

In conclusion, the monitoring system SensOor is very suitable to monitor the grazing time of cows. The literature study showed that dual-purpose cows have advantages over the Holstein Friesian (especially in milk composition, meat quality, health and longevity). Concerning their grazing behaviour, dual-purpose cows might spend more time on grazing and constitute more grass in their diet. This difference in behaviour was not shown by the output of SensOor. The absence of this difference might be explained by the excellent environment of the Friesian pastures. It is expected that the advantage concerning grass and feed intake of dual-purpose cows are better demonstrated in a harsh environment with pastures of low quality.

Content

Abstract	3
1. Introduction	6
2. Materials & Methods	8
2.1. Literature study.....	8
2.2. Participating farmers	8
Mts. Lozeman, Achterveld, Utrecht	8
Mts. Bosma, De Wilgen, Friesland	10
2.3. Grazing management on the farms.....	12
2.4. Validation of SensOor under grazing conditions	12
How does SensOor work?	12
SensOor: Support and value for farmers	12
Validation	12
2.5. Differences in grazing behaviour among breeds	14
2.6. Measuring herbage production.....	15
2.7. Impact of environmental factors.....	17
2.8. Impact of grazing on herbage production.....	19
3. Results	20
3.1. Literature study: Dual-purpose cattle vs. Holstein Friesian cattle.....	20
Cows in Europe: From local breeds to Holstein Friesian cows.....	20
Milk production.....	21
Milk composition	22
Meat quality.....	22
Fertility	23
Health	23
Grazing behaviour	24
Genotype × Environment interactions.....	24
Environmental factors	25
Conclusions	25
3.2. Validation of SensOor	25
3.3. Dual-purpose cows & grazing	27
Grazing behaviour: differences between breeds.....	28
Behaviour study during daytime grazing hours.....	28
3.4. Herbage production	32
4. Discussion	34
4.1. Literature study: Milk-type vs. dual-purpose cow.....	34
4.2. SensOor	34

4.3. Comparison of Holstein Friesian and Dutch Friesian cattle.....	35
4.4. Herbage production	36
5. Conclusions	37
Literature	39

1. Introduction

In Europe a growing demand for organic dairy products can be observed, which asks for an increase in organic production (Schaack et al., 2014). In the Netherlands the sales of organic food grew by 17% in 2011. Especially the demand for organic products from animal origin is growing in the Netherlands. 41% of all purchased organic products in the Netherlands during 2011 were of animal origin (Bakker and Brouwer, 2012). Possible reasons for this increase in demand are that organic products are perceived as more healthy and are better for animal welfare and the environment (Bakker and Brouwer, 2012; Lairon and Huber, 2014). Furthermore, the image of organic agriculture is positive as consumers know that organic systems accommodate the animal's expression of natural behaviour better, for example by grazing cows or free-range chickens.

While the market for organic products is growing at a high rate, the production of organic products in the Netherlands stays somewhat behind. Organic milk is imported from different European countries. This causes a much higher purchase price for organic milk (€52/100 kg milk) compared to the price of conventional milk (€30/100 kg milk) (Jacobsen, 2015). This price difference of €22/100 kg milk motivates conventional farmers to convert their farm into an organic system.

Organic dairy production is associated with pasture-based production, because one of the criteria for organic farming is to practice grazing (Isselstein et al., 2013). Therefore it is important that grazing cows utilize the grass as efficiently as possible. Nowadays 95% of the cows in Europe are of the breed Holstein Friesian (Van Arendonk and Liinamo, 2003), in the Netherlands this is even 98% (CRV, 2015). This commercial milk-typical cow breed is selected for its high milk yield. Selection for many years on high milk yield has led to an impressive high milk production and for this reason many farmers keep the Holstein Friesian breed on their dairy farm.

However, a high level of milk production is associated with a reduced fertility and health issues. Cows which are selected for their high milk yield tend to use the majority of their available energy for the lactation, which becomes evident after parturition (Dillon et al., 2006). After calving most cows experience a negative energy balance, because of the start of their lactation and its associated energy requirement. Holstein Friesian cows however are more susceptible to this negative energy balance. Because of this negative energy balance after parturition the fertility is low (Roche et al., 2006; Fulkerson et al., 2008; Delaby et al., 2009; Roche et al., 2009). Indications for a decreased fertility are a reduced appearance of oestrus and a longer calving interval.

When cows are kept in an organic (i.e. pasture-based) system, it is questionable whether the Holstein Friesian breed is the best option. To keep the milk production of the Holstein Friesian as high as possible in a grazing system, intensive supplemental concentrate feeding is necessary. In an extensive system such as a pasture-based system Holstein Friesians do not reach their production potential (McCarthy et al., 2007). Research shows that cattle breeds which have not been exclusively selected for a high milk yield keep a more stable production in a pasture-based system than the Holstein Friesian (Delaby et al., 2009). These cattle breeds are called dual-purpose cows, which means that they are kept for both meat and milk production (Peniche-González et al., 2014). The meat of dual-purpose cows is as profitable as their milk production, because the meat quality is high and farmers receive a better price for both calves and adult cows. Because dual-purpose breeds are not selected exclusively for milk production, dual-purpose cows have a more diverse genotype. In general they show better health, higher fertility and a better longevity than the Holstein Friesian (De Winter et al., 2010; Piccand et al., 2013).

For this reason the European project 2-ORG-COWS focuses on dual-purpose cattle breeds in Europe. The project aims to find novel genetic traits within these breeds which makes them advantageous compared to the Holstein Friesian. Instead of trying to adapt the environment to the requirements of the high-yielding Holstein Friesian, organic breeding focuses on finding a cow that fits in the local production environment. For this reason 2-ORG-COWS investigates the characteristics of the different

dual-purpose cattle breeds in Europe, and the environment of the local production systems (Kristensen, 2015).

Nine European partners are involved in the project and conduct research on farms with their local dual-purpose breeds. In the Netherlands, two farmers participate who keep the Dutch Friesian cow, a black-pied dual-purpose breed which is in part the ancestor of the Holstein Friesian. On the participating research farms a monitoring system is installed, called SensOor. This system was developed by the Dutch company AGIS.

SensOor monitors the behaviour of cows and gives an indication of animal health, fertility and nutrition. The system consists of a three-dimensional accelerometer which is placed in the cow's ear and detects the ear movements. Cows make specific head and ear movements when eating, resting or ruminating. The temperature of the individual cows and of the total herd are measured and sent to the receiver where the data is collected and stored. Most farmers use the SensOor system mainly to detect when their cows are more active (and thus might be in heat). The use of SensOor has decreased the calving interval on many farms considerably. Another important function is the early detection of ill cows by dropping temperatures or decreased eating or rumination time. The system has been tested comprehensively under indoors circumstances, but never in a grazing environment. Therefore in this thesis SensOor will be validated outdoors, and if reliable, be used as an indicator for the grazing behaviour of the dual-purpose cows on the two Dutch farms.

The hypotheses of the project 2-ORG-COWS are:

1. Dual-purpose cows are more robust than modern dairy cattle breeds, and therefore more suitable for varying harsh organic grassland systems.
2. Novel environmental descriptors combined with novel statistical methodology reveal G×E for 'conventional' and for novel functional traits.
3. Breeding goals and breeding strategies for dual-purpose cattle breeds need to be adapted for selection of suitable cattle for varying harsh organic grasslands systems, regarding to constitution, physiology, health, milk quality, greenhouse gas emissions, etc.
4. Economic evaluation criteria exhibit superiority of dual-purpose cattle over modern dairy cattle breeds in organic pasture-based production systems when using a systemic approach on the farm as well as on the population level.
5. Pasture-based production system need to be adapted (improved) to realize a better health status, less greenhouse gas emissions, and improved product quality of dual-purpose cows.

In this thesis hypothesis 1 shall be investigated. The performance of dual-purpose cows in a pasture-based system will be evaluated by answering the following questions:

- What similarities and differences exist between European dual-purpose breeds and milk-typical breeds such as the Holstein Friesian?

This question will be answered by carrying out a literature study in which the characteristics of the Holstein Friesian breed and the different dual-purpose breeds will be investigated.

- Is SensOor a reliable system to monitor the grazing behaviour of dual-purpose cows?

To answer this question the SensOor system has to be validated. In order to this cows will be observed during their daytime grazing hours and their behaviour will be noted. The observations will be compared with the output of the SensOor system.

- Does the grazing behaviour of dual-purpose cattle differ from the grazing behaviour of the Holstein Friesian?

The output of SensOor will be used to give an overview of the behaviour of the different cattle breeds. A comparison of their behaviour will show whether a difference exists in their grazing behaviour.

The aim of this study is to test the first hypothesis of 2-ORG-COWS and to investigate what differences exist between the high-yielding Holstein Friesian cow and local dual-purpose breeds.

2. Materials & Methods

2.1. Literature study

First the concept of a dual-purpose cow was defined and explained which dual-purpose cow breeds are most common in Europe. Using an explorative literature study dual-purpose cows were compared to the well-known commercial cow breed Holstein Friesian. Similarities and differences between the breeds were discussed. Many comparative studies about the characteristics and qualities of the breeds have been executed in the past. The results of these studies were shown and discussed. It was expected that dual-purpose breeds have certain advantages over the Holstein Friesian and that this will become apparent by a thorough literature study.

Literature was acquired by using Google Scholar and the online library of Wageningen University. Personal communication with experts and farmers was used as well. As less information exists on dual-purpose breeds, 'grey' literature such as non-scientific agricultural journals was consulted.

2.2. Participating farmers

Two Dutch farms participate in this research. They are situated in Utrecht and in Friesland. Both farmers have installed the monitoring system SensOor on their farm. The farmer in Friesland already uses SensOor since 2012. The farmer in Utrecht has the system since October 2015. Both farmers endorse the system as a positive asset for their management and observe an increased fertility and disease detection among their herds.

Mts. Lozeman, Achterveld, Utrecht

Farmer Lozeman in Utrecht runs an organic dairy farm of 30 ha with 56 dairy cows of the Dutch Friesian breed. During the grazing season (April-November) the cows have access to a grass-clover pasture throughout the whole day. They graze 200 days a year, 20 hours a day. In summer the cows also receive concentrates (on average 7 kg/cow/day) and in the barn there is additional grass silage available. In winter the cows are fed grass and triticale silage and some additional concentrates. The cows are milked at 5:45 and at 17:15. The milking takes approximately 1 hour. The production of the cows is 6500 kg milk per cow per year, and 19,6 kg milk per cow per day.

In Appendix I a map of the farm can be found. The black line delineates the parcels of the farm, which add up to a total of 29.7 ha. This year (2016) the parcels 11-13 were not available to the farmer as the municipality diverts the stream north of the farm for cultural purposes. All parcels of the farmer are on sandy soil on which a grass-clover mixture is grown. Each year the farmer ploughs 1 parcel where he sows triticale (a cross of rye and wheat) to produce part of the concentrate feed of the cows. This year the triticale was sown on parcel 2. Triticale is suitable for sandy soils because it is a drought-resistant crop.

In Table 1 an overview of the parcels regarding their acreage, altitude, soil type and utilization for the last 6 years is given. The numbers of the parcels correspond with the numbers on the map in figure 1. Taking into account the parcels which are not available due to excavation around the stream (11-13) and the triticale parcel (2), the total grass acreage available on the farm is $29.7 - 1.8 - 1.46 - 0.66 - 2.49 - 3.8 = 19.49$ ha.

The farmer fertilizes the pastures during the growth season with the manure of his own cows. He estimates that he uses 40 m³ animal manure per ha in total.

Table 1. Overview of the parcels on the organic farm.

#	Hectares	Crop	Soil type	Average altitude (m)
1	0.44	2009-2015: grass-clover	Sand	3.96
2	3.8	2009-2015: grass-clover	Sand	4.17
3	1.36	2009-2015: grass-clover	Sand	3.28
4	3.99	2009-2013: grass-clover 2014: triticale 2015: grass-clover	Sand	3.13
5	4.10	2009-2012: grass-clover 2013: triticale 2014-2015: grass-clover	Sand	3.33
6	2.13	2009-2015: grass-clover	Sand	3.57
7	0.75	2009-2015: grass-clover	Sand	4.08
8	1.36	2009-2015: grass-clover	Sand	3.67
9a	0.71	2009-2014: grass-clover 2015: triticale	Sand	4.02
9b	3.30	2009-2014: grass-clover 2015: triticale	Sand	4.64
10	1.19	2009-2015: grass-clover	Sand	3.98
11	1.80	2009-2013: maize 2014-2015: grass-clover	Sand	3.17
12a	1.46	2009-2013: maize 2014-2015: grass-clover	Sand	3.16
12b	0.66	2009-2015: grass-clover	Sand	3.27
13	2.49	2009-2013: maize 2014-2015: grass-clover	Sand	3.26

The organic farmer has kept the Dutch Friesian dual-purpose cows for decades. The farmer is part of a national breeders association in which knowledge is exchanged and bulls and sperm is traded among farms. The farmer tries to keep the bloodlines of his cows pure but to avoid inbreeding he needs to crossbreed. Usually he uses sperm from British Friesian bulls to inseminate his cows. The British Friesians descend from the Netherlands as well and are similar to Dutch Friesians regarding their physique and milk production.

In Table 2 an overview is given of the herd and their bloodline composition. There are 16 pure-bred DF cows, and 29 cows are cross-bred DF and BF. The remaining cows have HF blood as well, and some blood of which the origin is unknown.

Table 2. Overview of bloodline composition (%) of the herd on the organic farm.

Dutch Friesian	British Friesian	Holstein Friesian	Unknown	Number of cows
100	-	-	-	16
87.5	12.5	-	-	14
75	25	-	-	10
75	12.5	12.5	-	5
50	50	-	-	4
87.5	-	12.5	-	1
62.5	12.5	25	-	1
50	25	25	-	1
37.5	62.5	-	-	1
25	62.5	12.5	-	1
50	-	12.5	37.5	1
25	50	-	25	1
				56

30% of the herd is purebred Dutch Friesian. 68% of the cows have BF blood.

In Figure 1 the dispersion of the different bloodlines is shown.




Figure 1. Dispersion of the bloodlines of the organic herd.

Mts. Bosma, De Wilgen, Friesland

Farmer Bosma in Friesland has a conventional dairy farm and keeps 100 Dutch Friesian cows. They have 48 ha of grassland and 12 ha of maize, for the maize silage production. There has been some crossbreeding with Holstein Friesian. Approximately 15% of the herd is cross-bred with HF and 5% is purebred HF. In summer the cows spend on average 6 hours of grazing per day and are inside the barn overnight, where they are fed maize and some brewer's grains. Additionally they receive on average 4 kg concentrates per cow per day. In winter the feed ration is grass and maize silage, brewer's grains, some soy and minerals.

In Appendix II the conventional farm is shown with all grass and maize parcels. The total acreage is about 60 ha, of which 20% is used for production of maize. On 4 parcels maize is grown, on the remaining land grass is grown for grazing and silage. The parcels on the south-west side of the farm are mostly on peat soils (parcel 7-14), whereas the other parcels are on sandy soils.

The conventional farmer is situated in Friesland. The parcels of the farmer are shown in Table 3. Typical for Friesland peat soils can be found, which have in general a lower altitude than sandy soils. Besides this, peat soils have a lower water permeability compared to sandy soils (Bot, 2011). On half of the parcels the farmer is involved in nature conservation, such as protection of nesting birds and rest areas for meadow birds.

The pastures are fertilized with animal manure and artificial fertilizer. On average the pastures are fertilized with 22 m³ manure/ha. Additionally 40 kg nitrogen per ha is used.

Table 3. Overview of the parcels on the conventional farm

#	Hectares	Crop	Soil type	Average altitude (m)	Nature conservation
1	1,57	2009-2015: grass	Sand	0.24	
2	3,82	2009-2015: grass	Sand	0.04	
3	1,24	2009-2015: grass	Sand	-0.02	
2+3	1,44	2009-2010: grass 2011-2015: maize	Sand	-0.04	
4	2,75	2009-2015: grass	Sand + peat	0.19	
5	4,63	2009-2015: grass	Sand	0.01	
6	3,41	2009-2015: grass	Sand	-0.56	
7	4,08	2009-2015: grass	Peat	-0.93	Yes
8	3,67	2009-2015: grass	Sand + peat	-0.47	Yes
9	3,96	2009-2015: grass	Sand + peat	-0.73	Yes
10	2,75	2009-2015: grass	Peat	-0.91	Yes
11	4,76	2009-2015: maize	Peat	-0.77	Yes
12	1,26	2009-2013: maize 2014-2015: grass	Peat	-0.72	Yes
13	0,97	2009-2015: maize	Sand	0.18	Yes
14	5,76	2009-2015: grass	Sand + peat	-0.31	Yes
15	2,25	2009-2015: grass	Sand	-0.10	
16	2,70	2010-2015: maize	Sand + peat	-0.51	Yes

The conventional farmer has a large herd of 104 dairy cows. The majority of the cows is pure-bred Dutch Friesian, but in 2010 the farmer bought 10 pure-bred Holstein Friesians to extend the herd. This resulted in cross-bred animals. Most of the bulls that are used to inseminate the cows are DF, but HF is used as well to avoid inbreeding and to increase the milk production. This resulted in a diversified herd which is shown in Table 4 and Figure 4.

Table 4. Overview of bloodline composition (%) of the herd on the conventional farm

Dutch Friesian	Holstein Friesian	Number of cows
100		72
75	25	5
50	50	22
0	100	5
		104




Figure 4. Dispersion of the bloodlines of the conventional herd

2.3. Grazing management on the farms

On both farms the herds have access to grazing during summer. The grazing behaviour of the HF cows of the conventional farmer will be studied and compared to DF cows. To enable a fair comparison, DF cows will be selected which are comparable to the HF cows in their age and production stage. If possible, it will be determined whether there is a difference in grazing time between the breeds.

2.4. Validation of SensOor under grazing conditions

The performance of the SensOor system under indoor conditions with dairy cows was already evaluated in 2013 (Bikker et al., 2014). The behaviour of the cows was observed and these observations were compared to the SensOor output. The results proved that the output of SensOor has a very high reliability. The testing was done during winter time, when the cows are kept indoors.

Another study evaluating the performance of SensOor was done in 2013, Canada (Wolfger et al., 2015). The animals that were studied were castrated bulls in an outdoor dirt pen, which were fed at a feeding bunk. This study also supports the reliability of SensOor, although attention must be paid to the difference between the activities eating and ruminating. The study showed that SensOor was not always correct in discerning these particular activities, especially in warm environments where due to insects the cows use their ears a lot, thus compromising the data.

At the time of writing this thesis no other validation studies concerning SensOor have been published. The reliability of SensOor data of cows grazing on a pasture has not been tested so far.

How does SensOor work?

SensOor consists of a three-dimensional accelerometer which is placed in the ear of the cow. It detects the temperature of the cow, ear movements and typical head movements. From these movements the system can derive the behaviour of the cow, whether she is eating, ruminating or resting. The chip in the ear of the cow collects the data every minute and can save the data for 48 hours. Then the data is sent via a router in the barn to the computer where the data is interpreted and saved.

The data from the ear chip is interpreted by a proprietary model developed by AGIS. The data is expressed in a behavioural category, either the cow is eating, ruminating, resting or active. The minute-to-minute data is converted into percentages of behaviours per hour and per day. Through an online web application the farmer can access the information about the herd.

SensOor: Support and value for farmers

SensOor can support the farmers management by providing information about the health, fertility and nutritional status of the herd. Health issues can be detected by comparing the ear temperature of a specific cow to the average temperature of the herd. When there is a large deviation SensOor notices the farmer about this by sending a notification (Zevenbergen, 2012).

The activity of the cows is also measured by SensOor. When there is a peak in the cows activity this might indicate that the cow is in heat and can be inseminated. The system even shows the LH (Luteinizing Hormone) peak, which enables the farmer to inseminate the cow on precisely the right time (Buning, 2012).

Fluctuations in eating and rumination activity can also indicate health problems or issues with the feed of the cows. The graphs created by SensOor in the web application can provide a helpful overview of the time the cow (or total herd) spends on eating, ruminating and resting.

Validation

In this study SensOor was validated under grazing conditions. To achieve this cows were studied during their time spent in the pasture. These observations were compared to the output from SensOor to validate the performance of the system under a grazing environment. There is no research done so far on the performance of SensOor on cows while grazing outside on pasture. However, the dual-purpose cows on the farms that were studied are often outside and spend time on grazing. Therefore it is interesting to see whether the activity that SensOor registers is viable for cows that graze outdoors.

For the validation, the cows were selected at random and observed by 1 person. Due to logistical reasons the observations were made on the farm in Utrecht. 3 or 4 cows were studied at the same time.

The observations were done in the same time frames, from 11:00 to 15:00. The behaviour of the cows were noted every minute in an observation scheme. This scheme can be found in Appendix II. Later the observations were classified in one of the following categories: 'eating, ruminating, resting or active'. The observation protocol as described by Bikker et al. (2014) was followed. Using the statistical program IBM SPSS Statistics 22 (from now referred to as SPSS) the observations were analysed.

In May 2016, on the 10th and 12th, cows were observed in Achterveld, Utrecht. The weather conditions during these days are displayed in Table 5.

Table 5. Weather conditions on observation days (KNMI, 2016b).

Date	Temperature (°C)	Sun hours	Wind speed (m/s)	Humidity (%)
May 10 th	15.4-22.1	0.9	3.4	62
May 12 th	14.4-25.1	13.7	5.5	50

The cows that were observed are listed in Table 6. Their genetics, age and days in lactation on the day of observation are shown as well.

Table 6. Information on observed cows.

Observation date	Cow number	Genetics	Birth date (age)	Calving date (DIL)
May 10 th	19	87.5% DF 12.5% BF	21-02-2008 (8.02)	4-5-2016 (6)
May 10 th	33	50% DF 50% BF	20-12-2012 (3.04)	12-12-2015 (150)
May 10 th and 12 th	57	87.5% DF 12.5% BF	05-03-2011 (5.02)	06-01-2016 (125, 127)
May 12 th	43	87.5% DF 12.5% BF	21-02-2007 (9.02)	27-09-2015 (228)
May 12 th	56	62.5% BF 25% DF 12.5% HF	18-11-2013 (2.06)	24-11-2015 (170)

The observant was situated outside of the pasture which provided an overview of the cows without disturbing their natural behaviour. However due to the presence of trees and the possibility of the cows to go indoors, it was not easy to observe the selected cows for consecutive hours. For this reason the data of four of the observed cows covers 1 hour.

SensOor shows five different activities in its output, namely eating, ruminating, resting, active and highly active. It is unknown where the difference in active and highly active is based on. Therefore the output of these two activities was combined. To accommodate a comparison between the observations and the output of SensOor, the behaviour of the cows was classified into the activities that SensOor uses (Table 7).

Table 7. Observed behaviour classified into four SensOor activities.

Observed behaviour	SensOor Activity
Grazing standing	Eating
Grazing walking	Eating
Ruminating lying down	Ruminating
Ruminating standing	Ruminating
Standing	Active
Lying down	Resting
Walking	Active

The minutes they spent on grazing, ruminating, resting or activity were recalculated into percentages. This was done to enable a comparison between the output of the SensOor web application and the observations.




Figure 5. Output of SensOor for cow 57 on May 12th between 9:00 and 16:00. Blue stands for ruminating, green for eating, yellow for resting and orange and red for active.

Figure 5 shows the output of SensOor. The time that the cow spends on different activities are shown in percentages per hour. It is also possible to compute the output in days or weeks.

2.5. Differences in grazing behaviour among breeds

As it was found that SensOor is indeed reliable under grazing circumstances (see Results), the data of SensOor was used to examine whether dual-purpose cows (Dutch Friesian) spend more time on grazing than commercial, milk-typical breeds (Holstein Friesian). Within the cow herds of the farms variation exists in the bloodlines, which makes it possible to compare the grazing behaviour of the cows within the herds. Although the sample size is quite small, it will be examined whether the behaviour of a 100% DF cow is different from a 100% HF cow.

Cows were selected from the farm in Friesland, as on this farm 5 purebred HF cows are present. Four of the HF cows are in their 4th or 5th lactation. The other HF cow is a heifer. The behaviour of these cows was compared to 5 DF cows. These cows were selected based on their age and lactation stage, so they were comparable to the HF cows. For every HF cow of a certain age and lactation stage a DF cow was selected who had similar features. Information on age, yield and lactation stage of the cows can be found in Table 8. This information was collected on May 26th, 2016.

Table 8. Information on 10 cows (5 Holstein Friesian, 5 Dutch Friesian) whose behaviour was studied.

	100% HF	100% DF
Cow number:	33	81
Birthdate:	Unknown	10-02-2011
Age (years):	Unknown	5.03
Yield (kg/day):	39.6	31.8
Lactation number:	5	4
Calving date:	20-01-2016	19-01-2016
Days in lactation:	127	128
Concentrates (kg/day)	6	2
Concentrates/kg milk	0.15	0.06
Cow number:	63	42
Birthdate:	Unknown	23-02-2007
Age (years):	Unknown	9.03
Yield (kg/day):	28.7	18.1
Lactation number:	4	6
Calving date:	22-05-2015	03-06-2015
Days in lactation:	370	358
Concentrates (kg/day)	8	0.5
Concentrates/kg milk	0.28	0.03
Cow number:	75	80
Birthdate:	17-12-2013	07-12-2013
Age (years):	2.05	2.05
Yield (kg/day):	38	20.3
Lactation number:	1	1
Calving date:	09-02-2016	12-02-2016
Days in lactation:	107	104
Concentrates (kg/day)	7	4
Concentrates/kg milk	0.18	0.20
Cow number:	94	36
Birthdate:	02-08-2008	09-11-2009
Age (years):	7.09	6.06
Yield (kg/day):	27.3	19.9
Lactation number:	5	5
Calving date:	16-12-2015	08-12-2015
Days in lactation:	162	170
Concentrates (kg/day)	3	4
Concentrates/kg milk	0.11	0.20
Cow number:	106	43
Birthdate:	20-10-2008	10-10-2010
Age (years):	7.07	5.07
Yield (kg/day):	19.3	24
Lactation number:	5	4
Calving date:	27-09-2015	30-09-2015
Days in lactation:	242	239
Concentrates (kg/day)	2	1
Concentrates/kg milk	0.10	0.04

The behaviour of the breeds will be compared. If differences are observed, factors such as the weather conditions and the amount of fed concentrates will be considered as a possible explanation for deviations.

2.6. Measuring herbage production

To measure the total grass production of the farms grass cages were placed on parcels that are grazed during the summer. The cages were placed to prevent the cows from eating the grass. The grass is mown every 5 or 6 weeks. Hereafter the grass was weighed and analysed. Because the dimensions of the cages are known (1.5 by 4 meters or 1.25 by 4.2 meters) the total grass production of the parcels

can be calculated. The cages were placed in Friesland on April 12th, and in Utrecht on April 13th. On each farm 6 parcels were measured by 2 cages. In total 12 cages were placed per farm.

In Figures 6a and 6b the locations of the grass cages are shown. The red dot indicates where a grass cage was placed. On every parcel the two cages are marked with A and B to differentiate between the cages. On the organic farm (Figure 6a) the cages were placed on parcel 3, 4, 5, 6, 9b and 10. Parcel 3, 6 and 10 are grass-clover parcels since 2009. On parcel 4, 5 and 9b triticale was grown in respectively 2014, 2013 and 2015.

On the conventional farm (Figure 6b) the cages were placed on parcel 2, 4, 5, 6, 8 and 9. The parcels which are more to the east (2, 5 and 6) are on sandy soils, parcel 4, 8 and 9 are partly on peat soils. All these parcels have been used for grass production since 2009.



Figure 6a.



Figure 6b.

The grass was mown on both farms on May 17th and on June 30th, 2016. The mowing was done by lifting the cages and mowing a strip of 0.9 m wide along the entire length of the cage. Depending on the type of the cage this was either 4 or 4.2 meters. The surface that was mowed per cage was therefore 3.6 m² or 3.78 m². The grass that was mown from this strip was collected in baskets and weighed on a scale. A random sample was taken from the grass which was analysed later to determine the grass quality. The dry matter content was determined by drying these samples overnight on 70 °C. From the dry matter content the DM production in kg/ha was calculated. As the acreage of the parcels is known, the total DM production per parcel was calculated. The grass cages were not placed on every parcel on the farms. To estimate the production of the remaining parcels the average DM production of the grass cages was taken. Crude protein content, digestibility and fiber content was determined by a chemical analysis.

Forage quality is influenced by many aspects. Palatability, animal uptake, digestibility and nutrient content are the main factors which determine forage quality. A good palatability is associated with high feed intake by the animal. The digestibility of the grass that is taken up is determined by the maturity of the grass. Young leaves can be digested for 80-90%, while older plant materials such as the stem are digested for less than 50% (Ball et al., 2001).

Neutral detergent fibers (NDF) comprise the parts of the grass which are difficult to digest, mainly the fibers (Oba and Allen, 1999). The poor digestible fibers such as cellulose and lignine are ranked under the acid detergent fibers (ADF). When the NDF content of forage grass is below 50%, this is considered good quality. When NDF exceeds 60% the forage quality is below average. ADF is an appropriate indicator for grassland quality as well because it contains the especially poor digestible fiber cellulose and lignine. For a good quality forage an ADF content below 35% is desired (Van Saun, 2006).

2.7. Impact of environmental factors

The netto primary production of grass can be influenced by adequate farm management. However, climatic factors are beyond the control of the farmer. The potential impacts of the climatic environment on grass growth are elaborated below.

Temperature can be a limiting factor for grass growth. In an experiment done by Kummerow and Ellis (1984) it was shown that arctic grasses will grow at 2 °C, but their biomass production is enhanced when temperature is increased to 12 °C. This indicates that although grass survives low temperatures, a higher air temperature increases grass growth and therefore the biomass production.

Thornley and Cannell (1997) discovered that the net primary production of a temperate grassland increases with rising temperatures, as long as the sward is ungrazed. Pastures that are grazed respond differently to an increased temperature, because the leaf area index decreases fast as the grass leaves are eaten by livestock. A high leaf area index is associated with a higher primary production.

The production of the grass in the grass cages will not be influenced by grazing, so it is expected that a higher air temperature will have a positive effect on grass growth in the cages. However, the grass outside the cages will be subject to grazing. The fact that grazing affects primary production should be taken into account when estimating total forage production of the farms.

The amount of precipitation and the water holding capacity of the soil influences the grass production as well (Sala et al., 1988). The authors state that in cases where the precipitation per year exceeds 370 mm, a high water holding capacity increases the production of the grassland. Below 370 mm per year the water holding capacity of the soil has a negative effect on primary production. As the average annual precipitation in the Netherlands is 792 mm (KNMI, 2016a), the water holding capacity of the soils of the pastures will have a positive effect on grass production. The soils of the farm in Utrecht are all sandy soils. In Friesland some of the parcels have partly peat soils. According to Bot (2011) peat soils are less water permeable than sand soils (0.001-0.1 m vs. 1-10 m in sand). The water holding capacity is therefore higher in Friesland than in Utrecht. It is expected that the parcels which have a partly peat soil have a higher grass production than the sandy soils. These parcels are 4, 8 and 9.

Climatic factors such as rainfall, temperature and sunshine hours have shown to influence grass growth. Therefore the meteorological information of the farms was collected. Using maps of the Dutch meteorological institute (KNMI) the rainfall, temperature and sunshine was noted during February, March, April, May and June of 2016. For the assessment of rainfall on the farm in Utrecht the data from the precipitation station in Barneveld was used. Concerning temperature and sunhours, the data from weather station 'De Bilt' was used. In Friesland the weather stations in Drachten (rainfall) and Leeuwarden (temperature and sunhours) were consulted.

The data was entered in Excel and visualised in graphs. This meteorological data might be used to explain differences in grass growth and for differences in the behaviour of cows. In Figure 7, 8 and 9 the data of respectively rainfall, sunshine hours and temperature are shown.




Figure 7. Total rainfall in mm during February, March, April, May and June on research farms.

During these five months the total rainfall in Friesland was 450 mm. In Utrecht this was slightly more, 494 mm. The difference in rainfall between the two locations was not significant. During the month April more rain fell in Friesland (105 mm, vs. 91 mm in Utrecht). In all other months the rainfall was higher in Utrecht. The month June was very wet, both in Friesland and in Utrecht 37% of the precipitation during February-June fell in June (165 and 182 mm, respectively).




Figure 8. Sunshine in hours on average per day during February, March, April, May and June on research farms.

During these five months the total sunhours in Friesland were 893. In Utrecht this was less: 817 hours. The difference in hours of sunshine over the five months between the two locations was not significant. Except for the month April, the sun shone more in Friesland than in Utrecht. This might be connected to the amount of rainfall, as it rained more in Utrecht than in Friesland. Usually in case of rainfall the sky is cloudy and there is less sunshine.




Figure 9. Average daily temperature during February, March, April, May and June on research farms.

The graphs in Figure 9 look very similar. However, during the five months that were studied the temperature was significantly higher in Utrecht. During April and in the first half of June the higher temperature in Utrecht can be clearly observed. In Friesland the daily temperature during the five months was on average 9,2 °C, whereas in Utrecht this was 10 °C. This difference in temperature might partly explain the higher yield of grass on the pastures of Lozeman in Achterveld.

For the evaluation of the grazing behaviour of the Dutch Friesian and Holstein Friesian cows in Friesland, the weather conditions of 9-13 May were studied in more detail. Relative humidity data was collected from the KNMI website. Using the ambient temperature and relative humidity data the temperature humidity index (THI) was calculated. This was done with the same method used by Charlton et al. (2011):

$THI = (1.8T + 32) - [(0.55 - 0.0055RH) \times (1.8T - 26)]$ in which T is the ambient temperature and RH is the relative humidity.

2.8. Impact of grazing on herbage production

Grasses can cope with grazing very well, as their leaves are produced quickly and abundantly. However, leaves are necessary for photosynthesis which is crucial for plant growth and respiration. When due to grazing too much leafy material is removed, grass growth is substantially decreased (Trlica, 1992). Parsons et al. (1983) state that a sward under intensive grazing has lower photosynthesis. According to Milchunas et al. (1988) the impact of grazing on a grassland depends heavily on the type of grassland, the grass species and the grazing intensity. Some grasses are very resistant to trampling. Other grass species show increased tillering and formation of rhizomes and stolons under grazing circumstances. Without grazing species diversity is decreased because dominant species are allowed to flourish.

3. Results

3.1. Literature study: Dual-purpose cattle vs. Holstein Friesian cattle

In this chapter an overview will be provided of the main European dual-purpose breeds and how they compare to the production performances of the Holstein Friesian cow.

Cows in Europe: From local breeds to Holstein Friesian cows

For centuries cows have been kept as a husbandry multi-purpose animal. The cow provided milk, meat, draught and leather for the farmer and his family (Syrstad, 1993). After World War II the Netherlands faced the challenge of reconstruction of their society and economy. This induced Dutch farmers to modernize their farms and increase their production. This was achieved by upscaling and mechanization of the farms which lead to specialisation while focusing on milk production in cows. The authorities played an active role in this process by creating a steady milk price and investing in dairy advisors and research (Theunissen, 2012). Thus, dairy cows have been selected for their high milk yields, resulting in a shift from the original Friesian black-pied cow to the highly productive Holstein Friesian cow. These Holstein Friesians were imported from the United States or Canada and have an astonishing high milk production, up to 10,000 kg milk/cow/year. In Europe the Holstein Friesian is commonly kept in an intensive system. This means that the cows are kept indoors during winter and are fed supplemental feeds such as maize, brewers grains and concentrates. Furthermore they are almost all fertilized by artificial insemination and calve year round (Van Arendonk and Liinamo, 2003). The same development was observed in other western European countries. Due to the new possibility of artificial insemination many European farmers crossbred their local cattle breeds with the milk-typical Holstein Friesian, or with more beef-producing cattle breeds such as the Belgian Blue (Hiemstra et al., 2010). In this way the process of specialisation of farms progressed quickly. Nowadays, the majority of European dairy farmers keep the Holstein Friesian, as 95% of the European cows is predominantly Holstein Friesian (Van Arendonk and Liinamo, 2003).

Due to the unilateral selection for milk production, other important physiological aspects of the Holstein Friesian have been neglected or at least have been underexposed. Commercial dairy cows such as the Holstein Friesian are susceptible to claw and udder infections. The drop in their body reserves after calving (negative energy balance) is severe, resulting in a low fertility. Genetic research shows that selecting for high milk yields might be correlated with low fertility and an increased prevalence of mastitis, milk fever and claw problems (Pryce et al., 1997). Royal et al. (2002) showed that it is feasible to select for fertility traits as they are correlated to milk yield. Because of the widespread use of the Holstein Friesian, the genetic variation in the total dairy cattle population has become low. Medugorac et al. (2009) therefore emphasize the importance of the conservation of local cattle breeds, arguing that these breeds are well adapted to specific environmental conditions and have unique production features. These genetic traits could be used to diversify and improve the current European cattle population if needed.

Local cow breeds were originally not only selected for their milk yield, but also for their meat production. Cows which are kept for both milk and meat production are called dual-purpose cows (Syrstad, 1993). Usually these breeds are indigenous and are in general more robust, fertile and strong. Their milk production is much lower than commercial cow breeds but they are also much less susceptible to infections and diseases, due to their better health. Because dual-purpose cows are local breeds, they are different in every country. Their positive and negative traits differ therefore per breed (Medugorac et al., 2009). The most well-known dual-purpose breeds in the Netherlands are Dutch Friesian, Groninger Whitehead and MRY (Meuse, Rhine and Yssel) cattle. These breeds are shown in Figure 10.



Figure 10. The Dutch dual-purpose breeds from left to right: Dutch Friesian, Groninger Whitehead and MRY.

In this chapter the main differences between the high-yielding Holstein Friesian and the European dual-purpose breeds will be discussed.

Milk production

The main and most striking difference between the Holstein Friesian and the dual-purpose cow is their milk yield. In an Irish study by Dillon et al. (2003) the Holstein Friesian was compared to the dairy breed Irish Holstein and to the French dual-purpose breeds Normande and Montbeliarde. All cows were kept in a pasture-based system and were fed the same diet of concentrates and silage. The milk production of the HF was much higher (5994 kg/cow) than the Normande (4561 kg/cow) and Montbeliarde (5119 kg/cow).

Another study in which cow breeds in a pasture-based system were compared was executed in Austria by Horn et al. (2013). They compared the performance of the Austrian Brown Swiss (which has been selected for milk yield) and a Holstein strain which was bred for longevity and fitness under low-input conditions. This Holstein type (HFL) is used as a dual-purpose breed in Austria. The Brown Swiss had a higher milk yield of 6595 kg than the HFL breed which produced 5616 kg.

Haiger and Knaus (2010) compared conventional Holstein Friesians to the same HFL as mentioned above and to the dual-purpose Simmental breed. The Simmental had a production of 6027 kg and the HFL 6145 kg, whereas the Holstein Friesian produced 7567 kg.

It is clear that the breed Holstein Friesian excels in milk production compared to dual-purpose breeds. Even under low-input conditions in a mostly pasture-based system the milk production of the Holstein Friesian proves superior, although their full potential is not reached in that case. A comparative French study carried out by Delaby et al. (2009) reviewed the performances of Holstein Friesian cows and the dual-purpose breed Normande under different feeding strategies. The milk yield of the HF was again higher than those of the Normandes. However, it was shown that the milk production of HF is much more dependent on the quality of the feed than those of Normande. When the diet was changed from high to low supplementation of concentrates, the milk production and body condition score dropped dramatically for HF. This change was observed in the Normande cows as well, but not so strong. Reversed, when the amount of concentrates was increased, the milk production of the HF increased much faster than that of the Normande.

Due to the fact that HF are bred for their milk production, their body condition score tends to be low. Cows with high genetic merit for milk yield spend their energy on milk production, even at the expense of their own body reserves (Buckley et al., 2000; Dillon et al., 2003; Fulkerson et al., 2008). Especially after calving this genetic feature leads to a negative energy balance. This is usually solved by the farmer by strongly supplementing the diet of the HF with concentrates.

Milk composition

A high milk production is one of the most important aspects of a cow, in which the HF breed is superior over dual-purpose breeds. However, the quality of the milk is determined by the composition of the milk, for example the fat and protein content. It has been shown that protein and fat percentages in the milk are negatively correlated to milk yield (Syrstad, 1993; Fulkerson et al., 2008). Therefore dual-purpose breeds which have a lower milk yield produce milk of a higher quality. The breeds that on average have a higher protein content than the HF are the MRY breed (3.75%) (Hiemstra, 2015), Normande (3.60%) and Montbeliarde (3.49%) (Dillon et al., 2003).

The local Holstein strain of New Zealand is often used as a dual-purpose breed and was compared to the American Holstein Friesian by (Roche et al., 2006). They observed that the protein and fat content of the New Zealand Holstein is consistently higher throughout the lactation. Interestingly, by supplementing the diet of the cows with concentrates the fat content for both breeds was decreased. At the other hand, the protein content of the milk increased.

However, fat and protein contents are not necessarily higher in every dual-purpose breed. According to the research done by Renna et al. (2014) among the Italian red-pied Aosta breed the protein content ranged from 3.28-3.35%. This might be explained by the fact that the milk composition is not only genetically determined, but is also influenced by feed regimes (Roche et al., 2006), lactation stage, season (Palmquist et al., 1993) and health status of the cow (Auld et al., 1995). So although the negative relationship between high milk yield and functional components in the milk is proven, it is still possible to influence the milk quality by managing the ration carefully and assure proper cow health.

Meat quality

Besides milk production the dual-purpose cow is bred for meat production. Beef quality in dual-purpose breeds is determined by the daily weight gain of the animal, the SEUROP classification (European meat quality assessment system) and the meat percentage (Schild et al., 2003). In their study about the Dutch dual-purpose breed Groninger Whitehead De Winter et al. (2010) underline the superior carcass quality of dual-purpose cows when they are too old for milk production. Within the SEUROP classification the meat of dual-purpose cows fits on average in the U-class ('very good'). In contrast, the HF carcasses score usually in the O ('fair') or P class ('poor').

This difference can be explained by the composition of the animals. Where HF cows for years have been selected for their high milk production, dual-purpose cows were bred for their meat production as well. The German dual-purpose breeds Fleckvieh and Gelbvieh proved to have much better carcass and meat quality than the dairy types Braunvieh and HF.

Besides getting a better price for old dairy cows, farmers receive also a higher price for the bull calves when sold to a fattener because they grow faster (Geuder et al., 2012). This is confirmed by the Dutch farmers involved in this research, as they name the better price they get for their bull calves as a benefit of their dual-purpose cows. In Norway two genetic lines of the dual-purpose breed Norwegian Red were compared. One line was selected for high milk production while the other served as a control line. There were no differences in carcass quality, although it became clear that it is very hard to improve carcass quality while selecting for milk production traits (Aass and Vangen, 1997).

Monsón et al. (2005) compared the carcass and meat quality of 4 different cattle breeds. Among these breeds were the Holstein Friesian and the dual-purpose breed Brown Swiss. The slaughter live weight, dressing percentage and SEUROP score were all higher for the Brown Swiss. The Holstein Friesian had a much higher fat content which is not favourable in beef products. Overall consumers liked the

meat of the Brown Swiss better than the Holstein. In comparison however, the other breeds (which were bred for meat production only) ranked better than the Brown Swiss.

Fertility

One of the most important aspects of dairy cows is their fertility. Reproduction is crucial for the cow's milk production and the herd replacement. Optionally, the sale of young stock can be an additional form of income for the farmer. Economically, the fertility of the herd has a direct effect on the income of the farmer. A low pregnancy rate results in more inseminations and a longer calving interval, which gives higher costs and a lower yearly milk production. For example, the overall income difference between a farm with a conception rate of 45% and 60% can be 10% (Boichard, 1990)

It is well-known that there is a negative genetic correlation between high milk yield and fertility. When cows are intensively selected for milk yield, a decline in reproductive performance is to be expected. Although this correlation is strong, the cows diet has an important effect on fertility as well (Pryce et al., 2004). After parturition a reduced ability to resume cyclicity is observed in Holstein Friesians because of their high genetic merit for milk and functional components yield (Carthy et al., 2016). This ability to resume cyclicity is positively correlated with body condition score and carcass with better conformation.

In a comparative study of Piccand et al. (2013) of the HF with Fleckvieh in a seasonal calving system it was shown that the Fleckvieh cattle perform much better than the HF. Because it takes longer for the HF to get pregnant, the HF is unsuited for a seasonal calving system. (Piccand et al., 2013) suggests that the higher body condition score of the Fleckvieh cattle accounts for the increased fertility of the cows compared to the HF cows.

This is supported by Fulkerson et al. (2008) who state that the fertility of a cow is closely related to the animal's energy balance. When the energy balance is negative (which happens often immediately after calving), the fertility of the cow is greatly reduced. This drop in energy balance happens in all cows after parturition, but is exacerbated in cows with a high genetic merit for milk yield (Pryce et al., 2004).

To improve the fertility of cows it is advised to avoid exclusive selection for milk yield. Because fertility traits do not have a high heritability, it is possible to improve fertility indirectly by selecting for health traits in cows. Berry et al. (2003b) suggest selection for body condition score, as this trait is associated with a good fertility. After all, when cows are in a good physical condition, the conception and calving interval are in general shorter (Jílek et al., 2008).

Health

Dual-purpose cows are perceived as more robust than HF cows due to their extra muscle development. Dual-purpose cows usually are ranked with a high body condition score (BCS). According to (Roche et al., 2009) the BCS can be used as a measure for health. Cows with a good BCS are less susceptible to infectious diseases and are more fertile. In addition, Rössler et al. (2013) examined the breeding goals of German cattle breeders who kept either Braunvieh or Hinterwälder. For both breeds the breeders underlined the robustness of the breed which prevents leg and claw problems and disease in general.

De Winter et al. (2010) describe in their report on Groninger Whitehead cows the strong legs and low veterinarian costs of the Whitehead, due to their better health. Many farmers who keep a dual-purpose breed are pleased with the good resistance of their herd and the low treatment costs (Hiemstra, 2015)

The good health and associated good fertility of dual-purpose cows are an incentive for farmers to crossbreed their HF herds with dual-purpose breeds (Van Ginneken, 2010).

Cows that are more robust live longer. Ray et al. (1992) showed that cows in their fourth and fifth lactation have their highest milk production. Many Holstein Friesian cows in the Netherlands never reach this number of lactations. According to CRV (2015) the average age of a Holstein Friesian in the Netherlands is 3.10 years, whereas the average age of the dual-purpose breeds MRY, Dutch Friesian and Groninger Whitehead is respectively 4.03, 4.04 and 4.02 years.

Grazing behaviour

Much studies have been executed concerning the diet and grazing behaviour of cows. The substitution rate plays an important role, which is defined as 'the decrease in pasture intake per kilogram of supplemental feed' (Bargo et al., 2002). This substitution rate indicates to what extent cows acquire their energy requirements from concentrate feed or from pasture. Several studies on cattle's grazing behaviour show that when concentrates are increased in the diet, the grazing time and fresh grass intake decreases (O'Connell et al., 2000; Bargo et al., 2002; McCarthy et al., 2007). According to Gibb et al. (1999) the grazing time of cows increases when the sward surface height decreases. Furthermore lactating cows display a higher grass intake, which is explained by their higher energy requirements.

A comparative study in Ireland of 3 HF strains (highly productive, highly durable and New Zealand) in 3 different feeding systems showed that genetics are related to grazing behaviour (McCarthy et al., 2007). In all 3 feeding systems the New Zealand strain grazed longer than the other HF cows. This can be explained by the New Zealander breeding programs for their cows. NZ cows are selected for fat and protein concentrations in the milk, a good feed conversion and the potential to produce well on low quality pastures.

Holstein Friesians which have been selected for high milk production, show consistently the largest response to supplementation of their grass-based diet by concentrates. This suggests that milk-typical cows cannot achieve their full potential under exclusively grazing conditions (Kennedy et al., 2003; Dillon et al., 2006; McCarthy et al., 2007; Delaby et al., 2009). At the other hand, high milk yield is positively correlated to feed intake. Therefore, Holstein Friesian cows consume more grass or other roughages (Oldenbroek and Van Eldik, 1980).

In their report on the Groninger Whitehead De Winter et al. (2010) describe the ability of this dual-purpose breed to feed on nature grass (up to 40% of their diet) without excessive loss in milk production. This breed is particularly suitable for nature conservation or extensive dairy production systems. Other dual-purpose breeds are also known for their ability to cope with forage of low quality.

Genotype × Environment interactions

Evidence exists that the performance of a cow is influenced by the environment which is called Genotype×Environment (G×E) interaction. This G×E interaction means that animals with a certain genotype may react differently to a certain environment than animals with a different genotype would in that same environment (Berry et al., 2003a). A Holstein Friesian cow, which has high genetic merit for milk production, might not display a high milk production because of an unfavourable environment (Dillon 2006). The environment in this case can differ in variable ways. For example, a study on the topic of G×E interaction compared HF cows in both organic and conventional systems (Nauta et al., 2006). In the organic system the milk yield of the cows was lower, showing the G×E interaction. Another study compared different feeding regimes. No significant interaction was found between genotype and the feeding regimes in the study, however there are indications that the composition of the diet has substantial influence on the performance of high-producing HF cows (Veerkamp et al., 1994). In a study where American HF and New Zealand HF cows were fed two

different diets, namely a grass-based diet and a total mixed ration (TMR), it was shown that American HF cows perform much better on a TMR diet than New Zealand HF cows. Conversely, the New Zealand HF cattle performed better on a grass diet than the American cows did (Kolver et al., 2002). Another study from New Zealand (Macdonald et al., 2008) showed that although the genetic merit for milk yield and milk solids has been increased in the American Holstein Friesian breed during the years, their production performance in a pasture-based system is not superior over the local New Zealand breed. Moreover, under pasture-based conditions their body condition score and fertility is lower which makes the local New Zealand breed more fit for a pasture-based system. Only when supplemented with concentrate feed the genetic merit of the Holstein Friesian cow became evident. Other factors than production system and diet that may affect G×E interaction are a hot or cold climate and herd size (Zwald et al., 2003).

Environmental factors

Environmental factors influence the cows performance and production as well. It is well known that cows are susceptible to heat stress. When the ambient temperature surpasses 26 degrees, cows experience heat stress because they cannot cool down themselves. Heat stress leads to a severe reduction in production (Kadzere et al., 2002). However, Sharma et al. (1983) found that Jersey cows are better resistant to heat stress compared to Holstein cows. This could be due to their difference in size. In general dual-purpose cows are smaller than HF cows, which suggests that dual-purpose breeds are less affected by heat stress.

The slope of the pasture increases the energy that the cow needs for maintenance (Knaus, 2016). This can take 50% of their energy. Coulon and Pradel (1997) discovered that an increased walking of cows decreases their milk production. It is suggested that when cows are on a steep or sloped pasture they walk more and spend more of their energy on walking than on production.

Another factor which influences cows to go outside on the pasture is the distance to the meadow. The closer the pasture, the more likely the cows will go graze outside. However, in general cows prefer to graze outside. When given the choice, cows were outdoors 57% of their time (Charlton et al., 2013). Cows on pasture experience less claw problems, but lose more weight and produce slightly less compared to cows which are housed indoors all day long (Hernandez-Mendo et al., 2007).

Conclusions

In this chapter the differences between European dual-purpose breeds and the milk-typical Holstein Friesian have been discussed. Although the Holstein Friesian breed has a very high milk production, it has also been shown that concentrate supplementation is needed to actually achieve their high production potential. The occurrence of G×E interaction might explain this dependence on an optimal diet. Therefore, in a pasture-based system where feed quality differs along the season, Holstein Friesians cannot perform to their full potential. Dual-purpose cows show a lower but stable milk production with high fat and protein content, while also being fertile, healthy and able to cope with fodder of low quality. These traits make them more fit for a low input dairy production system.

3.2. Validation of SensOor

The minutes the observed cows spent on grazing, ruminating, resting or activity were recalculated into percentages. This was done to enable a comparison between the output of the SensOor web application and the observations. The observations and output of SensOor are shown in Table 7.

Table 7. Observation of grazing behaviour vs. output SensOor.

May 10th		Number of minutes				Percentages			
11:00-12:00		graz	rum	rest	active	Graz	rum	rest	active
Cow 19	Observed	9	27	22	2	15%	45%	37%	3%
	SensOor					22%	65%	6%	8%
Cow 33	Observed	43	16	0	1	72%	27%	0%	2%
	SensOor					73%	27%	0%	0%
12:00-13:00									
Cow 57	Observed	57	0	0	3	95%	0%	0%	5%
	SensOor					94%	3%	0%	3%
May 12th									
11:00-12:00		graz	rum	rest	active	Graz	rum	rest	active
Cow 56	Observed	30	30	0	0	50%	50%	0%	0%
	SensOor					63%	21%	0%	16%
Cow 43	Observed	0	8	52	0	0%	13%	87%	0%
	SensOor					3%	11%	80%	7%
12:00-13:00									
Cow 43	Observed	32	14	4	10	53%	23%	7%	17%
	SensOor					55%	23%	5%	17%
13:00-14:00									
Cow 57	Observed	60	0	0	0	100%	0%	0%	0%
	SensOor					98%	0%	0%	2%

All observations were entered in Excel and linked with the corresponding output from SensOor. This generated a scatter plot which shows the similarities between the visual observations and the detected behaviour by SensOor. The scatter plot of the grazing behaviour is shown in Figure 11.




Figure 11. Scatter plot of SensOor output and visual observations of the grazing behaviour of 6 different cows.

The trendline and its R^2 (0.97) in Figure 11 indicate that the visual observations have a strong correspondence with the output from SensOor. When in the field the cows were observed as grazing,

SensOor detected this activity as eating. However, to assess whether there exists a significant resemblance between the observations and SensOor, statistical tests are necessary. Therefore a 2-sided paired t-test was performed using SPSS. The output is shown in Table 8 and 9.

Table 8. [Output SPSS - Paired Samples t-test: Correlation] Correlations between the observations and output SensOor.

		Correlation (r)	Sig. (p)
Pair 1	Grazing Observed Grazing SensOor	0.99	0.00
Pair 2	Ruminating Observed Ruminating SensOor	0.77	0.05
Pair 3	Resting Observed Resting SensOor	0.94	0.00
Pair 4	Active Observed Active SensOor	0.50	0.25

A positive correlation exists between the observed and detected activities of grazing, ruminating and resting ($p \leq 0.05$, Table 8). For the behaviour 'Active' this positive correlation was not significant ($p > 0.05$). A significant positive correlation means that the two variables are associated with one another. When for example the variable 'Grazing Observed' has a high value, the variable 'Grazing Observed' will also have a high value.

Table 9. [Output SPSS - Paired Samples t-test] Comparison of observed behaviour to detected behaviour by SensOor.

		Mean (%)	Std. Error Mean	t	Df	Sig. (2-tailed)
Pair 1	Grazing Observed Grazing SensOor	-3.3	2.0	-1.7	6	0.15
Pair 2	Ruminating Observed Ruminating SensOor	1.1	5.4	0.2	6	0.84
Pair 3	Resting Observed Resting SensOor	5.7	4.3	1.3	6	0.24
Pair 4	Active Observed Active SensOor	-3.7	2.4	-1.5	6	0.18

The paired samples t-test tests whether the difference between the paired variables is significantly different. For all 4 activities the differences between observation and SensOor output were not significantly different ($p > 0.05$, Table 9). Therefore the observations in the field and the detection output of SensOor are well comparable. From this we can conclude that the monitoring system SensOor presents a reliable image of the behaviour of cows under grazing conditions. Especially regarding the connection between eating and grazing: when SensOor detects the behaviour 'eating' when the cow is on pasture, it is certain that the cow is grazing.

3.3. Dual-purpose cows & grazing

It was confirmed that SensOor is reliable under grazing conditions. To assess a difference in behaviour between Holstein Friesian (HF) and Dutch Friesian (DF) cows their SensOor output was collected and compared.

To accommodate comparison between the breeds, the individual cow data was averaged and the average differences between production and diet of the breeds were calculated. This is shown in Table 10. Although the cows are all comparable concerning their age (lactation number), it is shown that the DF cows overall produce less and also receive less concentrates. To achieve their milk yield, DF cows

need less concentrates per kg milk compared to the HF cows. This is shown by calculating the amount of concentrates that the cows need to produce 1 kg of milk.

Table 10. Average information on the 10 selected cows.

	Holstein Friesian	Dutch Friesian	Difference
Lactation number	4	4	0
Milk yield (kg/day)	30,6	22,8	-7,8
Concentrates (kg/day)	5,2	2,3	-2,9
Concentrates per kg milk (kg)	0,17	0,10	-0,07

Grazing behaviour: differences between breeds

The behaviour in percentages of the cows during 7 days (May 9th – 15th) was gathered from the SensOor web application and entered in Excel. An average of these 7 days for each behaviour for each cow was obtained. For example, on average HF94 spent 23.9% on eating, 39.4% on ruminating, 30.9% on resting and 6% on activity during the 7 studied days. In the same way an average of the behaviour of the other cows was obtained using Excel.

Using SPSS, an independent samples t-test was executed. The 4 test variables were the 4 different behaviours in percentages and the grouping variable was the breed (HF or DF). The output is shown in Table 11.

Table 11. [Output SPSS - independent samples t-test] Comparison of the behaviour of 5 HF and DF cows during 1 week.

	Breed	Mean (%)	Std. Error Mean	t	df	Sig. (2-tailed)
Eating	HF	24.1	2.2	0.17	8	0.87
	DF	23.5	2.9			
Ruminating	HF	38.6	1.6	0.79	8	0.45
	DF	36.7	1.9			
Resting	HF	29.0	2.3	-0.80	8	0.45
	DF	31.4	1.9			
Active	HF	8.4	1.8	-0.17	8	0.87
	DF	8.8	1.4			

Equal variances were assumed. The output of the test shows that there are no significant differences between the eating, ruminating, resting and activity behaviour of the cows between breeds during 1 week.

Behaviour study during daytime grazing hours

The test before studied the behaviour of the two cow breeds during a week. However, the cows on the farm in Friesland are only outside between 9:00 in the morning and 16:00 in the afternoon. Therefore their pasture behaviour can only be studied during these hours, as SensOor does not distinguish between grazing and eating. Eating behaviour outside of grazing hours means that the cows are eating at the feed bunk. The behaviour of the cows during their 7 hours on pasture was averaged and interpret in the same way as before. This was done for the days May 9th -13th. In the test for May 13th the output of cow HF75 was disregarded, as this cow was in heat and did not show her normal behaviour. The corresponding cow DF80 was also excluded. The output of the test of the cow's behaviour on May 12th is shown as an example in Table 12.

Table 12. [Output SPSS - independent samples t-test] Comparison of the grazing behaviour of 5 HF and DF cows during 7 hours of pasture on May 12th.

	Breed	Mean (%)	Std. Error Mean	t	df	Sig. (2-tailed)
Grazing	HF	33.5	2.2	-2.80	8	0.02
	DF	42.1	2.1			
Ruminating	HF	32.2	1.3	2.51	8	0.04
	DF	26.1	2.0			
Resting	HF	24.5	3.3	0.29	8	0.78
	DF	23.1	3.4			
Active	HF	10.0	2.4	0.42	8	0.69
	DF	8.8	1.5			

As a result, for May 9th, 10th and 11th, no significant difference in behaviour between the HF cows and the DF cows was found during their time on pasture. The mean differences between the behaviour percentages of the breeds were very small or not significant enough.

For the days May 12th and 13th the percentage of grazing behaviour of the DF cows was significantly increased (12th: 42.1%, 13th: 45.3%) compared to the HF cows (12th: 33.5%, 13th: 34.9%). Additionally, on May 12th the rumination behaviour of the DF cows was decreased (26.1%) compared to the HF cows (32.2%). On May 13th there was no difference in rumination between the breeds.

In Figure 12 the behaviour of the animals on pasture is shown in percentages. During the five days their behaviour demonstrates differences.




Figure 12: Average activities of the 5 HF cows and 5 DF cows during the week.

When grazing behaviour was increased, the cows spent less time on resting. This can be explained by the fact that when cows graze, they walk or stand and are not lying down, which would be registered as 'resting'. The behaviour 'active' stays stable over the week.

The most striking difference in the behaviour of the cows during this week is that the grazing behaviour of the DF cows increased from May 10th, while the grazing behaviour of the HF cows

decreased after May 11th. An explanation for this increased grazing behaviour of the DF cows is difficult to find. The cows were on the same two parcels (2 and 4) for all five days, although they probably received access to new parts of pasture each 2 days.

The weather conditions may have played a role, which is why the weather of these days are shown in Figure 13, together with the grazing behaviours of both breeds. Throughout the week the humidity of the air increased (from 50% to 72%), while there was a steady decrease in temperature (from 19.9 °C to 11.8 °C) . May 10th and 11th had less sunshine (6.5 and 9.2 hours) compared to the other days, which were extremely sunny (around 14 hours sunshine). To assess the effect of temperature and humidity on the grazing behaviour of cows, the temperature humidity index (THI) was calculated using the method of Charlton et al. (2011). According to Ehrenreich and Bjugstad (1966) the time spent on grazing by beef cattle decreases when the THI exceeds 60. When the THI is between 65 and 67, the average time less spent on grazing is 0.25 hours, between 73 and 75 this is 0.84 hours. According to Ravagnolo and Misztal (2000) heat stress in cattle occurs when THI exceeds 72.




Figure 13. Climatic conditions and the grazing behaviour of the studied cows.

During the week relative humidity increased, while temperature decreased. This resulted in a constant THI during the first 4 days, which was around 72 (range: 70 - 73). Therefore it is possible that during those days the cows suffered from heat stress which might have reduced their grazing time. The average grazing behaviour of the HF cows decreased during the week. In contrast, the grazing time of the DF did increase from May 10th. This suggests that the HF cows did suffer from heat stress whereas the DF cows are less susceptible to heat stress. This agrees with the findings of Sharma et al. (1983) who states that smaller cows are less affected by heat stress than larger breeds such as the Holstein Friesian.

Another explanation for the slightly higher grazing time of the DF cows is that they are on average fed less concentrates per kg milk they produce. This might have motivated the DF cows to graze more. Taking as an example the pair of HF63 and DF42 (Figure 14), it shows that DF42 spent more time on grazing during the week of May 9th – 13th. DF42 received 0.5 kg concentrates a day, whereas HF63 received 8 kg. These two cows are both at the end of their lactation.




Figure 14. Ruminating and grazing behaviour of pair HF 63 & DF42.

The two primiparous cows HF75 and DF80 have a high production. HF75 produced 38 liter milk per day, whereas DF80 produced 20.3 liters. Both were fed concentrates and received approximately 0.2 kg concentrates per kg milk produced. In this case it shows that when the concentrates per kg milk is the same, the DF cow spent less time on grazing than the HF did (Figure 15). HF75 in this case might be motivated to graze more, because she received per kg milk the same amount of concentrates, but produced twice as much as her counterpart DF80.




Figure 15. Ruminating and grazing behaviour of pair HF 75 & DF80.

3.4. Herbage production

On May 17th and June 30th 2016 the grass in the grasscages was mown and sampled. The grass was dried overnight at 70 °C. The DM proportion was determined and the yield of the pastures was calculated. In Table 13 the yield of the pastures of the first and second cut on both farms is shown.

Table 13. Yield of the grasscages on the pastures on May 17th and June 30th.

Farm in Utrecht (organic)				Farm in Friesland (conventional)			
Cage	DM (kg/ha)		Difference	Cage	DM (kg/ha)		Difference
	May 17 th	June 30 th			May 17 th	June 30 th	
3A	5236	3978	-24%	2A	3950	4242	+7%
3B	4686	3302	-30%	2B	3762	3705	-2%
4A	6806	2250	-67%	4A	4460	4001	-10%
4B	6330	3307	-48%	4B	4710	4004	-15%
5A	5497	2789	-49%	5A	2731	3095	+13%
5B	7060	2692	-62%	5B	3839	3521	-8%
6A	5494	2126	-61%	6A	4763	2806	-41%
6B	5435	2025	-63%	6B	4899	2337	-52%
9A	4742	2369	-50%	8A	4854	4276	-12%
9B	4749	2264	-52%	8B	4540	3912	-14%
10A	4873	3150	-35%	9A	5030	2921	-42%
10B	3804	3122	-18%	9B	4144	2974	-28%
Average	5393	2781	-48%		4307	3483	-19%

On May 17th, the first cut was made. In general this cut produces the highest yield. When the first cut is very heavy (high yield), a delay in regrowth occurs. According to R Emmelink et al. (2015) when the first cut was around 5000 kg DM/ha, the regrowth to 4000 kg DM/ha for the second cut is 12 days delayed. When this delay is not taken into account, it results in lower yields than expected. These lower yields can clearly be observed in both locations. Especially however in Utrecht, where the average yield is almost 50% lower in June than in May. To assess whether the heavy cut in May was the only reason for the reduced production in June, the production of the first and second cut of both farms were set in a scatter plot. Trendlines were constructed but resulted both in very low R²'s (Figure 16), which means that the heavy first cut was not the only reason for a much lower yield.




Figure 16. Scatter plot of the yields of first and second cut on both farms.

The average grass yield of the first cut of the organic farm was higher (5393 kg DM/ha) than the conventional farm (4307 kg DM/ha). The organic farm had the highest average yields on parcel 4 and 5. These parcels were re-sown in respectively 2015 and 2014. Parcel 9 was re-sown this year with a mixture of grass, clovers, plantain and chicory. The other parcels are all grass-clover pastures for six years. The lowest production was found on parcel 10. On June 30th the production was much lower as can be seen in Table 13. The average production of the organic farm was 2781 kg DM/ha, 48% less than on May 17th. All parcels produced less grass than before.

For the first cut, the conventional farm has highest yields on 6 and 8, and the lowest yields are observed on parcel 2 and 5. Parcel 8 has a peat soil, the other parcels are on sandy soils. The peat soil might explain the higher yield, as the peat retains water very well due to its low permeability. In June the DM production was lower on all parcels except for 2A and 5A. This might be the result of the lower production of the first cut of these parcels. The lowest production was on parcel 6 and 9. Those parcels are the wettest parcels (Bosma, personal communication) and taking into account the extreme rainfalls in June, an excess amount of water might have limited grass growth.

4. Discussion

In this chapter the results of the thesis are discussed and possible improvements for further research are mentioned.

4.1. Literature study: Milk-type vs. dual-purpose cow

The aim of the literature study was to compare the typical dairy cow to the dual-purpose cow in Europe. Knowledge on the typical dairy cow (the Holstein Friesian) was abundantly available, as there are so many Holstein Friesians used in dairy production. In contrast, dual-purpose cows represent many different breeds, often local and not well-researched. This resulted in a limited amount of articles on dual-purpose cows and the use of grey literature to obtain enough information on the breeds.

In this study the dual-purpose breed Dutch Friesian was evaluated. However, in the Netherlands just over 800 animals are registered (CRV, 2015). To state that this breed represents all dual-purpose cattle in Europe is untenable. However, the Dutch Friesian possesses traits which are found in all dual-purpose breeds, which makes it a typical dual-purpose cow. Although not of the same breed, the same grazing behaviour might be expected in other dual-purpose cows. Other factors, such as environment, farming system, temperament and diet should be considered as they might influence the grazing behaviour. Although dual-purpose cows are less susceptible to G×E interaction than Holstein Friesian cows, studies show that the quality of the diet influences their production performance.

The advantages of the dual-purpose cows which were discovered by literature study, were recognized by the participating farmers. The organic farmer was very appreciative of the fertility of his Dutch Friesian cows. The calving interval of 376 days is much lower than the average calving interval of dairy cows on Dutch farms which was 416 days in 2014 (CRV, 2015). The insemination number per cow was estimated by the organic farmer to be 1.4. The trait robustness was observable as well, as the percentage of lame cows was below 5%. The conventional farmer uses sporadically antibiotics for his herd and the occurrence of lameness is below 5%. The farmers both mentioned the higher protein and fat contents of the milk compared to HF cows as a positive aspect. Furthermore, the farmers find that their DF cows are smaller (less energy needed for maintenance) and need less additional concentrates to produce milk.

Although the two farms are different in management (organic vs. conventional), size (55 vs. 105 dairy cows) and location (Utrecht vs. Friesland), both farmers see the advantages of dual-purpose cows in their farming. This is consistent with the results of the literature study.

4.2. SensOor

This study showed that it is possible to show and follow the grazing behaviour of cows using the monitoring system SensOor. When the cow is grazing on the pasture, this behaviour is recorded as 'eating' very accurately ($R^2 = 0.97$). To monitor the behaviour of cows on the pasture the system SensOor is very suitable. However, the other observed behaviours were not as good a good match for the output of SensOor. This might be due to the lack of experience of the observant to correctly assess for example the rumination behaviour. Another explanation could be that multiple cows were observed at the same time, which increased the difficulty of observation.

At the time of writing the owning company of SensOor, AGIS, is researching new possibilities of the system. They are developing a new data storage system and are distinguishing between rumination while standing and lying down. In addition, the system will be tested under grazing conditions. It is expected that the results of AGIS will be consistent with the results of this study.

The use of monitoring systems on farms to track the behaviour of ruminants is becoming increasingly important. Around the topic Precision Dairy Farming a symposium was organised by the DairyCare

group (Rutter, 2016). As herds are becoming larger and farmers cannot keep track of all of their animal anymore, a monitoring system can help. During the conference it became clear that for this purpose many systems are already on the market and that the development of new systems and improvement of existing systems progresses quickly.

Besides SensOor, more systems which assess grazing behaviour exist and have been validated. An example is the RumiWatch system, designed to monitor the eating, drinking and rumination behaviour of the cow (Zehner et al., 2012). This system registers the movements of the jaw of the cow through a noseband, and counts the bites and chews. When a cow ruminates, she displays a very regular pattern of chewing and regurgitating a bolus. The RumiWatch system enables the researcher to create a precise overview of the cow's behaviour during rumination. This could be a valuable asset to researchers using SensOor, as SensOor only gives the time of rumination but not the pattern of biting and chewing.

4.3. Comparison of Holstein Friesian and Dutch Friesian cattle

The comparison of the Holstein Friesian and Dutch Friesian cows was done on the farm in Friesland. Only 5 purebred HF cows were available, and four of them were in their fourth or fifth lactation. The selected cows were therefore not a fair representation of the whole herd, as the herd consists of cows of many different lactation numbers.

To effectively compare the grazing behaviour of two different cow breeds, it would be desirable to have two research farms in the same region, with on each farm a predominant breed such as HF or DF. One farm with both breeds in an equal distribution would be even better. In this way the environment and a different management cannot influence the results of one of the farms.

The research described in this thesis was carried out on two different farms. As shown in the results the grass production of the pastures on the farms was very high. In general, grass production in the Netherlands is very good. Therefore it is questionable whether the outcome of these research can be applied to other countries in Europe. The pasture quality in other countries might be different from the Netherlands. Furthermore, the farming systems of the studied farms might not represent a true pasture-based system. The conventional farmer fed the herd concentrates and silage when the cows were indoors. The organic farmer supplemented the cows with concentrates as well. While the cows were outside for most of the day on the organic farm, they were only between 9:00 and 16:00 outside on the conventional farm. On the organic farm the cows could go indoors whenever they wanted. When using the results of this study, the pasture quality and the time cows spend on the pasture should be considered in order to achieve a representative result.

No significant difference in grazing behaviour between the HF and DF could be found. Although it was expected that DF cows spend more time on grazing than HF cows, this was not observed. The results of the literature study show that selection for a high milk yield is correlated with high feed intake which suggests that the HF cows graze more. Furthermore, HF cows are in general heavier than DF cows (650 kg vs. 575 kg) and therefore require more energy. This could explain why the time spent on grazing of both breeds is similar.

Another explanation for the lack of difference in grazing behaviour of the cows is that the farm has pastures of a high quality. Perennial ryegrass (*Lolium perenne*) is an excellent grass species for feeding lactating cows due to its high yield and nutritional quality (Smit et al., 2005). The literature study showed that dual-purpose cows are better able to cope with a harsh environment compared to HF cows. On this particular farm both breeds were grazed on a pasture of high quality. Therefore the advantage of the dual-purpose breed could not be observed. It is very well possible that if the cows were kept on a semi-natural pasture (of worse quality) a difference in grazing behaviour would show.

4.4. Herbage production

Although it was expected that the weather circumstances in Friesland would be very different from those in Utrecht this was not the case. The amount of rainfall was almost the same and also the total hours of sunshine. Only the average daily temperature was significantly higher in Utrecht, which explains the higher grass production of the first and second cut in Utrecht. Lantinga (personal communication) stated that a difference of 1 °C (of the average daily temperature) can make a 20% growth difference in grass. This only occurs between 5 and 10 °C. Above an average ambient daily temperature of 10 °C the temperature does not restrict grass growth anymore. The influence of the temperature could even be higher on a local scale, as the farm in Utrecht is quite secluded by trees and therefore less susceptible to hard wind and cold. In contrast, the parcels of the farm in Friesland are not sheltered at all and probably more vulnerable to extreme weather.

Both farms used manure of the own cows to fertilize the pastures. On the conventional farm artificial fertilizer was applied as well. The difference in grassland production can be caused by several factors. The soil composition of both farms is unknown, while the level of nutrients and minerals in the soil directly affects grass growth. The parcels of the organic farm are organic since 1977, therefore probably the organic matter in the soil is higher than the soils of the conventional farm. This would mean that the soils of the organic farm are more fertile (Maeder et al., 2002). On the other hand, the conventional farmer adds extra nitrogen in the form of artificial fertilizer.

The difference in production between the first and second cut is very high. For the organic farm the average production decreased with 52%. The production of the conventional farm was lower as well, but not as much as the organic farm: 19%. The large difference with the second cut might be partially explained by the high herbage production of the first cut. It is possible that because of the heavy first cut, the regrowth was seriously delayed and resulted in a much lower yield than expected.

The month June was extremely wet, KNMI never recorded such high precipitation during June before. This can partly explain the lower production of both farms because excessive water on pastures is unfavourable for plant growth. An explanation for the differences in production between the farms might be that on the organic farm the manure management has been different from the conventional farm. Precise numbers on how much and when the parcels were fertilized was not available. However, upon arrival on the organic farm on June 30th, the colour of the grass was light green. This might indicate that there was a nitrogen shortage in the pasture. The conventional farm uses additional artificial fertilizer to fertilize his pastures, which might explain the less severe drop in production.

The production of the grasslands was estimated by using grass cages. There was no influence of grazing on the grass in the cages, whereas the majority of the pastures of both farms is used for grazing. As grazing has a considerable influence on the production of a grassland the grass cages might not give an entirely honest picture of the actual production (Trlica, 1992). Grazed pastures have a lower net grass production than mown pastures (Lantinga et al., 1987).

5. Conclusions

The literature study in this thesis has shown that Holstein Friesian cows are not suited for pasture-based systems of low forage quality, especially when their performance is compared to that of dual-purpose cows. Holstein Friesians tend to show a large drop in their body reserves after calving, which is only solved by supplementing the cow with large amount of concentrates. When fed only grass her body condition decreases rapidly after calving and also her production will not reach its potential. A dual-purpose cow distributes her energy more gradually and therefore her body condition stays stable after calving. While her milk production increases, this is not at the cost of the energy required for her own maintenance.

This is the strongest advantage that dual-purpose cows have over Holstein Friesian. The first hypothesis formulated by the project 2-ORG-COWS ('Dual-purpose cows are more robust than modern dairy cattle breeds, and therefore more suitable for varying harsh organic grassland systems') is therefore endorsed by the results of this thesis.

Additionally, when a cow has a bad body condition score, the disease resistance and fertility of the cow decreases as well. Diseases cost the farmer money for treatment and losses in the milk. Fertility problems costs the farmer money because it takes more time to get the cow in milking again, and the costs of semen and artificial insemination.

Another great advantage of dual-purpose cows is their longevity. The milk production of a cow increases with the number of lactations the cow has had and due to their better health and robustness the dual-purpose breeds live longer than HF cows. The experiences of the participating farmers in this study with their own dual-purpose breed 'Dutch Friesian' emphasize these results. The farmer in Utrecht named the fertility of the Dutch Friesian as a particular important asset, and the farmer in Friesland their robustness and health.

In this thesis the cow behaviour monitoring system 'SensOor' was validated under grazing conditions. Especially for the monitoring of grazing behaviour the system is very well suited. A combination with other monitoring systems such as the RumiWatch could increase the reliability of the output. The SensOor system measures concerning grazing and rumination only the time the cows spend on this activity. RumiWatch gives more information on the frequency of bites and chews and shows a rumination pattern. This information could increase the knowledge on the differences in eating and rumination behaviour of cows and in particular dual-purpose cows. Currently the company of SensOor (AGIS) is researching the system under grazing conditions and developing improvements.

Regarding the comparison of the grazing behaviour of the dual-purpose breed Dutch Friesian to the Holstein Friesian on the farm in Friesland, no clear difference in behaviour was found. There was much variability between the individual cows. It is expected that if the sample size would be larger a more complete view could be obtained of the grazing behaviour of the cows. In this case one cow with aberrant behaviour could have a large impact on the results as every cow provides 10% of the results in the comparative behaviour study. The results of this study showed no significant difference in grazing behaviour of both breeds, but rather a similar pattern. The result that both breeds spend a considerable part of their pasture time on grazing, can be explained by the correlation between high milk yield and feed intake for the HF cows and the grazing preference of the DF cows. The pasture time of these cows was approximately 7 hours a day and the average milk yield of the DF cows was lower than the HF cows (22,8 liter/day vs. 30,6 liter/day).

On the two studied farms the herbage production was estimated using grass cages. A clear difference in yield was observed in the first cut in May, as the organic farm in Utrecht had an impressive production (average 5400 kg DM/ha) compared to the conventional farm in Friesland (average 4300 kg DM/ha). This was probably caused by the ambient temperature differences. During the spring the temperature in Utrecht was on average 1 °C higher than in Friesland. This difference could have been reinforced by the openness of the pastures in Friesland, compared to Utrecht where the pastures lie secluded between trees. The absence of trees in Friesland makes the pastures more vulnerable to extreme weather conditions and might have affected the perceived temperature as even more cold.

The second cut resulted in much lower yields which can be expected after a heavy first cut. The yield of the pastures of the conventional farm were higher than those of the organic farm during the second cut, which is explained by a better manure management on the conventional farm. Although precise information on the manure management is missing, it is estimated that the manure application per hectare on both farms was comparable. Additionally, the conventional farmer used artificial fertilizer on all his parcels.

Dual-purpose cows perform well in a pasture-based system. Those breeds are traditionally selected for their ability to produce on a diet of low quality and the time they spent on grazing. Farmers selected breeding bulls based on the performance and grazing behaviour of the mother of these animals. Compared to the popular Holstein Friesian, dual-purpose cows are better suited for a pasture-based system. The Dutch pasture-based systems might not be representative for other European countries because of their excellent grass production and suitable environment. The Dutch pastures inhabit in general favourable, nutritional grass species and are not steep or remote. If the behaviour study in this thesis was conducted in for example the Alps, the pasture conditions would be very different from the Dutch pasture conditions. The Alpine pastures are not suited for comprehensive cultivation, are in general steep and can be remote. This results in pastures of lower quality, and it is expected that the performance of dual-purpose breeds would be significantly better than that of Holstein Friesian cows. The good conditions of the Dutch pastures conceal the better grazing properties of the dual-purpose breeds with respect to the Holstein Friesian cows.

Literature

- Aass, L., and O. Vangen. 1997. Effects of selection for high milk yield and growth on carcass and meat quality traits in dual purpose cattle. *Livestock Production Science* 52: 75-86.
- Auldust, M., S. Coats, G. Rogers, and G. McDowell. 1995. Changes in the composition of milk from healthy and mastitic dairy cows during the lactation cycle. *Australian Journal of Experimental Agriculture* 35: 427-436.
- Bakker, J., and A. Brouwer. 2012. Monitor duurzaam voedsel 2011. In: L. e. I. Ministerie van Economische Zaken (ed.). p 1-40, The Hague, The Netherlands.
- Ball, D. M. et al. 2001. Understanding forage quality. American Farm Bureau Federation Park Ridge, IL.
- Bargo, F., L. D. Muller, J. E. Delahoy, and T. W. Cassidy. 2002. Milk Response to Concentrate Supplementation of High Producing Dairy Cows Grazing at Two Pasture Allowances. *Journal of Dairy Science* 85: 1777-1792.
- Berry, D. P. et al. 2003a. Estimation of genotype \times environment interactions, in a grass-based system, for milk yield, body condition score, and body weight using random regression models. *Livestock Production Science* 83: 191-203.
- Berry, D. P. et al. 2003b. Genetic Relationships among Body Condition Score, Body Weight, Milk Yield, and Fertility in Dairy Cows. *Journal of Dairy Science* 86: 2193-2204.
- Bikker, J. P. et al. 2014. Technical note: Evaluation of an ear-attached movement sensor to record cow feeding behavior and activity. *Journal of Dairy Science* 97: 2974-2979.
- Boichard, D. 1990. Estimation of the economic value of conception rate in dairy cattle. *Livestock Production Science* 24: 187-204.
- Bot, B. 2011. Grondwaterzakboekje. Bot, Raadgevend Ingenieur, Rotterdam, NL.
- Buckley, F., P. Dillon, S. Crosse, F. Flynn, and M. Rath. 2000. The performance of Holstein Friesian dairy cows of high and medium genetic merit for milk production on grass-based feeding systems. *Livestock Production Science* 64: 107-119.
- Buning, S. 2012. Precisielandbouw in de stal: observatiesysteem SensOor volgt elke hap en stap. *Melkvee magazine: Onafhankelijk magazine voor de melkveehouder en veefokker* 11: 30-31.
- Carthy, T., D. Ryan, A. Fitzgerald, R. Evans, and D. Berry. 2016. Genetic relationships between detailed reproductive traits and performance traits in Holstein-Friesian dairy cattle. *Journal of dairy science* 99: 1286-1297.
- Charlton, G., S. Rutter, M. East, and L. Sinclair. 2011. Effects of providing total mixed rations indoors and on pasture on the behavior of lactating dairy cattle and their preference to be indoors or on pasture. *Journal of dairy science* 94: 3875-3884.
- Charlton, G. L., S. M. Rutter, M. East, and L. A. Sinclair. 2013. The motivation of dairy cows for access to pasture. *Journal of Dairy Science* 96: 4387-4396.
- Coulon, J., and P. Pradel. 1997. Effect of walking in roughage intake and milk yield and composition of Montbeliarde and Tarentaise dairy cows. In: *Annales de zootechnie*. p 139-146.
- CRV. 2015. Jaarstatistieken. <https://www.crv4all.nl/downloads/prestaties/jaarstatistieken/> Accessed 23-5-2016.
- De Winter, M., T. Vogelzang, and J. Schaick. 2010. De blaarkop: ouderwets goed: inventarisatie van de mogelijkheden voor een dubbeldoelkoe in deze tijd. 9086154247, LEI Wageningen UR.
- Delaby, L., P. Faverdin, G. Michel, C. Disenhaus, and J. L. Peyraud. 2009. Effect of different feeding strategies on lactation performance of Holstein and Normande dairy cows. *Animal* 3: 891-905.
- Dillon, P., D. P. Berry, R. D. Evans, F. Buckley, and B. Horan. 2006. Consequences of genetic selection for increased milk production in European seasonal pasture based systems of milk production. *Livestock Science* 99: 141-158.
- Dillon, P., F. Buckley, P. O'Connor, D. Hegarty, and M. Rath. 2003. A comparison of different dairy cow breeds on a seasonal grass-based system of milk production: 1. Milk production, live weight, body condition score and DM intake. *Livestock Production Science* 83: 21-33.
- Ehrenreich, J. H., and A. J. Bjugstad. 1966. Cattle Grazing Time is Related to Temperature and Humidity. *Journal of Range Management Archives* 19: 141-142.

- Fulkerson, W. J. et al. 2008. Holstein-Friesian Dairy Cows Under a Predominantly Grazing System: Interaction Between Genotype and Environment. *Journal of Dairy Science* 91: 826-839.
- Geuder, U., M. Pickl, M. Scheidler, M. Schuster, and K.-U. Gótz. 2012. Growth performance, carcass traits and meat quality of Bavarian cattle breeds. *Zuchtungskunde* 84: 485-499.
- Gibb, M. J., C. A. Huckle, R. Nuthall, and A. J. Rook. 1999. The effect of physiological state (lactating or dry) and sward surface height on grazing behaviour and intake by dairy cows. *Applied Animal Behaviour Science* 63: 269-287.
- Haiger, A., and W. Knaus. 2010. A comparison of dual-purpose Simmental and Holstein Friesian dairy cows in milk and meat production: 1st comm. Milk production without concentrates. *Zuchtungskunde* 82: 131-143.
- Hernandez-Mendo, O., M. A. G. von Keyserlingk, D. M. Veira, and D. M. Weary. 2007. Effects of Pasture on Lameness in Dairy Cows. *Journal of Dairy Science* 90: 1209-1214.
- Hiemstra, A. 2015. 'Naar 10.000 kilo melk per koe willen we niet': Ton Spierings looft probleemloze karakter van MRIJ-ras. *Melkvee magazine: Onafhankelijk magazine voor de melkveehouder en veefokker* 14: 30-33.
- Hiemstra, S. J., Y. De Haas, A. Mäki-Tanila, and G. Gandini. 2010. Local cattle breeds in Europe. Development of policies and strategies for self-sustaining breeds. Wageningen Academic Publ., Wageningen, The Netherlands.
- Horn, M. et al. 2013. Suitability of different dairy cow types for an Alpine organic and low-input milk production system. *Livestock Science* 153: 135-146.
- Isselstein, J. et al. 2013. Qualität des Weidefutters in der ökologischen Milchviehhaltung.
- Jacobsen, S. 2015. Biologische melkprijs FrieslandCampina fors omhoog. <http://www.melkvee.nl/nieuws/8048/biologische-melkprijs-frieslandcampina-fors-omhoog> Accessed 8-2 2016.
- Jílek, F. et al. 2008. Relationships among body condition score, milk yield and reproduction in Czech Fleckvieh cows. *Czech Journal of Animal Science* 53: 357-367.
- Kadzere, C. T., M. R. Murphy, N. Silanikove, and E. Maltz. 2002. Heat stress in lactating dairy cows: a review. *Livestock Production Science* 77: 59-91.
- Kennedy, J. et al. 2003. Effect of genetic merit and concentrate supplementation on grass intake and milk production with Holstein Friesian dairy cows. *Journal of Dairy Science* 86: 610-621.
- Knaus, W. 2016. Perspectives on pasture versus indoor feeding of dairy cows. *Journal of the Science of Food and Agriculture* 96: 9-17.
- KNMI. 2016a. Archief maand/seizoen/jaaroverzichten. <http://www.knmi.nl/nederland-nu/klimatologie/maand-en-seizoensoverzichten> Accessed 8-6-2016.
- KNMI. 2016b. KNMI - Daggegevens van het weer in Nederland. <http://projects.knmi.nl/klimatologie/daggegevens/index.cgi> Accessed 18 May 2016.
- Kolver, E., J. Roche, M. De Veth, P. d. Thorne, and A. Napper. 2002. Total mixed rations versus pasture diets: Evidence for a genotype x diet interaction in dairy cow performance. In: *Proceedings-New Zealand Society of Animal Production*. p 246-251.
- Kristensen, H. 2015. 2-ORG-COWS. <http://coreorganicplus.org/research-projects/2-org-cows/> Accessed 1-8-2016.
- Kummerow, J., and B. A. Ellis. 1984. Temperature effect on biomass production and root/shoot biomass ratios in two arctic sedges under controlled environmental conditions. *Canadian Journal of Botany* 62: 2150-2153.
- Lairon, D., and M. Huber. 2014. Food quality and possible positive health effects of organic products *Organic Farming, Prototype for Sustainable Agricultures*. p 295-312. Springer, the Netherlands.
- Lantinga, E., J. Keuning, J. Groenwold, and P. Deenen. 1987. Distribution of excreted nitrogen by grazing cattle and its effects on sward quality, herbage production and utilization *Animal Manure on Grassland and Fodder Crops. Fertilizer or Waste?* p 103-117. Springer, the Netherlands.
- Macdonald, K. et al. 2008. A comparison of three strains of Holstein-Friesian grazed on pasture and managed under different feed allowances. *Journal of dairy science* 91: 1693-1707.
- Maeder, P. et al. 2002. Soil Fertility and Biodiversity in Organic Farming. *Science* 296: 1694-1697.

- McCarthy, S. et al. 2007. The influence of strain of Holstein-Friesian dairy cow and pasture-based feeding system on grazing behaviour, intake and milk production. *Grass and Forage Science* 62: 13-26.
- Medugorac, I. et al. 2009. Genetic diversity of European cattle breeds highlights the conservation value of traditional unselected breeds with high effective population size. *Molecular ecology* 18: 3394-3410.
- Milchunas, D., O. E. Sala, and W. K. Lauenroth. 1988. A generalized model of the effects of grazing by large herbivores on grassland community structure. *American Naturalist*: 87-106.
- Monsón, F., C. Sañudo, and I. Sierra. 2005. Influence of breed and ageing time on the sensory meat quality and consumer acceptability in intensively reared beef. *Meat Science* 71: 471-479.
- Nauta, W. J., R. F. Veerkamp, E. W. Brascamp, and H. Bovenhuis. 2006. Genotype by Environment Interaction for Milk Production Traits Between Organic and Conventional Dairy Cattle Production in The Netherlands. *Journal of Dairy Science* 89: 2729-2737.
- O'Connell, J. M., F. Buckley, M. Rath, and P. Dillon. 2000. The Effects of Cow Genetic Merit and Feeding Treatment on Milk Production, Herbage Intake and Grazing Behaviour of Dairy Cows. *Irish Journal of Agricultural and Food Research* 39: 369-381.
- Oba, M., and M. S. Allen. 1999. Evaluation of the Importance of the Digestibility of Neutral Detergent Fiber from Forage: Effects on Dry Matter Intake and Milk Yield of Dairy Cows. *Journal of Dairy Science* 82: 589-596.
- Oldenbroek, J., and P. Van Eldik. 1980. Differences in feed intake between holstein Friesian, Dutch Red and White and Dutch Friesian cattle. *Livestock Production Science* 7: 13-23.
- Palmquist, D. L., A. Denise Beaulieu, and D. M. Barbano. 1993. Feed and Animal Factors Influencing Milk Fat Composition1. *Journal of Dairy Science* 76: 1753-1771.
- Parsons, A. J., E. L. Leafe, B. Collett, and W. Stiles. 1983. The Physiology of Grass Production Under Grazing. I. Characteristics of Leaf and Canopy Photosynthesis of Continuously-Grazed Swards. *Journal of Applied Ecology* 20: 117-126.
- Peniche-González, I. N. et al. 2014. Milk production and reproduction of dual-purpose cows with a restricted concentrate allowance and access to an association of *Leucaena leucocephala* and *Cynodon nlemfuensis*. *Journal of Applied Animal Research* 42: 345-351.
- Piccand, V. et al. 2013. Production and reproduction of Fleckvieh, Brown Swiss, and 2 strains of Holstein-Friesian cows in a pasture-based, seasonal-calving dairy system. *Journal of dairy science* 96: 5352-5363.
- Pryce, J., R. Veerkamp, R. Thompson, W. Hill, and G. Simm. 1997. Genetic aspects of common health disorders and measures of fertility in Holstein Friesian dairy cattle. *Animal Science* 65: 353-360.
- Pryce, J. E., M. D. Royal, P. C. Garnsworthy, and I. L. Mao. 2004. Fertility in the high-producing dairy cow. *Livestock Production Science* 86: 125-135.
- Ravagnolo, O., and I. Misztal. 2000. Genetic Component of Heat Stress in Dairy Cattle, Parameter Estimation. *Journal of Dairy Science* 83: 2126-2130.
- Ray, D. E., T. J. Halbach, and D. V. Armstrong. 1992. Season and Lactation Number Effects on Milk Production and Reproduction of Dairy Cattle in Arizona1. *Journal of Dairy Science* 75: 2976-2983.
- Remmelink, G., J. van Middelkoop, W. Ouweltjes, and H. Wemmenhove. 2015. Handboek melkveehouderij 2015/16, Wageningen UR Livestock Research.
- Renna, M. et al. 2014. Milk yield, gross composition and fatty acid profile of dual-purpose Aosta Red Pied cows fed separate concentrate-forage versus total mixed ration. *Animal Science Journal* 85: 37-45.
- Roche, J. R., D. P. Berry, and E. S. Kolver. 2006. Holstein-Friesian Strain and Feed Effects on Milk Production, Body Weight, and Body Condition Score Profiles in Grazing Dairy Cows. *Journal of Dairy Science* 89: 3532-3543.
- Roche, J. R. et al. 2009. Invited review: Body condition score and its association with dairy cow productivity, health, and welfare. *Journal of Dairy Science* 92: 5769-5801.
- Rössler, R., P. Herold, A. Weidele, and A. V. Zärate. 2013. Definition of user-specific breeding goals in Brown Swiss and Hinterwälder cattle in Baden-Württemberg. *Zuchtungskunde* 85: 173-187.

- Royal, M. D., A. P. F. Flint, and J. A. Woolliams. 2002. Genetic and Phenotypic Relationships Among Endocrine and Traditional Fertility Traits and Production Traits in Holstein-Friesian Dairy Cows. *Journal of Dairy Science* 85: 958-967.
- Rutter, M. 2016. The on-farm use of automated measures of ruminant foraging behaviour. In: *Precision Dairy Farming: Activity measurement in ruminant research and beyond*, WTC, Leeuwarden, The Netherlands
- Sala, O. E., W. J. Parton, L. Joyce, and W. Lauenroth. 1988. Primary production of the central grassland region of the United States. *Ecology* 69: 40-45.
- Schaack, D., J. Lernoud, B. Schlatter, and H. Willer. 2014. The organic market in Europe 2012. The world of organic agriculture. *Statistics and emerging trends*: 200-206.
- Schild, H., E. Niebel, and K.-U. Götz. 2003. Across country genetic evaluation of beef traits in Middle European dual purpose breeds. *Interbull Bulletin*: 158.
- Sharma, A. K. et al. 1983. Climatological and Genetic Effects on Milk Composition and Yield. *Journal of Dairy Science* 66: 119-126.
- Smit, H., B. Tas, H. Taweel, S. Tamminga, and A. Elgersma. 2005. Effects of perennial ryegrass (*Lolium perenne* L.) cultivars on herbage production, nutritional quality and herbage intake of grazing dairy cows. *Grass and Forage Science* 60: 297-309.
- Syrstad, O. 1993. Evaluation of dual-purpose (milk and meat) animals. *World Animal Review* 77: 56-59.
- Theunissen, B. 2012. Breeding for nobility or for production? Cultures of dairy cattle breeding in the Netherlands, 1945–1995. *Isis* 103: 278-309.
- Thornley, J., and M. Cannell. 1997. Temperate grassland responses to climate change: an analysis using the Hurley pasture model. *Annals of Botany* 80: 205-221.
- Trlica, M. J. 1992. Grass growth and response to grazing. Colorado State University Cooperative Extension.
- Van Arendonk, J. A. M., and A.-E. Liinamo. 2003. Dairy cattle production in Europe. *Theriogenology* 59: 563-569.
- Van Ginneken, R. 2010. Zorgvuldigheid troef bij inkruisen ander ras: Interesse in crossbreeding neemt toe. *Melkvee magazine: Onafhankelijk magazine voor de melkveehouder en veefokker* 7: 12-14.
- Van Saun, R. J. 2006. Determining forage quality: Understanding feed analysis. *Lamalink.com* August 3: 18-19.
- Veerkamp, R. F., G. Simm, and J. D. Oldham. 1994. Effects of interaction between genotype and feeding system on milk production, feed intake, efficiency and body tissue mobilization in dairy cows. *Livestock Production Science* 39: 229-241.
- Wolfger, B. et al. 2015. Technical note: Accuracy of an ear tag-attached accelerometer to monitor rumination and feeding behavior in feedlot cattle. *Journal of Animal Science* 93: 3164-3168.
- Zehner, N. et al. 2012. Validation of a new health monitoring system (RumiWatch) for combined automatic measurement of rumination, feed intake, water intake and locomotion in dairy cows. In: *Proceedings of international conference of agricultural engineering CIGR-Ageng*. p 1-6.
- Zevenbergen, G. 2012. Gegevens uit het oor: Agis Sesnoor ziet zieke koe snel. *Veehouderij Techniek* 15: 33-35.
- Zwald, N. R., K. A. Weigel, W. F. Fikse, and R. Rekaya. 2003. Identification of Factors That Cause Genotype by Environment Interaction Between Herds of Holstein Cattle in Seventeen Countries. *Journal of Dairy Science* 86: 1009-1018.

Appendix I
Map of the organic farm.



Appendix II
Map of the conventional farm.



Appendix III

Observation scheme used for cow observations on May 10th and 12th, 2016.

Date:

Cow number:

	Grazing st	Grazing w	Ruminating l	Ruminating st	Standing	Resting	Active
0:00:00							
0:01:00							
0:02:00							
0:03:00							
0:04:00							
0:05:00							
0:06:00							
0:07:00							
0:08:00							
0:09:00							
0:10:00							
0:11:00							
0:12:00							
0:13:00							
0:14:00							
0:15:00							
0:16:00							
0:17:00							
0:18:00							
0:19:00							
0:20:00							
0:21:00							
0:22:00							
0:23:00							
0:24:00							
0:25:00							
0:26:00							
0:27:00							
0:28:00							
0:29:00							
0:30:00							
0:31:00							
0:32:00							
0:33:00							
0:34:00							
0:35:00							
0:36:00							
0:37:00							
0:38:00							
0:39:00							
0:40:00							
0:41:00							
0:42:00							
0:43:00							
0:44:00							
0:45:00							
0:46:00							
0:47:00							
0:48:00							
0:49:00							
0:50:00							
0:51:00							
0:52:00							
0:53:00							
0:54:00							
0:55:00							
0:56:00							
0:57:00							
0:58:00							
0:59:00							