

# Mechanism of endocarp-imposed constraints of germination of *Lannea microcarpa* seeds

Oblé Neya<sup>1,2</sup>, Folkert A. Hoekstra<sup>1</sup> and Elena A. Golovina<sup>1,3\*†</sup>

<sup>1</sup>The Graduate School 'Experimental Plant Sciences', Laboratory of Plant Physiology, Wageningen University, Arboretumlaan 4, Wageningen, The Netherlands; <sup>2</sup>Centre National de Semences Forestières, 01 BP 2682 Ouagadougou 01, Burkina Faso; <sup>3</sup>K.A. Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, Botanicheskaja 35, Moscow 127276, Russia

## Abstract

*Lannea microcarpa*, a multipurpose tree species from the dry African savanna, sheds seeds that often display inhibition of germination. The underlying mechanism was investigated using seeds processed from fully matured fruits collected from natural stands in Burkina Faso. Germination of fresh seeds was variable (16–28%), while they did not germinate after drying and rehydration. Mechanical scarification of the endocarp at the proximal end of the seeds increased germination to 83–94%. Scarification on the distal end led to delayed radicle emergence through the produced hole in c. 40% of the seeds. The endocarp was permeable to water and respiratory gases. Increased water content in scarified seeds was associated with radicle extension during germination. Intact and scarified non-germinated seeds displayed a moderate rate of respiration with respiratory quotient (RQ) values of c. 1. Respiration increased and RQ decreased to c. 0.7 with radicle emergence. Ethylene evolution peaked in both intact and scarified seeds at the beginning of incubation and then decreased to low values. Inhibition of ethylene production by 1–5 mM 2-amino-ethoxyvinylglycine (AVG) caused only a partial decrease of germination of the scarified seeds. Intact non-germinated seeds gradually lost viability during incubation at 30°C, but could be rescued by delayed scarification before day 15 of incubation. It is concluded that radicle emergence in dry *L. microcarpa* seeds is inhibited only mechanically. The mechanical properties of the endocarp are attributed

to irreversible structural changes of the lignin–hemicellulose complex, which occur during drying.

**Keywords:** endocarp-imposed constraints, ethylene, germination, *Lannea microcarpa*, mechanical scarification, respiration, viability loss

## Introduction

*Lannea microcarpa* (Anacardiaceae) is one of the important multipurpose tree species from the African savanna. Together with other savanna tree species, *L. microcarpa* is considered endangered as a result of over-exploitation and repetitive severe drought spells in recent decades. Regular propagation of local tropical trees species is important for successful reforestation. Propagation is generally carried out via seeds in nurseries. However, little research has been conducted on the seeds. The lack of knowledge of the seed physiology of *L. microcarpa* is a major problem for the sustainable management and successful implementation of reforestation programmes.

*L. microcarpa* inflorescences are racemes, in which fruits of different stages of maturity are found (Neya, 2006). Freshly depulped seeds have water contents in the range 30–40%. Seeds of *L. microcarpa* have been classified previously as recalcitrant (Hong *et al.*, 1996). However, recent studies showed that these seeds are desiccation tolerant (Daws *et al.*, 2004; Flynn *et al.*, 2004; Pritchard *et al.*, 2004), but often have low germination after drying (Neya, 2006), which led to an incorrect conclusion about their recalcitrant behaviour.

Germination of both fresh and dry seeds of *Lannea* is very variable. This variation was observed between seeds collected in different years, or in the same year, but from different maturity stages of fruits (Neya, 2006). The increase of germination to  $\geq 80\%$  by mechanical scarification indicates that the generally low germination of *L. microcarpa* seeds is caused by

\*Correspondence

Fax: + 31 317 482725

Email: elena.golovina@wur.nl

†Current address: Wageningen NMR Centre, Dreijenlaan 3, 6703 HA Wageningen, The Netherlands

coat-imposed constraints (Neya, 2006). Differences in the extent of such constraints are likely to be the cause of the variability in germination observed among and within seed accessions. Coat-imposed constraints can be eliminated by artificial treatments, such as mechanical or acid scarification, natural causes such as fire, high temperature or fluctuation in temperature, or biological actions, such as micro-organisms or passage through the gut (Baskin and Baskin, 1998, 2000; Jones, 1999; van Assche *et al.*, 2003; Manzano *et al.*, 2005).

The seed coat (testa) in the *Anacardiaceae* family is not well developed (undifferentiated), and its functional role is performed by the fruit coat, represented by the endocarp (Baskin *et al.*, 2000). The endocarp in the *Anacardiaceae* family develops from the inner epidermis of the ovary wall and contains four layers (Baskin *et al.*, 2000). The three innermost layers are palisade-shaped and may cause mechanical and permeability constraints (Li *et al.*, 1999a, b; Baskin *et al.*, 2000; Baskin and Baskin, 2004). A number of mechanisms have been proposed by which seed/fruit coats can inhibit germination (Bewley and Black, 1994; Adkins *et al.*, 2002). These include: (1) interference with water uptake; (2) mechanical constraints; (3) interference with gas exchange; (4) retention of inhibitors; and (5) inhibitor production. Often, several of the possible constraints act together to maintain the intact seed in a dormant state (Bewley and Black, 1994).

In the present study, the germination patterns of non-scarified and scarified seeds were used to test the possible mechanisms of endocarp-imposed constraints of the germination of dried *L. microcarpa* seeds.

## Materials and methods

### Plant material

Seeds of *Lannea microcarpa* (Engl. & K. Krause) were harvested from a natural stand in Ipélcé, Burkina Faso. They were collected from fully mature (purple-black) fruits and were, on the same day, soaked in water and depulped at the Centre National de Semences Forestières (CNSF) in Ouagadougou. The seeds were rubbed with sand to remove the mesocarp tissue and washed with water. Subsequently, they were dried at ambient conditions (c. 29°C and 53% relative humidity) in the laboratory for 2–3 h, and seeds surrounded by an intact endocarp were then selected and sent the next day to Wageningen (The Netherlands) in cotton bags, arriving 2 d after harvest. Water content of seeds on arrival was 0.32–0.43 g H<sub>2</sub>O g<sup>-1</sup> (DM, dry mass basis).

### Moisture content and germination

Three replicates of five seeds were used to assess seed moisture content (MC) throughout the study.

Seeds were weighed before and after drying at 103°C for 17 h (ISTA, 1999), and their MCs were calculated as g H<sub>2</sub>O g<sup>-1</sup> (DM). The MC during incubation was determined by temporary removal of the seeds from the germination medium and surface-drying with tissue paper before the fresh mass was weighed. From the increase in mass, the actual MC was calculated.

For each single germination assay, 25 seeds in duplicate (unless stated otherwise) were used. The seeds were sown on moist filter paper contained in transparent plastic boxes (11 cm × 17 cm × 5 cm) for germination and incubated at 30°C in an incubator. Seeds were scored for germination at least twice a week until all viable seeds had germinated. They were regarded as germinated when the radicle had emerged to at least 2 mm. Soft and rotten seeds were considered as non-viable and discarded.

### Mechanical scarification

Scarification consisted of making a small hole in the endocarp of seeds on the proximal (above the axis) or on the distal ends, with respect to the micropylar region of the seed, using a nail-clipper or a razor blade. Damage to the embryo during this process was avoided, and seeds with visibly damaged embryos were discarded. Seeds were scarified prior to sowing or later during incubation. In the latter case, they were first incubated non-scarified at 30°C for 3, 6, 9, 12 or 15 d, before being scarified at the proximal end and returned to the germination medium for further incubation at 30°C.

### Electron paramagnetic resonance spectroscopy

Electron paramagnetic resonance (EPR) spectra were recorded at room temperature with an X-band EPR spectrometer (Bruker, Rheinstetten, Germany, Model Elexsys 500). The water-soluble nitroxide radicle, perdeuterated TEMPONE (PDT; kindly provided by Professor I. Grigoriev, Institute of Organic Chemistry of the Russian Academy of Sciences, Novosibirsk, Russia), was used as the spin probe for testing the integrity of the plasma membranes of the cells in excised axes, according to the method described by Golovina and Tikhonov (1994). The seeds were first deoiled and rehydrated for 2 h or incubated intact at 30°C on moist filter paper for 15 d. Then axes were dissected and incubated in 1 mM PDT + 120 mM potassium ferricyanide for at least 20 min. EPR spectra of PDT in the isolated axes were recorded at 0.25 G modulation amplitude and 18 dB (3.17 mW). The spectra were recorded on individual axes, and the ratio of amplitudes of water/lipid (W/L) was calculated for each spectrum.

### **Respiration and ethylene measurements**

Respiration of seeds was investigated using headspace analysis. The individual seeds were allowed to respire in air-tight 7 ml vials for *c.* 5 h at 30°C. If seeds were incubated in water prior to respiration measurements, pieces of wet filter paper were placed at the bottom of the vials to avoid drying of the samples. After 5 h, 40  $\mu$ l headspace gas samples were taken from the vials and injected into a gas chromatograph (GC) equipped with a catharometer detector as described by Leprince and Hoekstra (1998), which allowed O<sub>2</sub> uptake and CO<sub>2</sub> release to be measured simultaneously. The sensitivity of the GC detection was enhanced by an additional amplifier, connected to an electronic integrator. The carrier gas was He. Calibration of the respiratory gases was performed with pure CO<sub>2</sub> and air, assuming a gas composition of the air at atmospheric pressure of 20.95 vol % O<sub>2</sub>, 0.93 vol % Ar, and 78.09 vol % N<sub>2</sub>. Accuracy of the quantification was on the basis that the sum of the injected gases (CO<sub>2</sub>, O<sub>2</sub>, Ar and N<sub>2</sub>) equalled 100%.

For ethylene measurements, samples of the headspace gas of intact and scarified seeds (proximal end) were taken from corresponding vials after 9 h of accumulation. One-ml samples were taken and analysed using a gas chromatograph equipped with an alumina column and a flame ionization detector. Temperatures were 180 and 200°C for the oven and the detector, respectively. Ethylene peaks were registered by an integrator coupled to the GC and quantified by comparison with peaks of ethylene standards.

### **2-Amino-ethoxyvinylglycine treatments**

Seeds were scarified on the proximal end and incubated at 30°C in Petri dishes containing filter paper soaked in 0.1, 1, 5 or 10 mM AVG solutions, plus a water control. At daily intervals, four seeds were removed from each Petri dish (with different AVG concentrations) and individually placed into air-tight 7 ml vials, where they were allowed to accumulate ethylene for *c.* 2 h. Subsequently, headspace gas samples were taken for ethylene analysis as described above. For germination tests, 25 seeds in duplicate were sown in transparent plastic boxes containing filter papers moistened with each of the AVG solutions and incubated at 30°C.

### **Fourier transform infrared microspectroscopy**

Transmission Fourier transform infrared (FT-IR) spectra were recorded at room temperature on a Perkin-Elmer 1725 IR-spectrometer (Perkin-Elmer, Beaconsfield, UK), equipped with a narrow-band, liquid nitrogen-cooled mercury/cadmium/telluride

detector and a Perkin-Elmer microscope. The optical bench was purged with dry CO<sub>2</sub>-free air (Balston, Maidstone, UK). The acquisition parameters were 4 cm<sup>-1</sup> resolution, 32 co-added interferograms, 2 cm<sup>-1</sup> moving mirror speed, and 3600 to 800 cm<sup>-1</sup> wavenumber range. Spectral analysis and display were carried out using Spectrum version 2.00 (Perkin-Elmer).

Samples scratched from the outer or inner layers of the endocarp by a scalpel were placed between two diamond windows that were mounted into a temperature-controlled holder. The sample was pressurized until sufficient transparency was obtained. To determine the components of the endocarp, recorded spectra were compared with spectra of the natural reference material kapok fibres, containing a mixture of cellulose, lignin and hemicellulose.

## **Results**


### **Water content and germination pattern**

Intact and scarified seeds have different patterns of increased water content during incubation in water (Fig. 1A). Water content in intact seeds levelled off after 3 d of incubation, whereas water contents of the scarified seeds continued to increase. The increase of water content was similar for both types of scarified seeds during the first 2 d. Later on, the increase of water content was faster in seeds scarified on the proximal end than on the distal end.

The range of germination for fresh seeds was 16–28%. After drying, intact seeds did not germinate during 2 weeks of incubation in water (Fig. 1B). When scarified on the proximal end, seeds began to germinate on the second day of incubation, with a maximum germination of 92% at day 9. Germination of seeds scarified on the distal end occurred through the hole made during scarification, i.e. opposite to the normal germination side. Radicle protrusion in these seeds started at the fifth day of incubation, and a maximum germination of 40% was reached at day 12. Delayed scarification of the proximal end at different times during incubation showed a gradual decrease of seed viability until day 15, when scarification caused only 2–4% of the seeds to germinate (Fig. 1C).


### **Membrane intactness in axes from non-germinated seeds**

Membrane intactness in axes of non-germinated seeds after 15 d of incubation was analysed by a spin-label technique. Briefly, the axes were placed in an aqueous solution of spin label (PDT) and broadening agent



**Figure 1.** Effect of scarification on water uptake (A) and germination (B and C) of *Lannea microcarpa* seeds incubated on moist filter paper at 30°C. Scarification of air-dried seeds was carried out on dry seeds (A and B), or 0, 3, 6, 9, 12 and 15 d after the start of incubation (C, arrows indicate the day of scarification). Scarification of dry seeds was carried out either on proximal or distal ends of the seed (A and B). Delayed scarification was carried out on the proximal end (C). Each point is the average water content of ten individual seeds  $\pm$  SE (A) or the average germination percentage of two batches of  $2 \times 25$  seeds each (B and C).

(ferricyanide ions) for labelling. The EPR spectrum of PDT in axis cells contained two superimposed components, originating from the spin label in the aqueous cytoplasm and lipid bodies (Fig. 2A and B, inserts). In leaky cells, the aqueous cytoplasmic signal is zero because of the broadening effect of ferricyanide ions that have entered the cells through damaged membranes, whereas the lipid signal remains unchanged. The upper part of the high field (right-hand-side of the spectrum) water line was used to estimate the aqueous component of the spectra (W),




**Figure 2.** Distribution of the W/L ratios calculated from the electron paramagnetic resonance (EPR) spectra (inserts) of perdeuterated TEMPONE (PDT) in individual axes from *Lannea microcarpa* seeds. Axes were excised from decoated seeds that were rehydrated at 30°C for 2 h ( $n = 20$ ; panel A), or from intact seeds after 15 d of incubation at 30°C ( $n = 50$ ; panel B). Inserts show representative EPR spectra of PDT in cells of an axis excised from a decoated seed after 2h of imbibition (top) or from a seed after 15 d of incubation (bottom). The spectra contain two superimposed components. Determination of the heights of the lines representing aqueous cytoplasm (W) and lipid bodies (L) is as indicated in the figure. The ratio W/L was used as an indicator of cellular integrity.

which is proportional to the total cytoplasmic volume of cells with intact membranes. The amplitude of the lipid component (L) is proportional to the total number of cells in a sample without respect to membrane intactness. By measuring W/L in the EPR spectra, the proportion of cells with intact plasma membranes was estimated (Golovina and Tikhonov, 1994; Golovina *et al.*, 1997; Neya *et al.*, 2004). Axes with higher W/L ratios (Fig. 2A, insert) contain more intact cells than those with low W/L values (Fig. 2B, insert). The extent of cell swelling during imbibition or cell extension during germination should increase the proportion of the water component in the spectra and thus W/L values.

Figures 2A and 2B show the distribution of W/L values of individual axes isolated from decoated, 2 h rehydrated seeds and from intact seeds after 15 d of incubation at 30°C, respectively. Axes from the 2 h incubated decoated seeds (control) had a distribution of W/L values ranging from 2 to 9. Because all the cells in the control samples are viable, the range of the distribution of W/L values is likely caused by the different extent of cell swelling. The distribution of W/L from the seeds incubated for 15 d was different. Thirty-eight percent of the axes from 15-d incubated seeds lost firmness and did not stain in tetrazolium chloride. EPR spectra of rotted axes did not have a water component (W/L = 0), indicating the absence of living cells. All other axes stained in tetrazolium chloride, but scarification on the proximal end did not induce their germination (Fig. 1C). The W/L value of 13 probably indicates cell extension, which occurred in a few cases (2–4% of which germinated, Fig. 1C). The W/L values below 2 are out of the range expected for viable axes. The rest of the axes had W/L values from 2 to 7, which was within the range of the viable control.

### Effect of drying and rehydration on seed respiration

Oxygen uptake and CO<sub>2</sub> evolution during drying of freshly depulped seeds from 0.43 g H<sub>2</sub>O g<sup>-1</sup> to the air-dried state are shown in Fig. 3A. Respiratory activity decreased with the loss of water to reach immeasurably low values *c.* 0.1 g H<sub>2</sub>O g<sup>-1</sup>. The respiratory quotient (RQ) was *c.* 0.7. On drying, there was no indication of unbalanced respiration caused by a possibly impeded O<sub>2</sub> uptake. During the first 2 d of rehydration of the air-dried seeds, respiration increased to values similar to those of the freshly depulped seeds at comparable water contents (Fig. 3B). Also, RQ values returned to approximately the same values. This indicates that the seeds were able to recruit sufficient O<sub>2</sub> to avoid fermentative metabolism, at least during the first 2 d of rehydration.




**Figure 3.** Effects of (A) drying of freshly depulped seeds and (B) rehydration of intact, air-dried seeds of *Lannea microcarpa* on respiratory gas exchange and respiratory quotient (RQ). Samples for gas analysis by gas chromatography were taken after 5 h of seed respiration in 7-ml flasks at 30°C. Conditions for the depulped seeds (A) were 8 d total drying time, 5 seeds per flask ( $n = 4$ )  $\pm$  SE, and for the rehydrating seeds (B) were 2 d rehydration, 1 seed per flask ( $n = 6$ )  $\pm$  SE.

### Effect of scarification on seed respiration

The gas exchange rate from scarified seeds (proximal end) was higher than that from intact seeds at day 1 of incubation, and continued to increase following radicle emergence at day 2 (Fig. 4A). The respiratory activity of the non-germinating intact seeds remained constant during days 1 through 6 after an initial increase due to rehydration, and then slightly increased, perhaps due to the decay of some of these seeds.

Seed scarification on the distal end also allows oxygen to have free access to the embryo. Because gas exchange measurements were performed on individual seeds, categories of non-germinating and germinating seeds were tested for respiratory activity separately. Those seeds that did not germinate in spite of scarification had the same amount of respiration as intact non-scarified seeds. The upsurge of respiration coincided with the emergence of the radicle (Fig. 4B).




**Figure 4.** Effect of scarification on respiratory gas exchange from rehydrated *Lannea microcarpa* seeds at 30°C. (A) Seeds were intact and did not germinate (open and closed circles) or were scarified on the proximal end (open and closed triangles), which allowed for emergence of the radicle from day 2 onwards; data are the average ( $\pm$  SE) of measurements of 6–10 individual seeds. (B) Seeds were scarified on the distal end of the seed and categorized according to whether they displayed radicle emergence at day 4–5 ( $n = 6$ ) or did not at day 5 ( $n = 4$ ). (C) Seeds were scarified on the proximal end at 3, 6 and 9 d of incubation; each point is the average value of 5 individual seeds. Samples for gas analysis by gas chromatography were taken after 5 h of seed respiration in 7 ml flasks at 30°C. Closed symbols, oxygen uptake; open symbols, carbon dioxide release.

Further support of these results was obtained in experiments with delayed scarification. Seed scarification on the proximal end at days 3, 6 and 9 of incubation led to an upsurge in respiration, but with decreasing intensity (Fig. 4C). The degree of respiration upsurge correlated with the percentage of germinated seeds (Fig. 1C). In all cases, the RQ was *c.* 1 for non-germinating seeds and *c.* 0.7 for germinating seeds (data not shown).


#### **Possible role of ethylene in the regulation of seed germination**

Figure 5 shows the evolution of ethylene from intact and scarified seeds (at their proximal ends) during incubation in water. The rate of ethylene production was determined by sampling headspace gas after 9 h of ethylene accumulation. After a peak in the production on day 1, the ethylene emanation by scarified seeds gradually decreased with time of incubation. The controls had high ethylene production at days 1 and 2, after which ethylene emanation fell to almost zero.

The effect of the 1-aminocyclopropane-1-carboxylic acid (ACC)-synthase inhibitor amino-ethoxyvinylglycine (AVG) on ethylene evolution is shown in Fig. 6A. Seeds scarified on the distal end were



**Figure 5.** Ethylene release by intact and scarified (proximal end) *Lannea microcarpa* seeds upon incubation at 30°C. Each point is the average value of the ethylene production of ten individual seeds  $\pm$  SE. Ethylene was allowed to accumulate in the headspace of 7 ml flasks for *c.* 9 h.



**Figure 6.** Effect of amino-ethoxyvinylglycine (AVG) on (A) ethylene biosynthesis and (B) percentage germination of *Lannea microcarpa* seeds upon incubation at 30°C. All seeds were scarified on the proximal end prior to rehydration in the AVG solutions. Each point in (A) is the average ethylene production of four individual seeds  $\pm$  SE. Ethylene was allowed to accumulate for c. 2 h. Each point in (B) is the average percentage germination of  $2 \times 25$  seeds.

incubated in solutions of a range of AVG concentrations. Ethylene production was determined in headspace gas after 2 h of accumulation. Inhibition of ethylene production by AVG was concentration dependent, and  $\geq 1$  mM AVG caused almost complete inhibition during the first 4 h of seed incubation. Inhibition of ethylene production decreased the rate and final percentage of germination (Fig. 6B). The inhibition of germination was concentration dependent; but even at high AVG concentrations, when ethylene production was almost zero (Fig. 6A), half the seeds still germinated (Fig. 6B).

### Seed coat composition

Using FT-IR, the main endocarp components were determined on the basis of specific IR-absorption

peaks. Figure 7A shows the IR-spectrum of material scraped from the inner layer of the endocarp. There was no evidence for the presence of lipids or proteins. Instead, a number of IR-absorption peaks could be linked to structural carbohydrates and lignin (Yu *et al.*, 2005).


The carbonyl C=O stretching absorbance at  $1737\text{ cm}^{-1}$  refers to the ester bonds of the hemicelluloses, lignins and cellulose, and the absorbance at  $1243\text{ cm}^{-1}$  points to the presence of hemicelluloses (Yu *et al.*, 2004, 2005). The peaks at 1595, 1505 and  $1461\text{ cm}^{-1}$  are associated with aromatic ring vibrations in lignin molecules (Dorado *et al.*, 2001; Sun *et al.*, 2004), whereas the main peak at  $1050\text{ cm}^{-1}$ , associated with C–O stretching, and another at  $1159\text{ cm}^{-1}$  can be assigned mainly to cellulose (Ilharco *et al.*, 1997; Sun *et al.*, 2004). The other peaks may be the result of both lignin and cellulose, since these compounds have overlapping peaks in this wavenumber region (Dorado *et al.*, 2001).

For comparison, the spectrum of kapok fibres is shown in Fig. 7A. It resembles to a large extent the spectrum of the inner endocarp layer of *L. microcarpa*. Kapok has a composition of c. 43% cellulose, 13–15% lignin and 32% hemicellulose (Brink and Escobin, 2003). From this similarity, it is assumed that the endocarp of *L. microcarpa* has approximately the same composition. Figure 7B shows representative spectra of material from the outer layer of the endocarp, isolated from fresh seeds ( $0.43\text{ g H}_2\text{O g}^{-1}$ ) and seeds after 10 months of dry storage ( $0.04\text{ g H}_2\text{O g}^{-1}$ ). There was no large difference in peak position and area under the peaks, as evidenced by an almost flat difference spectrum. FT-IR analysis was carried out with two or three specimens from three seeds, each scraped from the outer and inner layers of the endocarp. The endocarp of neem (*Azadirachta indica*) seed, removal of which promotes radicle emergence (Neya, unpublished data), exhibited an IR spectrum almost identical to that of the *L. microcarpa* endocarp (Fig. 7C). On the contrary, the IR-spectrum of the *Khaya senegalensis* seed coat differed considerably from that of the *L. microcarpa* endocarp, in that hemicelluloses were low and lignins were almost absent. Growing radicles of *K. senegalensis* always break through the seed coat easily (Neya, 2006).

## Discussion

### Endocarp impedes germination of *L. microcarpa* seeds

Germination of both fresh and dry seeds of *L. microcarpa* is very variable among and within seed accessions. Drying increases germination of seeds from green fruits, does not influence the germination



**Figure 7.** Fourier transform infrared spectra. (A) Material from the inner layer of the endocarp of air-dried *Lannea microcarpa* seeds after 10 months of storage (0% germination), with dry kapok fibres shown for comparison. The position of peaks associated with the carbonyl C=O stretching vibration ( $1737\text{ cm}^{-1}$ ), the lignin-specific aromatic ring vibrations ( $1461$ ,  $1505$ ,  $1595\text{ cm}^{-1}$ ) and hemicellulose ( $1243\text{ cm}^{-1}$ ) are indicated. (B) Comparison of material from the outer layer of endocarp from fresh *L. microcarpa* seeds ( $28 \pm 6\%$  germination) and after 10 months of storage, with the difference spectrum shown at the bottom of the panel. (C) Material scraped from the outer layer of the endocarp from *Azadirachta indica* and from the seed coat of *Khaya senegalensis*.

of seeds from green-red fruits and decreases the germination of seeds from purple-black fruits (Neya, 2006). The increase of seed germination with drying is related to seed maturation (Neya, 2006) or to loss of physiological dormancy (Pritchard *et al.*, 2004).

The decrease of seed germination with drying of mature seeds from purple-black fruits obviously relates to coat-imposed constraints, because mechanical scarification improved germination considerably (Fig. 1B). Differences in the extent of such constraints among seeds are likely to be the cause of the variability in germination.

The embryo of *L. microcarpa* seeds is enclosed by a seed coat (testa) and a fruit coat (endocarp). The seed coat is a thin, papery layer and can hardly impose any constraints on seed germination. However, the endocarp is a well-developed outer covering of the seeds (Baskin *et al.*, 2000), which is responsible for the constraints of germination. In spite of the fact that impermeability of the endocarp for water is common in the *Anacardiaceae* family that inhabits arid and semi-arid regions (Baskin and Baskin, 1998), the endocarp of *L. microcarpa* is water permeable, and the seed imbibes freely during incubation (Pritchard *et al.*, 2004; Neya, 2006; Fig. 1A). The production of inhibitors by coats is also unlikely, because the removal of only a small part of the endocarp during scarification allows seed germination (Fig. 1B). The possible involvement of three other mechanisms was tested: interference with gas exchange, retention of inhibitors and mechanical constraints.

### Interference with gas exchange

Permeability of the endocarp to water does not necessarily rule out its ability to interfere with gas exchange (Bewley and Black, 1994). Drying of *L. microcarpa* seeds decreased their subsequent germination from 16–28% to 0%. If the permeability of the endocarp to respiratory gases changes during their drying, respiration characteristics will also change, leading to the inhibition of respiration and stimulation of fermentation (Wager, 1974; Rolletschek *et al.*, 2002). The end products of such fermentation may be damaging to the seed, causing loss of viability (Fig. 1C). A high  $\text{CO}_2/\text{O}_2$  ratio (RQ), reaching values considerably above 1, is diagnostic for the lack of sufficient oxygen. From the low RQ values ( $<1$ ) in Fig. 3, it can be deduced that sufficient oxygen is available during rehydration of intact seeds. Moreover, the rates of gas exchange were almost the same at similar water contents in both dehydrating fresh seeds and rehydrating seeds (Fig. 3). Thus, the endocarp does not impede respiratory gas exchange in imbibed *L. microcarpa* seeds.

However, endocarp permeability for oxygen may be still not enough to allow radicle emergence, when seed demands for oxygen increase. Seeds, scarified on the proximal end attained much higher rates of respiration than intact seeds (Fig. 4A). This might be interpreted as the result of free access to oxygen due to

scarification. If this were the case, scarification on the distal end would also cause the immediate increase in respiration. However, the increase of respiration took place only after radicle protrusion (Fig. 4B). In general, high respiratory activity appears to be closely associated with radicle protrusion, but not with making holes in the endocarp during scarification (Fig. 4C). This is comparable with the results of Derkx *et al.* (1993), who found that dormancy breaking in seeds of *Sisymbrium officinale* does not cause an upsurge in O<sub>2</sub> uptake, but completion of germination does. Radicle protrusion as the result of completed germination also caused a decrease of the RQ from 1 (carbohydrate oxidation) to 0.7 (oil oxidation). This indicates the switch of metabolism from using carbohydrates (sugars) as substrates for respiration to oil, which is a sign of the mobilization of the major storage reserves during seedling growth.

### Ethylene production

The effect of the endocarp on seed germination may also be connected with insufficient ethylene production (Bewley and Black, 1994). Ethylene is produced by the embryo during the initial stages of germination and has a close relationship with cell elongation (Abeles, 1986; Sánchez-Calle *et al.*, 1989). In the ethylene biosynthesis pathway, the conversion of 1-aminocyclopropane-1-carboxylic acid (ACC) to ethylene is oxygen dependent (Adams and Yang, 1979). The release of seeds from dormancy is associated with enhanced consumption of oxygen, which is not only involved in respiration, but also in other metabolic processes playing a role in the breaking of dormancy. Insufficient oxygen supply may decrease ethylene production and influence cellular extension and radicle protrusion. Therefore, ethylene emanation from intact seeds of *L. microcarpa* was compared with that of seeds scarified on their proximal end. A peak in ethylene in the headspace gas over both intact and scarified seeds was noted early during incubation; after that the evolution fell to low values (Fig. 5). This demonstrates that the endocarp is permeable to ethylene and does not interfere with ethylene production by the seed.

Usually the peak of ethylene production is associated with radicle protrusion (Kepczynski and Kepczynska, 1997). In the case of scarified seeds, ethylene emanation decreased considerably at day 2, when radicles started to emerge (Fig. 1B and Fig. 5). The ethylene production by scarified seeds remained in the range of 0.5–1 nl h<sup>-1</sup>g<sup>-1</sup> at days 4 and 5, which could be associated with radicle growth occurring at the same time. On the contrary, intact seeds continued to produce ethylene on day 2 at the same rate as on day 1. However, at day 3 and after, intact seeds did not

produce ethylene, even though they were still viable, and scarification on this day led to 78% germination (Fig. 1C). At day 3, the increase in water content levelled off. Probably, both events were associated with the inability of the cells to elongate. Thus, the endocarp does not interfere with ethylene production.

While almost all seeds produce ethylene during germination, requirements for this hormone vary among species (Lalonde and Saini, 1992). To verify whether ethylene production is necessary for the germination of *L. microcarpa* seeds, the effect of amino-ethoxyvinylglycine (AVG), an ACC-synthase inhibitor, on ethylene evolution and seed germination was followed. In germinating seeds, ethylene is known to be synthesized exclusively from ACC produced *de novo* (Kepczynski and Kepczynska, 1997). The inhibition of the ethylene production by AVG (Fig. 6A) shows that ethylene in *L. microcarpa* seeds is also produced via *de novo* synthesized ACC and not by conversion of pre-existing ACC. Germination decreased with the increased concentration of AVG (Fig. 6B). However, even when the production of ethylene was almost abolished by a high concentration of AVG, 50% of seeds were still able to germinate. These data show that the requirements for endogenous ethylene during germination of *L. microcarpa* seeds vary among seeds.

### Mechanical constraints and retention of inhibitors

Because the endocarp in *L. microcarpa* seeds is hard, tough tissue, it is reasonable to expect mechanical constraints as the main mechanism of inhibition of seed germination. The radicle always emerged through the hole made during scarification, even when it was made on the end opposite from where the radicle normally emerges – evidence for a strong mechanical restriction of germination.

The continuing increase of water content in seeds scarified on the proximal end shows that this treatment allows cellular extension followed by radicle protrusion at day 2. In intact seeds the endocarp prevents cellular extension, and water content increases only due to osmotic forces, and thus levelled off after 3 d of incubation. However, seeds scarified on the distal end had the same pattern of water content increase as seeds scarified on the proximal end, despite the observation that radicle protrusion occurred 3 d later (Fig. 1A and B). In this case, radicle extension also took place early during incubation, but inside the seed, enabling the radicle to find the hole for visible protrusion. This occurred in only 40% of the seeds. Probably, only the most vigorous embryos can emerge from the hole at the distal end of the seeds.

The different patterns of germination of seeds scarified on the proximal and distal ends also suggest that the retention of inhibitors is not involved in lack of germination of intact seeds. If this were the case, the position of the hole made by scarification would not influence the pattern of germination because of inhibitor leakage through the hole.

### **Loss of viability of incubated intact seeds**

The percentage of radicle protrusion decreased with time of seed incubation in water before scarification (Fig. 1C). Germination fell to 2–4% when the seeds were scarified at day 15. Apart from 38% soft, obviously rotten seeds after 15 d of incubation at 30°C, the remaining 62% of seeds were still firm and looked viable. However, there was no sign of cellular extension. This follows from the fact that the majority of axes from the firm seeds had W/L values of 2–7 (Fig. 2B), which is within the range established for just-rehydrated, but not extended, viable axes (from 2–9) (Fig. 2A). Only in few cases (*c.* 4%) did W/L increase to 13 or decrease to 1 (Fig. 2B). W/L = 1 indicates a loss of membrane integrity in the majority of cells; W/L = 13 relates to cellular extension. If cellular extension occurred in all the axes, the distribution of W/L values would be well above 9 after 15 d of rehydration. The inability of cells to extend is also supported by the limited increase of water content in non-scarified seeds (Fig. 1A). Thus, endocarp-imposed mechanical constraints prevent not only radicle emergence, but also cell extension before protrusion.

The inability of seeds to germinate results in their slow decay, but the reason for this is unclear, because the endocarp does not interfere with water and gas exchange. Also, ethylene is not involved in the slow decay of the hydrated, intact seeds, because radicle emergence could be evoked by delayed scarification of the incubated, intact seeds (Fig. 1C), long after ethylene had dissipated. The loss of seed viability after long incubation without germination of intact seeds may be associated with the exhaustion of the sugar pool as a respiratory substrate for the axes and inability to use other substrates (oil), because the program of seedling growth is not switched on. The CO<sub>2</sub> production of *c.* 60 nmol min<sup>-1</sup> g<sup>-1</sup> at 30°C by hydrated, intact seeds (Fig. 4) leads to a (calculated) weight loss of *c.* 0.3% of the DM per day and consequently to 4.5–5% after 15 d.

### **Relationships between endocarp composition and hardness**

*In situ* Fourier transform infrared microspectroscopy (FTIR) was used to find the relationships between

endocarp composition and its hardness. The hard endocarps of *L. microcarpa* (Fig. 7A) and *A. indica* (Fig. 7C) contain lignin and hemicellulose. On the contrary, *K. senegalensis* seeds do not have such an endocarp; their seed coats are soft and do not contain lignin and hemicellulose (Fig. 7C).

The FTIR spectra of the endocarp of *L. microcarpa* are not changed upon drying and dry storage of seeds (Fig. 7B), while hardness increases and does not decrease after rehydration. There may be irreversible structural changes in the endocarp during drying. This may relate to covalent binding of lignin to hemicellulose, which confers mechanical strength to the cell wall. The cross-linking of polysaccharides by lignin is a potential obstacle for water absorption by the cell wall (<http://en.wikipedia.org/wiki/Lignin>), which may prevent endocarp swelling during rehydration.

### **Ecological and practical consequences of *L. microcarpa* seed germination behaviour**

*L. microcarpa* seeds are shed at the beginning of the rainy season (Arbonnier, 2002), i.e. favourable conditions for immediate germination. They are not subjected to maturation drying, but can withstand dehydration and maintain viability in dry state for 2 years or more (Neya, 2006). However, drying may cause the endocarp-imposed constraints to germination. It is debatable as to whether endocarp strengthening can be considered as secondary dormancy, which aids in the dispersal of *L. microcarpa* seeds in space and time, or a fatal constraint, such that dry seeds will never germinate in nature. Nevertheless, non-germinated dry *L. microcarpa* seeds can still be used for reproduction in forestry practice if they are scarified on the micropylar end of the seed before germination. The loss of viability of rehydrated seeds (Fig. 1C) within 15 d at 30°C is not a likely strategy followed in nature.

### **Acknowledgements**

The fellowship of the Dutch Organization for the Advancement of Tropical Research (WOTRO) for O.N. is gratefully acknowledged.

### **References**

- Abeles, F.B. (1986) Role of ethylene in *Lactuca sativa* cv 'Grand Rapids' seed germination. *Plant Physiology* **81**, 780–787.
- Adams, D.O. and Yang, S.F. (1979) Ethylene biosynthesis: identification of 1-aminocyclopropane-1-carboxylic acid as an intermediate in the conversion of methionine

- to ethylene. *Proceedings of the National Academy of Sciences, USA* **76**, 170–174.
- Adkins, S.W., Bellairs, S.M. and Loch, D.S.** (2002) Seed dormancy mechanisms in warm season grass species. *Euphytica* **126**, 13–20.
- Arbonnier, M.** (2002) *Arbres, arbustes et lianes des zones sèches d'Afrique de l'Ouest* (2nd edition). Paris, CIRAD, MNHN.
- Baskin, C.C. and Baskin, J.M.** (1998) *Seeds: Ecology, biogeography, and evolution of seed dormancy and germination*. San Diego, Academic Press.
- Baskin, J.M. and Baskin, C.C.** (2000) Evolutionary considerations of claims for physical dormancy-break by microbial action and abrasion by soil particles. *Seed Science Research* **10**, 409–413.
- Baskin, J.M. and Baskin, C.C.** (2004) A classification system for seed dormancy. *Seed Science Research* **14**, 1–16.
- Baskin, J.M., Baskin, C.C. and Li, X.** (2000) Taxonomy, ecology, and evolution of physical dormancy in seeds. *Plant Species Biology* **15**, 139–152.
- Bewley, J.D. and Black, M.** (1994) *Seeds. Physiology, development and germination* (2nd edition). New York, Plenum Press.
- Brink, M. and Escobin, R.P.** (2003) *Plant resources of south-east Asia (PROSEA) 17. Fibre plants*. Leiden, Backhuys Publishers.
- Daws, M.I., Gaméné, C.S., Sacandé, M., Pritchard, H.W., Groot, S.P.C. and Hoekstra, F.A.** (2004) Desiccation and storage of *Lannea microcarpa* seeds from Burkina Faso. pp. 32–39 in Sacandé, M.; Jøker, D.; Dulloo, M.E.; Thomsen, K.A. (Eds) *Comparative storage biology of tropical tree seeds*. Rome, International Plant Genetic Resources Institute.
- Derx, M.P.M., Smidt, W.J., Van der Plas, L.H.W. and Karszen, C.M.** (1993) Changes in dormancy of *Sisymbrium officinale* seeds do not depend on changes in respiratory activity. *Physiologia Plantarum* **89**, 707–718.
- Dorado, J., Almendros, G., Field, J.A. and Sierra-Alvarez, R.** (2001) Infrared spectroscopy analysis of hemp (*Cannabis sativa*) after selective delignification by *Bjerkandera* sp. at different nitrogen levels. *Enzyme and Microbial Technology* **28**, 550–559.
- Flynn, S., Turner, R.M. and Dickie, J.B.** (2004) Seed information database. Available at Web site <http://www.kew.org/data/sid> (release 6.0, October 2004).
- Golovina, E.A. and Tikhonov, A.N.** (1994) The structural differences between the embryos of viable and nonviable wheat seeds as studied by EPR spectroscopy of lipid-soluble spin labels. *Biochimica et Biophysica Acta* **1190**, 385–392.
- Golovina, E.A., Tikhonov, A.N. and Hoekstra, F.A.** (1997) An electron paramagnetic resonance spin-probe study of membrane permeability changes with seed aging. *Plant Physiology* **114**, 383–389.
- Hong, T.D., Linington, S. and Ellis, R.H.** (1996) *Seed storage behaviour: A compendium. Handbook for genebanks, No. 4*. Rome, International Plant Genetic Resources Institute.
- Illharco, L.M., Garcia, A.R., Lopes da Silva, J. and Vieira Ferreira, L.F.** (1997) Infrared approach to the study of adsorption on cellulose: influence of cellulose crystallinity on the adsorption of benzophenone. *Langmuir* **13**, 4126–4132.
- ISTA (International Seed Testing Association)** (1999) International rules for seed testing. Supplement. *Seed Science and Technology* **27**, 98–107.
- Jones, H.D.** (1999) Seeds get a wake-up call. Dormancy, a survival mechanism for seeds, or a useful tool for agriculture. *The Biologist* **46**, 65–69.
- Kepczynski, J. and Kepczynska, E.** (1997) Ethylene in seed dormancy and germination. *Physiologia Plantarum* **101**, 720–726.
- Lalonde, S. and Saini, H.S.** (1992) Comparative requirement for endogenous ethylene during seed germination. *Annals of Botany* **69**, 423–428.
- Leprince, O. and Hoekstra, F.A.** (1998) The responses of cytochrome redox state and energy metabolism to dehydration support a role for cytoplasmic viscosity in desiccation tolerance. *Plant Physiology* **118**, 1253–1264.
- Li, X., Baskin, J.M. and Baskin, C.C.** (1999a) Anatomy of two mechanisms of breaking dormancy by experimental treatments in seeds of two North American *Rhus* species (Anacardiaceae). *American Journal of Botany* **86**, 1505–1511.
- Li, X., Baskin, J.M. and Baskin, C.C.** (1999b) Seed morphology and physical dormancy of several North American *Rhus* species (Anacardiaceae). *Seed Science Research* **9**, 247–258.
- Manzano, P., Malo, J.E. and Peco, B.** (2005b) Sheep gut passage and survival of Mediterranean shrub seeds. *Seed Science Research* **15**, 21–28.
- Neya O.** (2006) Conservation of tree seeds from tropical dry-lands. PhD Thesis, Wageningen University, The Netherlands.
- Neya, O., Golovina, E.A., Nijse, J. and Hoekstra, F.A.** (2004) Ageing increases the sensitivity of neem (*Azadirachta indica*) seeds to imbibitional stress. *Seed Science Research* **14**, 205–217.
- Pritchard, W.H., Daws, M.I., Fletcher, B.J., Gaméné, C.S., Msanga, H.P. and Omondi, W.** (2004) Ecological correlates of seed desiccation tolerance in tropical African dryland trees. *American Journal of Botany* **91**, 863–870.
- Rolletschek, H., Borisjuk, L., Koschorreck, M., Wobus, U. and Weber, H.** (2002) Legume embryos develop in a hypoxic environment. *Journal of Experimental Botany* **53**, 1099–1107.
- Sánchez-Calle, I.M., Delgado, M.M., Bueno, M., Diaz-Miguel, M. and Matilla, A.** (1989) The relationships between ethylene production and cell elongation during the initial growth period of chick-pea seeds (*Cicer arietinum*). *Physiologia Plantarum* **76**, 569–574.
- Sun, X.F., Sun, R.C. and Sun, X.J.** (2004) Acetylation of sugarcane bagasse using NBS as a catalyst under mild reaction conditions for the production of oil sorption-active materials. *Bioresource Technology* **95**, 343–350.
- Van Assche, J.A., Debucquoy, K.L.A. and Rommens, W.A.F.** (2003) Seasonal cycles in the germination capacity of buried seeds of some *Leguminosae* (Fabaceae). *New Phytologist* **158**, 315–323.
- Wager, H.G.** (1974) The effect of subjecting peas to air enriched with carbon dioxide 2. Respiration and the metabolism of the major acids. *Journal of Experimental Botany* **25**, 338–351.

- Yu, P.Q., McKinnon, J.J., Christensen, C.R. and Christensen, D.A.** (2004) Using synchrotron transmission FTIR microspectroscopy as a rapid, direct, and nondestructive analytical technique to reveal molecular microstructural-chemical features within tissue in grain barley. *Journal of Agricultural and Food Chemistry* **52**, 1484–1494.
- Yu, P.Q., Christensen, C.R., Christensen, D.A. and McKinnon, J.J.** (2005) Ultrastructural-chemical makeup of yellow-seeded (*Brassica rapa*) and brown-seeded

(*Brassica napus*) canola within cellular dimensions, explored with synchrotron reflection FTIR microspectroscopy. *Canadian Journal of Plant Science* **85**, 533–541.

Received 31 May 2007  
accepted after revision 19 November 2007  
© 2008 Cambridge University Press

---