

increase of the above specified decomposition products accounted completely for the observed sabinene decrease, the total sum of the measured monoterpene compounds was not affected by this decomposition. To specify the composition of monoterpene compounds that are emitted from European beech, a correction factor was applied to the relevant compounds taking a cartridge storage time of more than 7 days into account. In contrast, previous experiments performed for the GC-FID cartridges did not indicate a similar decomposition of sabinene on the respective VOC samples. However, during the present study some of the potential decomposition products were not investigated by the GC-FID analysis and assuming a similar decomposition process as observed for the GC-MS cartridges may therefore lead to an underestimation of the total sum of monoterpenes measured by GC-FID of up to 20%.

2.1.4. Emission Algorithms

[15] The emission of monoterpenes from European beech was simulated by two different light and temperature dependent algorithms. The first algorithm applied (in the following referred to as G97) was developed for isoprene emission by *Guenther et al.* [1993, 1995] and *Guenther* [1997]. In previous studies, isoprene emission has been shown to be triggered by light as a result of the close link between its emission and production from photosynthetic precursor compounds. However, several authors demonstrated that the latter algorithm may also be used to calculate the emission of monoterpenes that are released directly upon their production [e.g., see *Kesselmeier et al.*, 1996; *BEMA-Project*, 1997; *Ciccioli et al.*, 1997; *Kuhn et al.*, 2002b; *Kuhn et al.*, 2004].

[16] The G97 algorithm assumes a hyperbolic increase of VOC emissions to light intensity, leading to a saturation effect. With respect to leaf temperature the algorithm assumes enzymatic processes leading to a temperature optimum of the VOC emission at temperatures of 39°C. Within the algorithm, the light dependent term of the G97 function (in the following referred to as C_L) is specified by Formula 1. The temperature dependent term (in the following referred to as C_T) is specified by Formula 2. To calculate the actual VOC emission, both factors (C_L and C_T) are linked by multiplication with a standard emission factor (in the following referred to as SEF) that describes the basal VOC emission at standard light and standard temperature conditions (1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 30°C, see Formula 3).

(1) Light dependent term of the G97 function:

$$C_L = \frac{\alpha \cdot C_{L1} \cdot L}{\sqrt{1 + \alpha^2 \cdot L^2}}$$

(2) Temperature dependent term of the G97 function:

$$C_T = \frac{\exp\left(\frac{C_{T1} \cdot (T - T_S)}{R \cdot T_S \cdot T}\right)}{C_{T3} + \exp\left(\frac{C_{T2} \cdot (T - T_M)}{R \cdot T_S \cdot T}\right)}$$

(3) Calculation of the actual VOC emission by the G97 function:

$$\text{VOC emission} = \text{SEF} \cdot C_L \cdot C_T$$

($\alpha = 0.0027$, $C_{L1} = 1.066$, C_L = light dependent term of the G97 function, C_T = temperature dependent term of the G97 function, $C_{T1} = 95,000 \text{ J mol}^{-1}$, $C_{T2} = 230,000 \text{ J mol}^{-1}$, $C_{T3} = 0.961$, L = actual light intensity ($\mu\text{mol m}^{-2} \text{s}^{-1}$), R = universal gas constant ($8.314 \text{ J K}^{-1} \text{mol}^{-1}$), SEF = standard emission factor, T = leaf temperature ($^{\circ}\text{K}$), $T_M = 314^{\circ}\text{K}$, $T_S = 303^{\circ}\text{K}$).

[17] The second algorithm (in the following referred to as S97) that was applied to the present data set was developed to describe the emission of monoterpenes from sunflower and European beech as a function of light and temperature by *Schuh et al.* [1997]. In comparison to the G97 function the S97 algorithm assumes also a saturation effect of VOC emission at high light intensities. However, in contrast to the G97 simulation which assumes a hyperbolic increase of VOC emission, the S97 algorithm assumes an allosteric enzyme regulation leading to a sigmoid increase of VOC emission at lower light intensities. This light dependent term (in the following referred to as $C_{L(S)}$) of the S97 algorithm is described by Formula 4. The temperature dependent term, describing the instantaneous emission of monoterpenes, is identical to the G97 function (see C_T , Formula 2). In addition to the instantaneous emission of monoterpenes, the original S97 function assumes also a release of VOCs from unspecific storage pools (Formula not shown). However, in the present study a release of monoterpenes from unspecific storage pools was not detectable. Therefore the storage pool term of the S97 algorithm was neglected, as recommended for sabinene emission from European beech by *Schuh et al.* [1997]. To calculate the actual VOC emission by the S97 function, the respective terms were linked by multiplication with a standard emission factor (in the following referred to as Φ_{LT}) as described by Formula 5, yielding a modified form of the original S97 algorithm that neglects a monoterpene release from storage pools.

(4) Light dependent term of the S97 function:

$$C_{L(S)} = C_{L1} \cdot \left(\frac{\alpha \cdot L}{\sqrt{1 + \alpha^2 \cdot L^2}} \right)^2$$

(5) Calculation of the actual VOC emission by the S97 function:

$$\text{VOC emission} = \Phi_{LT} \cdot C_{L(S)} \cdot C_T$$

($\alpha = 0.0027$, $C_{L1} = 1.066$, $C_{L(S)}$ = light dependent term of the S97 function, C_T = temperature dependent term of the G97 function, L = actual light intensity ($\mu\text{mol m}^{-2} \text{s}^{-1}$), Φ_{LT} = standard emission factor).

[18] As shown by Formulas 3 and 5, both standard emission factors (SEF or Φ_{LT}) can be calculated from the slope of linear regression of the measured VOC emission to the product of the light and temperature dependent terms of the respective algorithm (note that standard conditions correspond to 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 30°C for both algorithms). Since both algorithms assume a cessation of VOC emission in the dark, a linear regression without bias ($y = 0$) was used for the calculation of both emission factors.

- Kuhn, U., S. Rottenberger, T. Biesenthal, C. Ammann, A. Wolf, G. Schebeske, S. T. Oliva, T. M. Tavares, and J. Kesselmeier (2002a), Exchange of short-chain monocarboxylic acids by vegetation at a remote tropical forest site in Amazonia, *J. Geophys. Res.*, *107*(D20), 8069, doi:10.1029/2000JD000303.
- Kuhn, U., S. Rottenberger, T. Biesenthal, A. Wolf, G. Schebeske, P. Ciccioli, E. Brancaleoni, M. Frattoni, T. M. Tavares, and J. Kesselmeier (2002b), Isoprene and monoterpene emissions of Amazonian tree species during the wet season: Direct and indirect investigations on controlling environmental functions, *J. Geophys. Res.*, *107*(D20), 8071, doi:10.1029/2001JD000978.
- Kuhn, U., S. Rottenberger, T. Biesenthal, A. Wolf, G. Schebeske, P. Ciccioli, E. Brancaleoni, M. Frattoni, T. M. Tavares, and J. Kesselmeier (2004), Seasonal differences in isoprene and light-dependent monoterpene emission by Amazonian tree species, *Global Change Biol.*, *10*, 663–682.
- Kuhn, U., T. Dindorf, C. Ammann, S. Rottenberger, P. Guyon, R. Holzinger, S. Ausma, T. Kenntner, F. Helleis, and J. Kesselmeier (2005), Design and field application of an automated cartridge sampler for VOC concentration and flux measurements, *J. Environ. Monitor.*, *7*, 568–576.
- Lenz, R., R. Köble, and G. Seufert (2001), Species-based mapping of biogenic emissions in Europe: Case study Italy, paper presented at 8th European Symposium on the Physico-chemical Behaviour of Atmospheric Pollutants, Joint Res. Cent., Inst. for Environ. and Sustainability, Torino, Italy.
- Lichtenthaler, H. K. (1999), The 1-deoxy-D-xylulose-5-phosphate pathway of isoprenoid biosynthesis in plants, *Annu. Rev. Plant Phys.*, *50*, 47–65.
- Llusia, J., and J. Peñuelas (2000), Seasonal patterns of terpene content and emission from seven Mediterranean woody species in field conditions, *Am. J. Bot.*, *87*, 133–140.
- Loreto, F., P. Ciccioli, A. Cecinato, E. Brancaleoni, M. Frattoni, and D. Tricoli (1996), Influence of environmental factors and air composition on the emission of alpha-pinene from *Quercus ilex* leaves, *Plant Physiol.*, *110*, 267–275.
- Loreto, F., P. Ciccioli, E. Brancaleoni, M. Frattoni, and S. Delfino (2000), Incomplete C-13 labelling of alpha-pinene content of *Quercus ilex* leaves and appearance of unlabelled C in alpha-pinene emission in the dark, *Plant Cell Environ.*, *23*, 229–234.
- Monson, R. K., P. C. Harley, M. E. Litvak, M. Wildermuth, A. B. Guenther, P. R. Zimmerman, and R. Fall (1994), Environmental and developmental controls over the seasonal pattern of isoprene emission from aspen leaves, *Oecologia*, *99*, 260–270.
- Moukhtar, S., B. Bessagnet, L. Rouil, and V. Simon (2005), Monoterpene emissions from Beech (*Fagus sylvatica*) in a French forest and impact on secondary pollutants formation at regional scale, *Atmos. Environ.*, *39*, 3535–3547.
- Niinemets, U., and M. Reichstein (2002), A model analysis of the effects of nonspecific monoterpenoid storage in leaf tissues on emission kinetics and composition in Mediterranean sclerophyllous *Quercus* species, *Global Biogeochem. Cycles*, *16*(4), 1110, doi:10.1029/2002GB001927.
- Niinemets, U., K. Hauff, N. Bertin, J. D. Tenhunen, R. Steinbrecher, and G. Seufert (2002a), Monoterpene emissions in relation to foliar photosynthetic and structural variables in Mediterranean evergreen *Quercus* species, *New Phytol.*, *153*, 243–256.
- Niinemets, U., G. Seufert, R. Steinbrecher, and J. D. Tenhunen (2002b), A model coupling foliar monoterpene emissions to leaf photosynthetic characteristics in Mediterranean evergreen *Quercus* species, *New Phytol.*, *153*, 257–275.
- Niinemets, U., F. Loreto, and M. Reichstein (2004), Physiological and physicochemical controls on foliar volatile organic compound emissions, *Trends Plant Sci.*, *9*, 180–186.
- Olson, J. (1992), World ecosystems (WE1.4): Digital raster data on a 10 minute geographic 1080 × 2160 grid square, in *Global Ecosystem Database*, NOAA, National Geophysical Data Center, Boulder, Colo.
- Owen, S. M., P. Harley, A. Guenther, and C. N. Hewitt (2002), Light dependency of VOC emissions from selected Mediterranean plant species, *Atmos. Environ.*, *36*, 3147–3159.
- Peñuelas, J., and J. Llusia (1999), Seasonal emission of monoterpenes by the Mediterranean tree *Quercus ilex* in field conditions: Relations with photosynthetic rates, temperature and volatility, *Physiol. Plantarum*, *105*, 641–647.
- Peuke, A. D., C. Schraml, W. Hartung, and H. Rennenberg (2002), Identification of drought-sensitive beech ecotypes by physiological parameters, *New Phytol.*, *154*, 373–387.
- Rinne, H. J. I., A. B. Guenther, J. P. Greenberg, and P. C. Harley (2002), Isoprene and monoterpene fluxes measured above Amazonian rainforest and their dependence on light and temperature, *Atmos. Environ.*, *36*, 2421–2426.
- Roeckner, E., K. Arpe, L. Bengtsson, M. Christoph, M. Claussen, L. Dümenil, M. Esch, M. Gioletta, U. Schlese, and U. Schulzweida (1996), The atmospheric general circulation model ECHAM-4: Model description and simulation of the present-day climate, *Rep. 218*, Max Planck Inst. for Meteorol., Hamburg, Germany.
- Sabillon, D., and L. V. Cremades (2001), Diurnal and seasonal variation of monoterpene emission rates for two typical Mediterranean species (*Pinus pinea* and *Quercus ilex*) from field measurements—relationship with temperature and PAR, *Atmos. Environ.*, *35*, 4419–4431.
- Schäfer, L., J. Kesselmeier, and G. Helas (1992), Formic and Acetic acid emission from conifers measured with a “cuvette” technique, in *CeC Air Pollution Research 39: Field Measurements and Interpretation of Species Related to Photooxidants and Acid Deposition*, edited by G. Angeletti, S. Beilke, and J. Slanina, pp. 319–323, Eur. Comm., Brussels.
- Schuh, G., A. C. Heiden, T. Hoffmann, J. Kahl, P. Rockel, J. Rudolph, and J. Wildt (1997), Emissions of volatile organic compounds from sunflower and beech: Dependence on temperature and light intensity, *J. Atmos. Chem.*, *27*, 291–318.
- Seufert, G., et al. (1997), An overview of the Castelporziano experiments, *Atmos. Environ.*, *31*, 5–17.
- Simpson, D., et al. (1999), Inventorying emissions from nature in Europe, *J. Geophys. Res.*, *104*, 8113–8152.
- Solmon, F., C. Sarrat, D. Serca, P. Tulet, and R. Rosset (2004), Isoprene and monoterpenes biogenic emissions in France: Modeling and impact during a regional pollution episode, *Atmos. Environ.*, *38*, 3853–3865.
- Spirig, C., A. Neftel, C. Ammann, J. Dommen, W. Grabmer, A. Thielmann, A. Schaub, J. Beauchamp, A. Wisthaler, and A. Hansel (2005), Eddy covariance flux measurements of biogenic VOCs during ECHO 2003 using proton transfer reaction mass spectrometry, *Atmos. Chem. Phys.*, *5*, 465–481.
- Staudt, M., and N. Bertin (1998), Light and temperature dependence of the emission of cyclic and acyclic monoterpenes from holm oak (*Quercus ilex* L.) leaves, *Plant Cell Environ.*, *21*, 385–395.
- Staudt, M., and G. Seufert (1995), Light-dependent emission of monoterpenes by Holm Oak (*Quercus-Ilex* L.), *Naturwissenschaften*, *82*, 89–92.
- Staudt, M., S. Rambal, R. Joffre, and J. Kesselmeier (2002), Impact of drought on seasonal monoterpene emissions from *Quercus ilex* in southern France, *J. Geophys. Res.*, *107*(D21), 4602, doi:10.1029/2001JD002043.
- Staudt, M., R. Joffre, and S. Rambal (2003), How growth conditions affect the capacity of *Quercus ilex* leaves to emit monoterpenes, *New Phytol.*, *158*, 61–73.
- Steinbrecher, R., W. Schürmann, A. M. Schreiner, and H. Ziegler (1993), Terpenoid emissions from common oak (*Quercus robur* L.) and Norway spruce (*Picea abies* L. Karst), in *CeC Air Pollution Research 47: General Assessment of Biogenic Emissions and Deposition of Nitrogen Compounds, Sulfur Compounds, and Oxidants in Europe*, edited by G. Angeletti, J. Slanina, and S. Beilke, pp. 251–257, Eur. Comm., Brussels.
- Street, R. A., S. Owen, S. C. Duckham, C. Boissard, and C. N. Hewitt (1997), Effect of habitat and age on variations in volatile organic compound (VOC) emissions from *Quercus ilex* and *Pinus pinea*, *Atmos. Environ.*, *31*, 89–100.
- Thomas, F. M. (2000), Growth and water relations of four deciduous tree species (*Fagus sylvatica* L., *Quercus petraea* [MATT.] LIEBL., *Q.-pubescens* WILLD., *Sorbus aria* [L.] CR.) occurring at Central-European tree-line sites on shallow calcareous soils: Physiological reactions of seedlings to severe drought, *Flora*, *195*, 104–115.
- Tollsten, L., and P. M. Müller (1996), Volatile organic compounds emitted from beech leaves, *Phytochemistry*, *43*, 759–762.
- Weiss, A., and J. M. Norman (1985), Partitioning solar-radiation into direct and diffuse, visible and near-infrared components, *Agric. Forest Meteorol.*, *34*, 205–213.
- Went, F. W. (1960), Blue hazes in the atmosphere, *Nature*, *187*, 641–643.

P. Ciccioli, Istituto di Metodologie Chimiche, Area della Ricerca del CNR di Montelibretti, I-00019 Monterotondo Scalo, Italy.

T. Dindorf, J. Kesselmeier, U. Kuhn, and G. Schebeske, Department of Biogeochemistry, Max Planck Institute for Chemistry, Johann-Joachim Becher Weg 27, D-55128 Mainz, Germany. (dindorf@mpch-mainz.mpg.de)

L. Ganzeveld, Department of Atmospheric Chemistry, Max Planck Institute for Chemistry, Johann-Joachim Becher Weg 27, D-55128 Mainz, Germany.

C. Holzke, Research Center Jülich GmbH, Institute II: Troposphere, D-52425 Jülich, Germany.

R. Köble and G. Seufert, Joint Research Center Ispra, Institute for Environment and Sustainability, T.P. 051, I-21020 Ispra, Italy.