


Fulmar Litter EcoQO Monitoring in the Netherlands 1979-2007 in relation to EU Directive 2000/59/EC on Port Reception Facilities

J.A. van Franeker, A. Meijboom, M. de Jong
& H. Verdaat

Report nr: C032/09



Institute for Marine Resources and Ecosystem Studies

Wageningen *IMARES*

Location Texel

Client: Ministry of Transport, Public Works and Water Management (VenW)
Directorate-General for Civil Aviation and Maritime Affairs (DGLM)
PO Box 20904, 2500 EX Den Haag, The Netherlands
Contact: Drs. T.M. van Bommel-Reehoorn
tel: + 31 (0) 70 351 1889
@ Tineke.van.Bommel@minvenw.nl
contract nr: DGLM 08001 / 4500125534
'Graadmeter Stormvogel-Zwerfvuil 2007'

Publication Date: 23 april 2009

- Wageningen **IMARES** conducts research providing knowledge necessary for the protection, harvest and usage of marine and coastal areas.
- Wageningen **IMARES** is a knowledge and research partner for governmental authorities, private industry and social organisations for which marine habitat and resources are of interest.
- Wageningen **IMARES** provides strategic and applied ecological investigation related to ecological and economic developments.

Citation

Van Franeker, J.A., Meijboom A., De Jong, M. & Verdaat, H. (2009) Fulmar Litter EcoQO Monitoring in the Netherlands 1979-2007 in relation to EU Directive 2000/59/EC on Port Reception Facilities. *IMARES Report nr C032/09. Wageningen IMARES, Texel. 39pp*

Contact details:

Dr. J.A. (Jan Andries) van Franeker
Wageningen IMARES
PO Box 167, 1790 AD Den Burg (Texel), The Netherlands
tel +31 (0) 317 487 085
@: jan.vanfraneker@wur.nl

© 2009 Wageningen **IMARES**

Wageningen IMARES is registered in the Dutch trade record
Amsterdam nr. 34135929,
BTW nr. NL 811383696B04.

The Management of IMARES is not responsible for resulting damage, as well as for damage resulting from the application of results or research obtained by IMARES, its clients or any claims related to the application of information found within its research. This report has been made on the request of the client and is wholly the client's property. This report may not be reproduced and/or published partially or in its entirety without the express written consent of the client.

A_4_3_2-V5.1 *adapted*

Contents

- Summary 5
- Samenvatting 9
- 1 Introduction 13
- 2 The Fulmar as an ecological monitor of marine litter 15
- 3 Shipping, marine litter and policy measures 19
- 4 Material and methods 21
- 5 Results 25
 - 5.1 Monitoring in the Netherlands 1979-2007 and trends 25
 - 5.2 Results North Sea 34
- 6 Conclusions 35
- 7 Acknowledgements 36
- 8 References 37
- 9 Quality Assurance 39
- Justification 39




Photo 1: Northern Fulmar (*Fulmarus glacialis*) in flight



Photo 2: Example of an approximately average content of plastic in stomachs of Fulmars beached in the Netherlands (31 pieces of plastic, 0.2273 g is slightly below the current Dutch average).

Left - top: industrial pellets; **centre:** sheetlike plastics; **bottom:** threadlike plastics
Right - top: clay pellet (non plastic rubbish); **centre:** foamed plastic; **bottom:** fragments
(scale reference: industrial pellets have about 4 mm diameter)

Fulmar Litter EcoQO Monitoring in the Netherlands 1979-2007 *in relation to EU Directive 2000/59/EC on Port Reception Facilities*

Summary

Operational and cargo related wastes from ships are an important source of litter in the marine environment in the southern North Sea and cause serious economical and ecological damage. Inadequacies in the ship to shore waste delivery procedures are considered a major factor in illegal discharges. The European Union therefore addressed the problem with the Directive on Port Reception Facilities (Directive 2000/59/EC). Obligatory waste delivery to shore and indirect financing of the costs are key-elements of the Directive to stimulate and enforce proper disposal of shipwaste in harbours. Monitoring the effect of the EU Directive is required.

A marine litter monitoring program using plastic abundance in stomachs of a seabird, the Northern Fulmar, was already operational in the the Netherlands and was further developed also for international implementation by OSPAR as one of the 'Ecological Quality Objectives (EcoQOs)' for the North Sea (OSPAR 2008). Fulmars are purely oceanic foragers, ingest all sorts of litter from the sea surface, and do not regurgitate poorly degrading diet components, but slowly wear these down in the stomach. Accumulated hard plastic items in stomachs of beached Fulmars thus integrate marine litter levels encountered over a number of weeks in a particular area.

In the Netherlands, the *Ministry of VenW* commissions regular updates of Dutch data in the Fulmar-Litter monitoring database maintained by IMARES. This report summarizes the Dutch monitoring results up to 2007. North Sea wide monitoring was started in the 'Save the North Sea' project supported by the EU Interreg IIIB program, and is currently funded by Corporate and Social Responsibility awards from the *NYK Group Europe Ltd*. Jointly, these national and international efforts will merge in a complete update of the OSPAR North Sea Fulmar-Litter-EcoQO up to the year 2007, to be published in a scientific journal.

Monitoring in the Netherlands 1979-2007

Volunteers of the Dutch Seabird Group (NZG) and a range of other participating groups collect Fulmars found dead on Dutch beaches. In 2007, 67 beached Fulmars were collected and 61 of the corpses contained suitable intact stomachs. Samples of 40 or more birds are considered to result in reliable annual 'average' data.

Results 2007: Among the 61 stomachs, 56 contained plastic (92% incidence), with an overall average number of 36 items per bird and average mass of 0.37 gram plastic per bird. Industrial plastics represent a minority (3 'pellets' per bird, 0.07g) compared to discarded user plastics (33 items and 0.30g) (Table 1). In 2006 relatively high averages for that year were discussed as a potential error from the small sample size in that year; however, the properly sized sample of 2007 shows that the earlier results were not an artefact or incident.

Current levels 2003-2007: As a standard, because of variability and occasional years of lower sample size, it has been agreed that the 'current pollution level' is best described as the average situation over the most recent 5 years. Over the 2003-2007 period in a sample of 309 Fulmars, plastic incidence was 93% with an average number of over 26 pieces, and average mass of 0.28 gram plastic (further details in Table 2). Thus, in terms of number and mass of plastics, the year 2007 appears to be above 'average' for the situation over the past 5 years.




Figure i Trend in plastic mass in stomachs of Fulmars beached in the Netherlands 1980-2007 – represented by running arithmetic average over 5 year periods, i.e. bars shift one year ahead at a time; period and number of birds shown below each bar.




Figure ii Trend in plastic mass in stomachs of Fulmars beached in the Netherlands 1980-2007– Annual changes as suggested by 'geometric means' calculated from ln-transformed data (transformation reduces the impact of extremes). Sample sizes before 1997 were too small to calculate annual geometric means.

Trends: As convened in earlier reports, the metric for discussion of trends is based on the mass of plastics in stomachs, rather than on incidence or number of plastic particles. In trend discussions, a distinction is made between:

- 'long-term trend' which calculates the trend over all years in the dataset (now 1979-2007)
- 'recent trend' defined as trend over the past 10 years (now: 1998-2007)

Average figures for plastic mass may be strongly influenced by a few highly polluted stomachs, especially when sample sizes are small. Therefore, for a visual representation of trends over time, *Fig.i* uses a single overall average for the 1980's and as of 1995 averages over 5 year periods. As an alternative, *Fig. ii* does use annual data as off 1997, but shows 'geometric mean mass'. Geometric means are calculated from logarithmically transformed data, which reduces the importance of incidental higher values in the dataset.

Statistical calculations to test for significance of trends use neither averages nor annual geometric means. Test results are based on simple linear regressions of ln-transformed data for the mass of plastic in all individual stomachs against year of collection of the birds.

Long term trend 1979-2007

In spite of evident strong changes in Fig's *i* and *ii*, the 'all plastics' category (Table 3A) shows no clear long term linear trend. This has several reasons. Firstly, the overall mass of plastics strongly increased from the 1980s to the 1990s but has subsequently decreased to approximately the initial level. Linear analyses do not 'see' the variable components in non-linear trends. But trend calculations are also compromised because different types of plastic have shown strongly different trends. User plastics were responsible for the above described increase and later decrease. Industrial plastics on the other hand have strongly decreased since the early 1980s, resulting in a highly significant long-term reduction ($p < 0.001$). As a consequence of these mixed trends, the composition of plastic litter has strongly changed since the early 1980s, with nowadays a strongly reduced proportion of industrial plastics (reduced from about 50% to 20% of total mass) and an increased mass of user plastics from discarded waste.

Recent 10 year trend 1998-2007

Contrary to results in the earlier reports, the regression analyses for 10 year recent trends (Table 3B) no longer reveals a significant decrease in total plastic since the late 1990s. From *Fig.ii* the loss of significance can be explained by the exclusion of the high 1997 value from the start of the trend period, and replacement by elevated 2007 values at the end. Earlier decreases in plastics, both user (since 1990's) and industrial (from the start in 1980's) may still occur (negative signs of t values in table 3B), but not at a statistically relevant level. Geometric mean data for the most recent 5 years in *Fig.ii* may even suggest a reversal of the trend after 2003.

Ecological Quality Objective (EcoQO) for the North Sea

In OSPAR, the Dutch Fulmar monitoring study initiated the development of an Ecological Quality Objective for marine litter based on the amount of plastic in Fulmar stomachs. However, data are looked at in a slightly differently way, being 'the percentage of birds exceeding a critical value of plastic in the stomach'. This is an alternative approach to exclude strong influence of a few exceptional birds on the dataset. The OSPAR **Ecological Quality Objective** for marine litter currently formulates its target for acceptable ecological quality in the North Sea as:

"There should be less than 10% of Northern Fulmars having 0.1 gram or more plastic in the stomach in samples of 50-100 beachwashed fulmars from each of 5 different regions of the North Sea over a period of at least 5 years".

In 2007, 70% of Dutch Fulmars exceeded the critical value of 0.1g plastic in the stomach (Table 2). In 2006, an exceptionally high figure of 85% of birds exceeding 0.1g of plastic was suspected to be an artefact from small sample size, but the 70% score in the good sample size for 2007 confirms high levels of plastic pollution in recent years. Over the latest five years, an average of 61% of Fulmars beached in the Netherlands exceeds the critical level of 0.1 g of plastic in the stomach. Following the EcoQO target

definition, Fig. *iii* shows the Dutch data in terms of EcoQO performance over 5-year periods. The EcoQO graph supports the suggestion made under 'Short term trends' that after a period of decreasing plastic loads since the 1990's (in the order of 10% of EcoQO performance!), improvements seem to have halted or even reversed. Thus, the Fulmar study until now can not observe an effect from implementation of the EU Directive 2000/59/EC on Port Reception Facilities.




Figure *iii* EcoQO performance in the Netherlands 1980-2007 – Trend in percentage of beached Fulmars in the Netherlands having more than 0.1 gram plastic in the stomach (running average over 5 year periods, each bar shifting one year; number of birds in brackets below each bar). Note that for clarity, the Y-axis only shows the 50 to 70% range where the OSPAR target for the EcoQO is that less than 10% of birds has more than 0.1 g plastic in the stomach (see Fig.5).

Conclusions

- In 2007, 92% of Fulmars beached in the Netherlands (61 stomachs) had plastic in the stomach with an average number of 36 pieces and mass of 0.37 gram per bird. The critical level of 0.1 gram of plastic used in the OSPAR EcoQO was exceeded by 70% of the birds (the EcoQO target aims at less than 10%)
- Averaged over the past 5 years (2003-2007; 309 stomachs), 93% of Dutch Fulmars had plastic in the stomach with 27 pieces and 0.28g on average per bird (*cf photo 2 on page 4*). The critical EcoQO level of 0.1 g was exceeded by 61% of the birds. It is recommended to rely on the 5-year averages.
- Thus, the 2007 abundance of plastic exceeds the recent 5-year average, suggesting no recent improvements. Earlier improvements in EcoQO performance by about 10% seem to have stopped or may even have reversed (Fig *iii*).
- Statistical tests for significance of changes over the past 10 years show no significant trends for industrial plastics, user plastics or their combined total.
- These findings indicate that the implementation of the EU Directive on Port Reception Facilities since late 2004 has not yet resulted in detectable ecological improvement in the southern North Sea.

Stormvogel Zwerfvuil EcoQO monitoring in Nederland 1979-2007 in relatie tot Richtlijn 2000/59/EG betreffende havenontvangstvoorzieningen voor scheepsafval en ladingresiduen

Samenvatting

Het door schepen overboord zetten van operationeel en aan lading gerelateerd afval is één van de belangrijke bronnen van zwerfvuil in de zuidelijke Noordzee. Zulk afval heeft ernstige economische en ecologische gevolgen. Tekortkomingen in afgifteprocedures in havens worden gezien als een belangrijk achtergrond voor het illegaal overboord zetten van scheepsafval. De EU heeft dit probleem aangepakt met de 'Richtlijn betreffende havenontvangstvoorzieningen' (Richtlijn 2000/59/EC; de zogenaamde 'HOI-Richtlijn'). Verplichte afgifte van afval en indirecte financiering van de kosten vormen de kern van de maatregelen waarmee de Richtlijn correcte afvalafgifte wil stimuleren en afdwingen. Het monitoren van de effecten van de HOI-Richtlijn is noodzakelijk.

In Nederland worden trends in zwerfafval op zee reeds onderzocht in een monitoring programma dat is gebaseerd op de hoeveelheid plastic afval in magen van dood angespoelde zeevogels: de Noordse Stormvogel. Dit graadmeter instrument wordt ook internationaal verder ontwikkeld als een 'Ecological Quality Objective (EcoQO)' door OSPAR (OSPAR 2008). De Noordse Stormvogel fourageert uitsluitend op zee, eet geregeld afval, en hoopt slecht verteerbaar materiaal zoals plastic op in de maag. Daardoor geeft de maaginhoud een geïntegreerd beeld van de hoeveelheden afval die de vogels in voorgaande weken op zee zijn tegengekomen.

Het Ministerie van VenW heeft Wageningen IMARES opdracht gegeven de Nederlandse stormvogel graadmeter aan te vullen met gegevens van het jaar 2007. Het voorliggende rapport is daarvan het eindresultaat. Vergelijkbare internationale monitoring van stormvogels in de hele Noordzee is in 2002 opgestart als onderdeel van het Europese 'Save the North Sea' project, en daarna voortgezet met bedrijfsmaatschappelijke prijzen toegekend door de NYK Group Europe Ltd. Het resultaat van al gezamenlijke nationale en internationale inspanning zal later dit jaar worden geïntegreerd in een wetenschappelijke tijdschriftpublicatie over de OSPAR Stormvogel-Zwerfvuil-EcoQO in de Noordzee tot en met 2007.

Nederlands graadmeter onderzoek 1979-2007

Vrijwilligers van de Nederlandse Zeevogelgroep (NZG) en een reeks andere organisaties verzamelen Noordse Stormvogels die dood op de stranden aanspoelen. In 2007 konden aldus 67 stormvogels worden verzameld, waarvan er 61 nog een voor onderzoek geschikte intacte maag bevatten. Voor een betrouwbaar jaargemiddelde voor een bepaalde locatie wordt een bemonsterd aantal van 40 of meer vogels aanbevolen.

Resultaten 2007: Van de 61 magen bevatten er 56 (92%) plastic met een gemiddeld aantal van 36 stukjes en gemiddeld gewicht van 0.37 gram per vogel (Tabel 1). In 2006 kon een relatief hoog gehalte aan plastic mogelijk een vertekend beeld zijn ten gevolge van beperkt aantal vogels in het onderzoek, maar het grotere aantal vogels uit 2007 bevestigt het beeld.

Huidige situatie 2003-2007: omdat toevallige jaarfluctuaties of te kleine monstergrootte het beeld kunnen beïnvloeden, wordt geadviseerd om "de huidige situatie" te beschrijven als het gemiddelde over de vijf meest recente jaren. Gemeten over 2003-2007 periode had 93% van de 309 onderzochte stormvogels plastic in de maag. Het gemiddelde aantal stukjes was 26 per vogel en het gemiddeld gewicht 0.28 gram per vogel (Tabel 2). Binnen de recente 5-jaarsperiode was het jaar 2007 dus qua aantallen en gewicht van plastic een bovengemiddeld jaar.

Trends: Trendmatige ontwikkelingen worden geanalyseerd op basis van het gewicht van het plastic in de magen, waarbij

- 'lange termijn trends' verwijzen naar de complete periode 1979-2007
- 'recente trends' zijn gedefinieerd als de trends over de voorgaande 10 jaar (1998-2007)

Gemiddelde waarden voor het gewicht aan plastic kunnen sterk worden beïnvloed door een klein aantal sterk vervuilde magen, vooral bij een beperkte monsternamen. Derhalve toont Fig. i een totaal gemiddelde voor de jaren tachtig met zogenaamd 'lopende gemiddeldes' over latere 5 jaars periodes waarvoor de gebruikte gegevens telkens één jaar opschuiven. Als alternatief worden in Fig. ii. toch afzonderlijke jaargegevens getoond, maar dan 'geometrische gemiddeldes' die zijn berekend aan de hand van logaritmisch omgevormde getallen.

Logaritmische omvorming onderdrukt de invloed van uitschieterende extreem hoge waardes. Statistische berekeningen om de 'significantie' van bepaalde trends te toetsen zijn niet op gewone, lopende of geometrische gemiddeldes gebaseerd. De trends worden getoetst door lineaire regressie analyses van ln-getransformeerde plastic gewichten in individuele vogels tegen het jaar waarin de vogels werden verzameld.

Figuur i (blz. 6)

Trend in plastic gewicht in magen van in Nederland aangespoelde Noordse Stormvogels – getoond aan de hand van lopende 5-jaars gemiddeldes waarbij grafiek staven telkens één jaar verschuiven; de jaren en het aantal vogels in de berekening staan bij iedere staaf vermeld. Voor de jaren '80 is een enkel gemiddelde berekend.

Figuur ii (blz. 6)

Trend in plastic gewicht in magen van in Nederland aangespoelde Noordse Stormvogels – getoond aan de hand van geometrische gemiddeldes per jaar berekend uit logaritmisch getransformeerde data die de invloed van extreem hoge waardes onderdrukken. In de jaren voor 1997 was de het aantal monsters ontoereikend voor jaarlijkse gemiddeldes. Voor de jaren '80 is een enkel gemiddelde berekend.

Lange termijn trend 1979-2007

Ondanks de in de figuren *i* en *ii* zichtbare veranderingen, is voor alle plastic soorten tezamen ('all plastics' in Tabel 3A) over de volledige studieperiode geen significante trend waarneembaar. De eerste reden daarvoor is dat plasticgewicht tussen jaren '80 en '90 sterk toenam, maar daarna weer is teruggezak naar ongeveer het beginniveau. Lineaire regressies "missen" per definitie niet-rechthoekige verbanden. Maar daarnaast worden lange-termijn trends enigszins verdoezeld omdat verschillende vormen van plastic een afwijkende ontwikkeling hebben doorgemaakt. Gebruiks plastics zijn verantwoordelijk voor de hierboven omschreven toename en afname. Maar industrieel plastic heeft over de volle periode een significante afname vertoond ($p < 0.001$). Ten gevolge van deze afwijkende trends is de samenstelling van het plastic in de stormvogelmagen sterk veranderd van een aanvankelijk gelijk gewicht aan industrieel en gebruiksplastic naar een situatie waarin ca 80% van het plastic tot het van afval afkomstige gebruikers type behoort.

Recente 10 jaar trend 1998-2007

In tegenstelling tot eerdere rapportages, laat Tabel 3B geen significante trends meer zien over de laatste 10 jaar. De verklaring is te zien in Fig. ii, die toont dat zeer hoge waardes uit 1997 nu voor het eerst buiten de berekening vallen, en dat een patroon van afname over de jaren 1998-2003 wordt 'tegengewerkt' door een geleidelijke toename over de jaren 2003-2007, hetgeen tezamen resulteert in 'géén trend' voor de afgelopen tien jaar. Weliswaar is er nog steeds afname (negatieve t-waardes in tabel 3B) maar deze is niet langer significant. Het beeld voor industrieel en gebruiksplastics is over deze periode vergelijkbaar.

Ecologische Kwaliteitsdoelstelling voor de Noordzee

In de Ecological Quality Objective (EcoQO) voor zwerfvuil op zee bekijkt OSPAR dezelfde gegevens op een iets andere manier, namelijk als een percentage van de vogels waarbij de maaginhoud een grensgewicht aan plastic overschrijdt. OSPAR definieert de 'doelwaarde voor aanvaardbare ecologische kwaliteit' in de Noordzee als de situatie waarin:

"minder dan 10% van aangespoelde Noordse Stormvogels 0.1 gram of meer plastic in de maag heeft, in monsternames van 50 tot 100 vogels uit 5 verschillende Noordzee regio's over een periode van tenminste 5 jaar"

In 2007 overschreed 70% van de onderzochte stormvogels de kritische grens van 0.1g plastic in de maag. Dit is weliswaar een verbetering t.o.v. de situatie in 2006, maar steekt over een wat langere periode slecht af, passend bij de eerdere conclusies. Het 5-jaar gemiddelde voor de periode 2003-2007 is dat 61% van de Stormvogels van de Nederlandse kust uitkomt boven de grenswaarde van 0.1 gram plastic in de maag; onveranderd t.o.v. de periode 2002-2006..

De trend in EcoQO ontwikkelingen in Nederland wordt getoond in Fig.iii als lopend 5-jaars-gemiddelde, dwz de ook in de EcoQO doelstelling verwoorde periode. De figuur ondersteunt de eerder opmerking (Recente trend) dat na

een periode van afname in vervuiling (in de orde van 10% in de EcoQO!) in de allerlaatste jaren de afname is gestopt of zelfs is omgeslagen in een geleidelijke toename. Dit betekent dat het stormvogelonderzoek tot dusverre nog géén effect kan aantonen van de invoering van de EU Richtlijn voor havenontvangstvoorzieningen 2000/59/EG.

Figuur i (blz. 8)

Ontwikkeling in de Stormvogel-Zwerfvuil Ecologische Kwaliteitsdoelstelling 1980-2007. Trend in het percentage Noordse Stormvogels dat méér dan 0.1 gram plastic in de maag heeft, als lopend gemiddelde op basis van 5-jaar-periodes. Voor de jaren '80 is een enkel gemiddelde berekend. Let op dat voor het helder tonen van veranderingen de y-as is beperkt tot een 50-70% bereik, waardoor de EcoQO-doelwaarde van 10% niet is getoond (zie daarvoor Fig.5).

Conclusies

- In 2007 had 92% van de in Nederland aangespoelde Noordse Stormvogels (61 magen) plastic in de maag, met een gemiddeld aantal van 36 stukjes en gemiddeld gewicht van 0.37 gram per vogel. De in de OSPAR EcoQO gebruikte grenswaarde van 0.1g plastic werd door 70% van de vogels overschreden (EcoQO doelstelling is < 10%).
- Gemiddeld over de afgelopen 5 jaar (2003-2007; 309 magen) had 93% van de Nederlandse stormvogels plastic in de maag en lag het gemiddelde op 27 stukjes en 0.28 gram per vogel. De EcoQO grenswaarde van 0.1 gram werd door 61% van de vogels overschreden. Gebruik van 5-jaar gemiddeldes wordt aanbevolen.
- Bovenstaande houdt in dat de hoeveelheid plastic in 2007 'bovengemiddeld' is voor de laatste vijf jaar en duidt op gebrek aan verbetering in deze korte periode. Een eerdere verbetering in termen van de EcoQO (ca 10% reductie) lijkt te zijn gestopt of zelfs omgekeerd (Fig. iii).
- Statistische toetsen voor veranderingen over de laatste tien jaar laten geen significante veranderingen zien voor industrieel plastic, gebruiksplastic of hun gecombineerde totaal.
- Deze bevindingen houden in dat de implementatie van de EG Richtlijn voor Havenontvangstvoorzieningen vanaf eind 2004 nog géén merkbare verbetering heeft opgeleverd in de ecologische toestand van de zuidelijke Noordzee.

1 Introduction

Marine litter, in particular plastic waste, represents an environmental problem in the North Sea with wide ranging economical and ecological consequences.

Economic consequences of marine litter are suffered by coastal municipalities who find themselves confronted with excessive costs for beach clean-ups. Tourist business suffers damage because guests stay away from polluted beaches, especially when various types of litter are a health-risk for tourists. Fisheries are confronted with a substantial by-catch of marine litter which causes loss of time and sometimes necessitates discarding of tainted catch. All sorts of shipping suffer financial damage and more importantly, safety-risks from fouled propellers or blocked water-intakes. Coastal litter blowing inland is even seriously affecting farmers. The economical damage from marine litter is difficult to estimate, but a detailed study in the Shetlands with additional surveys elsewhere indicates that extrapolated costs for the whole North Sea area may exceed one billion Euro per year (Hall 2000; pers.inf.).

The most pronounced ecological consequence of marine litter is the suffering and death of marine wildlife. Entanglement of seabirds and marine mammals regularly attracts public attention. However, only a small proportion of such mortality becomes visible among beached animals. Even less apparent are the consequences from the ingestion of plastics and other types of litter. Ingestion is extremely common among a wide range of marine organisms including many seabirds, marine mammals and sea-turtles. It does cause direct mortality but the major impact may well occur through reduced fitness of many individuals. Sub-lethal effects on animal populations remain largely invisible. In spite of spectacular examples of mortality from marine litter, the real impact on marine wildlife remains difficult to estimate (Laist 1987, 1997; Derraik 2002). Plastics gradually break down to microscopically small particles, but even these may pose serious problems to marine ecosystems (Thompson *et al.* 2004). Concern about microplastics attracts more and more attention as evidence is growing that plastics in seawater strongly bind organic pollutants and microplastics may enter the base of the food-web by ingestion by filterfeeding marine organisms (Endo *et al.* 2005; Teuten *et al.* 2007; Browne *et al.* 2008; Moore 2008; Arthur *et al.* 2009). Thus, in addition to the toxic substances incorporated into plastics in the manufacturing process, plastics may concentrate much more pollutants from the environment and act as a pathway boosting their accumulation in marine organisms that ingest plastic. Evidently, this same mechanism operates at all levels of organisms and sizes of ingested plastic material, from small zooplankton filterfeeders to large marine birds and mammals, but it is the microplastic issue and their ingestion by small filterfeeders that has emphasized the potential scale and urgency of the problem.

Recognizing the negative impacts from marine litter, a variety of international policy measures has attempted to reduce input of litter. Examples of these are the London Dumping Convention 1972; Bathing Water Directive 1976; MARPOL 73/78 Annex V 1988; Special Area status North Sea MARPOL Annex V 1991; and the OSPAR Convention 1992. In the absence of significant improvements, political measures have been intensified by for example the EU-Directive on Port Reception Facilities (Directive 2000/59/EC) and the Declaration from the North Sea Ministerial Conference in Bergen, March 2002.

Recent policy initiatives have recognized that policy aims need to be quantifiable and measurable. Therefore, the North Sea Ministers in the 2002 Bergen Declaration have decided to introduce a system of Ecological Quality Objectives for the North Sea (EcoQO's). A number of these EcoQO's is implemented in an immediate pilot program. For example, the oil pollution situation in the North Sea will be measured by the rate of oil-fouling among Guillemots (*Uria aalge*) found on beaches (Camphuysen 2005). The ecological quality target is set at a level in which less than 10% of beached Guillemots have oil on the plumage.

Another set of EcoQO's has to be developed for future implementation. Among this latter group is an EcoQO for marine litter, to be measured by the abundance of plastic in stomachs of seabirds, *in casu* the Northern Fulmar (*Fulmarus glacialis*). Working Groups in ICES and in OSPAR are involved in the further development and implementation of the EcoQO system including the advice on realistic target levels (OSPAR 2005). For convenience this EcoQO for marine litter is referred to as the 'Fulmar-Litter-EcoQO' and is approaching final formal approval in the OSPAR system, as expressed in the publication of the 'Background Document' (OSPAR

2008). The EcoQO approach has also been included as an element in the approach for the intended European Marine Strategy (EC 2005).

Within the Netherlands, the Ministry of Transport, Public Works and Water Management (VenW) has a coordinating role in governmental issues related to the North Sea environment. As such, VenW is involved in the development of environmental monitoring systems ("graadmeters") for the Dutch continental shelf area. As a part of this activity, VenW have commissioned several earlier projects by IMARES working towards a Fulmar-Litter-EcoQO. The first pilot project considered stomach contents data of Dutch Fulmars up to the year 2000 and made a detailed evaluation of their suitability for monitoring purposes (Van Franeker & Meijboom 2002). A series of later reports (see 'References') have provided annual updates on the Dutch time-series up to the year 2006 (van Franeker et al 2008), paying special attention to shipping issues and EU Directive 2000/59/EC.

As of 2002, the Dutch Fulmar research was also expanded to all countries around the North Sea as a project under the **Save the North Sea (SNS)** program. SNS was co-funded by EU Interreg IIIB over period 2002-2004 and aimed to reduce littering in the North Sea area by increasing stakeholder awareness. The Fulmar acted as the symbol of the SNS campaign. SNS project results and issues related to the development of the Fulmar-Litter-EcoQO were published in Van Franeker *et al.* 2005 (Alterra-rapport 1162). Findings strongly supported the important role of shipping (incl. fisheries) in the marine litter issue. For further publications of the SNS Fulmar study see e.g. Save the North Sea 2004, Van Franeker 2004b and 2004c, Edwards 2005, Guse *et al* 2005, Olsen 2005, van Franeker *et al* 2008).

Building upon this earlier work, the current assignment from the Dutch Ministry of Transport included the following tasks:

- To update the time series on litter in stomach contents of Dutch Fulmars with the data from 2007, and publish the result in a new report;
- To continue co-ordination of the beached bird sampling in the Netherlands into 2009

The fulmar study additionally received a 'Corporate and Social Responsibility Award from the NYK Group Europe Ltd. This funding made it possible:

- To analyse samples from other North Sea locations collected in the year 2007.
- To continue international co-ordination of EcoQO sampling in the North Sea area into 2009, including the organisation of an international workshop. Collected samples from 'foreign' Fulmars to be stored frozen, awaiting future sources of funding).
- To promote expansion of the litter monitoring network to new groups and locations

This report provides the details of the addition of data from the Netherlands over the year 2007. Integrated results from the combined national and international projects will be used to prepare an article in a scientifically reviewed journal to ensure public and scientific 'quality control' on the Fulmar Litter EcoQO research.

2 The Fulmar as an ecological monitor of marine litter

The interpretation of monitoring information presented in this report requires a summary of earlier findings.

Van Franeker & Meijboom (2002) discussed the feasibility of using stomach contents of beached Northern Fulmars to measure changes in the litter situation off the Dutch coast in an ecological context. Samples of Fulmars available for the feasibility study mainly originated from the periods 1982 to 1987 and 1996 to 2000, with smaller number of birds from the years in between.

Reasons for selection of the Fulmar out of a list of potential monitoring species, are of a practical nature:

- Fulmars are abundant in the North Sea area (and elsewhere) and are regularly found in beached bird surveys, which guarantees supply of adequate samples for research.
- Fulmars are known to consume a wide variety of marine litter items.
- Fulmars avoid inshore areas and forage exclusively at sea (never on land).
- Fulmars do not normally regurgitate indigestible items, but accumulate these in the stomach (digestive processes and mechanical grinding gradually wear down particles to sizes that are passed on to the gut and are excreted).
- Thus, stomach contents of Fulmars are representative for the wider offshore environment, averaging pollution levels over a foraging space and time span that avoids bias from local pollution incidents.
- Historical data are available in the form of a Dutch data series since 1982 (one earlier 1979 specimen); and literature is available on other locations and related species worldwide (Van Franeker 1985; Van Franeker & Bell 1988).
- Other North Sea species that ingest litter either do not accumulate plastics (they regurgitate indigestible remains); are coastal only and/or find part of their food on land (e.g. Larus gulls); ingest litter only incidentally (e.g. North Sea alcids) or are too infrequent in beached bird surveys for the required sample size or spatial coverage (e.g. other tubenoses or Kittiwake *Rissa tridactyla*).

Beached birds may have died for a variety of reasons. For some birds, plastic accumulation in the stomach is the direct cause of death, but more often the effects of litter ingestion act at sub-lethal levels, except maybe in cases of ingestion of chemical substances. For other birds, fouling of the plumage with oil or other pollutants, collisions with ships or other structures, drowning in nets, extremely poor weather or food-shortage may have been direct or indirect causes of mortality.

At dissection of birds, their sex, age, origin, condition, likely cause of death and a range of other potentially relevant parameters are determined. Standardized dissection procedures for EcoQO monitoring have been described in detail in a manual (Van Franeker 2004b). Stomach contents are sorted into main categories of plastics (industrial and user-plastics), non-plastic rubbish, pollutants, natural food remains and natural non food-remains. Each of these categories has a number of subcategories of specific items. For each individual bird and litter category, data are recorded on presence or absence ("incidence"), the number of items, and the mass of items (see methods). For efficiency/economy reasons, some of the details in manual and earlier reports have been reduced.

The pilot study undertook extensive analyses to check whether time-related changes in litter abundance were susceptible to error caused by bias from variables such as sex, age, origin, condition, cause of death, or season of death. If any of these would substantially affect quantities of ingested litter, changes in sample composition over the years could hamper or bias the detection of time-related trends.

A very important finding of the pilot study was that no statistical difference was found in litter in the stomach between birds that had slowly starved to death and 'healthy' birds that had died instantly (e.g. because of collision or drowning). This means that our results, which are largely based on beached starved birds, are representative for the 'average' healthy Fulmar living in the southern North Sea.

Only age was found to have an effect on average quantities of ingested litter, adults having less plastic in their stomachs than younger birds. Possibly, adults lose some of the plastics accumulated in their stomach when

they feed chicks or spit stomach-oil during defence of nest-sites. Another factor could be that foraging experience may increase with age. Understanding of the observed age difference in plastic accumulation is still fairly poor, and further study is required. With financial support from Chevron Upstream Europe, we have started a cooperative project with the Faroese Fisheries Laboratory using Fulmars from the Faroe Islands, where birds are hunted for consumption and large numbers of samples are easily obtained. Additional samples have been obtained from fisheries by-catch in the area. Stomach contents are analysed for both normal diet (in a Faroese study) and for accumulated litter (in a Dutch study). Samples have been obtained from all months of the year over the period 2003-2006. Detailed analyses are still to be conducted, but overall averaged data have already been used in this new report.

Although age has been shown to affect absolute quantities of litter in stomach contents, changes over time follow the same pattern in adults or non-adults. As long as no directional change in age composition of samples is observed, trends may be analysed for the combined age groups. However, background information for the presentation of results and their interpretations always requires insight in age composition of samples.

Significant long term trends from 1982 to 2000 were detected in incidence, number of items and mass of industrial plastics, user plastics and suspected chemical pollutants (often paraffine-like substances). Over the 1982-2000 period, only industrial plastics decreased while user plastics significantly increased. When comparing averages in the 1980s to those in the 1990s, industrial plastics decreased from 6.8 granules per bird (77% incidence; 0.15g per bird) to 3.6 granules (64%; 0.08g). User-plastics increased from 7.8 items per bird (84%; 0.19g) to 27.6 items (97%; 0.52g). An analysis for shorter term recent trends over the period of 1996 to 2000 revealed continued significant decrease in industrial plastics and suggested stabilization or slight decreases in other litter categories.

Analysis of variability in data and Power Analysis revealed that reliable figures for litter in stomachs in a particular region are obtained at a sample size of about 40 birds per year and that reliable conclusions on change or stability in ingested litter quantities can be made after periods of 4 to 8 years, depending on the category of litter.

Mass of litter, rather than incidence or number of items, should be considered the most useful unit of measurement in the long term. Mass is also the most representative unit in terms of ecological impact on organisms. Incidence loses its sensitivity as an indicator when virtually all birds are positive (as is the case in Fulmars). In regional or time-related analyses, mass of plastics is a more consistent measure than number of items, because the latter appears to vary with changes in plastic characteristics.

The pilot study concluded that stomach content analysis of beached Fulmars offers a reliable monitoring tool for (changes in) the abundance of marine litter off the Dutch coast. By its focus on small-sized litter in the offshore environment such monitoring has little overlap with, and high additional value to beach litter surveys of larger waste items. Furthermore, stomach contents of Fulmars reflect the ecological consequences of litter ingestion on a wide range of marine organisms and create public awareness of the fact that environmental problems from marine litter persist even when larger items are broken down to sizes below the range of normal human perception. As indicated there is an increasing awareness of the dangers from microplastics, but monitoring quantities and effects in these species is more difficult than that of intermediate sized plastics in seabirds.

The pilot study concluded that the formal indicators which would be recommended in future Dutch Fulmar-Litter monitoring were abundances by mass of industrial plastic, user plastic, their combined total and suspected chemicals. Each of these represents different sources of pollution, and thus specific policy measures aimed at reduced inputs. Because no specific funding was raised the suspected chemicals plastics have become the main focus. Addition of further formal indicators from other litter (sub-)categories would produce little added value in the current situation. However, data-recording procedures are such that at the raw data-level, these categories continue to be recorded and can be extracted from databases, should the need arrive.

After publication of the pilot study, the Dutch monitoring has continued annually and has resulted in a series of reports (Van Franeker et al 2003 to 2008). These studies confirmed a decrease of industrial plastics and also gradually provided evidence for significant decreases in user plastics after the late 1990s, a trend however that more recently has become doubtful.

In 2002, the Fulmar Litter monitoring was boosted by participation in the 'Save the North Sea' project. The Save the North Sea ('SNS') campaign, co-funded by EU Interreg III B, aimed at increasing awareness among stakeholders so as to reduce littering behaviour. Expanding the Dutch Fulmar study to locations all around the North Sea was one of the project components. Co-operation was established with interested groups in all countries around the North Sea. In 2005 the final project report (Van Franeker et al 2005) showed that Fulmars from the southern North Sea had almost two times more plastic in the stomach than Fulmars from the Scottish Islands, and almost four times as much as that in a small sample from the Faroe Islands. Location differences and relative abundances of different types of litter suggested a major role of shipping, and show that the bulk of the litter problem in the North Sea region is of local origin.

In 2002, North Sea Ministers in the Bergen Declaration, decided to start a system of Ecological Quality Objectives for the North Sea. One of the EcoQO's to be developed was for the issue of marine litter pollution, using stomach contents of a seabird, the Fulmar, to monitor developments, and set a target for 'acceptable ecological quality'. OSPAR was requested to look after implementation of the ecological quality objectives. Since then, a number of steps have been taken, based on reports from the Dutch studies and the Save the North Sea project. The preliminary wording of the EcoQO on target level for ecological quality is that:

"There should be less than 10% of Northern Fulmars having 0.1 gram or more plastic in the stomach in samples of 50-100 beachwashed fulmars from each of 5 different regions of the North Sea over a period of at least 5 years".

So, the basis of the EcoQO monitoring system is mass of plastics as recommended from the Dutch studies. But rather than using average plastic mass for the target definition, a combination is used of frequency of occurrence of plastic masses above a certain critical level. The background of this is that a few exceptional outliers can have a strong influence on the calculated average. The wording of the target level basically excludes influence of exceptional outlying values. A background document on the EcoQO has now been formally accepted and has been published on the OSPAR website (OSPAR 2008). The EcoQO approach is also considered in discussions on the European Marine Strategy Directive.

Anticipating further development of the EcoQO approach in OSPAR and EU, the international Save the North Sea Fulmar study group was kept active after completion of the 2002-2004 EU-project. Funding from OSPAR partners has not yet been achieved. Fortunately, in recent years CSR awards (Corporate Social Responsibility) from NYK Europe Ltd made it possible to analyse stomachs from outside the Netherlands collected after 2004, to continue international coordination and outreach and the publication of integrated results. Dutch government funding, plus the support from NYK Europe, has ensured a North Sea EcoQO update covering data from the period 2002-2006 (van Franeker 2008). The new international data up to 2007 will be used in a scientific publication on the EcoQO in a refereed journal.

3 Shipping, marine litter and policy measures

In historic times, all waste products from ships were discarded almost anywhere and at any time. The relatively low intensity of shipping and generally decomposable nature of wastes allowed such practice to continue for centuries without significant problems except inside harbour areas. However, exponential population growth and global industrialization has boosted marine transports by fast mechanically-powered ships with ever increasing quantities of poorly decomposable and toxic wastes from fuel, cargo and household practises. Old habits are hard to change, particularly if such change involves costs in an extremely competitive international industry such as shipping. For example, the dramatic environmental consequences of oil discharges from ships were already known in the early 1900s. More than a century later, under continuous public pressure and a continuous sequence of policy measures, the oil pollution problem is to some extent under control, but definitely not solved. Compared to the problems from dumping of oil or toxic wastes, the issue of disposal of 'garbage' into the marine environment has long been considered of minor importance. It might still be considered that way if not for plastics. Plastics, although known since the early 1900s, started their real development only after 1960. Since then, they have found their way into almost every application, replacing old materials in existing products, and creating new use in for example an endless array of 'disposable' packaging products.

Unfortunately, the same factors that made plastics such a popular product have resulted in them becoming an environmental problem. Low production costs have promoted careless use and low degradability leads to accumulation in the environment. In 2007, the world production of synthetic polymers amounted to about 315 million metric tons, over 40% of which is used for packaging (PlasticsEurope2008). Annual growth rates are between 5 to 10%!

At the same time, intensity of shipping has increased. Between 1994 and 2008, the world's active merchant fleet grew from 437 to 742 million gross tons. Fleets grow faster and faster. Over the 1994-2003 decade, tonnage showed 30% growth, matched by a similar growth over the 5 year period 2003-2008. The tonnage of new merchant ships (>100 gtons) leaving shipyards was 18 million gross tons in 1994 and reached an all time peak of 57 million gross tons over 2007. (Department for Transport 2008).

Marine litter originates from a variety of sources, including merchant shipping, fisheries, offshore industry, recreational boating, coastal tourism, influx from rivers or direct dumping of wastes along seashores. The relative importance of various sources differs strongly in different parts of the world, and is almost impossible to quantify. Dutch Coastwatch studies (e.g. Stichting de Noordzee 2003) score litter into categories 'from sea' (shipping, fisheries, offshore); 'beach-tourism'; 'dumped from land'; and 'unknown'. In the Netherlands, the 'from sea' category consistently represents in the order of 40% of litter items recorded. The 'unknown' category scores a similar percentage. Considerable uncertainties are linked to this categorization. More specific information may come from the OSPAR initiative for monitoring litter on beaches in a somewhat more systematic approach. In a first German report (Fleet 2003), ten years of Coastwatch-like surveys, plus two years of the more detailed OSPAR pilot project, were evaluated. From both studies it is concluded that shipping, fisheries and offshore installations are the main sources of litter found on German North Sea beaches. The larger proportion of litter certainly originates from shipping, with a considerable proportion of this originating in the fisheries industry. In the Netherlands, data to this effect were collected in a large beach litter study on Texel (van Franeker 2005) suggesting that up to 90% of plastic litter originates from shipping and fisheries in the Dutch area.

So, although there may be uncertainties in details, there is little doubt that waste disposal by ships is one of the important sources of marine litter worldwide, a fact also recognized by the International Maritime Organization (IMO) in a specific 'garbage-annex' to the MARPOL Convention.

The International Convention for the Prevention of Marine Pollution from Ships (MARPOL 73/78) entered into force on 2nd October 1983 for Annexes I (oily wastes) and II (bulk liquid chemicals), but its Annex V, covering garbage, only achieved sufficient ratifications to enter into force on 31st December 1988. MARPOL Annex V contains the following main prohibitions for discharge of solid wastes:

- No discharge of plastics.
- No discharge of buoyant dunnage, lining or packaging material within 25 nautical miles (nm).
- No discharge of garbage within 12 nm. Food waste may be discharged if ground to pieces smaller than one inch.
- No discharge of any solid waste, including food waste, within 3 nm.

Unfortunately, control of compliance with Annex V regulations on ships is difficult. During Port State Inspections, garbage-related issues will definitely not receive the strongest attention. Nevertheless, in the year 2002, 13% of deficiencies recorded related to Annex V garbage regulations (OECD-MTC 2003).

In the European region, and especially the North Sea area, the sheer intensity of merchant shipping and fisheries makes them an undisputed source of marine litter. From that background, North Sea states promoted that the North Sea received the status of MARPOL Special Area for its annexes I (oil) and V (garbage). Amendments to that effect were made in 1989, and the Special Area status for the North Sea entered into force in February 1991. "Special Areas" under MARPOL Annex V have a more restrictive set of regulations for the discharge of garbage, with the main additions being:

- No discharge, not only of plastics, but also of any sort of metal, rags, packing material, paper or glass.
- Discharge of food wastes must occur as far as practicable from land, and never closer than 12 nm.

Within the European Union, progress under worldwide MARPOL regulations was considered insufficient. In the port of Rotterdam, approximately 5 to 10% of visiting ships used port reception facilities. Clearly not every ship needs to discharge wastes at every port visit, but the level of waste delivery was clearly too low. High costs of proper disposal in combination with low risk of being fined for violations are a clear cause. Poor functioning of available reception facilities definitely plays a role as well. Compliance with MARPOL regulations is hard to enforce at sea, especially when many ships fall under jurisdiction of cheap flag-states with little concern for environmental issues. Compliance can only be promoted by measures that can be enforced when ships visit the harbour. From this perspective, the European Commission and parliament have installed the EU-Directive on Port Reception Facilities for ship-generated waste and cargo residues (Directive 2000/59/EC). Key elements of the Directive are:

- Obligatory disposal of all ship-generated waste to reception facilities before leaving port. Ship-generated waste includes operational oily residues, sewage, household and cargo-associated waste, but not residues from holds or tanks.
- Indirect financing, to a 'significant' degree, of the delivery of ship-generated waste. Finances for such 'free' waste reception should be derived from a fee system on all ships visiting the port. Delivery of cargo residues remains to be paid fully by the ship
- Ports need to develop and implement a 'harbour waste plan' that guarantees appropriate reception and handling of wastes

The term 'Significant' was later identified as meaning 'in the order of at least 30%'. Implementation date for the Directive was December 2002, but unfortunately suffered some delay in several countries. In the Netherlands, the Directive became implemented in late 2004, operating at or above the minimum level of indirect financing depending on the harbour. On an annual basis, results are evaluated by the Minister of Transport, Public Works and Water Management.

The Netherlands Ministry of Transport, Public Works and Water Management wants to measure whether implementation of the EU Directive for Port Reception Facilities has the intended effect. As far as litter is concerned, the Fulmar-Litter-EcoQO approach can be used. This tool complements surveys of quantities of litter delivered in ports, or beach surveys for quantities of waste washing onto beaches. These approaches have their specific merits but do not measure residual levels of litter in the marine environment itself. The Fulmar-Litter-EcoQO does look at this marine environment and at the same time places such information in the context of ecological effects.

4 Material and methods

In 2007 Wageningen IMARES has continued the collection of beached Fulmars from Dutch beaches with the assistance of the Dutch Seabird Group (Nederlandse Zeevogelgroep NZG) through its Working Group on Beached Bird Surveys (Nederlands Stookolieslactoffer Onderzoek - NSO). Also several coastal bird rehabilitation centres support the collection program. Since the start of the **Save the North Sea** project in 2002, IMARES co-ordinates similar sampling projects at a range of locations in all countries around the North Sea. Organisations involved differ widely, and range from volunteer bird groups to governmental beach cleaning projects.

Bird corpses are stored frozen until analysis. Standardized dissection methods for Fulmar corpses have been published in a dedicated manual (Van Franeker 2004b). Stomach content analyses were described in full detail in Van Franeker & Meijboom (2002) as were the methods for data analysis and presentation of results. For convenience, some of the methodological information from earlier reports is repeated here in a condensed form.

At dissections, a full series of data is recorded that is of use to determine sex, age, breeding status, likely cause of death, origin, and other issues. Age, the only variable found to influence litter quantities in stomach contents, is largely determined on the basis of development of sexual organs (size and shape) and presence of *Bursa of Fabricius* (a gland-like organ positioned near the end of the gut which is involved in immunity systems of young birds; it is well developed in chicks, but disappears within in the first year of life or shortly after). Further details are provided in Van Franeker 2004b. In the near future, an updated version of the manual should be published to improve details and maximize efficiency of methods.

After dissection, stomachs of birds are opened for analysis. Stomachs of Fulmars have two 'units': initially food is stored and starts to digest in a large glandular stomach (the *proventriculus*) after which it passes into a small muscular stomach (the *gizzard*) where harder prey remains can be processed through mechanical grinding. Initially, data for the two individual stomachs were recorded separately, but for the purpose of reduction in monitoring costs, the contents of proventriculus and gizzard are now combined.

If oil or chemical types of pollutants are present, these may be sub-sampled and weighed before rinsing the remainder of stomach content. Although this was a standard component at the start of our studies, requirements for the Dutch "graadmeter" and international EcoQO have a focus on plastic or at best MARPOL Annex V litter types. Thus, for financial efficiency, potential chemical pollutants in the stomachs are no longer part of the project. If sticky substances hamper further processing of the litter objects, hot water and detergents are used to rinse the material clean as needed for further sorting and counting under a binocular microscope.

The following categorization is used for plastics and other rubbish found in the stomachs:

1 PLASTICS (PLA)

1.1 Industrial plastic pellets (IND). These are small, often cylindrically-shaped granules of ± 4 mm diameter, but also disc and rectangular shapes occur. Various names are used, such as pellets, beads or granules. They can be considered as "raw" plastic or a half-product in the form of which, plastics are usually first produced (mostly from mineral oil). The raw industrial plastics are then usually transported to manufacturers that melt the granules and mix them with a variety of additives (fillers, stabilizers, colourants, anti-oxidants, softeners, biocides, etc.) that depend on the user product to be made. For the time being, included in this category is a relatively small number of very small, usually transparent spherical granules, also considered to be a raw industrial product.

1.2 User plastics (USE) (all non-industrial remains of plastic objects) differentiated in the following subcategories:

1.2.1 sheetlike user plastics (she), as in plastic bags, foils etc., usually broken up in smaller pieces;

1.2.2 threadlike user plastics (thr) as in (remains of) ropes, nets, nylon line, packaging straps etc. Sometimes 'balls' of threads and fibres form in the gizzard;

1.2.3 foamed user plastics (foa), as in foamed polystyrene cups or packaging or foamed polyurethane in mattresses or construction foams;

- 1.2.4 **fragments (fra)** of more or less hard plastic items as used in a huge number of applications (bottles, boxes, toys, tools, equipment housing, toothbrushes, lighters etc);
- 1.2.5 **other (oth)**, for example cigarette filters, rubber, elastics etc., so items that are 'plastic-like' or do not fit into a clear category.

2 RUBBISH (RUB) other than plastic:

- 2.1 **paper (pap)** which besides normal paper includes silver paper, aluminium foil etc, so various types of non-plastic packaging material;
- 2.2 **kitchenfood (kit)** for human food wastes such as fried meat, chips, vegetables, onions etc, probably mostly originating from ships' galley refuse;
- 2.3 **various rubbish (rva)** is used for e.g. pieces of timber (manufactured wood); paint chips, pieces of metals etc.;
- 2.4 **fish hook (hoo)** from either sportfishing or longlining.

Further optional categories of stomach contents

- 3 POLLUTANTS (POL)
for items indicating industrial or chemical waste remains such as slags (the remains of burning ovens, eg remains of coal or ore after melting out the metals); tar-lumps (remains of mineral oil); chemical (lumps or 'mud' of paraffine-like materials or sticky substances arbitrarily judged to be unnatural and of chemical origin) and featherlumps (indicating excessive preening by the bird of feathers sticky with oil or chemical pollutants).
- 4 NATURAL FOOD REMAINS (FOO)
Numbers of specific items may be recorded in separate subcategories (fish otoliths, eye-lenses, squid-jaws, crustacean remains, jelly-type prey remains, scavenged tissues incl feathers, insects, other).
- 5 NATURAL NON-FOOD REMAINS (NFO)
Numbers of subcategories eg plant-remains, seaweed, pumice, stone and other may be recorded.

For the main categories 1 (plastic) and 2 (rubbish) we record for each stomach and each (sub)category:

- incidence (Presence or absence) and
- abundance by number (count of Number of items)
- abundance by mass (Weight in grams) using Sartorius electronic weighing scale after a one to two day period of air drying at laboratory temperatures. For marine litter (categories 1 to 3 above), this is done separately for all subcategories. In the early Fulmar study we also weighed the natural-food and natural-non-food categories as a whole, but this was discontinued in 2006 to reduce costs. Weights are recorded in grams accurate to the 4th decimal (= tenth of milligram).

To be able to sort out items of categories 1 and 2, all other materials in the stomachs described in categories 3 to 5, have to be cleaned out. However in these latter categories, further identification, categorization, counting, weighing and data-processing is not essential for the EcoQO. Whether details are recorded depends of the interest of the participating research group and their reasons to collect beached Fulmars.

In addition to the acronyms used for (sub)categories as above, further acronyms may be used to describe datasets. Logarithmic transformed data are initiated by 'ln'; mass data are characterised by capital G (gram) and numerical data by N(number). For example lnGIND refers to the dataset that uses ln-transformed data for the mass of industrial plastics in the stomachs; acronym NUSE refers to a dataset based on the number of items of user plastics.

Analysis

Data from dissections and stomach content analysis are recorded in Excel spreadsheets and next stored in Oracle relational database. GENSTAT 8 was used for statistical tests. As concluded in the pilot study (Van Franeker & Meijboom 2002) and later reports, statistical analysis of data for presence of trends over time is conducted using mass-data. Tests for trends over time are conducted using linear regressions fitting ln-transformed plastic mass values for individual birds on the year of collection. Logarithmic transformation is needed because the original data are strongly skewed and need to be normalized for the statistical procedures. Tests for 'long term' trends use the full data set; 'recent' trends only use the past ten years of data.

For earlier Dutch reports, the tests on significance of trends on the chosen indicators of 'total plastic', 'industrial plastic' and 'user plastic' were the final main output. Focus was on significance of trends in specified categories without defining the final target. However, the wording of the Fulmar-Litter-EcoQO as now proposed in OSPAR is:

"There should be less than 10% of Northern Fulmars having 0.1 gram or more plastic in the stomach in samples of 50-100 beached fulmars from each of 5 different regions of the North Sea over a period of at least 5 years".

Thus, the information requested now focuses more on the information on 'total plastic' and 5-year averages for mass of the combined plastics in the bird stomachs. Such information is already incorporated in the Dutch approach, and merely requires a simplified data-presentation for EcoQO purposes. In the background however, tests using individual data as described above, and data collection on specified main litter categories, continue to play an important role for correct interpretation of the EcoQO metric.

5 Results

5.1 Monitoring in the Netherlands 1979-2007 and trends

Over the year 2007 a total of 67 Fulmars was collected from Dutch beaches, of which 61 proved to have an undamaged stomach adequate for the EcoQO research. This good sample size somewhat compensated the lower number of stomachs obtained in the previous year (27). A sample size of 40 or more is recommended to reliably characterize the pollution level in a particular time-frame and area (i.e. the annual 'average'). However, for multi-year trends a lower sample size in a particular year is not a problem, as analyses are not based on annual averages but on individual data for each bird.

In the year 2007, 56 out of 61 stomachs (92%) contained plastic, with an overall average number of nearly 36 items and mass of 0.37 gram per bird (Table 1). Non-plastic rubbish was found in about one in five stomachs, most frequently being galley food wastes. Numerical and mass abundance of plastics in 2007 were in line with relatively high occurrence of plastics observed in the smaller 2006 sample.

As convened in earlier reports, the metric for discussion of trends focuses on the mass of plastics in stomachs, in which the

- 'current situation' is described by the last 5-year average as above;
- 'long-term trends' refer to the full dataset (now 1979-2007)
- recent trends' are defined as trends over the past 10 years (now 1998-2007); and
- Trends are tested for significance by linear regressions of ln-transformed plastic data of individual birds against year.

The **current situation** as shown in the 5 year average shown in the bottom of table 2 fits in with the pattern of reduced plastic loads in Fulmar stomachs after peak levels in the 1990s. Mean values over the most recent 5 years (2003-2007; 309 Fulmars) are that 93% of birds had plastic in the stomach, with an average number of 27 pieces, and average mass of 0.28 gram plastic. Thus, the year 2007 was above 'average' for the situation over the past 5 years, which implies that improvements seen around the turn of the century seem to have stopped, and the trend in amounts of ingested plastic over the most recent years seems upward rather than downward.

The **long term 1979-2007 trend** analysis for "all plastics" ignores the 1990s peak in pollution levels and sees no significant change (Table 3A), i.e. indicates comparable levels in the 1980s and recent period. However, compared to the 1980s, the composition of plastic litter has strongly changed, with a significantly reduced proportion of industrial plastics and a somewhat increased mass of user plastics from discarded waste (Table 3A).

Statistical tests for **recent trends over the past 10 years (1998-2007)** (Table 3B) no longer show significant changes. As was feared from data interpretation in the 2006 report, the significant downward trends over earlier 10 year periods have ceased in the tests for the 1998-2007 time frame. Both industrial and user plastics contributed to this change.

A general overview of the various trends in plastic pollution since the 1980s is best obtained from Fig.1, which is based on the data from Table 2, recalculated to 'stable' 5-year means, each time shifting one year ahead. Shown are trends in plastic incidence, average number and average mass of plastic per bird, and specifies industrial and user plastics in those figures. In all three aspects, the increase in plastic pollution between the 1980s and the second half of the 1990s is visible, and is completely caused by increased user plastics, masking substantial decreases in industrial plastic over that period. In the late 1990s nearly 100% of Fulmars had plastic in the stomach, approaching 30 particles and 0.6 gram mass of plastic per bird. The graphs show that these late 1990s figures represented peak levels and that since then, on top of the continued decrease of industrial plastic, user plastics also started a downward trend. Remarkably, this is not the case when looking at the average number of plastic items, which has remained at a more or less constant high level of near 30 pieces per bird. Apparently, characteristics of user plastics are changing with smaller fragments becoming more dominant.

However, Fig.1 clearly illustrates that plastic ingestion continues to occur at a very high level, and that decreases in plastic mass seen around the change of the century have slowed down and so far provide no evidence for improvement following implementation of the EU Port Reception Facilities Directive in late 2004.

Table 1

Summary of sample characteristics and stomach contents of Fulmars collected for Dutch marine litter monitoring in the year 2007. For each litter-(sub)category the table lists: Incidence, representing the proportion of birds with one or more items of the litter category present; the average number of items per bird stomach; the average mass per bird stomach; and the maximum mass observed in a single stomach. The final column shows the geometric mean mass, which is calculated from ln-transformed values as used in trend-analyses. The bottom note explains sample composition in terms of age, sex, origin (by colourphase; darker phases are of distant Arctic origin), death cause oil, and the average condition-index (which ranges from emaciated condition=0 to very good condition=9). Although only age is currently considered relevant in affecting litter contents in stomachs of Dutch birds, this is not necessarily true in other comparisons.

Year 2007 (n=61 *)		incidence	average number of items	average mass of litter (g/bird) ± standard deviation	max. mass recorded	geometric mean mass (g/bird)
ALL PLASTICS		92%	35.6	0.366 ± 0.399	2.3	0.1287
1.1	INDUSTRIAL PLASTIC	70%	3.1	0.070 ± 0.090	0.4	0.0182
1.2	USER PLASTIC	90%	32.5	0.296 ± 0.355	2.0	0.0957
1.2.1	sheets	61%	3.8	0.009 ± 0.020	0.1	0.0028
1.2.2	threads	49%	1.5	0.032 ± 0.110	0.6	0.0023
1.2.3	foamed	69%	6.9	0.019 ± 0.035	0.2	0.0048
1.2.4	fragments	85%	19.1	0.203 ± 0.271	1.7	0.0611
1.2.5	other plastic	34%	1.2	0.033 ± 0.068	0.3	0.0028
OTHER RUBBISH		33%	2.7	0.052 ± 0.147	0.8	0.0027
2.1	paper	0%	0.0	0.000 ± 0.000	0.0	0.0000
2.2	kitchenwaste (food)	23%	2.6	0.039 ± 0.135	0.8	0.0015
2.3	rubbish various	11%	0.2	0.013 ± 0.062	0.5	0.0005
2.4	fishhook	0%	0.0	0.000 ± 0.000	0.0	0.0000

* of which 42% adult, 46% male, 90% LL (white colourphase); 3% death caused by oil; and 1.3 average conditionscore

Table 2

Annual details for plastic abundance in Fulmars from the Netherlands

For separate and combined plastic categories, incidence (%) represents the proportion of birds with one or more items of that litter present; number (n) abundance by average number of items per bird; and mass (g) abundance by average mass per bird in grams. The column on the far right indicates level of performance in relation to the OSPAR EcoQO, viz. the percentage of birds having more than the critical level of 0.1 gram of plastic in the stomach. The bottom line of the table shows the 'current' situation as the average over the past 5 years. *Note sample sizes (n) to be very low for particular years implying low reliability of the annual averages for such years, not to be used as separate figures. Also note erratic variability in age proportions of birds in samples, where age is known to influence amount of litter in the stomach. Trend analyses (table 3) are not based on annual averages, but on values from all individual birds, together and in age-groups, to overcome problems of years of poor sample size or variable age composition.*

YEAR	n	% adult	INDUSTRIAL PLASTICS			USER PLASTICS			ALL PLASTICS (industrial + user)			EcoQO > 0.1 g
			%	n	g	%	n	g	%	n	g	
1979	1	0%	100%	2.0	0.07	100%	3.0	0.17	100%	5.0	0.24	100%
1980												
1981												
1982	3	0%	100%	5.0	0.11	67%	6.0	0.50	100%	11.0	0.61	100%
1983	19	37%	84%	8.8	0.19	89%	7.2	0.31	100%	16.0	0.49	89%
1984	20	40%	70%	9.6	0.19	90%	8.4	0.17	90%	17.9	0.35	55%
1985	3	33%	100%	5.3	0.14	100%	5.0	0.14	100%	10.3	0.28	100%
1986	4	25%	50%	0.8	0.02	75%	4.8	0.06	75%	5.5	0.08	25%
1987	15	67%	80%	3.9	0.11	67%	8.9	0.09	80%	12.7	0.20	53%
1988	1	0%	0%	0.0	0.00	100%	2.0	0.04	100%	2.0	0.04	0%
1989	4	50%	75%	5.3	0.14	100%	11.0	0.16	100%	16.3	0.29	75%
1990												
1991	1	0%	0%	0.0	0.00	100%	11.0	0.14	100%	11.0	0.14	100%
1992												
1993												
1994												
1995	2	50%	100%	1.5	0.02	100%	3.5	0.03	100%	5.0	0.06	0%
1996	8	63%	75%	2.9	0.07	100%	24.5	0.19	100%	27.4	0.26	63%
1997	31	16%	74%	5.9	0.13	97%	29.8	0.60	97%	35.8	0.73	84%
1998	74	45%	69%	3.1	0.07	95%	25.9	0.88	96%	29.0	0.95	72%
1999	107	69%	58%	3.4	0.06	97%	31.8	0.38	98%	35.3	0.44	61%
2000	38	58%	61%	3.4	0.08	100%	18.6	0.27	100%	22.0	0.35	61%
2001	54	37%	63%	2.6	0.06	96%	20.4	0.18	96%	22.9	0.24	48%
2002	56	54%	68%	4.6	0.09	96%	47.2	0.41	98%	51.8	0.50	68%
2003	39	56%	51%	2.3	0.05	92%	26.3	0.12	95%	28.5	0.17	54%
2004	131	79%	54%	2.6	0.06	91%	20.8	0.22	91%	23.4	0.27	60%
2005	51	67%	53%	2.0	0.05	96%	15.8	0.22	98%	17.8	0.27	47%
2006	27	59%	78%	3.5	0.08	93%	30.4	0.23	93%	33.9	0.30	85%
2007	61	39%	70%	3.1	0.07	90%	32.5	0.30	92%	35.6	0.37	70%
03-07 *	309	64%	59%	2.6	0.06	92%	23.8	0.22	93%	26.5	0.28	61%

* Five-year data were averaged over all individual birds in the five year period (so not from annual averages)

Table 3**Details of linear regression analyses of the selected litter indicators.**

Analysis of trends was conducted by linear regression, fitting ln-transformed litter mass values for individual birds on the year of collection. Tests were conducted over the full time period 1979-2007 (Table 3A) and the most recent 10 years of data (Table 3B). *The regression line ('trend') is described by $y = \text{Constant} + \text{estimate} * x$ in which y is the calculated value of the regression-line for year x . When the t -value of a regression is negative it indicates a decreasing trend in the tested litter-category; a positive t -value indicates increase. A trend is considered significant when the probability (p) of misjudgement of data is less than 5% ($p < 0.05$). Significant trends in the table have been labelled with positive signs in case of increase (+) or negative signs in case of decrease (-). Significance at the 5% level ($p < 0.05$) is labelled as - or +; at the 1% level ($p < 0.01$) as - or ++; and at the 0.1% level ($p < 0.001$) as - or +++.*

**A. LONG TERM TRENDS 1979-2007
for plastics in Fulmar stomachs, the Netherlands**

INDUSTRIAL PLASTIC (lnGIND)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	750	103.9	-0.0540	0.0137	-3.94	<0.001	---
adults	418	63.6	-0.0341	0.0208	-1.64	0.103	n.s.
non adults	320	102.9	-0.0534	0.0186	-2.88	0.004	--
USER PLASTICS (lnGUSE)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	750	-31.6	0.0145	0.0126	1.15	0.249	n.s.
adults	418	-22.6	0.0099	0.0186	0.53	0.596	n.s.
non adults	320	-77.6	0.0377	0.0155	2.43	0.016	+
ALL PLASTICS COMBINED (lnGPLA)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	750	27.5	-0.0148	0.0115	-1.29	0.198	n.s.
adults	418	9.4	-0.0059	0.0184	-0.32	0.748	n.s.
non adults	320	10.7	-0.0063	0.0143	-0.44	0.662	n.s.

**B. RECENT 10-year TRENDS (1998-2007)
for plastics in Fulmar stomachs, the Netherlands**

INDUSTRIAL PLASTIC (lnGIND)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	638	-1.4	-0.0015	0.0300	-0.05	0.960	n.s.
adults	378	-30.6	0.0130	0.0407	0.32	0.750	n.s.
non adults	250	43.5	-0.0237	0.0444	-0.53	0.594	n.s.
USER PLASTICS (lnGUSE)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	638	66.9	-0.0347	0.0255	-1.36	0.174	n.s.
adults	378	102.8	-0.0527	0.0363	-1.45	0.147	n.s.
non adults	250	23.3	-0.0127	0.0347	-0.37	0.714	n.s.
ALL PLASTICS COMBINED (lnGPLA)	<i>n</i>	Constant	estimate	s.e.	t	p	
all ages	638	82.3	-0.0422	0.0249	-1.69	0.091	n.s.
adults	378	94.3	-0.0483	0.0358	-1.35	0.177	n.s.
non adults	250	68.3	-0.0350	0.0329	-1.07	0.287	n.s.




Fig. 1 Visual summary of Fulmar-Litter monitoring results in the Netherlands 1982-2007, comparing average data for incidence, number of items and mass in the 1980's with running 5-year averages for the more recent period.

Annual data: geometric means

As explained in methods, the statistical tests (Table 3) for trends over time do not use annual or multi-year averages, but are based on stomach contents data from individual birds and year of collection. This allows greater detail and the inclusion of data from years where only small samples of birds were collected. Values for plastic contents are logarithmically transformed, because data are not normally distributed with a few high values obscuring trend analysis. Logarithmic transformation normalises the distribution of data and reduces the influence of the exceptionally high values.

However, annual figures are more convenient for regular annual updates in a monitoring program and since 1997 the Dutch annual sample sizes have usually been large enough to calculate annual means. Logarithmic transformation of data is still needed, but the average of logarithmic values can be transformed back into a 'normal' value, which is then known as the 'geometric mean'. Geometric means are appropriate to make comparisons between groups of samples (years, but also regions), but it has to be kept in mind that they can be very different from normal averages ('arithmetic means'). Since logarithmic transformation reduces higher values, the geometric mean is usually considerably lower than the arithmetic mean for the same data. In mass data for plastics in the Fulmar stomachs, geometric means are only about one third of the arithmetic means (see table 1). Annual geometric means for total plastic mass in the Fulmar stomachs since 1997 and the combined figure for the early 1982-1990 period are shown in Fig. 2. Graphs illustrate the trends also found by regressions in Table 3, and clearly illustrate the effects of age. Differences between the age groups are mostly consistent between annual samples indicating that summarized monitoring results can be expressed as the figure for all ages combined (Summary Fig. *ii*). As shown in tables 1-3 and Fig.1, recent years including the year 2007 show no evidence for continued improvement in the marine litter situation but suggests stabilization or even a weak increase in pollution levels after 2003.




Figure 2 Annual geometric means for mass of plastics in stomachs of beached Fulmars from the Netherlands 1982-2007 for all age groups combined (including birds of unknown age), adult birds and non-adults, with sample sizes in brackets in the x-axis labels. Data illustrate the trends and consistency in age-differences that allow usage of the all-age trendline in summaries.

Geometric mean masses are also an appropriate basis to compare the separate trends in abundance of industrial and user plastics (Fig.3). The remarkable jump in industrial plastic mass in the small 2006 sample was not fully confirmed in the larger 2007 sample, but nevertheless relatively high within the recent years. The high level of user plastics in 2006 was confirmed in the 2007 samples. Figure 3 suggests that user plastics irregularly decreased from the mid 1990's to 2003, but seem to be steadily increasing since then. This is particularly disappointing since the turning point into an increase, more or less coincides with the start of the implementation of the EU Directive on Port Reception Facilities, which clearly had the opposite intention. In the 1980s about equal masses of both types of plastic were present in the stomachs of Fulmars, but nowadays user plastics represent \pm 80% of the plastic mass in Fulmar stomachs and exceed levels seen the 1980s.




Figure 3 Annual geometric means for mass of industrial and user plastics in stomachs of beached Fulmars from the Netherlands 1982-2007 (all ages)

Dutch data in terms of the OSPAR EcoQO metric

ICES working groups, followed by OSPAR, have always described the EcoQO metric for marine litter in terms of a percentage of birds exceeding a critical value of plastic in the stomach. At first sight, one might argue that it would be easier to use a definition based on for example only the average mass of plastics. However, whether intentional or not, the ‘percentage plus critical value’ definition represents a sort of simplified procedure that avoids the mathematical problems caused by a few excessive stomach contents distorting comparative analyses. In the testing procedures and geometric means used above, such problems are overcome by logarithmic transformation of data. And although this is a standard statistical procedure, it is not always easily conveyed to the general public, and differences between means (arithmetic versus geometric) can be confusing. The EcoQO metric avoids such problems by using classes of birds in which the exceptional stomach contents lose their influence. Currently, the target for acceptable ecological quality has been defined as the situation in which “less than 10% of Northern Fulmars has 0.1 gram or more plastic in the stomach; in all North Sea regions; for a consecutive period of at least five years”.

In most earlier years, the simplified mode of presentation of annual datapoints for the EcoQO metric as in Fig. 4 has shown fairly good similarity to the more sophisticated use of annual figures for using geometric means for mass as in Fig.2. However, in last years report the EcoQO figure for 2006 (van Franeker et al 2008) made an abrupt change upward (from 47% to 85% of birds exceeding the critical 0.1g level). This was very much “out of proportion” with the more moderate changes observed in the underlying data. The explanation for the disproportionate response was a combination of a small sample size and the fact that in the current situation, many individuals are close to the critical level of 0.1g of plastics in the stomach (van Franeker et al 2008 Appendix C), which may cause that relatively small changes lead to a strong response in the EcoQO metric.




Figure 4 EcoQO performance of Fulmars from the Netherlands 1980s to 2007. Annual percentages of beached Fulmars having more than 0.1g plastic in the stomach for adult birds, non adults and for all age groups combined (sample sizes in brackets in x-axis labels). Target level for acceptable ecological quality as in preliminary OSPAR documentation.

Insufficient sample size in 2006 also showed in the unusual data for different age groups (Fig.4). Fortunately, the 2007 sample was larger but confirms that the amounts of plastic in the birds stomachs tends to be on the increase. Instability in annual results as in Fig.4 re-emphasizes that evaluations should not be based on a year to year comparisons, but must be based on averages and trends over a larger number of years! In line with the wording for the EcoQO target, it is thus sensible to present EcoQO data for periods of five years. Calculated from individual data, 61% of Dutch Fulmars over the past 2003-2007 period exceeds the 0.1g critical EcoQO level. Figure 5 shows a graph of time trends in the 5-year average EcoQO performance of Fulmars found in the Netherlands. With the Y-axis scaled to 100%, the wide gap from the 10% EcoQO target is strongly visualised and emphasizes the need for further improvement. However, at this scale the graphs insufficiently shows the changes since the mid 1990's. The same data at a finer scale can be seen in Fig iii of the summary, showing gradual improvements in EcoQO performance (from 67% down to 57% exceeding 0.1g level) until the 2001-2005 period, but a reversed pattern since then.




Figure 5 EcoQO performance of Fulmars from the Netherlands 1980s to 2007. Running 'five-year-averages' for the percentage of beached Fulmars having more than 0.1g plastic in the stomach (all ages). Samples sizes in brackets in x-axis labels. Target level for acceptable ecological quality as in preliminary OSPAR documentation. See also Figure *iii* in the summary.

5.2 Results North Sea

Funding from the NYK Group Europe Ltd has made it possible to process stomach samples from other North Sea locations for the year 2005-2007 and to continue the international coordination of sampling for the Fulmar project into 2009.

The combination of the Dutch and the international efforts has effectively permitted continuation of the EU funded Save the North Sea Fulmar study 2002-2004.

All projects combine implement the OSPAR litter EcoQO as requested by North Sea Ministers at their Bergen meeting in 2002.

Recent integrated results of the international North Sea EcoQO project are being prepared for a scientific publication. Quantities of plastics decrease from the southern North Sea to the north: Fulmars from the Netherlands have about twice as much plastic in their stomach than Fulmars from the Scottish Islands. For earlier reports on the North Sea situation, see van Franeker et al 2008.




Figure 6 Fulmar-Litter-EcoQO study sites
(colours of stars indicate regional groups)

6 Conclusions

With an increasing number of study years after the initial pilot study (Van Franeker & Meijboom 2002), the Fulmar-Litter monitoring program has strongly matured. Good annual samples for the Netherlands for most years since 1997, and international expansion of the project since 2002, have delivered a wealth of data and firmly established the approach of plastic abundance in stomachs of the Northern Fulmar as being suitable for monitoring marine litter in the framework of Ecological Quality Objectives (EcoQO's) for the North Sea. At the request of OSPAR, a *Background Document* has been prepared which has been published on the OSPAR website. (OSPAR 2008). The EcoQO approach may also become an element in the European Marine Strategy.

This report updates Dutch long term monitoring information for the Netherlands up to the year 2007. Combined with new data for all countries around the North Sea, data from this report will be used to prepare a scientific publication on the Fulmar-Litter-EcoQO, which is required to firmly establish the quality of the research and to further stimulate similar monitoring and awareness raising work in other regions.

- In 2007 in the Netherlands, 56 out of 61 Fulmar stomachs contained plastic (incidence 92%) with an overall average of 36 items per bird and average mass of 0.37 gram per bird. The sample size was adequate as a sample of 40 or more stomachs is considered sufficient for reliable annual averages.
- In terms of mass, the 2007 data suggest a slight increase for plastic abundance in stomachs of Dutch Fulmars from 2006 (0.30g per bird) to 2007.
- In terms of the metric used in the Fulmar-Litter-EcoQO, viz. the percentage of birds with more than 0.1 gram of plastic in the stomach, 70% of the Dutch Fulmars exceeded the critical level in 2007.
- The 'current situation' calculated as the five year average over years 2003-2007 (309 birds) is that 61% of beached Fulmars exceeds the critical EcoQO level of 0.1gram of plastic in the stomach. Incidence of plastics was 61%, with an average number of 26.5 items and average mass 0.28 g per bird.
- Thus, mass and EcoQO data for 2007 are above the current 5 year average for the Netherlands, and seem to indicate a gradual increase over the past five years
- Tests for long-term trends 1979-2007 do not reveal significant change because total plastic mass in Fulmar stomachs peaked in the late 1990s, but have since then returned to a level similar to that in the 1980s.
- However, over the years from the early 1980's to the 2000's, the composition of the ingested plastic has strongly changed from about equal mass proportions of industrial and user plastics in the to about 80% of user plastics in the current situation.
- Tests over the recent 10-year period 1998-2007 show that earlier decreases have come to a halt, and that no significant change can be detected at the moment. Ingested plastics seem to have decreased from high levels in the mid 1990's up to 2003, but that trend has certainly stopped and has maybe even reversed after year 2003.

7 Acknowledgements

In several stages of this long-term project, the coordinating activities, dissection work, data analysis and writing of reports have been financially supported by the Netherlands Ministry of Transport, Public Works and Water Management (VenW). Internationally, the Fulmar project truly expanded during the ***Save the North Sea*** project 2002-2004, which was co-funded by the EU Interreg IIIB program for the North Sea. Continuation of the international North Sea work has later been supported by awards from the NYK Group Europe. Faroe Island data are derived from a study on diets and litter funded by Chevron Upstream Europe.

Beached Fulmars are mostly collected by volunteers associated with the coordinators from a wide range of organisations. There are too many persons involved to be named. However, without that huge amount of volunteer effort, a project like this would be totally impossible.

8 References

- Arthur, C.; Baker, J.; & Bamford, H. (eds) 2009. Proceedings of the International Research Workshop on the Occurrence, Effects and Fate of Microplastic Marine Debris. Sep 9-11, 2008.. NOAA Technical Memorandum NOS-OR&R-30. NOAA, Silver Spring 530pp.
- Breen, H. 2002. Plastics discarded at sea. MARMAM Network message of 29 Jul 2002
- Browne, M.A.; Sissanayake, A.; Galloway, T.S.; Lowe, D.M.; & Thompson, R.C. 2008. Ingested microscopic plastic translocates to the circulatory system of the Mussel, *Mytilus edulis* (L.). *Environmental Science and Technology* 42: 5026-5031
- Camphuysen, C.J. 2005. Oiled-guillemot-EcoQO - Implementation Document. North Sea Pilot Project on Ecological Quality Objectives, Issue 4. Seabirds, EcoQO element (f). NIOZ Report, Royal Netherlands Institute for Sea Research, Texel. 12pp.
- Department for Transport 2008. Transport Statistics Report Maritime Statistics 2007.. DfT, London, pp 173. (www.dft.gov.uk/pgr/statistics/datatablespublications/maritime/compendium/maritimestatistics2007).
- Derraik, J.G.B 2002. The pollution of the marine environment by plastic debris: a review. *Marine Pollution Bulletin* 44: 842-852.
- EC 2000. Directive 2000/59/EC of the European Parliament and of the Council of 27 November 2000 on port reception facilities for ship-generated waste and cargo residues. *Official Journal of the European Communities* L 332: 81-90 (28 Dec 2000).
- EC (Commission of the European Communities) 2005. Proposal for a Directive of the European Parliament and of the Council establishing a framework for community action in the field of marine environmental policy (Marine Strategy Directive). COM(2005) 505 final, 24-10-2005 (*incl communication 504 final and Impact Assessment Annex SEC(2005)1290*).
- Edwards, R. 2005. Litter at sea means a bellyful of plastic. *New Scientist* 185(2481): 11.
- Endo, S.; Takizawa, R.; Okuda, K.; Takada, H.; Chiba, K.; Kanehiro, H.; Ogi, H.; Yamashita, R.; & Date, T. 2005. Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets: Variability among individual particles and regional differences. *Marine Pollution Bulletin* 50: 1103-1114.
- Fleet, D.M 2003. Untersuchung der Verschmutzung der Spülsäume durch Schiffsmüll and der deutschen Nordseeküste. (UFOPLAN) FAZ 202 96 183, im Auftrag des Umweltbundesamtes, Hochschule Bremen. 194pp.
- Guse, N., Fleet, D., van Franeker, J. & Garthe, S. 2005. Der Eissturmvogel (*Fulmarus glacialis*)- Mülleimer der Nordsee? *Seevögel* 26(2): 3-12.
- Hall, K. 2000. Impacts of marine debris and oil: economic and social costs to coastal communities. KIMO, c/o Shetland Islands Council, Lerwick. 104 pp.
- Laist, D.W. 1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. *Mar. Pollut. Bull.* 18(6B): 319-326.
- Laist, D.W. 1997. Impacts of marine debris: entanglement of marine life in marine debris including a comprehensive list of species with entanglement and ingestion records. pp 99-140 in: Coe, J.M. and Rogers, D.B. (eds.). *Marine debris sources, impacts and solutions*. Springer Series on Environmental Management. Springer Verlag, New York. 432pp.
- Mallory, M.L.; Roberston, G.J.; & Moenting, A. 2006. Marine plastic debris in northern fulmars from Davis Strait, Nunavut, Canada.. *Marine Pollution Bulletin* 52: 813-815.
- Moore, C.J. 2008. Synthetic polymers in the marine environment: a rapidly increasing, long-term threat. *Environmental Research* 108: 131-139.
- OECD-MTC (Organisation for Economic Co-operation and Development - Maritime Transport Committee) 2003. Cost savings stemming from non-compliance with international environmental regulations in the maritime sector. MEPC 49/INF.7. 14 March 2003. 53pp.
- Olsen, K. 2005. Havhesten - en flygende søppelbøtte. *Var Fuglefauna* 28: 28-32.
- OSPAR 2004. OSPAR Commission Summary Record. OSPAR 04/23/1-E, Annex 17.
- OSPAR Commission BDC 2005. Plastic particles in the stomachs of seabirds. BDC 05/3/1 Add.1 : 39-40.
- OSPAR 2008. Background Document for the EcoQO on plastic particles in stomachs of seabirds. OSPAR Commission, Biodiversity Series. ISBN 978-1-905859-94-8 Publication Number: 355/2008.

- Save the North Sea. 2004. Reduce marine litter: 'Save the North Sea' project results. Keep Sweden Tidy Foundation, Stockholm, 17pp.
- Stichting De Noordzee 2000. HOI's: beschikbaarheid en rendement. Inzamelfaciliteiten voor oliehoudend- en vast afval in Nederlandse zeehavens. Verslag Stichting de Noordzee, Utrecht.16pp.
- Stichting De Noordzee 2003. Coastwatch Onderzoek 2002. Stichting de Noordzee, Utrecht. 17pp + addenda.
- Teuten, E.L.; Rowland, S.J.; Galloway, T.S.; & Thompson, R.C. 2007. Potential for Plastics to Transport Hydrophobic Contaminants. *Environmental Science and Technology* 41: 7759-7764
- Thompson, R.C., Olsen, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D. & Russell, A.E. 2004. Lost at sea: Where is all the plastic?. *Science* 304 (5672): 838-838.
- Van Franeker, J.A. 1985. Plastic ingestion in the North Atlantic Fulmar. *Marine Pollution Bulletin* 16: 367-369.
- Van Franeker, J.A. & Bell, P.J. 1988. Plastic ingestion by petrels breeding in Antarctica. *Marine Pollution Bulletin* 19: 672-674.
- Van Franeker, J.A. & Meijboom, A. 2002. Litter NSV - Marine litter monitoring by Northern Fulmars: a pilot study. ALTERRA-Rapport 401. Alterra, Wageningen, 72pp.
- Van Franeker, J.A. & Meijboom, A. 2003. Marine Litter Monitoring by Northern Fulmars: progress report 2002. Alterra-rapport 622. Alterra, Wageningen, 49 pp.
- Van Franeker, J.A. 2004a. The Fulmar-Litter-EcoQO in relation to EU Directive 2000/59/EC on Port Reception Facilities. Rapportage in opdracht van Directoraat Generaal Goederenvervoer, Ministerie van Verkeer en Waterstaat. Contract DGG-27020138. DGG-VenW, Den Haag. 47pp.
- Van Franeker, J.A. 2004b. Save the North Sea - Fulmar Study Manual 1: Collection and dissection procedures. Alterra Rapport 672. Alterra, Wageningen. 38pp.
- Van Franeker, J.A. 2004c. Fulmar wreck in the southern North Sea: preliminary findings. *British Birds* 97: 247-250.
- Van Franeker, J.A., Meijboom, A. & De Jong, M.L. 2004. Marine litter monitoring by Northern Fulmars in the Netherlands 1982-2003. Alterra Rapport 1093. Alterra, Wageningen. 48pp.
- Van Franeker, J.A. 2005. Schoon strand Texel 2005: onderzoeksresultaten van de schoonmaakactie van het Texelse strand op 20 april 2005. Alterra speciale uitgave 2005/09. Alterra, Texel. 23pp.
- Van Franeker, J.A., Heubeck, M., Fairclough, K., Turner, D.M., Grantham, M., Stienen, E.W.M., Guse, N., Pedersen, J., Olsen, K.O., Andersson, P.J. & Olsen, B. 2005. 'Save the North Sea' Fulmar Study 2002-2004: a regional pilot project for the Fulmar-Litter-EcoQO in the OSPAR area. Alterra-rapport 1162. Alterra, Wageningen. 70pp.
- Van Franeker, J.A. & Meijboom, A. 2006. Fulmar Litter EcoQO Monitoring in the Netherlands 1982-2004 in relation to EU Directive 2000/59/EC on Port Reception Facilities.. Report for the Ministry of Transport, Public Works and Water Management (VenW), contract nr DGTL/ZH/2.53.2.5012. Alterra, Texel. 41pp..
- Van Franeker, J.A.; & Meijboom, A. 2007. Fulmar Litter EcoQO Monitoring in the Netherlands 1982-2005 in relation to EU Directive 2000/59/EC on Port Reception Facilities.. Wageningen IMARES Report Nr C019-07. IMARES Texel, 40pp
- Van Franeker, J.A.; & the SNS Fulmar Study Group 2008. Fulmar Litter EcoQO monitoring in the North Sea - results to 2006. IMARES Report nr C033/08. Wageningen IMARES, Texel. 53pp.
- Vlietstra, L.S. & Parga, J.A. 2002. Long-term changes in the type, but not the amount, of ingested plastic particles in Short-tailed Shearwaters in the southeastern Bering Sea. *Marine Pollution Bulletin* 44: 945-955.

9 Quality Assurance

IMARES utilises an ISO 9001:2000 certified quality management system (certificate number: 08602-2004-AQ-ROT-RvA). This certificate is valid until 15 December 2009. The organisation has been certified since 27 February 2001. The certification was issued by DNV Certification B.V. Furthermore, the chemical laboratory of the Environmental Division has NEN-AND-ISO/IEC 17025:2005 accreditation for test laboratories with number L097. This accreditation is valid until 27 March 2009 and was first issued on 27 March 1997. Accreditation was granted by the Council for Accreditation, with the last inspection being held on the 5th of October 2007.

Justification

Rapport nr	C032/09
IMARES Project Number:	430.61001.04
Client project reference	DGLM 08001 / 4500125534

The scientific quality of this report has been peer reviewed by a colleague scientist and the head of the department of Wageningen IMARES.

Approved:




Dr. N.M.J.A. Dankers
Senior Scientist

Date:

16-apr-2009

Approved:



Dr. H.J. Lindeboom – *Science Director IMARES*

Date:

16-apr-2009

Number of copies:	30
Number of pages	39
Number of tables:	3
Number of graphs:	iii + 6 + 2 photo's
Number of appendix attachments:	0