

Chromosome numbers in *Begonia* 2*

R. A. H. Legro¹ and J. Doorenbos²

¹ Laboratory of Plant Taxonomy, Agricultural University, Wageningen, the Netherlands

² Laboratory of Horticulture, Agricultural University, Wageningen, the Netherlands

Received: 7 May 1971

Summary

The somatic chromosome numbers of 90 species of *Begonia* were established. The following numbers were found: 16, 22, 26, 28, 30, 32, 34, 36, 38, 44, 48, 52, 56, 60, 64 and 76. *B. gigantea* Wall. ($2n = 16$) has the lowest chromosome number reported so far in *Begonia*. The conclusion of a previous paper that 22, 28 and 56 are the most common numbers within the genus was confirmed, but it was found that there is also a considerable number of species with $2n = 38$.

Introduction

At the conclusion of a previous publication (Legro and Doorenbos, 1969) in which the chromosome numbers of 100 species of *Begonia* were given, the prospect was held out that shortly the chromosome numbers of another 100 species would be made known. The present article only partly fulfils this promise. In many species, the cytological observations presented more difficulties than expected. Other plants have not yet been identified with certainty. As a result, only ninety chromosome counts, not all of them conclusive, can be added to those of our previous paper.

In the course of the present study, a few imperfections of the first paper have come to light, viz a few cases of incorrect nomenclature and one case of an incorrect chromosome number. The species concerned have been listed again and are mentioned separately in the discussion.

Material and methods

Much of the plant material came from the sources mentioned in the first paper. An important contribution, mainly of Mexican species, was made by R. Ziesenhenne of Santa Barbara, California. Mrs. J. Neal of Worthing, Surrey, England, sent us several species, such as *B. prismatocarpa*, *B. venusta* and *B. pavonina*, the latter two collected in the Cameron mountains in Malaya. J. J. Bos gave us a number of species collected in the Cameroons, such as *B. elatostemmoides*, *B. mannii* and *B. quadrialata*, and in Natal, such as *B. princeae*. *B. ficicola* was brought by M. C. G. Middelburg from the type locality (Ekumbe) in the Cameroons. *B. keniensis* was collected by J. J. F. E. de

* Publication 356, Laboratorium voor Tuinbouwplantentuinbouw, Landbouwhogeschool, Wageningen, the Netherlands.

Wilde in Ethiopia. J. F. M. Zieck sent us several species from New Guinea, such as *B. physandra*.

Correct identification is a perennial problem in this genus, of which no modern monograph exists. The species have been compared with all available descriptions, but not with type material. Of the plants studied, voucher specimens have been deposited in WAG, the Herbarium of the Department of Plant Taxonomy of the Agricultural University at Wageningen.

Some authors have used a wider species concept than is used in the present study. Smith and Schubert in particular consider several of the specific names used here to be synonymous with others. In many cases the present authors agree with them, but nevertheless it seemed advisable to use a narrow species concept in this study, as cytological differences between morphologically related plants might well have come to light.

The chromosomes were studied in squash preparations of root tips according to the methods described in the previous paper.

Results

The results are presented in Table 1, which is set up in the same way as Table 1 of the previous article. Species marked with an asterisk have been mentioned already in the

Table 1. Survey of somatic chromosome numbers in *Begonia* species in comparison with numbers mentioned by previous authors.

	2n	Literature data		
		2n	authors ¹	year
<i>African species</i>				
Squamibegonia				
<i>B. haulevilleana</i> De Wild.	38			
Augustia				
<i>B. caffra</i> Meissn.	26			
<i>B. princeae</i> Gilg	26			
<i>B. suffruticosa</i> Meissn.	26			
Scutobegonia				
<i>B. ficicola</i> Irmsch.	34 + + + +			
<i>B. elatostemmioides</i> Hook. f.	(38)			
Loasibegonia				
<i>B. prismatocarpa</i> Hook. f.	32			
Tetraphila (Fusibegonia)				
<i>B. mannii</i> Hook. f.	38			
<i>B. polygonoides</i> Hook. f.	36 + +			
* <i>B. spraguei</i> C. M. Weber (<i>B. parva</i> Sprague)	36 + +			
Rostrobegonia				
<i>B. johnstonii</i> Oliv.	26			
<i>B. keniensis</i> Gilg	(38)			
<i>Asiatic species</i>				
Haagea				
<i>B. dipetala</i> Grah.	30	ca. 28	Heitz	1927
Reichenheimia				
<i>B. flocicifera</i> Bedd.	28 + + + + or 32	30	S. & B.	1957
	<i>B. rajah</i> Ridl.	30 + +		

Table 1 (Continued)

		Literature data		
	2n	2n	authors ¹	year
Spenanthera				
<i>B. roxburghii</i> A.DC.	22			
Monopteron				
<i>B. gigantea</i> Wall.	16		28, 32, 34, S. & B.	1961
<i>B. griffithiana</i> (A.DC.) Warb.	22		40, 42	
Platycentrum				
<i>B. pavonina</i> Ridl.	22			
<i>B. tenuifolia</i> Dryand.	22			
<i>B. venusta</i> Ridl.	44			
<i>B. versicolor</i> Irmsch.	22			
Diploclinium				
<i>B. subnummularifolia</i> Mer.	(32)			
Parvibegonia				
<i>B. crenata</i> Dryand.	56			
Petermannia				
<i>B. brevirimosa</i> Irmsch.	44			
<i>B. sp.</i> New Guinea	30			
Knesbeckia				
<i>B. cyclophylla</i> Hook. f.	22			
Section unknown				
<i>B. physandra</i> Merr. & Perry	28			
<i>American species</i>				
Doratometra				
<i>B. franconis</i> Liebm.	26			
Scheidweileria				
<i>B. luxurians</i> Scheidw.	56			
Steineria				
<i>B. hookeriana</i> Gard.	(56)			
<i>B. oxyphylla</i> A.DC.	(56)			
Pritzelia				
<i>B. crispula</i> Brade	(38)			
<i>B. epipsila</i> Brade	(56)			
<i>B. hispida</i> Schott var. <i>cucullifera</i> Irmsch.	56			
<i>B. hugelii</i> A.DC.	56			
<i>B. itaguassiensis</i> Brade	38++			
<i>B. listida</i> hort.	(76)			
<i>B. olsoniae</i> Sm. & Schub. (<i>yellogoana</i> Brade)	56			
<i>B. paleata</i> A.DC.	(38)			
<i>B. paranaensis</i> Brade	(56)			
<i>B. princeps</i> (Kl.) A.DC.	(48)	48	Piton	1962
<i>B. pulchella</i> Raddi	(56)			
<i>B. sarmentacea</i> hort.	(38)			
<i>B. scabrida</i> A.DC.	48			
<i>B. scharffiana</i> Regel	(56)			
<i>B. tomentosa</i> Schott	(56)			
<i>B. valida</i> Goebel	(38)	36, 38	Heitz	1927
<i>B. vitifolia</i> Schott (<i>acerfolia</i> hort. non HBK)	38		?	
<i>B. vitifolia</i> Schott (<i>reniformis</i> Hook. f. non Dry.)	38	36	Hamel	1937
Solananthera				
<i>B. solananthera</i> A.DC.	56			
Weilbachia				
<i>B. purpusii</i> Zies.	28			

CHROMOSOME NUMBERS IN BEGONIA. 2

Table 1 (Continued)

	2n	Literature data		
		2n	authors ¹	year
<i>B. pustulata</i> Liebm.	38			
<i>Huszia</i>				
* <i>B. micranthera</i> Griseb.				
var. <i>hieronymii</i> (Lind.) Sm. & Schub.	28			
var. <i>fimbriata</i> hort.	27			
var. <i>foliosa</i> Sm. & Schub.	28			
var. <i>venturia</i> hort.	27			
<i>Giroudia</i>				
<i>B. barkeri</i> Knowles & Westc.	28			
<i>B. chivatoa</i> Zies.	28			
<i>B. cobana</i> C.DC.	28			
<i>B. daedalea</i> Lemoine	28			
<i>B. kortsae</i> Zies.	28			
<i>B. hypolipara</i> Sandw.	28			
<i>B. lindleyana</i> Walp.	28			
<i>B. megaphylla</i> A.DC.	28			
<i>B. plebeja</i> Liebm.	28			
<i>B. pringlei</i> S. Watson	28			
<i>B. sericoneura</i> Liebm.	28			
<i>B. squamosa</i> C.DC.	28			
<i>B. squarrosa</i> Liebm. (<i>hepaticamaculata</i> Zies.)	28			
<i>B. villipetiola</i> C.DC.	28			
<i>Latistigma</i>				
<i>B. aconitifolia</i> A.DC.	60			
<i>B. platanifolia</i> Schott	60			
<i>Begoniastrum</i>				
<i>B. crinita</i> Oliv.	(34)			
<i>B. cucullata</i> Willd.	(64)			
<i>B. cucullata</i> var. <i>hookeri</i> (A.DC.) Sm. & Schub.	34			
<i>B. fischeri</i> Schrank var. <i>tovarensis</i> Irmsch.	34			
<i>B. hirtella</i> Willd.	34			
<i>B. leptotricha</i> C.DC.	34			
<i>B. leptotricha</i> 'Woolly-bear'	34			
<i>B. mollicaulis</i> Irmsch. (<i>subvillosa</i> hort. non Klotzsch)	34			
<i>B. parvifolia</i> Schott	32			
<i>B. porteriana</i> Fisch.	60			
<i>B. rotundifolia</i> Lam.	52			
<i>B. semperflorens</i> Link & Otto	(34)			
<i>Knesebeckia</i>				
<i>B. bulbiflora</i> Link & Otto	28			
<i>B. biserrata</i> Lindl. (<i>palmaris</i> A.DC.)	28			
<i>B. cavum</i> Zies.	28			
* <i>B. ignea</i> Warsz. (<i>palmaris</i> hort. non A.DC.)	56			
<i>B. kellermannii</i> C.DC.	28 + + +			
<i>B. martiana</i> Link & Otto	56			
<i>B. uniflora</i> S. Watson (<i>taylori</i> hort.)	28			
<i>B. viscida</i> Zies.	28			
<i>B. wollnyi</i> Herzog	28			
<i>Tetrachia</i>				
<i>B. egregia</i> N. E. Br.	(52)			

¹ S. & B. = Sharma and Bhattacharyya, 1961.

latter, but are repeated here because it appeared necessary to reconsider nomenclature or chromosome number.

Some results have been included here that are not yet fully conclusive. These tentative chromosome counts appear in the Table between brackets. The authors are fairly certain that they are correct, but have not been able to find sufficient cells showing distinct metaphase chromosomes which could be counted without any doubt.

Discussion

To the 18 different chromosome numbers listed in our previous article, the present results add only four new ones: 16, 32, 64 and 76, the last two being tentative counts. The recurrence of the same numbers and the fact that some appear to be characteristic of certain sections lead to the emergence of a pattern; the cytology of this genus may not be so complicated after all.

The chromosome numbers of West African species studied so far lie between 32 and 38. The determination of the exact number is hampered by the occurrence of bits of chromatin smaller than regular chromosomes. If these are not fragments, as has been assumed here, but chromosomes, the prevalent chromosome number among West African species would appear to be 38, at least within the large sections *Scutobegonia* and *Tetraphila*. The dainty *B. prismatocarpa*, however, which belongs to *Loasibegonia*, has 32 chromosomes.

In East Africa the number 26 is prevalent among the sections *Augustia* and *Rostrobegonia*. So far, *B. keniensis* ($2n = 38$) is the only exception. As this is a typical representative of *Rostrobegonia*, it may be of triploid origin. It is to be regretted that none of the 20 *Begonia* species of Madagascar (sections *Mezierea*, *Erminea*, *Quadrilobaria*, etc.) was available for study.

Among the Asiatic species studied, *B. gigantea* is the most interesting from a cytological point of view. Not only does it have the smallest number of chromosomes of all 190 species studied, but it is also the only species in which the individual chromosomes of the genome can be recognized (Fig. 1). The count of $2n = 16$ for this species leads to the speculation that the number of $2n = 22$, found in the Asiatic sections *Platycentrum* and *Sphenanthera*, may have arisen from a cross between a species with $2n = 16$ and one with $2n = 28$. The fact that *B. griffithiana*, morphologically very similar to *B. gigantea* and belonging to the same section, has 22 chromosomes supports this theory. Sharma and Bhattacharyya (1961) counted between 28 and 42 chromosomes for *B. gigantea*. These counts are very tentative (one wonders why they were published at all) but nevertheless render it highly unlikely that these authors had the same material as used for the present study.

B. dipetala of the small section *Haagea* is often treated as a subspecies of *B. malabarica* Lam. Irmscher (1961) however, considers it a species in its own right, and the present chromosome counts support this opinion. *B. venusta*, which has 3-5-fid placentae while those of the other species of *Platycentrum* are bifid, appears to be tetraploid in relation to these other species.

Knowledge of the large Asiatic sections *Reichenheimia*, *Diploclinium* and *Petermannia* remains scanty. Most species of *Petermannia* from New Guinea appear to have 44 chromosomes, but one unidentified species was found to have 30. Perhaps the species with 44 chromosomes arose from triploids of species with 30, rather than as tetraploids of species with 22 chromosomes..




Fig. 1. Mitotic metaphase of *Begonia gigantea* Wall. Note the two very large chromosomes. Magnification $\times 3100$.

The tuberous *B. physandra* has never been cultivated before. The ovaries (not described previously) were found to be two-celled. This probably makes it necessary to erect a new section for this species.

Among American sections, Pritzelia is one of the largest. Further study of the species of this section has confirmed that they can be divided into a group with 38 and a group with 56 chromosomes. So far, no correlation was found between these chromosome numbers and the morphological characteristics of the plants. The pretty *B. listida*, undoubtedly a true species but as far as we know not properly described yet, appears to have 76 chromosomes. This means it is tetraploid in relation to those species of Pritzelia which have 38.

B. princeps ($2n = 48$) may have originated as a hybrid between a species with 38 and one with 56 chromosomes. A plant grown as *B. scabrida* A.DC. in botanic gardens, and in fairly good agreement with A. de Candolle's description (although not all placentae are entire, as they should be), was also found to have 48 chromosomes.

B. vitifolia stands badly in need of taxonomic study. This species as interpreted by De Candolle, is grown in collections under the erroneous name *B. acerifolia* (non HBK). Although even Irmscher maintains that *B. reniformis* Hook. (non Dryander) is synonymous with *B. vitifolia*, it is a different species. *B. valida*, on the other hand, appears




Fig. 2. Mitotic early metaphase of *Begonia plumieri* A. DC., demonstrating the difficulties (small, very numerous chromosomes in a granular protoplasm) which have made it impossible to arrive at a conclusive chromosome count for this species. Magnification $\times 3000$.

to be closely related to *B. vitifolia* (sensu A. DC.). It is, however, beyond the scope of this article to solve these taxonomic puzzles.

The section Giroudia appears to be very homogeneous cytologically. The chromosomes of these species are relatively easy to count, with the exception of those of *B. pringlei*. In this species, one has to find exactly the right stage of the metaphase. Otherwise, one gets the impression that there are more than 28, possibly because of strong constrictions of the chromosomes. Within the related section Weilbachia, *B. pustulata* ($2n = 38$) is a notable exception among species with 28 chromosomes.

B. micranthera var. *hieronymii* is the same plant as *B. micranthera* of the previous publication. It is a dioecious plant with large pink flowers, and of much interest for hybridizing purposes. The var. *foliosa* is also dioecious, but the small white flowers have no ornamental value. From the USA we received two monoecious forms with showy, orange flowers, which are grown under the names *B. micranthera* var. *fimbriata* and *B. micranthera* var. *venturia*. Neither confirms to the original description of these varieties. Both were found to have 27 chromosomes. We regard the 'var. *fimbriata*' as a hybrid between *B. cinnabrina* ($2n = 26$) and, presumably, a form of *B. micranthera* with $2n = 28$, and the 'var. *venturia*' as a similar hybrid, although its parentage is less clear.

Further study of Begoniastrum has shown that 34 is a common chromosome number in this section. The material of *B. leptotricha* was imported from two sources in Brazil

as an unidentified species. The Begonia known among growers in the USA as 'Woolly-bear' differs from these imports in a more compact habit, broader, larger leaves and a denser fulvous tomentum.

B. cucullata was imported as an unidentified species from Venezuela. It has a stiff, erect and succulent stem, persistent foliaceous denticulate stipules and acute seeds. It agrees in all respects with the old picture of Loddiges' *B. spathulata*, reproduced by Smith and Schubert (1941, p. 102). It appears to be tetraploid, although 64 chromosomes were counted rather than the expected 68 (the number 56, given in the previous paper, was an administrative error). The plant called here *B. cucullata* var. *hookeri* has a much more spreading habit, stipules that are less broad and less persistent, and blunt seed. This may be the wild ancestor of *B. semperflorens*, although in the material studied there were several differences between the two.

There are still about 30 species in the collection of which the chromosome numbers could not yet be established with a reasonable degree of certainty. The reasons for this are threefold. In some species, the number of metaphase plates found is too small to permit definitive conclusions. Examples of such species are *B. dietrichiana* Irmsch., *B. fernandoi-costae* Irmsch., *B. obscura* Brade, *B. inciso-serrata* A.DC., *B. venosa* Skan and many others. Other species do yield sufficient metaphase plates, but the attainment of conclusive results is rendered very difficult because the chromosomes are very small, and the protoplasm is full of other granules, which are hard to distinguish from true chromosomes. This group includes species as *B. quadrialata* Warb., *B. angularis* Raddi and *B. lindeniana* A.DC. A third group comprises such species in which the sheer number of the very small chromosomes (well over 100) makes counting an almost impossible task. Examples of this (often complicated by granular protoplasm) are *B. convallariodora* C.DC., *B. serratifolia* C.DC. and *B. plumieri* A.DC. (Fig. 2). Presumably, these species have to be studied by more sophisticated methods than the ones used in the present study.

Literature

Bowden, W. M., 1945. A list of chromosome numbers in higher plants. I. Acanthaceae to Myrtaceae. *Am. J. Bot.* 32: 81-92.

Hamel, J. C., 1937. Etudes caryologiques sur quelques Begoniacées. *Revue Cytol. Biol. vég.* 2: 392-413.

Heitz, E., 1927. Über multiple und aberrante Chromosomenzahlen. *Abh. Naturw. Ver. Hamburg* 21: 48-58.

Irmscher, E., 1961. Begoniaceae, Schiebblattgewächse. Parey's Blumengärtnerie, 2nd ed., Vol. 2 B. 67-98; 838-839. Parey, Berlin.

Legro, R. A. H. & J. Doorenbos, 1969. Chromosome numbers in *Begonia*. *Neth. J. agric. Sci.* 17: 189-202.

Piton, F., 1962. Contributions à l'étude cytologique des Begoniacées. *Rev. Cyt. Biol. vég.* 24: 165-216.

Sharma, A. K. & U. C. Bhattacharyya, 1956-67. Cytological studies in *Begonia*. 1. *Cellule* 58: 307-330.

Sharma, A. K. & U. C. Bhattacharyya, 1961. Cytological studies in *Begonia*. 2. *Caryologia* 14: 279-301.

Smith, L. B. & B. G. Schubert, 1941. Revisión de la especies Argentinas del género *Begonia*. *Darwinia* 5: 78-117.

Zeilinga, A. E. 1962. Cytological investigations of hybrid varieties of *Begonia semperflorens*. *Euphytica* 126-136.