

Microbial degradation of amygdalin of bitter apricot seeds (*Prunus armeniaca*).

International Journal of Food Microbiology

Nout, M.J.R.; Tunçel, G.; Brimer, L.

[https://doi.org/10.1016/0168-1605\(94\)00115-m](https://doi.org/10.1016/0168-1605(94)00115-m)

This publication is made publicly available in the institutional repository of Wageningen University and Research, under the terms of article 25fa of the Dutch Copyright Act, also known as the Amendment Taverne. This has been done with explicit consent by the author.

Article 25fa states that the author of a short scientific work funded either wholly or partially by Dutch public funds is entitled to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed under The Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' project. In this project research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and / or copyright owner(s) of this work. Any use of the publication or parts of it other than authorised under article 25fa of the Dutch Copyright act is prohibited. Wageningen University & Research and the author(s) of this publication shall not be held responsible or liable for any damages resulting from your (re)use of this publication.

For questions regarding the public availability of this publication please contact openscience.library@wur.nl

Short Communication

Microbial degradation of amygdalin of bitter apricot seeds
(*Prunus armeniaca*)

M.J.R. Nout ^{a,*}, G. Tunçel ^b, L. Brimer ^c

^a Department of Food Science, Agricultural University, Bomenweg 2, 6703 HD Wageningen,
The Netherlands

^b Faculty of Engineering, Ege University, 35100 Bornova-Izmir, Turkey

^c Department of Pharmacology and Pathobiology, The Royal Veterinary and Agricultural University,
13 Bülowsvej, DK-1870 Frederiksberg C, Copenhagen, Denmark

Received 20 September 1993; revised 19 April 1994; accepted 8 August 1994

Abstract

Amygdalin is a cyanogenic glycoside occurring among others in almonds and bitter apricot seeds with interesting levels of dietary protein. Utilization of seeds for human or animal nutrition requires adequate detoxification. In the present paper, selected filamentous fungi (*Mucor circinelloides*, *Penicillium nalgiovense*) and yeasts (*Hanseniaspora valbyensis*, *Endomyces fibuliger*) were tested for their in-situ ability to decompose amygdalin. The latter (*Endomyces fibuliger*) was best able to grow on autoclaved bitter apricot seeds and detoxify them from 30 $\mu\text{Mol CN/g}$ dry matter to less than 1 $\mu\text{Mol CN/g}$ dry matter after 48 h of incubation at 27°C.

Keywords: Apricot seeds; Amygdalin; Cyanogenic; Detoxification; Fungi; Yeasts; *Endomyces fibuliger*

1. Introduction

Bitter apricot seeds (*Prunus armeniaca*) are by-products of the apricot processing industry. The seeds have a high protein content (20–25%, dry weight basis) and could be of interest as a food or feed ingredient. However, depending on variety, they contain approximately 50–150 $\mu\text{Mol/g}$ (dry weight basis) of the potentially toxic cyanogenic glycosides amygdalin and prunasin (Tunçel et al., 1990; Brimer et al., 1993). We reported earlier (Tunçel et al., 1990) on prospects for detoxification

* Corresponding author.

by biotechnological means, i.e. by using the endogenous enzymes combined with microbial fermentation. The tempe fermentation process using the fungus *Rhizopus oligosporus* as inoculant enabled a removal of around 70% of total cyanide potential. As we considered this inadequate from toxicological point of view, we developed a simple method based on the picrate test (Guignard, 1906), as used by Brimer and Moelgaard (1986), to screen microbial strains for their ability to degrade amygdalin by hydrolysis (Brimer et al., 1993). The results obtained can be summarized as follows: of 29 strains of lactic acid bacteria, 16 were unable to degrade amygdalin while the maximum extent of degradation was 27% of the initial level after overnight incubation; of 21 yeast strains, six could not degrade amygdalin, and maximum degradation was 98%; of 25 strains of filamentous fungi, only 1 could not degrade amygdalin while several strains were found able to degrade up till 99% of the initial level of amygdalin (Brimer et al., 1993). This paper describes detoxification trials with those strains able to degrade > 95% of initial levels of amygdalin, i.e. strains of *Hanseniaspora valbyensis*, *Endomyces fibuliger*, *Mucor circinelloides* and *Penicillium nalgiovense*.

2. Materials and methods

2.1 Materials.

Bitter apricot seeds were obtained from Izmir, Turkey. *Hanseniaspora valbyensis* strain LU 263 (LU codes refer to the collection maintained at the laboratory of food microbiology, Agricultural University, Wageningen, the Netherlands), *Endomyces fibuliger* strain LU 677, *Mucor circinelloides* strain LU M40 and *Penicillium nalgiovense* strain LU 520 originate from fermented foods. All were grown and maintained on Malt Extract Agar (Oxoid CM 59) slants. Incubation was at 28°C for 3–5 days, and grown cultures were stored at 5°C. *Bacillus* sp. strains B4 and B6 had been isolated from apricot seeds previously (Brimer et al., 1993). They were grown (30°C for 2–3 days) and maintained (5°C) on Plate Count Agar (Oxoid CM325) slants.

2.2. Reagents.

D-Amygdalin (Sigma St.Louis USA, A6005) was used for standard as a 1 mM solution in distilled water. Pectinase (EC 3.2.1.15) from *Rhizopus* sp. (Sigma St.Louis USA, P2401) was used as a source of hydrolytic enzymes (Brimer and Rosling, 1993) for degradation of amygdalin standards, as a 0.2% w/v solution in distilled water in the quantification of total cyanide potential. Picrate reagent sheets were prepared according to Brimer et al. (1983).

2.3. Methods.

Extraction of amygdalin and degradation products: 10 g sample (seed dry weight) was homogenized with 90 ml 0.1 M ortho-phosphoric acid (Cooke, 1978) in

a glass blender jar (Braun Multimix MX32 type 4207 Germany) as follows: 15 s speed 1, 1 min speed 3, 1 min rest, 1 min speed 3. Homogenates (A) were found stable for 7 days at 4°C as also found for cassava extracts by Cooke (1978).

Amygdalin degradation. To 5 ml homogenate (A), 15 ml phosphate buffer (0.1 M $\text{Na}_3\text{PO}_4/\text{H}_3\text{PO}_4$, pH 7) is added to give homogenate (B) with pH 6.5. In a microtiter plate, the following wells are prepared: (a) amygdalin standard (6 wells/plate): 10, 20 or 30 μL amygdalin + 100 μL pectinase + distilled water totalling 200 μL ; (b) concentrated sample for degradation products (HCN and cyanohydrins) (2 wells/plate): 100 μL sample homogenate (B) + 100 μL dist. water; (c) diluted sample for degradation products (2 wells/plate): 10 μL sample homogenate (B) + 190 μL dist. water; (d) concentrated sample for total cyanide potential (intact amygdalin + degradation products) (2 wells/plate): 100 μL sample homogenate (B) + 100 μL pectinase; (e) diluted sample for total cyanide potential (2 wells/plate): 10 μL sample homogenate (B) + 90 μL dist. water + 100 μL pectinase. A picrate reagent sheet (plastic side up) is placed on top of the microtiter plate and weighed down gently with a flat object of approx. 100 g, and incubated at 25°C overnight. In this way, one microtiter plate accommodates a 3-level amygdalin calibration and three different samples.

Assessment. The density of the red-brown spots on yellow background typical of positive reactions, was assessed by measuring the absorption of transmitted light at 550 nm using a microtiter plate reader (EAR 400, SLT-Labinstruments, A-5082 Groedig, Austria) (Brimer et al., 1993). Using the amygdalin standards for calibration in each picrate sheet, extinctions of either dilute or concentrated sample which were closest to the standard, were used for calculations. Degradation products labeled as 'cyanohydrins' were based upon wells type (b) or (c), and the difference of total cyanide potential (wells type 'd' or 'e') and degradation products was labeled as 'amygdalin'. Both were expressed as $\mu\text{mol CN/g}$ sample dry weight. The detection level was 1 $\mu\text{mol HCN/g}$ dry weight.

Seed processing. Apricot seeds were rinsed 3 times with tapwater, and ground coarsely in a Fritsch mill (Fritsch Pulverisette type 14.702, Germany) (fine rotor, no sieve, speed 1). Particle size distribution of the grits was: 24% < 1.0 mm; 20% 1.0–2.0 mm; 36% 2.0–4.8 mm; 20% > 4.8 mm. The ground seeds were soaked overnight at 22°C in tapwater at a rate of seeds: liquid = 1:3 w/v. Subsequently, the material was recovered on a sieve and washed twice with tapwater, and either (a) cooked in fresh tapwater (seeds: water = 1:3 w/v) at 100°C for 30 min with open pan or (b) dispensed without further addition of water in screw-capped glass jars and autoclaved at 121°C for 30 min. After cooling, the autoclaved material was inoculated with the microbial strains tested at a rate of approximately 10^5 – 10^6 cfu/g, and incubated with loose-fitting screw-caps in order to avoid anaerobiosis.

Microbial enumeration. The growth of bacteria was monitored using pour-plates with Plate Count Agar (Oxoid CM325) incubated at 30°C for 3 days; yeasts were

counted after 3–5 d at 25°C in pour-plates of Yeast Extract Glucose Agar (Oxoid CM545) to which 0.01% (w/v) sterile oxytetracycline (Oxoid SR73) was added after autoclaving; and filamentous fungi were enumerated using surface-inoculated plates with Rose Bengal Agar modified according to Baggerman (1981), incubated at 28°C for 5 days.

3. Results and discussion

Table 1 shows the effect of preparatory treatments (grinding, soaking, heating) on cyanide compounds in bitter apricot seeds. Although the initial cyanide content of the present batch of seeds is much higher, these data show a similar trend as observed earlier (Tunçel et al., 1990), indicating that endogenous β -glycosidase activity causes significant degradation of amygdalin in soaked ground seeds. The resulting cyanohydrins volatilize during cooking, whereas the remaining amygdalin is practically unaffected. Autoclaved soaked seeds still contained detectable cyanohydrins, probably due to restricted volatilization in the closed autoclave vessel. When non-inoculated cooked (100°C, 30 min) seeds were kept in jars with loose-fitted screw-caps at 4°C for 48 h, no microbial growth occurred and the cyanide content did not change. However, when kept under similar conditions but incubated at 30°C for 48 h, profuse bacterial growth occurred ($\text{Log}_{10} \text{N/g} = 9.6$ on plate count agar) and cyanide was removed till below detection level ($1 \mu\text{mol/g dm}$) (data not shown). In similar investigations carried out previously, six out of 10 tested purified isolates all belonging to *Bacillus* sp. were able to degrade amygdalin in aqueous solution (Brimer et al., 1993). In these investigations, strains B4 and B6 resembling *B. subtilis* degraded 85% and 100%, respectively, of amygdalin overnight. These strains were included in the detoxification trials. When non-inoculated autoclaved (121°C, 30 min) seeds (Table 1) were incubated at 27°C for 72 h, no microbial growth occurred and the cyanide content remained constant. The latter treatment was therefore chosen to prepare seeds for detoxification trials with microbial pure cultures.

In Table 2, the effect of selected microbial cultures on cyanide levels in prepared bitter apricot seeds is shown. Not all cultures which can degrade

Table 1
Effect of pre-fermentation treatments on cyanide content of bitter apricot seeds.

	Cyanohydrins ^a	Amygdalin ^a
Raw seeds	6 (0.5) ^b	118 (29)
Ground, soaked overnight	24 (4)	30 (2)
Soaked + Cooked in open pan (100°C, 30 min)	ND	28 (2)
Soaked + Autoclaved (121°C, 30 min)	9 (1)	32 (3)

^a $\mu\text{mol HCN equivalents/g dry weight}$; ^b Duplicates from two independent batches of seeds (mean and standard deviations; $n = 4$); ND, not detected ($< 1 \mu\text{mol HCN equivalents/g dry weight}$).

Table 2
Effect of aerobic incubation at 27°C with selected microorganisms on cyanide content of soaked and autoclaved bitter apricot seeds.

Strain Code		Incubation period (h)			
		0	24	48	72
<i>Bacillus</i> sp.					
LU B4	CFU ^a	6.86	7.49	8.82	9.65
	Cyanohydrins ^b	9	5	2	9
	Amygdalin ^b	32	39	13	10
<i>Bacillus</i> sp.					
LU B6	CFU	6.81	7.48	9.05	9.90
	Cyanohydrins	9	4	2	11
	Amygdalin	32	30	11	7
<i>Hanseniaspora valbyensis</i>					
LU 263	CFU	5.45	5.40	6.21	6.36
	Cyanohydrins	9	8	6	10
	Amygdalin	31	30	30	28
<i>Mucor circinelloides</i>					
LU M40	CFU	5.10	4.84	4.67	4.42
	Cyanohydrins	10	6	9	7
	Amygdalin	32	32	28	27
<i>Penicillium nalgiovense</i>					
LU 520	CFU	4.94	4.20	4.12	4.40
	Cyanohydrins	9	10	8	4
	Amygdalin	30	31	29	26
<i>Endomyces fibuliger</i>					
LU 677	CFU	5.52	5.54	7.64	6.90
	Cyanohydrins	9	21	13	9
	Amygdalin	32	20	2	ND ^c
None (control)					
	CFU	< 1.7			< 1.7
	Cyanohydrins	9			1
	Amygdalin	30			31

^a Colony Forming Units expressed as Log₁₀N/g fresh sample; ^b μmol HCN equivalents/g dry weight;

^c ND, not detected (< 1 μmol HCN equivalents/g dry weight).

amygdalin in aqueous solution in a screening test (Brimer et al., 1993) can do so in-situ. The unsatisfactory results with *H.valbyensis*, *M.circinelloides* and *P.nalgiovense* are due to their poor growth on the substrate. Agar diffusion and auxanogram tests (data not shown) revealed that this was neither caused by water-soluble inhibitory compounds in the seeds, nor by C or N deficiency. Other possible reasons (e.g. poorly soluble phenolic compounds with antimicrobial effect) were not investigated. On the other hand, *Bacillus* sp. B4 and B6 grew well and caused significant removal of cyanide. However, detectable residues remained. The best results were obtained with *Endomyces fibuliger* LU 677 which showed good growth and resulted in complete removal of cyanide after 3 days of incubation. This yeast was isolated from Indonesian 'Ragi', a traditional inoculum for the fermentation of glutinous rice into the product 'Tape Ketan'. *E.fibuliger* is

considered as functional flora during this fermentation and contributes to the aroma of the fermented rice (Hesseltine et al., 1988). Further investigation of the effect of *E.fibuliger* on the chemical and organoleptic properties of bitter apricot seeds will be undertaken.

Acknowledgement

The financial support by NATO Collaborative Research Grant 920009 is gratefully acknowledged.

References

- Baggerman, W.I. (1981) A modified rose bengal medium for the enumeration of yeasts and moulds from foods. *Eur. J. Appl. Microbiol. Biotechnol.* 12, 242–247.
- Brimer, L., Brøgger Christensen, S., Mølgaard, P. and Nartey, F. (1983) Determination of cyanogenic compounds by thin-layer chromatography. 1. A densitometric method for quantification of cyanogenic glycosides, employing enzyme preparations (β -glucuronidase) from *Helix pomatia* and picrate-impregnated ion-exchange sheets. *J. Agric. Food Chem.* 31, 789–793.
- Brimer, L. and Mølgaard, P. (1986) Simple densitometric method for estimation of cyanogenic glycosides and cyanohydrins under field conditions. *Biochem. System. Ecol.* 14, 97–103.
- Brimer, L. and Rosling, H. (1993) Microdiffusion method with solid state detection of cyanogenic glycosides from cassava in human urine. *Food Chem. Toxicol.* 31, 599–603.
- Brimer, L., Tunçel, G. and Nout, M.J.R. (1993) Simple screening procedure for microorganisms to degrade amygdalin. *Biotechnology Techniques* 7, 683–687.
- Cooke, R.D. (1978) An enzymatic assay for the total cyanide content of cassava (*Manihot esculenta* Crantz). *J. Sci. Food Agric.* 29, 345–352.
- Guignard, M.L. (1906) Le Haricot à acide cyanhydrique, *Phaseolus lunatus* L. C. R. Hebd. Seances Acad. Sci. 142, 545–553.
- Hesseltine, C. W., Rogers, R. and Winarno, F.G. (1988) Microbiological studies on amylolytic oriental fermentation starters. *Mycopathology* 101, 141–155.
- Tunçel, G., Nout, M.J.R., Brimer, L. and Gökten, D. (1990) Toxicological, nutritional and microbiological evaluation of tempe fermentation with *Rhizopus oligosporus* of bitter and sweet apricot seeds. *Int. J. Food Microbiol.* 11, 337–344.