

LiGAPS-Beef, a mechanistic model to explore potential and feed-limited beef production 1: model description and illustration

van der Linden, A., van de Ven, G. W. J., Oosting, S. J., van Ittersum, M. K., & de Boer, I. J. M.

This is a "Post-Print" accepted manuscript, which has been published in "Animal"

This version is distributed under a non-commercial no derivatives Creative Commons  (CC-BY-NC-ND) user license, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited and not used for commercial purposes. Further, the restriction applies that if you remix, transform, or build upon the material, you may not distribute the modified material.

Please cite this publication as follows:

van der Linden, A., van de Ven, G. W. J., Oosting, S. J., van Ittersum, M. K., & de Boer, I. J. M. (2018). LiGAPS-Beef, a mechanistic model to explore potential and feed-limited beef production 1: model description and illustration. Animal, 1-11. DOI: 10.1017/S1751731118001726

You can download the published version at:

https://doi.org/10.1017/S1751731118001726

LiGAPS-Beef, a mechanistic model to explore potential and feed-limited beef production 2. Sensitivity analysis and evaluation of sub-models

A. van der Linden[1,2], G.W.J. van de Ven[2], S.J. Oosting[1], M.K. van Ittersum[2], and I.J.M. de Boer[1]

[1] *Animal Production Systems group, Wageningen University & Research, P.O. Box 338, 6700 AH Wageningen, The Netherlands*

[2] *Plant Production Systems group, Wageningen University & Research, P.O. Box 430, 6700 AK Wageningen, The Netherlands*

Corresponding author: Aart van der Linden. Email: aart.vanderlinden@wur.nl

Short title: LiGAPS-Beef 2. Sensitivity analysis and evaluation

Abstract

The model **LiGAPS-Beef** (Livestock simulator for Generic analysis of Animal Production Systems – Beef cattle) has been developed to assess potential and feed-limited growth and production of beef cattle in different areas of the world and to identify the processes responsible for the yield gap. Sensitivity analysis and evaluation of model results with experimental data are important steps after model development. The first aim of this paper, therefore, is to identify which parameters affect the output of LiGAPS-Beef most by conducting sensitivity analyses. The second aim is to evaluate the accuracy of the thermoregulation sub-model and the feed intake and digestion sub-model with experimental data. Sensitivity analysis was conducted using a one-at-a-time approach. The upper critical temperature (**UCT**) simulated with the thermoregulation sub-model was most affected by the body core temperature and parameters affecting latent heat release from the skin. The lower critical temperature (**LCT**) and UCT were considerably affected by weather variables, especially ambient temperature and wind speed. Sensitivity analysis for the feed intake and digestion sub-model showed that the digested protein per kg feed intake was affected to a larger extent than the metabolisable energy (**ME**) content. Sensitivity analysis for LiGAPS-Beef was conducted for ¾ Brahman × ¼ Shorthorn (**B×S**) cattle in Australia and Hereford cattle in Uruguay. Body core temperature, conversion of digestible energy (**DE**) to ME, net energy (**NE**) requirements for maintenance, and several parameters associated with heat release affected feed efficiency at the herd level most. Sensitivity analyses have contributed, therefore, to insight which parameters are to be investigated in more detail when applying LiGAPS-Beef. Model evaluation was conducted by comparing model simulations with independent data from experiments. Measured heat production in experiments corresponded fairly well to the heat

production simulated with the thermoregulation sub-model. Measured ME contents from two datasets corresponded well to the ME contents simulated with the feed intake and digestion sub-model. The relative mean absolute errors (**MAEs**) were 9.3% and 6.4% of the measured ME contents for the two datasets. In conclusion, model evaluation indicates the thermoregulation sub-model can deal with a wide range of weather conditions, and the feed intake and digestion sub-model with a variety of feeds, which corresponds to the aim of LiGAPS-Beef to simulate cattle in different beef production systems across the world.

Keywords: beef cattle, mechanistic modelling, production ecology, sensitivity analysis, yield gap

Implications

A generic model for beef cattle, named LiGAPS-Beef, has been described and illustrated in a companion paper (Van der Linden *et al.*, 2018a). This mechanistic model aims to assess the potential (*i.e.* theoretical maximum) and feed-limited growth and production of cattle in different beef production systems across the world. In this paper, we conducted sensitivity analyses and evaluated parts of LiGAPS-Beef with independent experimental data. Our results contribute to the evidence that LiGAPS-Beef can be used to simulate a broad range of beef production in systems with different climates and feeding strategies.

Introduction

The increasing demand for animal-source food calls for insight to what extent livestock production can be increased in different parts of the world. The biophysical scope to increase livestock production is the difference between the potential (*i.e.* maximum theoretical) production or feed-limited production and the actual production realized in

practice, which is also referred to as the yield gap (Van de Ven *et al.*, 2003, Van der Linden *et al.*, 2015). Identifying geographical regions with large yield gaps contributes to insight where food production can be increased per unit of land, which is generally regarded as a better strategy than expanding agricultural land at the expense of nature (Lobell *et al.*, 2009, Van Ittersum *et al.*, 2013).

Yield gaps of arable crops are widely assessed with mechanistic crop growth models, which simulate potential and water-limited production in different farming systems and in different regions of the world (Jones *et al.*, 2003, Keating *et al.*, 2003). Yield gaps of livestock have not been assessed with mechanistic models yet, since models simulating potential and feed-limited livestock production were hardly available at the start of this research. A generic, mechanistic model was developed, therefore, to assess potential and feed-limited beef production in different beef production systems and in different regions of the world (Van der Linden *et al.*, 2018a). This model is named **LiGAPS-Beef** (Livestock simulator for Generic analysis of Animal Production Systems – Beef cattle), and its results may eventually contribute to the identification of regions with a large biophysical scope to increase beef production.

Mechanistic models include the most important processes and mechanisms in systems, but still consist of multiple empirical elements and parameters that can considerably affect model output, and subsequently the conclusions based on the models' output (Thornley and France, 2007). Sensitivity analysis provides insight in how model output is affected by changes in model input. This method ranks input parameters based on their effect on model output (Pianosi *et al.*, 2016). Ranking parameters can be used to prioritize which parameters need to be estimated more precisely (Zuidema *et al.*, 2005). Sensitivity analysis is of particular importance if models are applied outside conditions they were calibrated for (Prisley and Mortimer,

2004). Since LiGAPS-Beef is designed to be applicable to a broad range of beef production systems, conducting sensitivity analysis is essential.

Furthermore, key processes in the model must be simulated in sufficient detail to ensure applicability of the model under a wide range of agro-ecological conditions and beef production systems. If key processes are simulated in sufficient detail, model output must resemble experimental data. Hence, model evaluation with experimental data is an essential and necessary step after model development to investigate whether model output is accurate. Model evaluation is conducted with experimental data not used for model calibration, so experimental data for model calibration and evaluation are independent (Bellocchi *et al.*, 2010). Model evaluation with independent experimental data is also referred to as model validation or testing, but we will use the term model evaluation consistently throughout this paper. Given the relevance of sensitivity analysis and model evaluation, the first aim of this paper is to assess which model parameters affect model output most. The second aim is to evaluate the performance of LiGAPS' sub-models on thermoregulation and feed intake and digestion with independent experimental data. The performance of the complete model LiGAPS-Beef in different beef production systems is evaluated in a companion paper (Van der Linden *et al.*, 2018b).

Materials and methods

Structure of LiGAPS-Beef

LiGAPS-Beef consists of a thermoregulation sub-model, a feed intake and digestion sub-model, and an energy and protein utilisation sub-model (Van der Linden *et al.*, 2018a) (Fig. 1). The thermoregulation sub-model simulates heat release, based on existing thermoregulation models (McGovern and Bruce, 2000, Turnpenny *et al.*,

2000a). This sub-model requires daily weather data if cattle are kept outdoors and climate conditions in stables if cattle are housed (Fig. 1). Genetic parameters and heat production from metabolic processes are inputs too. Minimum and maximum heat release are outputs of this sub-model. Cold conditions can increase feed intake, whereas hot conditions can decrease feed intake. The thermoregulation sub-model increases energy requirements under hot conditions, because energy is spent on panting (Fig. 1). Inputs for the feed intake and digestion sub-model are the energy requirements of cattle and the quality and quantity of the available feeds (Fig. 1). Feed intake is an output of this sub-model. Feed digestion is simulated based on a rumen model of Chilibroste *et al.* (1997), and yields metabolisable energy (**ME**) and digested protein as major outputs, which are used as input for the energy and protein utilisation sub-model. Energy and protein are distributed over the metabolic processes maintenance, physical activity, growth, gestation, and lactation (Van der Linden *et al.*, 2018a). Energy and protein for growth are allocated to different tissues (non-carcass tissue, and bone, muscle and fat tissue in the carcass). Beef is defined as deboned carcass. Feed efficiency of individual animals (**FE**, g beef kg^{-1} DM feed) is calculated from their beef production and feed intake (Fig. 1). Results for individual animals can be scaled up to the herd level.

Sensitivity analysis

Thermoregulation sub-model. Sensitivity analysis was used to assess the effect of changing parameters and weather data on the lower critical temperature (**LCT**) and upper critical temperature (**UCT**) simulated by the thermoregulation sub-model. In total, 31 parameters were investigated (23 cattle-specific; 8 breed-specific). These parameters were decreased and increased by 10%, while all other parameters were kept at their original values, according to the one-at-a-time approach (Pianosi *et al.*,

2016). Two exceptions were the body temperature (which was changed by 1 ºC, or 2.6%) and the standard temperature used in the formula to calculate the latent heat release from the skin (which was changed by 1%), because changing these two parameters by 10% caused excessive heat stress, which resulted in a reduction of feed intake, and eventually a complete depletion of the body fat reserves. We concluded, therefore, that changes of 10% were beyond the feasible biological range. Chemical and physical parameters were not included in the sensitivity analysis, since they were considered constants. In the baseline scenario for sensitivity analysis, solar radiation was set at 10 MJ m^{-2} coat day^{-1}, relative humidity at 50%, wind speed at 4 ms^{-1}, precipitation at 0 mm day^{-1}, and cloud cover at 4 Ω. The total body weight (**TBW**) was 450 kg in the baseline scenario, and heat production was 1.36 times maintenance heat production for *B. taurus* cattle, which corresponds to a situation where approximately half of the ME is allocated to maintenance, and half to growth. In addition, we investigated the LCT and UCT within a range of temperatures (-40°C to 40°C) combined with a range of solar radiation levels (0-30 MJ m^{-2} day^{-1}), relative humidity levels (10-100%), wind speeds (0.1-8.0 m s^{-1}), precipitation levels (0-30 mm day^{-1}), and cloud cover levels (0-8 Ω). In addition, the range of temperatures was combined with a range of TBWs (50-1300 kg), and heat production levels (1.0-2.0 × maintenance heat production).

Feed intake and digestion sub-model. Feed intake is dependent on the genotype of the animal, the climate, feed quality, and the available feed quantity, and is, therefore, an output of the joint sub-models of LiGAPS-Beef (Fig. 1). Feed digestion can be investigated with the feed intake and digestion sub-model only. The output of this sub-model is the ME content (MJ kg^{-1} DM) and digestible protein content (g kg^{-1} DM) of particular feeds and diets, using feed constituents as model inputs. Feed constituents

159 investigated were soluble, non-structural carbohydrates (**SNSC**), insoluble, non-
160 structural carbohydrates (**INSC**), digestible NDF (**DNDF**), soluble crude protein (**SCP**),
161 digestible crude protein (**DCP**) and total CP (Chilibroste *et al.*, 1997). In addition,
162 digestion (3×) and passage rates (2×) were included, as well as the slope and intercept
163 of a Lucas equation (Eq. 1) (Lucas *et al.*, 1961, Van Soest, 1994). Feed constituents
164 of thirteen feeds were decreased by 10% to investigate the effect on ME and digestible
165 protein content using the one-at-a-time approach.

166 Eq. 1 Digestible protein (g kg^{-1} DM) = 0.9 × CP (g kg^{-1} DM) - 32

167 *LiGAPS-Beef.* Sensitivity analysis was conducted to assess the effect of changing
168 parameters on FE at the herd level. Sensitivity analysis (one-at-a-time approach) was
169 conducted for all parameters of LiGAPS-Beef, including the 31 parameters from the
170 thermoregulation sub-model, and the slope and intercept of the Lucas equation (Eq.
171 1). Parameters were decreased and increased by both 5% and 10%. The arbitrary
172 changes of 5% and 10% were chosen because the standard deviations of parameters
173 or their expected range are unknown for most parameters. The disadvantage of this
174 approach is that the decrease or increase of parameters can be outside their
175 biologically feasible range, and consequently no meaningful model output is obtained.
176 Three parameters were changed by less than 5%, since biological limits did not allow
177 a change of 5% and 10%. The standard temperature used in the formula to calculate
178 the latent heat release from the skin and a parameter to calculate body area were
179 changed by 1%, and the body core temperature was changed by 0.1°C. Parameters
180 of the Gompertz curve were changed together because they are interrelated, except
181 for the rate constant. The sensitivity of model output was represented by the sensitivity
182 coefficient, which is the ratio of change in model output to the change in the parameter
183 value (Hamby, 1994).

Sensitivity analysis was conducted at the herd level for ¾ Brahman × ¼ Shorthorn (**B×S**) cattle, adapted to a tropical climate, and for Hereford cattle, adapted to a temperate climate. Four hypothetical baseline scenarios were used for the sensitivity analysis: B×S cattle in Australia under potential production; B×S cattle in Australia under feed quality limited production; Hereford cattle in Uruguay under potential production; and Hereford cattle in Uruguay under feed quality limited production. Under potential production, cattle were permanently housed, and the diet consisted of wheat (65%) and good quality hay (35%). Under feed quality limitation, the ME content of the diet was set at 11.1-12.2 MJ kg^{-1} DM in Australia, and 10.7-11.8 MJ kg-1 DM in Uruguay. Weather data used were from the year 1992 in Australia and 2002 in Uruguay. Weaning age was set at 210 days in both countries. The culling rate for a cohort of cows after birth of the first calf was set at 50% per year (Van der Linden *et al.*, 2015, Van der Linden *et al.*, 2018a). As cows were assumed to conceive up to an age of ten years, each cow gives, on average, birth to two calves. The female calf is used as a replacement for the reproductive cow and is not part of the herd unit, but gives rise to the next one (Van der Linden *et al.*, 2015, Van der Linden *et al.*, 2018a). Hence, one herd unit consists of a reproductive cow and one male calf. Slaughter weights of male B×S and Hereford calves were optimized to maximize FE at the herd level (Van der Linden *et al.*, 2018a).

Evaluation of sub-models

The thermoregulation sub-model and the feed intake and digestion sub-model were each evaluated with independent experimental data. The energy and protein utilisation sub-model is the largest and central sub-model, and it requires a significant amount of inputs from the thermoregulation and feed intake and digestion sub-model (Fig. 1). For this reason, evaluation of the energy and protein utilisation sub-model was not

209 conducted in this paper. Evaluation of this large sub-model is, however, included
210 indirectly in the evaluation of LiGAPS-Beef as a whole, which is reported in a
211 companion paper (Van der Linden *et al.*, 2018b).

212 *Thermoregulation sub-model.* The thermoregulation sub-model was calibrated, since
213 its daily time step was much coarser than the time step used in the thermoregulation
214 models of McGovern and Bruce (2000) and Turnpenny *et al.* (2000a). Model
215 simulations included an animal of 450 kg TBW kept outdoors. Solar radiation levels
216 were set at 15 MJ m^{-2} day^{-1} (horizontal surface), which was assumed to correspond to
217 7.5 MJ m^{-2} coat day^{-1}. Cloud cover was set at 4 Ω, and the level of precipitation at 0
218 mm day^{-1}. Parameters for respiration and latent heat release from the skin were
219 adjusted to fit to temperature-humidity indices (Eqs 2 and 3) (Mader *et al.*, 2006).

220 Eq. 2 $THI = 0.8 \times T + RH / 100 \times (T - 14.4) + 46.4$

221 Eq. 3 $THI_{adj.} = THI + 4.51 - 1.992 \times WS + 0.0068 \times SR$

222 Where THI is the temperature-humidity index, T is the temperature (ºC), RH is the
223 relative humidity (%), $THI_{adj.}$ is the temperature-humidity index adjusted for wind speed
224 and solar radiation, WS is wind speed (m s^{-1}), and SR is the level of solar radiation (W
225 m^{-2}). Threshold values for THI and $THI_{adj.}$ were adopted from Mader *et al.* (2006).
226 After calibration, simulated heat release was compared with measured heat release
227 from two experiments, which were also used to calibrate the model of Turnpenny *et al.*
228 (2000a). In the first experiment, heat release of Aberdeen Angus × Shorthorn steers
229 (323-361 kg TBW) was measured at low temperatures (-1.1 to 3.1°C), with low (<7
230 mm) and high coat lengths (>24 mm) (Blaxter and Wainman, 1964). In the second
231 experiment, heat release of Friesian calves (initial TBW 34.6 kg) and Jersey calves
232 (initial TBW 27.8 kg) was measured for a range of temperatures (3-20°C) and two wind

speeds (0.22 and 1.56 m s^{-1}) (Holmes and McLean, 1975). Coat length was not measured in this experiment, but it was assumed to be 25 mm. In both experiments, animals were expected to be below their LCT in most of the experimental treatments, and hence their measured heat release should correspond to the minimum heat release simulated with the thermoregulation sub-model.

Feed intake and digestion sub-model. We used the seven feed constituents and their digestion and passage rates specified by Chilibroste *et al.* (1997) as input to simulate the ME content of 13 feed types (MJ kg^{-1} DM). Simulated ME contents were compared with measured ME contents from MAFF (1986) and Kolver (2000). The mean absolute error (**MAE**) (Eq. 4), mean square error (**MSE**), and the RMSE (Eq. 5) reflect the deviation of simulated ME contents from the measured ME contents. The MSE was decomposed into the bias, slope, and random component (Bibby and Toutenburg, 1977). The bias component indicates systematic errors in the model, and the slope component indicates the models' ability to replicate the variability in the measured data. The random component is the remaining variation after accounting for the bias and slope components (Bibby and Toutenburg, 1977). A perfect fit of the regression line between simulated and measured data means that the bias and slope components explain 0% of the MSE, and the random component 100% (Bellocchi *et al.*, 2010).

Eq. 4 $$MAE = \frac{\Sigma \mid O - S \mid}{n}$$

Eq. 5 $$RMSE = \sqrt{\frac{\Sigma (O - S)^2}{n}}$$

Where O is the observed value, S is the simulated value, and n is the number of observations. The measured and simulated digested protein were not compared to

each other, because the CP content of feeds given in Chilibroste *et al.* (1997) was often different from the CP content given in MAFF (1986).

Results

Sensitivity analysis

Thermoregulation sub-model. The LCT was affected by more than 1.0°C for parameters used to calculate the body area and the minimum conduction between body core and skin (3 parameters) (Table 1). The UCT was affected by more than 1.0°C for parameters used to calculate the body area, body temperature, exhaled air temperature, maximum conduction between body core and skin, and latent heat release from the skin (2 parameters) (Table 1). The LCTs and UCTs decreased with increasing solar radiation, relative humidity, TBW, and heat production, whereas they increased with increasing wind speed and precipitation (Fig. 2). The ranges used for wind speed, TBW, and heat production resulted in considerable shifts in the LCTs and UCTs (10ºC or more for the LCT). Changes in relative humidity mainly affected the UCT, and hardly the LCT (Fig. 2). The shifts in LCT and UCT within the ranges specified were generally larger than the changes in LCT and UCT after changing parameters by 10% (Fig. 2, Table 1).

Feed intake and digestion sub-model. Reducing the content of SNSC, INSC, DNDF, SCP, DCP, and total CP by 10% resulted in a lower ME and digestible protein content for all feed types (Table 2). The ME content increased upon a 10% reduction in the passage rate in the rumen, the passage rate for DNDF, and the intercept of the Lucas equation (Eq. 1). The SNSC content affected the ME content of molasses (-10.4%), wheat (-5.3%), barley (-4.4%), and concentrates (-3.2%) most (Table 2). The DNDF content affected the ME content of cereal straw (-6.9%), hay (up to -5.9%), and grass

(up to -5.5%) most. Decreasing the slope of the Lucas equation had the same effect on the amount of digestible protein as decreasing the total CP content of the feed (Table 2). The digestible protein content of all feeds was negatively affected by a decrease in the slope of the Lucas equation, and positively by a decrease in its intercept. For molasses, the amount of protein digested in the baseline was negative, because the Lucas equation is negative at low levels of CP (4 g kg^{-1} DM for molasses). Its intercept and slope affected the digestible protein content of feeds with low CP contents (+80% and -90% for cereal straw) to a larger extent than feeds with high CP contents (+1% and -11% for soybean meal) (Table 2).

LiGAPS-Beef. For the baseline scenario, the FE of B×S cattle in Australia was 77.0 g beef kg^{-1} DM (65% wheat, 35% good quality hay) under potential production, and 40.8 g beef kg^{-1} DM (pasture) under feed quality limited production. The FE of Hereford cattle in Uruguay was 71.4 g beef kg^{-1} DM under potential production, and 37.1 g beef kg^{-1} DM under feed quality limited production. Changing parameter values by 5% or 10% hardly affected the FE at the herd level for most of the parameters under potential production. The sensitivity coefficient was only higher than one for the body temperature of B×S cattle in Australia, whereas sensitivity coefficients were below one for Hereford cattle in Uruguay (Table 3). Six parameters in the top ten parameters affecting model output most were found both in Australia and Uruguay under potential production. The net energy (**NE**) for maintenance and its multiplier were in the top ten parameters for each of the four scenarios. Sensitivity coefficients were higher under feed quality limited production than under potential production. Changing parameters in the top ten by 10% often did not result in meaningful output under feed quality limited production, due to simulated heat stress, the consequent reductions in feed intake, depletion of body fat reserves, and eventually mortality (Table 3).

Parameters related to heat release were listed more often in the top ten under feed quality limited production than under potential production. Latent heat release (Australia only), standard respiration rate (Australia only), maximum conduction between body core and skin, and the temperature of exhaled air (Uruguay only) were found in the top ten under feed quality limited production, but not under potential production (Table 3). Sensitivity coefficients were similar for changes of 5% and 10% in parameters under potential production in Australia, which suggests rather linear relations between parameters values and model output. The same holds for Uruguay, except for the adult weight used in the Gompertz curve, where sensitivity coefficients differ for a 5% change and a 10% change (Table 3).

Evaluation of sub-models

Thermoregulation sub-model. After calibration, the climate conditions resulting in heat stress in the thermoregulation sub-model corresponded to the climate conditions classified as alert, danger, and emergency by the temperature-humidity indices (Eqs 2 and 3) (Fig. 3). Measured heat release and simulated minimum heat release for the experiment of Blaxter and Wainman (1964) were in agreement for steers with high coat lengths, but simulations underestimated the minimum heat release for steers with low coat lengths (Fig. 4A). Measured heat release and simulated minimum heat release of Friesian and Jersey calves for the experiment of Holmes and McLean (1975) corresponded to each other at a heat release of approximately 90 W m^{-2} and higher (Fig. 4A). Treatments at 20°C and at 12°C with a wind speed of 0.22 m s^{-1} resulted in a heat release below 90 W m^{-2}. Latent and sensible heat release for the experiment of Blaxter and Wainman (1964) were simulated well for steers with high coat lengths, whereas sensible heat release was underestimated for steers with low coat lengths (Fig. 4B). Simulated and measured skin temperatures for the steers were assessed

329 reasonably well by the thermoregulation sub-model (Fig. 4C). Skin temperature was
330 underestimated considerably for one animal having low coat lengths (measured 23.7
331 and 22.0°C; simulated 16.5 and 15.4°C).

332 *Feed intake and digestion sub-model.* Simulated and measured ME contents of MAFF
333 (1986) generally corresponded to each other (RMSE = 1.28 MJ ME kg^{-1} DM, MSE =
334 1.64 MJ^2 ME kg^{-2} DM). The MAE was 1.06 MJ ME kg^{-1} DM, or 9.3% of the average
335 measured ME content. The intercept of the regression line was not significantly
336 different from zero (P = 0.79) and its slope was not significantly different from one (P
337 = 0.09). The bias component accounted for the largest part of the MSE (68.3%). The
338 slope component was 0.3% of MSE, and the random component was 31.4%.
339 Simulated and measured ME contents of Kolver (2000) generally corresponded also
340 to each other (RMSE = 0.87 MJ ME kg^{-1} DM, MSE = 0.76 MJ^2 ME kg^{-2} DM). The MAE
341 was 0.69 MJ ME kg^{-1} DM, or 6.4% of the measured ME content. The intercept of the
342 regression line was not significantly different from zero (P = 0.38) and its slope was
343 not significantly different from one (P = 0.38) (Fig. 5). The random component
344 accounted for the largest part of the MSE (56.1%). The bias component was 43.3% of
345 the MSE, and the slope component was 0.6%. The average difference in ME content
346 of the same feeds in the data of MAFF (1986) and Kolver (2000) was 0.58 MJ ME kg^{-1}
347 DM, or 5.3% of the mean measured ME content in MAFF (1986).

348 **Discussion**

349 *Sensitivity analysis*

350 *Thermoregulation sub-model.* The identification of parameters affecting the simulated
351 LCT and UCT prioritizes the parameters to be investigated in more detail. Such an
352 investigation may increase the accuracy of the sub-model further. Priority should be

given also to parameters with a large variability. For example, the maximum conduction between body core and skin was assumed to be constant for beef cattle, but the parameter value was 67% higher for dairy cattle than for beef cattle (Turnpenny *et al.*, 2000b). This suggests a considerable variability in parameter values among different cattle breeds. Hence, the LCT and UCT may be affected even more if the actual variability is larger than the 10% simulated. An opposite example is a parameter for calculating the body area (Table 1). The body area of a 400 kg animal decreases by 41% upon a 10% decrease in one parameter used to calculate body area from TBW (Thompson *et al.*, 2011). In comparison, the body area of *B. indicus* cattle is approximately 10% larger than for *B. taurus* cattle at the same weight (NRC, 2000). The effect of this particular parameter on LCT and UCT is, therefore, likely to be lower than with the 10% change simulated. Hence, investigating the ranges or standard deviations of parameters is important also to prioritize which parameters to measure more precisely or to investigate in more detail.

Changing weather variables in the ranges specified generally affected the LCT and UCT to a larger extent than changing parameter values by 10% (Fig. 2, Table 1). These results highlight the need for accurate weather data as input for the thermoregulation sub-model. Effects of weather variables on the LCT and UCT were in line with expectations. An increasing wind speed and precipitation increased heat release and hence increased both the LCT and UCT, whereas the reverse is true for increasing levels of solar radiation. Precipitation affected the simulated LCT and UCT by evaporation of water from the coat and an increase in heat conduction of the coat layer. Changes in relative humidity affected the UCT, but hardly the LCT (Fig. 2). This is explained by the latent heat release from the skin, which is maximized under hot conditions and minimized under cold conditions. Increasing TBW decreased the LCT

and UCT, which is mainly explained by a corresponding decrease in the ratio coat area to TBW. The range in TBW (50-1300 kg) and heat production (1.0-2.0 × maintenance) affected the LCT and UCT considerably (Fig. 2). Hence, heat production and TBW are important inputs for the thermoregulation sub-model that have to be simulated accurately.

Feed intake and digestion sub-model. The results of the sensitivity analysis suggest that the ME content is less sensitive to changes of input parameters than the digested protein content (Table 2). The ME content is determined by all parameters in Table 2, whereas the digested protein content is determined by fewer parameters (SCP, DCP, CP, and the slope and intercept of the Lucas equation). In addition, the intercept of the Lucas equation (-32 g CP kg^{-1} DM) amplifies the decrease in digested protein after a decrease in CP content, especially for feeds with a low CP content. As expected, the ME content of feed types with high SNSC contents was reduced most when the SNSC content was decreased by 10%, and the same holds for DNDF (Table 2).

LiGAPS-Beef. The identification of parameters affecting model output most prioritizes which parameters should be investigated in more detail for increasing the models' accuracy (Hamby, 1994, Zuidema *et al.*, 2005). The body core temperature affected model output most, except for Hereford cattle in Uruguay under potential production (Table 3). A higher body core temperature results in a larger temperature gradient between the body core and the ambient environment, which increases heat release, and reduces heat stress. The body core temperature is, however, fairly stable in cattle, but may be investigated further when simulating feed-limited production in (sub-)tropical climates. The conversion of digestible energy (**DE**) to ME ranked high in the top ten parameters under potential production (Table 3). Increasing the efficiency of the DE to ME conversion increases also the NE available for metabolic processes,

403 such as growth, which explains why this parameter affected the FE to a large extent.
404 Values of 0.81 or 0.82 are generally accepted for DE to ME conversion, and a value of
405 0.85 may be appropriate for diets containing high percentages of cereal grains
406 (CSIRO, 2007). Given the sensitivity coefficient of approximately one for the DE to ME
407 conversion, the maximum deviation in model output due to an imprecise estimation of
408 this parameter is approximately 5%.

409 The parameters affecting model output most in each of the four scenarios were NE
410 requirements for maintenance and the multiplier of NE requirements for maintenance
411 (Table 3). Decreasing these parameters increases the NE available for growth and
412 consequently the FE. Model users should thus aim to estimate the breed-specific NE
413 for maintenance, since this parameter is approximately 10% higher for *B. taurus* cattle
414 than for *B. indicus* cattle (NRC, 2000). Several parameters in the top ten affect heat
415 release, which affects the occurrence of heat stress, and consequently the FE.
416 Increasing the body area (or its multiplier), the conduction between body core and skin,
417 and the temperature of exhaled air increases heat release. Parameters associated with
418 heat release were more abundant under feed quality limited production than under
419 potential production. The average sensitivity coefficients were larger under feed quality
420 limited production than under potential production (Table 3). These results are partly
421 explained by the higher heat production during digestion of the grass-based diet under
422 feed quality limited production compared to the diet consisting of 65% wheat and 35%
423 hay under potential production. The higher heat production under feed quality limited
424 production makes thermoregulation and heat release more important than under
425 potential production.

426 Apart from three exceptions, parameters were changed by 5% and 10% using the one-
427 at-a-time approach, which is a structured procedure if standard deviations are

unknown, like in this study. The one-at-a-time approach has two major limitations. First, one parameter was changed at a time while the others were kept constant. We did not investigate effects of changing combinations of parameters, except for parameters of the Gompertz curve. Thus, investigating the joint effects of parameters is a direction for future research. Second, the one-at-a-time approach conducts a local sensitivity analysis and relies on the assumption of model linearity, which is often not justified (Saltelli and Annoni, 2010). The sensitivity coefficients of parameters affecting model output most generally did not differ for a 5% change and a 10% change, which suggests linearity (Table 3). Still, non-linear and non-additive interactions are expected for several parameters, since non-linear equations are used in LiGAPS-Beef. For example, the average sensitivity coefficients for Hereford cattle in Uruguay differed for a change of 5% and 10% in the values for the maximum body weight used in the Gompertz curve, which suggests non-linearity (Table 3). Global sensitivity methods account for non-linearity and non-additivity (Saltelli and Annoni, 2010). We partly addressed the issue of non-linearity by investigating changes in model output at four points (-10%, -5%, 5%, and 10%). Nevertheless, a global sensitivity analysis would provide more information than the one-at-a-time approach. Conducting a global sensitivity analysis is, therefore, another direction for future research.

Evaluation of sub-models

Thermoregulation sub-model. In the experiment of Blaxter and Wainman (1964), simulated and measured heat release generally corresponded to each other, but the sensible heat release with low coat lengths was underestimated (Figs 4A and 4B). A reduction in coat length by shaving might have resulted in a higher conduction of the remaining coat structure. Changing parameters related to coat structure did not decrease the deviation of measured and simulated sensible heat release with low coat

lengths. In addition, simulated skin temperatures were underestimated for one animal with low coat lengths (Fig. 4C). This animal had higher skin temperatures with a low coat length (average 22.9°C) than with a high coat length (average 21.7°C), which is opposite to expectations and measurements for the other animals. Changing parameters related to the coat structure did not decrease the average deviation of measured and simulated skin temperatures either.

Measured heat release and simulated minimum heat release of Friesian and Jersey breeds corresponded to each other at a heat production of 90 W m^{-2} and higher (Fig. 4A). Below 90 W m^{-2}, measured heat release and simulated minimum heat release did not corresponded to each other (treatments at 20°C and at 12°C with a wind speed of 0.22 m s^{-1}). An explanation for the deviations below 90 W m^{-2} is that calves might have been within the thermal neutral zone. The milk-fed calves had a ME intake equivalent to 125 W m^{-2} and a heat production of approximately 95 W m^{-2}, based on their growth rates and an assumed energy retention of 16 MJ kg^{-1} TBW. Hence, the expected heat release within the thermal neutral zone is at least approximately 95 W m^{-2}, which explains why measured heat release and simulated minimum heat release deviated below 90 W m^{-2}. All in all, evaluation of the thermoregulation sub-model indicates that simulated and measured results correspond fairly well to each other. Hence, we assume this sub-model is sufficiently capable of simulating thermoregulation within the model LiGAPS-Beef.

A limitation of the thermoregulation sub-model is its inability to simulate heat flows throughout the day, since it has a daily time step, just like the other two sub-models of LiGAPS-Beef. Evaluation of the thermoregulation sub-model was conducted, therefore, with experiments where climate conditions were kept constant. Nevertheless, climate conditions vary throughout the day for animals kept outdoors or

478 in open stables. For example, body core temperature is a constant in our sub-model,
479 whereas it is known to vary throughout the day under hot conditions (Parkhurst, 2010).
480 Still, an evaluation of LiGAPS-Beef in a companion paper shows that the occurrence
481 of heat stress is simulated fairly well with the daily time step (Van der Linden *et al.*,
482 2018b). The thermoregulation sub-model is calibrated to simulate the average cattle
483 behaviour at a time step, and behaviour throughout the day is not simulated. For
484 example, cattle may move to shaded areas during the warmest periods of the day to
485 mitigate heat stress, and shift their grazing pattern towards cooler periods.

486 *Feed intake and digestion sub-model.* Evaluation of the feed intake and digestion sub-
487 model aimed to investigate whether ME contents could be predicted from the feed
488 constituents specified by Chilibroste *et al.* (1997). Simulated and measured ME
489 contents were not significantly different for a range of feed types. The relative MAEs
490 were 9.3% for the dataset of MAFF (1986) and 6.4% for the dataset of Kolver (2000),
491 respectively. In our opinion, this performance meets the precision required in LiGAPS-
492 Beef sufficiently. As a comparison, the ME contents given by MAFF (1986) and Kolver
493 (2000) differed 5.3% for the same feed types, which may be caused by differences in
494 feed composition. In addition, minimum and maximum ME contents of feed types listed
495 by MAFF (1986) differ considerably as well (Fig. 5). The sub-model captured the
496 variability in simulated ME contents well, since the slope component contributed to less
497 than 1% of the MSE. The ME contents of feeds were generally underestimated (Fig.
498 5). This result corresponds to the result that the bias component accounted for 68.3%
499 of the MSE for the dataset of MAFF (1986), and for 43.3% of the MSE for the dataset
500 of Kolver (2000). Future research may focus, therefore, on fine-tuning parameters of
501 the feed intake and digestion sub-model to simulate the ME contents even more
502 accurately. The ME contents of particular feed types were simulated in detail by using

503 data on the seven feed constituents. If these data are not available, the ME contents
504 of feed types can be obtained from literature, and used as input for LiGAPS-Beef.
505 Further model comparison with regard to digestible protein may not be urgent, as the
506 conversion of CP to digestible protein is calculated via a well-established Lucas
507 equation (Van Soest, 1994).

508 **Conclusions**

509 LiGAPS-Beef aims to assess potential and feed-limited production of beef cattle in
510 different beef production systems across of the world. The first aim of this paper was
511 to assess which parameters affect the output of LiGAPS-Beef most. Sensitivity
512 analyses showed that model output was affected most by body core temperature,
513 conversion of DE to ME, NE requirements for maintenance, and several parameters
514 associated with heat release. Results of the sensitivity analyses can be used to
515 determine which parameters are to be investigated in more detail to increase the
516 accuracy of model simulations. The second aim of the paper was to evaluate the
517 performance of the thermoregulation sub-model and the feed intake and digestion
518 model. Simulated and measured heat release corresponded fairly well to each other.
519 Simulated ME contents of different feed types differed on average by 9.2% and 6.3%
520 from the measured ME contents of two datasets. In conclusion, the performance of
521 both sub-models was considered to be well enough to meet the aim of LiGAPS-Beef,
522 which provides scope to evaluate the complete model further at the animal level.

523 **Acknowledgements**

524 This research is part of the Wageningen University & Research strategic programme
525 ‘Mapping for sustainable intensification’, 2012-2016, funded by the strategic funds of

Wageningen University & Research, and the PE&RC and WIAS graduate schools of Wageningen University.

Declaration of interest

The authors declare they have no conflict of interests.

Software and data repository resources

The source code of LiGAPS-Beef is freely accessible at https://doi.org/10.18174/442973 and the model portal of the Plant Production Systems group of Wageningen University & Research, the Netherlands (http://models.pps.wur.nl/content/ligaps-beef). Updates and model applications will be published on the model portal.

References

Bellocchi G, Rivington M, Donatelli M and Matthews K 2010. Validation of biophysical models: issues and methodologies. A review. Agronomy for Sustainable Development 30, 109-130.

Bibby J and Toutenburg H 1977. Prediction and improved estimation in linear models. John Wiley & Sons, London, UK.

Blaxter KL and Wainman FW 1964. The effect of increased air movement on heat production and emission of steers. Journal of Agricultural Science 62, 207-214.

Chilibroste P, Aguilar C and Garcia F 1997. Nutritional evaluation of diets. Simulation model of digestion and passage of nutrients through the rumen-reticulum. Animal Feed Science and Technology 68, 259-275.

Commonwealth Scientific and Industrial Research Organisation (CSIRO) 2007. Nutrient requirements of domesticated ruminants. CSIRO Publishing, Collingwood, Australia.

Hamby DM 1994. A review of techniques for parameter sensitivity analysis of environmental-models. Environmental Monitoring and Assessment 32, 135-154.

Holmes CW and McLean NA 1975. Effects of air temperature and air movement on heat produced by young Friesian and Jersey calves, with some measurements of effects of artificial rain. New Zealand Journal of Agricultural Research 18, 277-284.

Jones JW, Hoogenboom G, Porter CH, Boote KJ, Batchelor WD, Hunt LA, Wilkens PW, Singh U, Gijsman AJ and Ritchie JT 2003. The DSSAT cropping system model. European Journal of Agronomy 18, 235-265.

Keating BA, Carberry PS, Hammer GL, Probert ME, Robertson MJ, Holzworth D, Huth NI, Hargreaves JNG, Meinke H, Hochman Z, McLean G, Verburg K, Snow V, Dimes JP, Silburn M, Wang E, Brown S, Bristow KL, Asseng S, Chapman S, McCown RL, Freebairn DM and Smith CJ 2003. An overview of APSIM, a model designed for farming systems simulation. European Journal of Agronomy 18, 267-288.

Kolver E, 2000. Nutrition guidelines for the high producing dairy cow. Proceedings of the Ruakura farmers conference 52, 17-28.

564 Lobell DB, Cassman KG and Field CB 2009. Crop yield gaps: their importance, magnitudes,
565 and causes. Annual Review of Environment and Resources 34, 179-204.
566 Lucas HL, Smart WWG, Cipolloni MA and Gross HD 1961. Relations between digestibility
567 and composition of feeds and foods, S-45 Report. North Carolina State College,
568 Raleigh, NC, USA.
569 Mader TL, Davis MS and Brown-Brandl T 2006. Environmental factors influencing heat
570 stress in feedlot cattle. Journal of Animal Science 84, 712-719.
571 Ministry of Agriculture, Fisheries & Food (MAFF) 1986. Feed composition - UK tables of feed
572 composition and nutritive value for ruminants. Chalcombe Publications, Marlow, UK.
573 McGovern RE and Bruce JM 2000. A model of the thermal balance for cattle in hot
574 conditions. Journal of Agricultural Engineering Research 77, 81-92.
575 National Research Council (NRC) 2000. Nutrient Requirements of Beef Cattle, 7th revised
576 edition, National Academy Press, Washington, DC, USA.
577 Parkhurst AM 2010. Model for understanding thermal hysteresis during heat stress: a matter
578 of direction. International Journal of Biometeorology 54, 637-645.
579 Pianosi F, Beven K, Freer J, Hall JW, Rougier J, Stephenson DB and Wagener T 2016.
580 Sensitivity analysis of environmental models: A systematic review with practical
581 workflow. Environmental Modelling & Software 79, 214-232.
582 Prisley SP and Mortimer MJ 2004. A synthesis of literature on evaluation of models for policy
583 applications, with implications for forest carbon accounting. Forest Ecology and
584 Management 198, 89-103.
585 Saltelli A and Annoni P 2010. How to avoid a perfunctory sensitivity analysis. Environmental
586 Modelling & Software 25, 1508-1517.
587 Thompson VA, Barioni LG, Oltjen JW, Rumsey T, Fadel JG and Sainz RD 2011.
588 Development of a heat balance model for cattle under hot conditions. In Modelling
589 nutrient digestion and utilization in farm animals (ed. D Sauvant, Van Milgen, J,
590 Faverdin, P, Friggens, N), pp. 243-251, Wageningen Academic Publishers,
591 Wageningen, The Netherlands.
592 Thornley JHM and France J 2007. Mathematical models in agriculture: quantitative methods
593 for the plant, animal and ecological sciences, 2nd edition. CABI, Wallingford, UK.
594 Turnpenny JR, McArthur AJ, Clark JA and Wathes CM 2000a. Thermal balance of livestock
595 1. A parsimonious model. Agricultural and Forest Meteorology 101, 15-27.
596 Turnpenny JR, Wathes CM, Clark JA and McArthur AJ 2000b. Thermal balance of livestock
597 2. Applications of a parsimonious model. Agricultural and Forest Meteorology 101,
598 29-52.
599 Van de Ven GWJ, de Ridder N, van Keulen H and van Ittersum MK 2003. Concepts in
600 production ecology for analysis and design of animal and plant-animal production
601 systems. Agricultural Systems 76, 507-525.
602 Van der Linden A, Oosting SJ, Van de Ven GWJ, De Boer IJM and Van Ittersum MK 2015. A
603 framework for quantitative analysis of livestock systems using theoretical concepts of
604 production ecology. Agricultural Systems 139, 100-109.
605 Van der Linden A, Van de Ven GWJ, Oosting SJ, Van Ittersum MK and De Boer IJM 2018a.
606 LiGAPS-Beef, a mechanistic model to explore potential and feed-limited beef
607 production 1. Model description and illustration. Animal.
608 Van der Linden A, Van de Ven GWJ, Oosting SJ, Van Ittersum MK and De Boer IJM 2018b.
609 LiGAPS-Beef, a mechanistic model to explore potential and feed limited beef
610 production 3. Model evaluation Animal.
611 Van Ittersum MK, Cassman KG, Grassini P, Wolf J, Tittonell P and Hochman Z 2013. Yield
612 gap analysis with local to global relevance - A review. Field Crops Research 143, 4-
613 17.
614 Van Soest PJ 1994. Nutritional ecology of the ruminant, 2nd edition. Cornell University Press,
615 Ithaca, NY, USA.
616 Zuidema PA, Leffelaar PA, Gerritsma W, Mommer L and Anten NPR 2005. A physiological
617 production model for cocoa (*Theobroma cacao*): model presentation, validation and
618 application. Agricultural Systems 84, 195-225.

619

620 **Table 1**. Changes in lower critical temperature (LCT) and upper critical temperature (UCT) of
621 beef cattle after changing parameters by 10%. The baseline LCT is -1.0°C and the baseline
622 UCT is 30.5°C. Changes (only 1°C or more) are given in degrees Celsius, relative to the
623 baseline.

Parameter determining:	LCT		UCT	
	-10%	+10%	-10%	+10%
Body area 2[1]	2.1	-1.6	2.3	-1.6
Body temperature[2]	-1.0	0.9	-4.8	9.5
Exhaled air temperature	-0.6	0.5	-1.2	1.5
Conduction core-skin 1[3]	-3.0	2.4	0.0	0.0
Conduction core-skin 2[3]	2.6	-2.7	0.0	0.0
Conduction core-skin 3[3]	4.8	-5.9	0.0	0.0
Max. conduction body core – skin	0.0	0.0	-1.4	1.6
Latent heat release 2[4]	0.0	0.0	-1.3	1.7
Latent heat release 3[4,5]	0.0	0.0	3.5	-2.1

624 [1] Body area (m^2) = body area multiplier × body area 1 × total body weight $^{\text{body area 2}}$ (McGovern and Bruce,
625 2000).

626 [2] Body temperature has been changed by 1°C.

627 [3] Min. conduction core-skin ($W\ m^{-2}\ K^{-1}$) = Conduction core-skin 1 / (Conduction core-skin 2 ×
628 $TBW^{\text{Conduction core-skin 3}}$)

629 [4] Maximum latent heat release ($W\ m^{-2}$) = minimum heat release + latent heat release 1 × $e^{(\text{latent heat}}$
630 $^{\text{release 2 × (skin temperature - latent heat release 3))}}$ × latent heat of water vapour

631 [5] This parameter has been changed by 1%.

Table 2. Effect of a 10% decrease in feed components on metabolisable energy (ME) and digested protein ($P_{dig.}$) per kg DM feed. Baseline ME and $P_{dig.}$ indicate the whole-tract digestibility for beef cattle. Other values indicate the relative change in ME and $P_{dig.}$ compared to the baseline (%).

	Baseline		SNSC	INSC	DNDF	SCP		DCP		Total CP		kdDNDF	kdPass	Dig. INSC	Dig. DNDF	Pass. DNDF	Lucas slope		Lucas intercept	
Feed type	ME	$P_{dig.}$	ME	ME	ME	ME	$P_{dig.}$	ME	$P_{dig.}$	ME	$P_{dig.}$	ME	ME	ME	ME	ME	ME	$P_{dig.}$	ME	$P_{dig.}$
	MJ kg^{-1} DM	g kg^{-1} DM	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
Barley	12.7	92	-4.4	-2.3	-1.6	-0.4	-3.4	-0.9	-8.1	-1.5	-13.5	-0.2	0.1	-1.8	-0.1	0.1	-1.9	-13.5	0.5	3.5
Concentrates	11.7	132	-3.2	-2.1	-2.3	-1.1	-5.0	-1.3	-6.0	-2.7	-12.4	-0.7	0.5	-2.1	-0.2	0.2	-2.7	-12.4	0.5	2.4
Hay (good quality)	9.6	123	-1.5	-2.2	-3.6	-0.8	-3.5	-1.3	-5.4	-2.9	-12.6	-1.4	1.0	-1.2	-0.4	0.4	-3.2	-12.6	0.7	2.6
Hay (poor quality)	7.8	31	-1.3	-1.3	-5.9	-0.5	-5.9	-3.6	-43.3	-1.6	-20.3	-2.3	1.6	-0.7	-0.6	0.6	-1.6	-20.3	0.8	10.3
Grass (spring)	11.0	207	-1.7	-0.4	-3.2	-2.4	-2.9	-3.5	-4.2	-8.8	-11.5	-1.3	0.9	-0.2	-0.3	0.3	-4.2	-11.5	0.6	1.5
Grass (summer)	8.8	130	-1.6	-0.9	-4.2	-1.0	-3.4	-1.5	-5.3	-3.5	-12.5	-1.6	1.2	-0.5	-0.4	0.5	-3.6	-12.5	0.7	2.5
Grass (dry summer)	7.4	72	-1.0	-1.1	-5.5	-0.4	-2.9	-1.3	-8.7	-2.2	-14.5	-2.1	1.5	-0.6	-0.6	0.6	-2.7	-14.5	0.8	4.5
Maize grain	13.3	89	-2.2	-5.5	-0.8	-0.2	-2.0	-0.9	-8.8	-1.4	-13.6	-0.3	0.2	-5.5	-0.1	0.1	-1.8	-13.6	0.5	3.6
Maize silage	10.1	42	-1.4	-4.8	-2.4	-0.9	-11.8	-0.4	-5.0	-1.4	-17.7	-0.9	0.6	-2.7	-0.2	0.2	-1.4	-17.7	0.6	7.7
Molasses	11.6	-28	-10.4	0.0	0.0	-0.1	-1.2	0.0	-0.1	-0.1	-1.3	0.0	0.0	0.0	0.0	0.0	-0.1	-1.3	0.6	11.3
Soy bean meal	11.6	424	-1.3	0.0	-1.6	-3.1	-4.3	-3.7	-5.2	-7.7	-10.8	-0.2	0.1	0.0	-0.1	0.1	-7.7	-10.8	0.5	0.8
Straw (cereals)	5.8	4	-0.3	-1.9	-6.9	-0.4	-22.5	-0.2	-11.2	-1.7	-90.0	-2.7	1.9	-1.0	-0.7	0.7	-1.2	-90.0	1.1	80.0
Wheat	12.8	88	-5.3	-2.3	-0.8	-0.4	-4.1	-0.7	-7.2	-1.4	-13.6	-0.1	0.1	-2.1	0.0	0.0	-1.8	-13.6	0.5	3.6

SNSC = Soluble, non-structural carbohydrates; INSC = Insoluble, non-structural carbohydrates; DNDF = digestible neutral detergent fibre; SCP = soluble crude protein; DCP = digestible crude protein; CP = crude protein. kdDNDF = digestion rate digestible NDF; kdPass = passage rate; Dig. INSC = digestion rate of insoluble, non-structural carbohydrates for the whole digestive tract; Dig. DNDF = digestion rate of degradable NDF in the intestines; Pass. DNDF = passage rate of degradable neutral detergent fibre in the intestines; Lucas slope and intercept = slope and intercept of a Lucas equation (Eq. 1, Lucas *et al.* 1961; van Soest, 1994).

Table 3. Average sensitivity coefficient (ASC) of the top-10 parameters affecting the feed efficiency of beef cattle at the herd level most. Sensitivity analysis was conducted with LiGAPS-Beef by increasing and decreasing parameters values by 5% (ASC 5%) and 10% (ASC 10%).

	B×S cattle, Australia, potential			B×S cattle, Australia, feed quality limited			Hereford cattle, Uruguay, potential			Hereford cattle, Uruguay, feed quality limited		
Rank	Parameter	ASC 5%	ASC 10%	Parameter	ASC 5%	ASC 10%	Parameter	ASC 5%	ASC 10%	Parameter	ASC 5%	ASC 10%
1	Body core temperature[1]	1.32	NA	Body core temperature[1]	25.64	NA	DE to ME conversion	0.98	0.98	Body core temperature[1]	12.90	NA
2	DE to ME conversion	0.94	0.94	NE for maintenance	1.80	NA	Maximum adult total body weight (Gompertz curve)	0.74	0.51	NE for maintenance	1.89	NA
3	NE for maintenance	0.60	0.60	Maintenance multiplier	1.80	NA	NE for maintenance	0.61	0.61	Maintenance multiplier	1.86	NA
4	Maintenance multiplier	0.60	0.60	Latent heat release 3[2]	1.45	NA	Maintenance multiplier	0.61	0.61	Maximum adult total body weight (Gompertz curve)	1.03	0.92
5	Maximum adult total body weight (Gompertz curve)	0.59	NA	Body area 1[3]	1.43	NA	Maximum adult total body weight	0.53	0.53	Body area 1[3]	1.02	NA
6	Slope Lucas equation[4]	0.29	0.29	Body area multiplier[3]	1.43	NA	Growth rate constant (Gompertz curve)	0.31	0.31	Body area multiplier[3]	1.02	NA
7	Carcass fraction	0.28	0.28	Latent heat release 2[2]	1.19	NA	Carcass fraction	0.31	0.31	Body area 2[3]	0.77	NA
8	Body area 2[3]	0.28	0.38	Maximum conduction body core - skin	1.17	NA	Slope Lucas equation[4]	0.24	0.24	Temperature exhaled air 1[5]	0.65	0.60
9	Efficiency of protein accretion	0.27	0.27	Body area 2[3]	1.12	NA	Lipid fraction fat tissue	0.23	0.23	Maximum conduction body core - skin	0.64	0.60

10	Lipid fraction fat tissue	0.26	0.26	Standard respiration rate	1.00	0.91	Constant of integration (Gompertz curve)	0.22	0.22	Growth rate constant (Gompertz curve)	0.61	0.52

642 B×S = Brahman × Shorthorn crossbred cattle; DE = digestible energy; ME = metabolisable energy; NA = no model output; NE = net energy.

643 [1] Body core temperature was decreased and increased by 0.1°C.

644 [2] Maximum latent heat release (W m^{-2}) = minimum heat release + latent heat release 1 × $e^{(\text{latent heat release 2} \times (\text{skin temperature - latent heat release 3}))}$ × latent heat of water

645 vapour. Latent heat release 3 was changed by 1%.

646 [3] Body area (m^2) = body area multiplier × body area 1 × total body weight $^{\text{body area 2}}$ (McGovern and Bruce, 2000). Body area 2 was changed by 1%.

647 [4] For the Lucas equation, see equation 1.

648 [5] Temperature exhaled air = temperature exhaled air 1 + temperature exhaled air 2 × air temperature + $e^{(\text{temperature exhaled air 3} \times \text{relative humidity} + \text{temperature exhaled air 4} \times \text{air}}$

649 $^{\text{temperature)}}$

Figures




Figure 1 Representation of LiGAPS-Beef (Livestock simulator for Generic analysis of Animal Production Systems – Beef cattle) and the connections among the three sub-models. Solid arrows indicate flows of material or energy in beef cattle, dashed arrows indicate a flow of information. ME = metabolisable energy; NE = net energy. Source: van der Linden *et al.* (2018a).






Figure 2. Effects of temperature in combination with solar radiation, relative humidity, wind speed, precipitation, cloud cover, total body weight, and heat production on the simulated thermal neutral zone (in white) of a bovine animal. The lower critical temperature of the cattle is the left edge of the thermal neutral zone (TNZ); the upper critical temperature the right edge.



663


664 **Figure 3**. Combined temperature and relative humidity to compare the occurrence of
665 heat stress in beef cattle simulated by the thermoregulation sub-model of LiGAPS-Beef
666 after calibration (A) with the temperature-humidity index of Mader *et al.* (2006) (B) and
667 the temperature-humidity index of Mader *et al.* (2006) accounting for wind speed and
668 solar radiation (C). Dashed lines indicate the simulated temperature at which heat
669 stress occurs with a relative humidity of 20% and 100%.



670

671 **Figure 4**. Simulated and measured total heat release of beef cattle for experiments of
672 Blaxter and Wainman (1964) and Holmes and McLean (1975) (A), together with
673 sensible and latent heat release (B) and skin temperature (C) for the experiment of
674 Blaxter and Wainman (1964). Dashed lines indicate y = x. CL = coat length.

675



676

677 **Figure 5**. Simulated and measured metabolisable energy (ME) content of feed types
678 consumed by beef cattle, which are given by MAFF (1986) and Kolver (2000).
679 Horizontal bars indicate the minimum and maximum simulated ME contents. Vertical
680 bars data indicate the minimum and maximum ME contents listed by MAFF (1986).