

Relationship between metabolic status and behavior in dairy cows in week 4 of lactation

R. J. van Hoeij¹, A. Kok^{1,2}, R. M. Bruckmaier³, M. J. Haskell⁴, B. Kemp¹
and A. T. M. van Knegsel^{1†}

¹Adaptation Physiology Group, Department of Animal Sciences, Wageningen University, PO Box 338, 6700 AH Wageningen, The Netherlands; ²Animal Production Systems Group, Department of Animal Sciences, Wageningen University, PO Box 338, 6700 AH Wageningen, The Netherlands; ³Veterinary Physiology, Vetsuisse Faculty, University of Bern, Bremgartenstrasse 109a, CH-3001 Bern, Switzerland; ⁴Scotland's Rural College (SRUC), West Mains Road, Edinburgh EH9 3JG, UK

(Received 22 September 2017; Accepted 22 May 2018; First published online 26 July 2018)

Blood metabolite and hormone concentrations are indicative of metabolic status, but blood sampling and analysis is invasive and time-consuming. Monitoring behavior can be done automatically, and behaviors may also be used as indicators of metabolic status. The aim of this study was to analyze the relationships between metabolic status and feeding behavior, lying behavior, motion index and steps of dairy cows in week 4 postpartum. Behavioral data from 81 Holstein-Friesian cows were collected using computerized feeders and accelerometers, and blood samples were collected for analysis of free-fatty acid (FFA), β -hydroxybutyrate (BHB), glucose, insulin, IGF-1 and growth hormone (GH) concentrations. First, cluster analysis was performed to categorize cows as having poor, average, good or very good metabolic status based on their plasma FFA, BHB, glucose, insulin, IGF-1 and GH concentration. Subsequently, the performance and behavior of cows in clusters with poor, average and good metabolic status were compared using GLM. Cows with a poor or average metabolic status tended to have greater fat-and-protein-corrected milk yield than cows with good metabolic status. Furthermore, cows with a poor metabolic status had a lower energy balance and dry matter intake (DMI) than cows with an average or good metabolic status and had a lower number of meals than cows with good metabolic status. Daily number of visits to the feeder and lying time tended to be positively related with metabolic status. Feeding rate (kg/min), daily meal time (min/day), number of lying bouts per day, steps and motion index were not related with metabolic status. In conclusion, better metabolic status in dairy cows in early lactation was associated with a greater DMI, increased feeding activity and a tendency to more time spent lying, compared with poor metabolic status. These results suggest that compromised metabolic status is reflected in altered cow's behavior in week 4 of lactation.

Keywords: dry period length, continuous milking, feeding behavior, lactogenic hormones, sensor technology

Implications

Blood metabolite and hormone concentrations are indicative of metabolic status, but blood sampling and analysis is invasive and time-consuming. Monitoring behavior can be done automatically, and behaviors may also be used as indicators of metabolic status. A better insight into relationships between metabolic status and behavior may improve the interpretation of the behavior of dairy cows in early lactation. Moreover, such associations can facilitate the interpretation of automatically recorded behavior to monitor the health of individual cows in early lactation.

Introduction

Diseases such as ketosis, retained placenta, metritis, mastitis, milk fever, displaced abomasum, lameness and impaired fertility result in a high disease incidence in the early lactation of dairy cows (Drackley, 1999; Ingvarsen *et al.*, 2003). In addition, disease events in early lactation may affect milk production performance in later lactation including total lactation yield (Fourichon *et al.*, 1999). Diseases in early lactation are often related with the negative energy balance (EB) of dairy cows during this period (Collard *et al.*, 2000; Ingvarsen *et al.*, 2003). The negative EB in early lactation is caused by insufficient energy intake to support the high milk yield. A negative EB is related with an altered metabolic status, indicated by lower glucose, insulin and IGF-1 concentrations in plasma, and greater free-fatty acid (FFA),

[†] E-mail: ariette.vankneggel@wur.nl

β -hydroxybutyrate (BHB) and growth hormone (GH) concentrations in plasma (Fenwick *et al.*, 2008).

Plasma metabolites and hormones can be indicators of metabolic status (Butler *et al.*, 2003), but blood sampling is invasive and time-consuming. Monitoring behavior is less invasive. Moreover, behavior such as feeding time and feeding rate, rumination time and steps are increasingly collected using sensors to indicate, for instance, heat or the moment of calving (Reith *et al.*, 2014). Changes in lying, walking and feeding behavior of dairy cows have been associated with diseases (Edwards and Tozer, 2004) such as lameness (González *et al.*, 2008), metritis (Urton *et al.*, 2005) or a displaced abomasum (Van Winden *et al.*, 2003). It can be hypothesized that lying, walking or feeding behavior is not only indicative for disease, but also for metabolic status of dairy cows in early lactation. A previous study reported that cows with a lower plasma FFA concentration had a greater walking activity *postpartum*, compared with cows with a greater plasma FFA concentration (Adewuyi *et al.*, 2006). In addition, cows with ketosis had a lower feed intake, feeding time, meal time and fewer meals and visits to the feeder than cows without ketosis (González *et al.*, 2008). To our knowledge, relationships between behavior and other plasma metabolites and metabolic hormones have not been described previously.

Shortening or omitting the dry period improves the EB and metabolic status through a lower milk yield and an increased or similar feed intake after calving (Rastani *et al.*, 2005; Van Knegsel *et al.*, 2014a). A previous study showed that a better EB after short or no dry period was weakly correlated with longer lying time, more feeding behavior and higher feed intake (Kok *et al.*, 2016). It can be hypothesized, however, that metabolic status may better reflect cow health and feelings than the EB, because metabolic status better reflects the degree to which the animal can cope with the catabolic state that is typical for early lactation (Van Knegsel *et al.*, 2014b).

The aim of this study was to analyze the associations between metabolic status, based on plasma metabolites and hormones, and lying and feeding behavior, motion index and steps of dairy cows. To address this aim, metabolic status and behavior were recorded in an experiment with dairy cows in early lactation after a 0-day or 30-day dry period. The current study monitored dairy cows in week 4 after calving to limit the direct effect of the calving process and start of lactation on behavior, and maximize the contrast in EB of cows with different dry period lengths (Van Knegsel *et al.*, 2014a). Moreover, lying time was expected to be lowest in week 4 of lactation (Maselyne *et al.*, 2017).

Material and methods

Animals and housing

The Institutional Animal Care and Use Committee of Wageningen University & Research approved the experimental protocol in compliance with the Dutch law on Animal

Experimentation (protocol no. 2014125). The experiment was originally designed to study the effects of dry period length on metabolic status (Van Hoeij *et al.*, 2017). The experiment was conducted at the Dairy Campus research herd (Lelystad, The Netherlands) between 27 January 2014 and 26 August 2015. The research herd was composed of 400 lactating Holstein cows. Cows were selected based on: (1) being bred with a Holstein sire, (2) expected calving interval <490 days, (3) daily milk yield >16 kg at 90 days before the expected calving date and (4) no clinical mastitis or high SCC ($\geq 250\,000$ cells/ml) at the final two test-days before drying off. All cows were housed in the same free-stall barn, which had concrete slatted floors, and cubicles (1.25 $\times$ 2.20 m) fitted with rubber mattresses (4 cm thick) covered with sawdust. Stocking density was maintained at 7 m² per cow, with one cow per cubicle. Cows were milked twice daily at ~0600 and 1700 h. Lying behavior, steps and motion index of cows were recorded for 6 complete days (Friday till Wednesday) in the 4th week after calving (i.e. week 4 after calving), because of limited sensor availability and changing of sensors on Thursdays.

Experimental design

In total, 130 cows entered the experiment, including six cows that entered twice. To obtain a balanced distribution of cows across treatments, cows were blocked according to the expected calving date, milk yield in the previous lactation and parity (2, ≥ 3) in the subsequent lactation. Within each group of three cows, two cows were assigned randomly to a dry period length treatment of 0 days (0-day DP) and one cow to a dry period length treatment of 30 days (30-day DP). Within the group of cows with a 0-day DP, cows were assigned randomly to either a low level of concentrate based on the energy requirement for their expected milk yield (LOW) or a standard (STD) level of concentrate based on the energy requirement for the expected milk yield of cows after a 30-day DP (Van Knegsel *et al.*, 2014). Cows with a 30-day DP were fed an STD level of concentrate, based on the requirement for their expected milk yield. This resulted in the following three treatment groups: cows with a 30-day DP fed the STD level of concentrate required for their expected milk yield (30-day DP(STD)), cows with a 0-day DP fed the same STD concentrate level as cows with a 30-day DP (0-day DP(STD)) and cows with a 0-day DP fed a LOW concentrate level (0-day DP(LOW)). Preliminary statistical analyses showed no effect of concentrate level within the 0-day DP treatment on feeding behavior, lying behavior, steps or motion index. Concentrate level was therefore excluded from further analyses in this study. Lying behavior, steps and motion index were only measured for 81 of 130 cows. The final data set for this study consisted of milk production, EB, BW, feed intake, plasma metabolite and metabolic hormone concentrations, and lying behavior, steps and motion index data of 81 unique cows in week 4 *postpartum* ($n = 53$ for 0-day DP and $n = 28$ for 30-day DP). Basal lactation ration, concentrate composition and feeding strategy were reported previously (Van Hoeij *et al.*, 2017).

Measurements

Behaviors. Measurement of behaviors was described earlier (Kok *et al.*, 2016). In short, feeding behavior was measured in week 4 *postpartum*. Basal lactation ration was provided and its daily intake was measured individually using roughage intake control (RIC) feeders (Insentec, Marknesse, the Netherlands). The stocking density was two cows per trough. The actual quantity of concentrate dispensed (kg/day) was recorded by the computerized feeder (Manus VC5; DeLaval, Steenwijk, the Netherlands). For each visit to a feeder, RIC feeders recorded cow identity, the start time and end time (hh:mm:ss) of the visit and the start weight and end weight of the feed in the feeder to the nearest 0.1 kg. Visits were clustered into meals based on the interval length between visits (Yeates *et al.*, 2001; Tolkamp *et al.*, 2002), with a threshold of 20.9 min between meals (Kok *et al.*, 2016). Feeding behaviors used for analyses were the average daily duration for 6 days of meals (meal time, min/day), average daily number of visits (visits, *n*/day), the average daily number of meals (meals, *n*/day) and the secondary variables daily feed intake (kg dry matter (DM)/day) and feeding rate (kg/min) that were derived from these variables.

Lying behavior, steps and motion index were recorded in week 4 *postpartum* with triaxial accelerometers (IceQube; IceRobotics, South Queensferry, UK). Lying behavior is recorded when the hind leg is in a horizontal position; the step count measures the number of times the animal lifts its leg up and places it back down again. Lying bouts <33 s were discarded as erroneous (Kok *et al.*, 2015). The step count was used as an indicator for walking activity. Motion index is a measure of the overall acceleration measured by the sensor in all three axes. Sensors were attached and detached to the hind leg on Thursdays between 1000 and 1200 h. Means of lying bouts, lying time, steps and motion index per cow per day over 6 consecutive days were used for the analysis in week 4 of lactation.

Milk yield and milk composition. Milk yield was recorded daily in week 4 *postpartum*. Milk samples for fat, protein and lactose analysis (ISO 9622; Qlip, Zutphen, the Netherlands) were collected four times per week (Tuesday afternoon, Wednesday morning, Wednesday afternoon and Thursday morning) and were analyzed as a pooled sample per cow per week. Fat-and-protein-corrected milk (FPCM) was calculated as:

$$\text{FPCM} = (0.337 + 0.116 \times \text{fat percentage} + 0.06 \times \text{protein percentage}) \times \text{milk yield}$$

(Centraal Veevoederbureau, 2016)

Body weight and energy balance. Body weight was recorded daily before each milking and averaged for week 4 of lactation. Energy balance was calculated per week, according to the Dutch net energy (NE) system for lactation (VEM) (Van Es, 1975), as the difference between energy intake and energy requirements for maintenance and milk yield (1000 VEM = 6.9 MJ of NE). According to the VEM system, the daily

requirement for maintenance is 42.4 VEM/kg^{0.75} BW per day and the requirement for milk yield is 442 VEM/kg FPCM (Van Es, 1975). Energy intake and EB are expressed in kJ/kg^{0.75} BW per day (Van Es, 1975).

Blood collection and analysis. Blood was collected on Thursday in week 4 *postpartum*. Blood was collected after the morning milking, between 3 and 1 h before the morning feeding. Blood (10 ml) was collected from the coccygeal vein into evacuated EDTA tubes (Vacuette; Greiner BioOne, Kremsmunster, Austria). Blood samples were kept on ice before centrifugation for plasma isolation (3000 × *g* for 15 min, 4°C). Plasma samples were stored at -20°C. Concentrations of FFA and BHB were measured enzymatically using kit no. 994-75409 from Wako Chemicals (Neuss, Germany) and kit no. RB1007 from Randox Laboratories (Ibach, Switzerland), respectively (Graber *et al.*, 2012). Plasma glucose concentration was measured using kit no. 61269 from BioMerieux (Marcy l'Etoile, France) (Graber *et al.*, 2012). Plasma insulin concentration was measured using kit no. PI-12K from EMD Millipore Corporation (Billerica, MA, USA). Plasma IGF-1 concentration was measured using kit no. A15729 from Beckman Coulter (Fullerton, CA, USA). Plasma GH concentration was measured by radioimmunoassay as described previously (Vicari *et al.*, 2008).

Statistical analyses

The natural logarithm of the plasma FFA, BHB and GH concentration was calculated to approximate normal distribution of these variables and was used in all statistical analyses. To evaluate the normality of residuals, a normality test was used in data distribution (PROC UNIVARIATE), and skewness between -1 and 1, kurtosis between -2 and 2 and a non-significant Shapiro-Wilk test were used as criteria for normality. To analyze correlations of plasma metabolites and hormones (FFA, BHB, glucose, insulin, IGF-1 and GH) with dry matter intake (DMI), feeding behavior, lying behavior, steps and motion index, a Pearson correlation was used (PROC CORR; SAS 9.3, SAS Institute Inc. (2011)). Pearson's correlation analysis was also used to analyze correlations among the different plasma metabolites and hormones.

Because in the current study most metabolites and hormones were correlated (Table 1b), as a second step, cluster analysis with the Ward method was performed using plasma FFA, BHB, glucose, insulin, IGF-1 and GH concentration as explanatory variables (PROC CLUSTER). Based on the cubic clustering criterion, pseudo F statistic and pseudo *t*-squared, cows were clustered in four clusters using the tree procedure (PROC TREE). The cluster with the very good metabolic status included only six cows that had a 0-day DP, a 12 kg lower average milk yield and lower feed intake than other clusters. Because of the limited number of animals and the extremely low milk production in this cluster, the cluster with a very good metabolic status was excluded from the analysis. Clusters with poor, average and good metabolic status were compared.

Table 1 Pearson's correlation coefficients ($P < 0.05$) (a) of plasma metabolites with fat-and-protein corrected milk (FPCM) yield, dry matter intake (DMI), energy balance (EB), feeding rate, number of meals and visits, meal time, lying time and bouts, steps and motion index; (b) among plasma free-fatty acids (FFA), β -hydroxybutyrate (BHB), glucose, insulin, IGF-1 and growth hormone (GH) concentrations in dairy cows

	FFA (mmol/l)	BHB (mmol/l)	Glucose (mmol/l)	Insulin (μ U/ml)	IGF-1 (ng/ml)	GH (μ g/l)
(a) Plasma metabolites with						
FPCM (kg/day) ¹	0.49**	0.31**	-0.37**	-0.47**	-0.41**	0.26*
DMI (kg DM/day) ¹	-0.49**	Ns	0.23*	Ns	0.22*	-0.34**
EB (kJ/kg ^{0.75} •per day) ¹	-0.78**	-0.41**	0.48**	0.33*	0.55**	-0.45**
Feeding rate (kg/min)	Ns	Ns	0.30**	Ns	Ns	-0.25*
Meals (n/day)	-0.34**	Ns	Ns	Ns	0.34**	-0.24*
Visits (n/day)	-0.39**	-0.2*	0.23*	Ns	0.22*	Ns
Meal time (min/day)	-0.38**	Ns	Ns	Ns	Ns	Ns
Lying time (h/day)	-0.43**	Ns	Ns	Ns	0.32**	Ns
Lying bouts (n/day)	Ns	Ns	Ns	Ns	Ns	Ns
Steps (n/day)	-0.32**	-0.25*	Ns	Ns	Ns	Ns
Motion index	-0.37**	-0.28*	Ns	Ns	Ns	Ns
(b) Plasma metabolites						
FFA ¹		0.33**	-0.48**	-0.40**	-0.52**	0.40**
BHB ²			-0.52**	-0.24*	Ns	Ns
Glucose				0.39**	0.53**	-0.27*
Insulin					Ns	Ns
IGF-1						-0.39**

Ns = not significant.

¹FFA is analyzed using the natural logarithm of FFA, BHB is analyzed using the natural logarithm of BHB and GH is analyzed using the natural logarithm of GH.* $P < 0.05$, ** $P < 0.01$.

To evaluate the plasma concentration of metabolites and hormones, FPCM yield (kg/day), EB (kJ/kg^{0.75}•per day) and BW for cows with different metabolic status, a GLM was used (PROC GLM). The independent variable was cluster:

$$Y_i = \text{cluster}_i + \varepsilon_i$$

where cluster_{*i*} indicates the mean of cluster *i* (*i* = good, average or poor) and ε_i indicate the random residual. To evaluate DMI, feeding behavior, lying time, steps and motion index for cows with different metabolic status, the GLM was extended with dry period length (0-day DP or 30-day DP) and parity (2 or ≥ 3) as dependent variables.

$$Y_{ijkl} = \text{cluster}_i + \text{DP}_j + \text{parity}_k + \varepsilon_{ijk}$$

where DP_{*j*} indicates the DP length (*i* = 0-, or 30- DP) and parity_{*k*} indicates the parity of the cow (*j* = parity 2, or ≥ 3).

The model was analyzed using a backward elimination procedure with a stay-in *P*-value of < 0.05 in the type III Wald test.

Results

Correlations between plasma metabolites and hormones and behavioral traits

The average daily FPCM yield was negatively correlated with plasma glucose, insulin and IGF-1, and positively with plasma FFA, BHB and GH ($P < 0.05$) (Table 1a). Dry matter intake was negatively correlated with plasma FFA and GH, and positively with plasma glucose and IGF-1 ($P < 0.05$). Energy balance was negatively correlated with the plasma FFA, BHB and GH, and positively with plasma glucose, insulin and IGF-1 ($P < 0.05$). Feeding rate was negatively correlated

with plasma GH, and positively with plasma glucose ($P < 0.05$). A number of meals per day was negatively correlated with plasma FFA and GH. Number of visits to the feeder was negatively correlated with plasma FFA and BHB, and positively with plasma glucose and IGF-1 ($P < 0.05$). Meal time and lying time were negatively correlated with plasma FFA, and lying time was positively correlated with plasma IGF-1 ($P < 0.01$). Steps and motion index were negatively correlated with plasma FFA and BHB ($P < 0.05$).

The plasma FFA concentration was negatively correlated with the plasma glucose, insulin and IGF-1, and positively with the plasma BHB and GH ($P < 0.01$) (Table 1b). The plasma BHB concentration was negatively correlated with the plasma glucose and insulin concentration ($P < 0.05$). The plasma glucose concentration was negatively correlated with the plasma GH concentration, and positively with the plasma insulin and IGF-1 concentration ($P < 0.05$). The plasma IGF-1 concentration was negatively correlated with the plasma GH concentration ($P < 0.01$).

Relation between metabolic status and feeding and lying behavior, steps and motion index

Cows were clustered for metabolic status based on their plasma FFA, BHB, glucose, insulin, IGF-1 and GH concentration in week 4 *postpartum*. This resulted in four groups of cows with a poor, average, good or very good metabolic status. Cows with a poor, average or good metabolic status had a 0-day or 30-day DP length, and parity 2 or ≥ 3 (Table 2). All six cows with a very good metabolic status had a 0-day DP length and were not included in further analyses. Cows with a good metabolic status had lower plasma FFA concentration (Figure 1a), and greater plasma glucose

Table 2 Number of cows in different clusters based on the plasma free-fatty acid (FFA), β -hydroxybutyrate (BHB), glucose, insulin, IGF-1 and bovine growth hormone (GH) concentration

Dry period length	Parity	Metabolic status			
		Poor	Average	Good	Very good
0-day DP	2	0	10	14	4
0-day DP	≥ 3	3	14	6	2
30-day DP	2	2	6	3	0
30-day DP	≥ 3	4	11	2	0
Total cows (<i>n</i>)		9	41	25	6

DP = dry period.

(Figure 1c) and IGF-1 concentration (Figure 1e) than cows with average or poor metabolic status ($P < 0.01$) (Table 3). Cows with an average metabolic status had lower plasma FFA and greater plasma glucose and IGF-1 concentrations than cows with poor metabolic status. Cows with a good metabolic status tended to have a lower plasma BHB (Figure 1b) and greater plasma insulin (Figure 1d) than cows with a poor metabolic status ($P < 0.10$). Plasma IGF-1 increased with metabolic status (adjusted R^2 : 0.82, $P < 0.01$; Figure 1e). Energy balance increased with metabolic status ($P < 0.01$). Cows with a good metabolic status tended to have lower FPCM yield than cows with a poor or average metabolic status ($P < 0.10$). Body weight did not differ between cows with different metabolic status.


**Figure 1** The plasma (a) free-fatty acids (FFA), (b) β -hydroxybutyrate (BHB), (c) glucose, (d) insulin, (e) IGF-1 and (f) growth hormone (GH) concentration for cows clustered for metabolic status based on plasma FFA, BHB, glucose, insulin, IGF-1, and GH concentration. Values represent the minimum, first quartile, median, second quartile and maximum. ^{a,b,c}Values with different symbols differ ($P < 0.05$).

Table 3 Plasma metabolite and metabolic hormone concentrations in week 4 postpartum for cows with different metabolic status after a 0- or 30-day dry period

	Adjusted R^2	Metabolic status			SEp	P-value
		Poor	Average	Good		
Cows		9	41	25		
FFA (mmol/l) ¹	0.30	0.46 (0.28 to 0.75) ^c	0.16 (0.13 to 0.20) ^b	0.10 (0.07 to 0.13) ^a		< 0.001
BHB (mmol/l) ¹	0.07	1.00 (0.72 to 1.39)	0.72 (0.62 to 0.84)	0.63 (0.52 to 0.77)		0.06
Glucose (mmol/l)	0.23	3.21 ^a	3.72 ^b	3.96 ^c	0.18	< 0.001
Insulin (μ U/ml)	0.07	7.7	15.0	14.9	3.8	0.09
IGF-1 (ng/ml) ¹	0.82	52 ^a	101 ^b	151 ^c	6.5	< 0.001
GH (μ g/l) ¹	0.04	4.68 (3.48 to 6.29)	4.52 (3.94 to 5.19)	3.81 (3.15 to 4.60)		0.29
FPCM	0.07	38.8	39.0	35.6	2.5	0.06
EB (kJ/kg ^{0.75} •per day)	0.22	-242 ^a	-124 ^b	6 ^c	66	< 0.001
BW (kg)	0.03	638	674	66	27	0.30

FFA = free-fatty acids; BHB, β -hydroxybutyrate; GH = growth hormone; FPCM = fat-and-protein corrected milk; EB = energy balance.

Metabolic status was based on the plasma FFA, BHB, glucose, insulin, IGF-1 and GH concentration. Values are presented as LSM and pooled SE (SEp).

^{a,b,c}Values with different superscripts differ ($P < 0.05$).¹ FFA, BHB and GH were log-transformed for analyses, but are shown as actual values with confidence intervals.**Table 4** Behavior in week 4 postpartum of cows with different metabolic status after a 0- or 30-day dry period

	Adjusted R^2	Metabolic status				P-value		
		Poor	Average	Good	SEp	C	DP	P
Cows		9	41	25				
Dry matter intake (kg DM/day)	0.25	19.5 ^b	22.0 ^a	22.4 ^a	0.9	< 0.01	< 0.01	0.44
Basal ration intake (kg DM/day)	0.30	12.5 ^b	15.1 ^a	15.3 ^a	0.9	< 0.01	< 0.01	0.41
Feeding rate (kg/min)	0.22	0.20	0.22	0.23	0.03	0.44	0.11	< 0.01
Meals (n/day)	0.15	7.17 ^b	7.38 ^b	8.24 ^a	0.49	0.02	0.62	0.64
Visits (n/day)	0.33	23.7	28.5	31.7	4.1	0.08	0.04	< 0.01
Meal time (min/day)	0.25	200	224	225	22	0.39	0.57	< 0.01
Lying time (h/day)	0.12	10.3	10.9	11.9	0.8	0.06	0.26	0.32
Lying bouts (n/day)	0.07	11.6	12.0	13.2	1.5	0.40	0.84	0.16
Steps (n/day)	0.18	1100	1231	1324	134	0.21	0.73	0.01
Motion	0.20	4409	4945	5386	542	0.15	0.51	0.01

C = cluster; DP = dry period length; P = parity; DM = dry matter.

Metabolic status was based on the plasma free-fatty acids, β -hydroxybutyrate, glucose, insulin, IGF-1 and bovine growth hormone concentration. Values are presented as LSM and pooled SE (SEp).^{a,b,c}Values with different superscripts differ ($P < 0.05$).

Cows with good or average metabolic status had greater DMI and basal ration intake than cows with a poor metabolic status ($P < 0.01$) (Table 4). Cows with a good metabolic status had more meals per day than cows with an average or poor metabolic status ($P < 0.05$). Cows with a good metabolic status tended to have a longer lying time than cows with a poor metabolic status ($P < 0.10$). Irrespective of the metabolic cluster, cows with a 0-day DP had greater DMI (21.9 ± 0.3 v. 20.5 ± 0.5 kg/day), basal ration intake (15.0 ± 0.3 v. 13.5 ± 0.4 kg DM/day) and visits to the feeder (29.0 ± 1.5 v. 24.2 ± 1.9 /day) than cows with a 30-day DP ($P < 0.05$). Cows of parity 2 had a lower feeding rate (0.20 ± 0.01 v. 0.24 ± 0.01 kg/min), and more visits to the feeder (29.7 ± 1.7 v. 23.1 ± 1.6 /day), longer meal times (233 ± 9 v. 195 ± 9 min/day) and more steps

(1280 ± 57 v. 1085 ± 56 /day) and motion index (5234 ± 233 v. 4464 ± 230) ($P < 0.01$) than cows with parity ≥ 3 .

Discussion

The aim of this study was to analyze the relationships between metabolic status, based on plasma metabolites and metabolic hormones, and feeding behavior, lying behavior, steps and motion index of dairy cows in week 4 *postpartum*. First, Pearson's correlation analysis was used to correlate metabolites and metabolic hormones with behavior. In particular, plasma FFA concentration was related to the behavioral indicators. Dry matter intake, number of meals and

visits to the feeder, meal times, lying times, steps and motion index were all lower at a greater plasma FFA concentration. Plasma BHB, glucose, insulin, IGF-1 and GH concentration were related to only three or fewer behavioral indicators, and the strength of these correlations was generally lower than between FFA concentration and behavioral indicators. This seems to indicate that cows with a greater FFA concentration, in particular, have a lower feed intake, partly explained by lower feed intake related behavior and lower activity. Our results are in-line with previous studies that found that cows with a greater plasma FFA concentration have lower feed intake (Waterman *et al.*, 1972; Lean *et al.*, 1992) and lower walking activity (Adewuyi *et al.*, 2006). Because in the current study most metabolites and hormones were correlated (Table 1b), as a second step, cluster analysis was performed to cluster cows for metabolic status based on their plasma FFA, BHB, glucose, insulin, IGF-1 and GH concentration. Cows were clustered in four groups for poor, average, good or very good metabolic status. Plasma IGF-1 concentration seemed to have the largest impact on clustering ($R^2 = 0.82$), followed by plasma FFA concentration ($R^2 = 0.30$) and glucose ($R^2 = 0.23$). Correlations with individual metabolites and metabolic hormones and behavior, however, indicated that plasma FFA concentration had more and stronger correlations with feeding behavior, lying behavior, steps and motion index than IGF-1. This might indicate that high plasma FFA is more related to behavior and feelings in week 4 of lactation than plasma IGF-1, possibly because cows did not recover from the catabolic state at this time.

To our knowledge, no other studies have directly related metabolic status with behavioral parameters. However, previous studies evaluated the relationships between disease events in early lactation and metabolic status and reported plasma metabolites for the diseased and non-diseased cows. These studies reported that cows with a metabolic disease, metritis or mastitis had lower metabolic status as reflected by a greater plasma FFA, BHB and haptoglobin, and lower plasma calcium concentration, than non-diseased cows (Soriani *et al.*, 2012). In addition, cows with metabolic disease in previous studies had lower daily feeding time (Urton *et al.*, 2005), decreased activity and rumination (Soriani *et al.*, 2012, Stangaferro *et al.*, 2016), compared with non-diseased cows. In another study, cows with ketosis had lower feed intakes, feeding times, meal times and fewer meals and feeder visits than cows without ketosis (González *et al.*, 2008). The lower reported DMI, and number of meals and visits to the feeder in these earlier studies are in accordance to our findings. In contrast, in our study, steps, motion index and meal time were not different among cows with different metabolic status. It should, however, be noted that in previous studies cows were diseased and likely showed sickness behavior, whereas cows in our study were not clinically diseased. The current study illustrates how behavior measures like a number of meals per day and lying time can be an indicator for metabolic status, even in case of no clinical disease.

On average, cows with a good metabolic status had a lower plasma FFA concentration and greater plasma glucose and IGF-1 concentrations compared with cows with an average or poor metabolic status. Better metabolic status was associated with greater DMI, basal ration intake and a number of meals, and tended to be associated with a longer daily lying time. These behaviors can be automatically recorded and may be reliable indicators of metabolic status. A number of visits to the feeder can be measured as proximity to the feed bunk using sensors, and can be processed into a number of meals. However, these sensors are unlikely to be used at present in commercial settings. Meanwhile, lying time can be recorded with commercially available sensors that are widely used for estrus detection (Rutten *et al.*, 2017). Lying time may, therefore, be a useful indicator for metabolic status, although the relation should be studied further, as well as potential interactions with other factors such as stocking density. Reduced lying time may be indicative of restlessness or discomfort, for example, because of udder pressure (Huzzey *et al.*, 2005; Bertulat *et al.*, 2017).

In the current study, plasma FFA concentration had stronger correlations and plasma IGF-1 concentration had similar correlations with behavior in week 4 of lactation, compared with correlations between EB and behavior in our previous study (Kok *et al.*, 2016). These results imply that plasma FFA has a stronger relation with behavior than calculated EB. It could be hypothesized, therefore, that plasma metabolites like FFA may better reflect cow health and feelings than the EB itself, because metabolic status better reflects the degree to which the animal can cope with the catabolic state typical in early lactation. In addition, it could be hypothesized that the relation between behavior and plasma metabolites may even be stronger in cows with a conventional dry period due to a more severe NEB and poorer metabolic status, than in cows with a 0-day or 30-day dry period.

Relations between metabolic status and behavior can be different for other weeks in early lactation. Daily lying time, for example, was shown to decrease in the weeks after calving, with the lowest lying time in week 4, after which lying time gradually increased and then leveled off at about 12.5 h per day (Maselyne *et al.*, 2017). Plasma FFA concentration, in contrast, peaks in the 1st week after calving (Chen *et al.*, 2015). Together, these two patterns imply that associations between plasma FFA concentration and daily lying time are different over time. Variable relations between metabolic status and behavior may be explained by the duration of the catabolic state and interactions with other motivations. For example, when cows were given access to feed and a lying place for only 12 h/day for a period of 2 weeks, they prioritized lying time over feed intake and consequently lost weight (Munksgaard *et al.*, 2005). The authors hypothesized that, if the experiment had lasted longer, a more severe weight loss could shift these behavioral time budgets toward feeding at the cost of lying time.

The application of different dry period lengths was used to create variation in metabolic status between cows in early lactation. A potential drawback of this method could be that short or no dry period affect behavior independent of metabolic status. However, the model to evaluate the effect of metabolic status on behavior corrected for effects of dry period length and parity. Correlations between metabolites and metabolic hormones and behavior were performed without dry period length and parity included as covariates in the model. Including dry period length and parity in a GLM to evaluate the effects of a plasma metabolite on behavior (results not shown) resulted in similar relations between plasma metabolites and behavior compared with the correlations presented in Table 1a.

The direction of causation between plasma metabolites or metabolic status with behavioral traits in dairy cows is unclear. For example, less feed-directed behavior and activity may result in lower DMI and more body fat mobilization, which is related to a greater plasma FFA and BHB concentration and a lower glucose and insulin concentration (Lean *et al.*, 1992). Vice versa, greater FFA concentration is related with metabolic acidosis and a feeling of discomfort that may decrease appetite and DMI (Waterman *et al.*, 1972; Allen *et al.*, 2009).

In conclusion, better metabolic status, as indicated by plasma metabolites and metabolic hormones, in dairy cows in week 4 of lactation after a short or no dry period was associated with a greater DMI, increased feeding activity and a tendency to more time spent lying, compared with poor metabolic status. A compromised metabolic status was reflected in altered cow's behavior in week 4 of lactation.

Acknowledgments

The authors thank DairyNL (Zuivel-NL; organization of the Dutch dairy supply chain, the Netherlands) and the Dutch Ministry of Economic Affairs (EZ, the Netherlands) for financing this study, which is part of the Public–Private Partnership 'Sustainable Dairy Chain' (Duurzame Zuivelketen), the Netherlands. The authors also thank Dairy Campus (Lelystad, the Netherlands), Rudi Koopmanschap and the Veterinary Physiology Group of the University of Bern (Switzerland) for their technical support and laboratory analyses during the experiment. This study was conducted as part of the PhD theses of the first two authors.

Declaration of interest

None.

Ethics statement

The Institutional Animal Care and Use Committee of Wageningen University and Research approved the experimental protocol in compliance with the Dutch law on Animal Experimentation (protocol no. 2014125).

Software and data repository resources

None.

References

- Adeuyi AA, Roelofs JB, Gruys E, Toussaint MJM and van Eerdenburg FJCM 2006. Relationship of plasma nonesterified fatty acids and walking activity in postpartum dairy cows. *Journal of Dairy Science* 89, 2977–2979.
- Allen MS, Bradford BJ and Oba M 2009. Board-invited review: the hepatic oxidation theory of the control of feed intake and its application to ruminants. *Journal of Animal Science* 87, 3317–3334.
- Bertulat S, Isaka N, de Prado A, Lopez A, Hetreau T and Heuwieser W 2017. Effect of a single injection of cabergoline at dry off on udder characteristics in high-yielding dairy cows. *Journal of Dairy Science* 100, 3220–3232.
- Butler S, Marr A, Pelton S, Radcliff R, Lucy M and Butler W 2003. Insulin restores GH responsiveness during lactation-induced negative energy balance in dairy cattle: effects on expression of IGF-I and GH receptor 1A. *Journal of Endocrinology* 176, 205–217.
- Chen J, Gross JJ, van Dorland HA, Remmelink GJ, Bruckmaier RM, Kemp B and van Knegsel ATM 2015. Effects of dry period length and dietary energy source on metabolic status and hepatic gene expression of dairy cows in early lactation. *Journal of Dairy Science* 98, 1033–1045.
- Collard BL, Boettcher PJ, Dekkers JCM, Petitclerc D and Schaeffer LR 2000. Relationships between energy balance and health traits of dairy cattle in early lactation. *Journal of Dairy Science* 83, 2683–2690.
- Centraal Veevoederbureau 2016. Tabellenboek Veevoeding. Voedernormen Rundvee, Schapen en Geiten, en voederwaarden voermiddelen herkauwers (in Dutch). The Hague, The Netherlands.
- Drackley JK 1999. Biology of dairy cows during the transition period: the final frontier? *Journal of Dairy Science* 82, 2259–2273.
- Edwards JL and Tozer PR 2004. Using activity and milk yield as predictors of fresh cow disorders. *Journal of Dairy Science* 87, 524–531.
- Fenwick MA, Llewellyn S, Fitzpatrick R, Kenny DA, Murphy JJ, Patton J and Wathes DC 2008. Negative energy balance in dairy cows is associated with specific changes in IGF-binding protein expression in the oviduct. *Reproduction* 135, 63–75.
- Fourichon C, Seegers H, Bareille N and Beaudeau F 1999. Effects of disease on milk production in the dairy cow: a review. *Preventive Veterinary Medicine* 41, 1–35.
- Graber M, Kohler S, Muller A, Burgermeister K, Kaufman T, Bruckmaier RM and Van Dorland HA 2012. Identification of plasma and hepatic parameters related to metabolic robustness in dairy cows. *Journal of Animal Physiology and Animal Nutrition* 96, 75–84.
- González LA, Tolkamp BJ, Coffey MP, Ferret A and Kyriazakis I 2008. Changes in feeding behavior as possible indicators for the automatic monitoring of health disorders in dairy cows. *Journal of Dairy Science* 91, 1017–1028.
- Huzzey JM, Von Keyserlingk MAG and Weary DM 2005. Changes in feeding, drinking and standing behavior in dairy cows during the transition period. *Journal of Dairy Science* 88, 2454–2461.
- Ingvartsen KL, Dewhurst RJ and Friggens NC 2003. On the relationship between lactational performance and health: is it yield or metabolic imbalance that cause production diseases in dairy cattle? A position paper. *Livestock Production Science* 83, 277–308.
- Kok A, van Knegsel ATM, van Middelaar CE, Hogeveen H, Kemp B and de Boer IJM 2015. Technical note: Validation of sensor-recorded lying bouts in lactating dairy cows using a 2-sensor approach. *Journal of Dairy Science* 98, 7911–7916.
- Kok A, Van Hoesel RJ, Tolkamp BJ, Haskell MJ, Van Knegsel ATM, De Boer IJM and Bokkers EA 2016. Behavioural adaptation to a short or no dry period in dairy cows. *Applied Animal Behaviour Science* 186, 7–15.
- Lean IJ, Farver TB, Trout HF, Bruss ML, Galland JC, Baldwin RL, Holmberg CA and Weaver LD 1992. Time series cross-correlation analysis of postparturient relationships among serum metabolites and yield variables in Holstein cows. *Journal of Dairy Science* 75, 1891–1900.
- Maselyne J, Pastell M, Thomsen PT, Thorup VM, Hänninen L, Vangoyte J, Van Nuffel A and Munksgaard L 2017. Daily lying time, motion index and step frequency in dairy cows change throughout lactation. *Research in Veterinary Science* 110, 1–3.
- Munksgaard L, Jensen M.B., Pedersen L.J., Hansen S.W. and Matthews L. 2005. Quantifying behavioural priorities – effects of time constraints on behaviour of dairy cows, Bos *taurus*. *Applied Animal Behavior Science* 92, 3–14.
- Rastani RR, Grummer RR, Bertics SJ, Gümen A, Wiltbank MC, Mashek DG and Schwab MC 2005. Reducing dry period length to simplify feeding transition cows: milk production, energy balance, and metabolic profiles. *Journal of Dairy Science* 88, 1004–1014.

- Reith S, Brandt H and Hoy S 2014. Simultaneous analysis of activity and rumination time, based on collar-mounted sensor technology, of dairy cows over the peri-estrus period. *Livestock Science* 170, 219–227.
- Rutten CJ, Kamphuis C, Hogeveen H, Huijps K, Nielen M and Steeneveld W 2017. Sensor data on cow activity, rumination, and ear temperature improve prediction of the start of calving in dairy cows. *Computers and Electronics in Agriculture* 132, 108–118.
- Soriani N, Trevisi E and Calamari L 2012. Relationships between rumination time, metabolic conditions, and health status in dairy cows during the transition period. *Journal of Animal Science* 90, 4544–4554.
- Stangaferro ML, Wijma R, Caixeta LS, Al-Abri MA and Giordano JO 2016. Use of rumination and activity monitoring for the identification of dairy cows with health disorders: Part I. Metabolic and digestive disorders. *Journal of Dairy Science* 99, 7395–7410.
- Tolkamp BJ, Friggens NC, Emmans GC, Kyriazakis I and Oldham JD 2002. Meal patterns of dairy cows consuming mixed foods with a high or a low ratio of concentrate to grass silage. *Animal Science* 74, 369–382.
- Urton G, von Keyserlingk MAG and Weary DM 2005. Feeding behavior identifies dairy cows at risk for metritis. *Journal of Dairy Science* 88, 2843–2849.
- Van Es AJH 1975. Feed evaluation for dairy cows. *Livestock Production Science* 40, 95–107.
- Van Hoeij RJ, Dijkstra J, Bruckmaier RM, Gross JJ, TJGM Lam, Rummelink GJ, Kemp B and Van Knegsel ATM 2017. The effect of dry period length and postpartum level of concentrate on milk production, energy balance, and plasma metabolites of dairy cows across the dry period and in early lactation. *Journal of Dairy Science* 100, 5863–5879.
- Van Knegsel ATM, Hammon HM, Bernabucci U, Bertoni G, Bruckmaier RM, Goselink RMA, Gross JJ, Kuhla B, Metges CC, Parmentier HK, Trevisi E, Tröschner A and Van Vuuren AM 2014b. Metabolic adaptation during early lactation: key to cow health, longevity and a sustainably dairy production chain. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 9, 1–15.
- Van Knegsel ATM, Rummelink GJ, Jorjone S, Fievez V and Kemp B 2014a. Effect of dry period length and dietary energy source on energy balance, milk yield, and milk composition of dairy cows. *Journal of Dairy Science* 97, 1499–1512.
- Van Winden SCL, Jorritsma R, Müller KE and Noordhuizen JPTM 2003. Feed intake, milk yield, and metabolic parameters prior to left displaced abomasum in dairy cows. *Journal of Dairy Science* 86, 1465–1471.
- Vicari T, van den Borne JJGC, Gerrits WJJ, Zbinden Y and Blum JW 2008. Postprandial blood hormone and metabolite concentrations influenced by feeding frequency and feeding level in veal calves. *Domestic Animal Endocrinology* 34, 74–88.
- Waterman R, Schwalm JW and Schultz LH 1972. Nicotinic acid treatment of bovine ketosis I. Effects on circulatory metabolites and interrelationships. *Journal of Dairy Science* 55, 1447–1453.
- Yeates MP, Tolkamp BJ, Allcroft DJ and Kyriazakis I 2001. The use of mixed distribution models to determine bout criteria for analysis of animal behaviour. *Journal of Theoretical Biology* 213, 413–425.