

The Vegetation of Jamaica

G. F. ASPREY AND R. G. ROBBINS
University College of West Indies

[Reprinted from ECOLOGICAL MONOGRAPHS, 23: 359-412. October, 1953]

Scanned from original by ISRIC – World Soil Information, as ICSU World Data Centre for Soils. The purpose is to make a safe depository for endangered documents and to make the accrued information available for consultation, following Fair Use Guidelines. Every effort is taken to respect Copyright of the materials within the archives where the identification of the Copyright holder is clear and, where feasible, to contact the originators. For questions please contact soil.isric@wur.nl indicating the item reference number concerned.

THE VEGETATION OF JAMAICA

G. F. ASPREY & R. G. ROBBINS
University College of the West Indies




TABLE OF CONTENTS

	PAGE		PAGE
INTRODUCTION	359	Alluvium	388
GENERAL DESCRIPTION OF JAMAICA	360	Vegetation of the Alluvial Plains	388
History and Land Settlement	360	Cultivated areas and induced savanna	388
Geology and Soils	361	Secondary communities	389
Climate	362	Thorn Scrub	389
FLORISTIC STUDIES	363	Swamps and Marshes	390
Analysis of the Flora	363	Herbaceous Swamp	390
Phytogeography	365	Mangrove Woodland	391
COMMUNITY STUDIES	370	Palm-Sedge Marsh	392
Introduction	370	Marsh Forest	392
Coastal Communities	371	Shale	393
Marine Vegetation	371	Vegetation on the Lower Shale Hills	393
Strand Communities	372	Montane Communities	394
Sand Spits	374	Lower Montane Rain Forest	394
Cays	376	Montane Sclerophyll	397
Lowland Communities	379	Montane Mist Forest	398
Limestone	379	Elfin Woodland	401
Dry Limestone Scrub Forest	380	SUMMARY AND DISCUSSION	403
Wet Limestone Forest	384	LITERATURE CITED	406
Vegetation of the bauxite plateaus	386	APPENDIX	
		Check-List of Plants mentioned in Text	408

INTRODUCTION

Plant ecology in the Caribbean has scarcely entered the broad phase of general descriptive accounts and the stage of intensive local analysis and experimental approach has not yet been reached. This is only to be expected since there are few ecologists available and often the taxonomy of whole floras is incomplete. The first step, then, is often one of descriptive phytogeography.

A number of the Central American Republics either possess recent floras or have been briefly covered in ecological papers, for example those of Miranda & Sharpe (1950), Standley (1941), Stevenson (1942) and Steyermark (1950). A general account of the area may be obtained from contributions to "Plants and Plant Science in Latin America" (Verdoorn 1945).

Of the West Indian Islands, Trinidad and Tobago have been described by Beard (1944a, 1946) but, geographically and botanically, they belong to the mainland (Venezuela). The same author (1949) covered the Lesser Antilles while the French Islands

(Guadeloupe and dependencies) had already received attention by Stehlé (1935-1945). Howard (1952) has given an account of the vegetation of the Grenadines.

Turning to the Greater Antilles which form a unit of closer affinities, the plant life of Cuba has received attention from Marie-Victorin & Leon (1942), Leon (1946), Leon & Alain (1951), Carabia (1943, 1945a), and Seifriz (1943).

Porto Rico forms the subject of a complete scientific survey to which Britton & Wilson (1923-1926) contributed the flora and Gleason & Cook (1927) the plant ecology.

Hispaniola, including Haiti and Dominican Republic, appears virtually neglected except for the short report by Holdridge (Verdoorn 1945).

Jamaica, which forms the subject of the present paper, has received only sporadic and localized study. It is obvious that a more complete evaluation of the vegetation of Jamaica is long over-due. The establishment of the new University College of the

ISN 27141

West Indies has now added a further incentive as well as being a centre for such studies.

In Jamaica, as indeed in the whole of the West Indies, there are two major aspects to be studied; the flora and the plant communities. It is under these headings that the present study is presented. While primarily concerned with describing the vegetation of Jamaica as such, some attempt is made to fit the various aspects into the whole Caribbean field by discussing the plant geographical and ecological affinities.

The ecological aspects are presented on a broad physiognomic basis and, whilst a detailed investigation of Jamaican vegetation cannot be claimed, the units described should form a sound and well-defined framework for future ecological studies of a more intensive nature. Where possible, such future lines of research have been suggested.

Acknowledgment for information and assistance is made to the following: Mr. I. B. Williams, Botany Department, University College of the West Indies, Jamaica; Mr. C. B. Lewis & Mr. G. R. Proctor of the Institute of Jamaica; Mr. K. Vernon, Soil Survey, Jamaica; Mr. W. J. Fowler, Meteorological Office, Jamaica; Mr. E. M. Brown, Forestry Department, Jamaica; Dr. L. Chubb, Geological Survey, Jamaica; Mr. C. W. Hewitt, Dept. of Agriculture, Jamaica; The Director of Surveys, Survey Dept., Jamaica; Dr. Archie Carr, University of Florida (Fig. 18, 36, 37); Mr. A. Packer, Kingston, Jamaica (Fig. 21, 29). Thanks are also due to Dr. A. R. Loveless for reading through the manuscript.

GENERAL DESCRIPTION OF JAMAICA

Lying in the tropical zone some 18 degrees north of the equator, Jamaica forms part of the string of Caribbean Islands stretching in an arc from Florida to Venezuela. It is the largest of the British West Indian Islands, being 144 mi (231 km) in length running from east to west, and some 30-40 mi (48-64 km) wide. It has an area of 4,411 sq mi (11,740 sq km). With Cuba, Hispaniola (Haiti and the Dominican Republic) and Porto Rico, Jamaica forms




FIG. 1. Tropical America and the Caribbean Area.

a natural group of islands known as the Greater Antilles (Fig. 1).

The island is dominated by its mountains which reach their maximum height of 7,402 ft. (2,256 m) at Blue Mountain Peak. More than half of the country is above 1,000 ft. in elevation (Fig. 2). The greater part of this area is taken up by the central upland plateaus at 2,000-3,000 ft., and the Blue Mountain range in the east. In the south, the low-lying coastal fringe includes broad, flat, alluvial plains.




FIG. 2. Topographical regions of Jamaica. The Blue Mountain Range in the east reaches a height of 7,402 ft.

HISTORY AND LAND SETTLEMENT

Four hundred years of European occupation have wrought a tremendous change in the vegetation. Forests now only occur in the wet limestone country of the interior, the higher elevations of the Blue Mountains in the east, the dry limestone hills and the mangrove swamps (Fig. 3).

Occupied by the British in 1665, there followed a period of agricultural expansion which was intensified during the sugar boom of the 18th. century. The lowlands were cleared for sugar cane and the slaves given cultivation rights on marginal lands and foothills. Seasonal burning by temporary tenants for their shifting patchwork cultivation has reduced much of the forest land to a state of second growth scrub, locally known as ruinate. In early colonial times, mahogany was exported in large quantities followed by such cabinet and dye woods as fustic (*Chlorophora tinctoria*), satinwood (*Zanthoxylum flavum*), sandalwood (*Amyris balsamifera*) and lignum vitae (*Guaiacum officinale*). Today, Jamaica provides but a fraction of its own lumber requirements. Early logging methods had no regard to conservation and appalling forest destruction has been wrought by the removal of bark from valuable trees for such minor products as dye chips and tannin.

Jamaica now has 18% of land area in forests and 47% in agricultural use. The remainder, 35%, consists of second growth scrub (ruinate), thorn scrub, mangrove swamp and so on. Of the agricultural areas, 8% can be classified as tree crops (cocoa, coffee, pimento, citrus, coconuts and bananas) and 39% as food crops, plantation crops (sugar and rice) and pasture. (Summarized from returns to F.A.O. 1951.)




FIG. 3. Vegetation map of Jamaica showing the distribution of forests and major plantation crops. Place names mentioned in the text are also included; the parishes are underlined.

GEOLOGY AND SOILS

The geological history of Jamaica began in upper Cretaceous times when volcanic activity occurred. This was probably submarine as the volcanic tuffs and lavas, the Trappian Series (Sawkins 1869) are interbedded with shallow-water, marine shales and reef limestones. At the end of the Cretaceous period, uplift converted Jamaica into land. Pronounced earth-movements with folding, thrusting and faulting, accompanied the uplift and continued into lower Eocene times but had ceased by middle Eocene. During Eocene times, gradual subsidence supervened and, by middle Eocene, large areas of the island were submerged beneath a shallow sea. The sea extended progressively until, by the Oligocene, probably the whole island was submerged. Renewed uplift began in the lower Miocene but probably no land appeared until the middle Miocene when there was considerable uplift so that only the margins of the present island remained beneath the sea. This uplift was accompanied by renewed folding and faulting on a less pronounced scale, with localized exceptions where folding was intense. Later, uplift brought the island to its present form.

Broadly speaking, Jamaica has an igneous and metamorphic core, covered, for the greater part, by a limestone mantle deposited during several marine submergences. The surface consists of approximately two-thirds limestone with the other third of igneous rock, sedimentary shales and alluvium.

Hard white limestone up to 2,000 ft. thick forms some three-quarters of the surface rock and may reach an elevation of 3,000 ft. Softer yellow limestone and marl occur in some areas and are often present as a narrow bordering strip around shale areas.

Soil-forming processes and erosion are active in Jamaica. On the one hand, there is natural geological erosion and weathering associated with the relatively youthful topography of the island. These factors are still active and have produced the sharp ridge topography characteristic of the shale hills of St. An-

drew. On the other hand, there is the steady loss of topsoil by leaching and erosion brought about by man's activities. Jamaica shares this common problem with other tropical countries where agriculture has been established on previous forest lands. Erosion is amplified by the steep topography, high temperature and torrential rains. Where the forest is left intact, an equilibrium is established and the soil type remains more or less stable. Due to slope and the porosity of the soil, drainage is rapid but the forest cover, so long as it protects the profile, remains. Once the forest cover is removed, however, steady leaching begins in the upper horizons with loss of nutrients. Exposure brings further erosion and often the whole of the upper horizon is missing, leaving a truncated profile. Here, cultivation is literally carried out on the lower horizons of weathering parent rock. Indeed, it may not be an overstatement that, in many parts of Jamaica, the soils approximate very nearly to the underlying geological rock pattern because of these facts. The condition is particularly prevalent in the shale areas. The eroded material is no longer deposited in valleys or flood plains but scoured out to sea during the sudden rainy seasons by swollen rivers passing through narrow valleys.

The major soil types of Jamaica will be described




FIG. 4. The distribution of the major soil types of Jamaica and the geological formation from which they have been derived.

under three headings: Soils of the Highlands, Soils of the Upland Plateaus and Soils of the Alluvial Plains. Their distribution is shown in Fig. 4.

SOILS OF THE HIGHLANDS

In Jamaica, the highlands are made up of ancient, acid, igneous and volcanic rocks, the greater part of which are above 3,000 ft., and form the shale areas, the most notable being the Blue Mountains. The soils which may be called lithosols are very young so that they can be classified according to the type of geological formation. Their porosity is great so that leaching is heavy, leading to acid conditions and a paucity of nutrients. Under forest conditions, a high organic content is built up in the surface layers and a soil profile is developed of which the following is typical: The first 6 in. consists of a dark brown, gravelly, sandy loam. This horizon is missing when the forest is denuded. From 6-33", a dark brown, sandy, gravelly loam with a high percentage of gravel. The latter increases until solid rock is reached.

Of more sporadic occurrence, are clays produced by the weathering of fine-grained, sedimentary rocks. Drainage here is poor and, on the whole, the soils are less acid. Here the first 15" consists of dark-brown, clay loam, richly stained with humus. This overlies a yellow-red, gravelly, sandy clay loam which may extend to a depth of 40" or more and gives way to a lighter, gravelly loam and finally to the weathered parent Richmond shale.

Both types of soil are liable to rapid surface erosion following removal of the forest, leading to a truncated profile with very little other than subsoil which is extremely poor in nutrients. Such soils may be found in both the western and central shale areas. The igneous area shown on the map is composed of granodiorite and reaches an elevation of 1,500 ft. Clearing of the original forest vegetation has led to rapid erosion and the surface is now comprised of rotten rock debris, low in available nutrients and extremely porous.

SOILS OF THE UPLAND PLATEAUS

These soils are derived from limestone and fall into two categories: Terra Rossa or Red Limestone Soils and Rendzinas or Black Marl Soils.

Terra Rossa is the name applied to the residual bauxitic soils which occur mainly on the upland plateaus of Manchester and St. Ann at an elevation of 2,000-3,000 ft. They are developed from the weathering of hard white limestone through solution and are typically coarse and porous in texture. Although under conditions of poor drainage, degraded forms of these bauxitic soils occur, they are usually much leached, acid, well oxidised and dehydrated. The red colour is due to ferric oxide and their depth over the limestone plateaus varies greatly. There is no distinctive profile but the surface has a high content of organic matter on which the agricultural value of the land depends.

The black marl soils or rendzinas are developed over soft, yellow limestones and marls. Such soils are of sporadic occurrence. They develop a typical,

mature profile in which the A horizon (0-6") is a heavy black or brown clay followed by a transitional zone (6-12") leading to a B horizon of light brown or yellow calcareous clay. This layer finally grades into the parent material. The soil is fine and heavy with poor drainage. High calcium status and wetness prevents development of the red colour typical of the bauxitic soils. Phosphorous is firmly bound but potash is adequate and available.

SOILS OF THE ALLUVIAL PLAINS

These soils are found on the extensive, southern, coastal plains. They have been deposited by rivers on the low-shelving coast, and are made up of loam, sand and gravel. Areas also occur where the surface alluvium is of marine origin. Here, heavy clays from 3-4 ft. deep, have been deposited over the normal riverine alluvium. The most useful agricultural land in Jamaica is found on the alluvial plains. For further information on the soils of the Caribbean, reference should be made to Hardy (1945).

CLIMATE

Lying 18½ degrees north of the equator, Jamaica is in the tropical zone. Climatic conditions are typical of a Caribbean island with a rainy, windward coast, a dry leeward coast and a cool, central, montane region. Local climate is insular with changes in land and sea breezes during the day and night and an equable temperature throughout the year.

Due to the varying altitude from sea-level to over 7,000 ft., in the eastern end, there is a wide difference in mean temperature amounting to 23°F. The mean annual, average temperature at Kingston on the dry, south coast is 79°F and, on Blue Mt. Peak, 56°F. Generally, the coast has a greater range (17°F) than the elevated inland (12°F at 5,000 ft.). Kingston has the coolest month in February (aver. 75.7°F) and the hottest month in July (aver. 81.4°F). The highest maximum is 97.8°F and the lowest minimum 56.7°F.

Rainfall is the most important single factor affecting the vegetation. The island lies in the path of the moisture-laden, easterly trade winds which blow consistently throughout the year. Striking first the high limestone John Crow Mountains and then the northern flanks of the Blue Mountains, they are forced upwards and cooled when excess moisture is precipitated. Hence, Portland in the north east is one of the wettest parishes with over 150 in. in contrast to the dry, southern St. Andrew parish with 35 in.

The rainfall distribution map (Fig. 5) shows that about three-quarters of the island receives an annual rainfall ranging above 60 in. Seventy year records for the island as a whole, give the average, total, annual rainfall as 77 in. during an average of 123 wet days.

A feature of the rainfall is its seasonal periodicity. The island mean shows that the wettest months are October and May. The major dry period is from January to March.

Charter (1941) and Beard (1944a) emphasize the importance of the number of months in the year in




FIG. 5. Rainfall distribution in Jamaica compiled from the 70-year records. The graphs show seasonal periodicity of the rainfall resulting in maximum in May and October and a major dry period from January to March.

which excessive evaporation over precipitation leads to drought. For the Caribbean area, this point is reached with a monthly fall of less than 4 in. (100 mm) with soil of normal porosity. Seasonal distribution for Jamaica upon such a basis is given in Fig. 6.




FIG. 6. Seasonal rainfall in Jamaica showing the four types of distribution. A rainfall of less than 4" per month is considered to produce drought conditions. Compiled by C. W. Hewitt, Dept. of Agriculture, Jamaica, B. W. I.

The mountainous nature of Jamaica results in a variety of local winds. Easterly Trade Winds blow constantly and constitute 89% of the winds over 15 m.p.h. recorded at Kingston. At the end of the year, north winds sweep down from Cuba. Hurricanes, which originate in the area of the Lesser Antilles to the east, develop during August, September and October and, on the average, Jamaica may expect to be on the path of a hurricane once in 8 years.

The shortest day (Dec. 21st.) is 11 hrs., and the longest (June 21st.), 13 hours. Jamaica records 60% sunshine at Kingston compared with 33% for London. Some areas inland which are low-lying (1,000 ft.) are subject to night and early morning fogs, while the Blue Mountains have daily mists down to 3,000 ft.

FLORISTIC STUDIES

The long list of distinguished botanists that have visited Jamaica shows that the island was one of the earliest and best botanised areas in the Caribbean.

The first records of the plants of Jamaica were published by Sir Hans Sloane (1669, 1707, 1725). Then followed contributions by Brown (1756), Long (1774) Lunan (1814) and MacFadyen (1837). A flora of the British West Indies published by Grisebach (1864) remained for many years the most complete study of the island's plants. It is still an important reference point for many groups. Five volumes of the Flora of Jamaica by Fawcett & Rendle appeared over the period from 1910-1936. The monocotyledons are represented by a single volume on the orchids, while some 21 families belonging to the Metachlamydeae have still to be described. Such a brief review cannot be left without mentioning Urban's *Symbolae Antillanae* (1898). The location of the important Jamaica herbaria are listed by J. Lanjouw (1945). The important collection by Harris is now in the possession of the University College of the West Indies. In spite of the numerous collectors, there are still comparatively large areas unknown botanically. This has been borne out in the recent work of Proctor (1953). Working on the Pteridophytes, he has collected abundantly of species listed as rare or known only from the type specimen.

ANALYSIS OF THE FLORA

Excluding the fungi and the lichens it is estimated that Jamaica has a total flora of some 4,000 plants. Cryptogams number about 1,500 (algae 350, bryophytes 600, pteridophytes 525). Phanerogams are just over 2,500 divided up as follows: gymnosperms 5, angiosperms 2,500 with 600 monocotyledons and 1,900 dicotyledons.

The monocotyledons number about 29 families and 213 genera with 600 species. The orchids and grasses alone account for about 400 of these species. Many of the 200 odd grasses are introductions. Of the orchids, embracing 61 genera, only one genus is endemic. In all, there are 194 species (Cuba 270), 31 of which belong to the genus *Epidendrum*. Sedges are well represented in Jamaica but are in need of revision. Bromeliaceae, listed in Grisebach as 26 species, now include 66 species in 12 genera. There are 7 genera with 13 species of indigenous palms which may be compared with Cuba's 80 species. Araceae has about 15 species in 7 genera while all the other monocotyledon families are small.

Of the 76 dicotyledon orders given in Hutchinson (1926), 62 are present in Jamaica, and contain 128 families, 647 genera and 1907 species (Tables 1, 1A, & 2).

TABLE 1. Analysis of the Flora. Orders of Dicotyledons.

Orders	Families	Genera	Species
Annonales	1	3	14
Apocynales	2	21 & 2*	49 & 3
Aristolochiales	1	1	5
Asterales	1	50 & 9	126 & 13
Berberidales	1	2	4
Bixales	4	10	22
Boraginales	1	7	36
Cactales	1	6	15
Campanales	3	4	13
Capparidales	2 & 1	6 & 1	16 & 2
Caryophyllales	4	12	18
Celestiales	5	10	25
Chenopodiales	5	23	38
Cruciales	1	6 & 2	7 & 3
Cucurbitales	3	9 & 4	16 & 6
Cunoniales	2	2	2
Dilleniales	1	2	2
Ebenales	2	8	27
Ericales	2	3	5
Euphorbiales	1	30 & 2	111 & 8
Garryales	1	1	1
Gentianales	1	7	13
Geraniales	4	6	7
Guttiferales	3	7	14
Hamamelidales	1	1	4
Lamiales	2	26 & 3	67 & 4
Laurales	2	8	20 & 1
Leguminosae	3	50 & 16	149 & 54
Loasales	2	3	5
Loganales	2	6	14
Lythrales	2	7	19 & 4
Malpighiales	2	10 & 1	31 & 1
Malvales	1	15	69 & 9
Meliaceles	1	4 & 1	9 & 1
Myricales	1	1	3
Myrtales	5	33	143 & 4
Myrsinales	1	5	28
Olacales	1	2	5
Passiflorales	1	1	14 & 3
Personales	5	54 & 3	113 & 7
Piperales	3	4	55
Plantaginales	1	1	3
Polemoniales	1	2	3
Polygonales	1	3 & 1	26 & 2
Polygales	1	2	6
Primulales	2	2	2
Ranales	3	5	9
Rhamnales	2	10	15 & 2
Rhoeadales	1	2	2
Rosales	1	5 & 1	9 & 6
Rubiales	2	42 & 4	147 & 7
Rutales	3	14 & 3	33 & 10
Santalales	2	7	22
Sapindales	4	17 & 3	36 & 4
Saxifragales	1	1	1
Solanales	2	19 & 1	98 & 1
Styracales	1	1	2
Theales	3	7 & 1	16 & 1
Thymelaeales	2	6	14
Tiliiales	3	13 & 1	30 & 3
Umbelliferae	2	6	21
Urticales	3	16	78

*Introductions are shown as additions. Introduced orders such as Violales, Proteales, Casuarinales, Pittosporales, Salicales are omitted.

TABLE 1A. Summary of the analysis of the Dicotyledons.

	Orders	Families	Genera	Species
Archichlamydeae .	45	97 & 1	389 & 37	1,161 & 124
Metachlamydeae .	17	31	258 & 22	746 & 35
Total	62	128 & 1	647 & 59	1,907 & 159

TABLE 2. Analysis of the Flora. Families of Dicotyledons. (C = Cosmopolitan, Tr = Predominantly tropical, T = Predominantly temperate).

Families	Genera	Species
Acanthaceae (Tr)	18*	35
Amarantaceae (C,Tr)	11	24
Ampelidaceae (Tr)	3	6
Anacardiaceae (Tr)	5 & 1	14 & 2
Annonaceae (Tr)	3	14
Apocynaceae (Tr)	11	26 & 1
Aquifoliaceae (C)	1	9
Araliaceae (Tr)	3	15
Aristolochiaceae (C)	1	5
Asclepiadaceae (Tr)	10 & 2	23 & 2
Balanophoraceae (Tr)	1	1
Basellaceae (Tr)	2	2
Batidaceae (Tr)	1	1
Begoniaceae (Tr)	1	6
Bignoniaceae (Tr)	8	17
Bixaeeae (Tr)	1	1
Bombacaceae (Tr)	2	2
Boraginaceae (C)	7	36
Brunelliaceae	1	1
Burseraceae (Tr)	2	4
Buxaceae (Tr)	1	4
Cactaceae	6	15
Caesalpiniaceae (Tr)	7 & 4	37 & 12
Campanulaceae (C,T)	2	2
Canellaceae	2	2
Capparidaceae (Tr)	5	15 & 1
Caprifoliaceae (C)	1 & 2	3 & 2
Caricaceae	1	2
Caryophyllaceae (C,T)	6	7
Celastraceae (C,T)	6	13
Ceratophyllaceae (C)	1	1
Chenopodiaceae (C,T)	3	4
Chloranthaceae	1	2
Clethraceae	2	4
Combretaceae (Tr)	6	8 & 1
Compositae (C)	50 & 9	126 & 13
Connaraceae (Tr)	1	2
Convolvulaceae (C)	6	50
Crassulaceae	1	1
Cruciferae (C,T)	6 & 2	7 & 3
Cucurbitaceae (C,Tr)	7 & 4	8 & 6
Cunoniaceae	1	1
Cyrillaceae	1	1
Dilleniaceae (Tr)	2	2
Ebenaceae (Tr)	1	1
Erythroxylaceae (Tr)	1	5
Euphorbiaceae (C,Tr)	30 & 2	111 & 8
Ficoidaceae	2	2
Flacourtiaceae (Tr)	1	3
Garryaceae	1	1
Gentianaceae (C,T)	7	13
Geraniaceae (C)	1	1
Gesneriaceae (Tr)	10	32
Goodeniaceae	1	1
Guttiferae (Tr)	5	12
Hernandiaceae (Tr)	1	2
Hippocrateaceae (Tr)	1	1
Hydrophyllaceae	2	3
Hypericaceae (C,T)	1	1
Iacinaceae (Tr)	1	1
Labiatae (C,T)	10 & 3	27 & 4
Lacistemaceae	1	1
Lauraceae (Tr)	7	18 & 1
Lecythidaceae (Tr)	1	1
Lentibulariaceae (C)	2	3
Linaceae (C)	1	1
Loasaceae	1	1
Lobeliaceae (C,Tr)	1	10
Loganiaceae (Tr)	2	3
Loranthaceae (C,Tr)	6	21
Lythraceae (C,T)	3	8
Malpighiaceae (Tr)	9 & 1	26 & 1
Malvaceae (C,Tr)	15	69 & 9

TABLE 2. Continued.

Families	Genera	Species
Maregraviaceae	1	2
Melastomaceae (Tr)	18	69
Meliaceae (Tr)	4 & 1	9 & 1
Menispermaceae (Tr)	2	4
Mimosaceae (Tr)	13 & 2	25 & 12
Molluginaceae (Tr)	2	3
Moraceae (Tr)	7	18
Moringaceae	0 & 1	0 & 1
Myricaceae	1	3
Myrsinaceae (Tr)	5	28
Myrtaceae (Tr)	6	60 & 3
Nyctaginaceae (Tr)	4	11
Nymphaeaceae	2	4
Ochnaceae (Tr)	2	5
Olacaceae (Tr)	2	5
Oleaceae	4	11
Onagraceae (C)	4	11 & 4
Oxalidaceae (C)	1	2
Papaveraceae (C)	2	2
Papilionaceae (C)	30 & 10	87 & 30
Passifloraceae (Tr)	1	14 & 3
Phytolaccaceae (Tr)	6	7
Piperaceae (Tr)	2	52
Plantaginaceae (C)	1	3
Plumbaginaceae (C)	1	1
Polygalaceae (C)	2	6
Polygonaceae (C)	3 & 1	26 & 2
Portulacaceae (C)	2	6
Primulaceae (C,T)	1	1
Quiinaceae	1	1
Ranunculaceae (C,T)	2	4
Rhamnaceae (C,T)	7	9 & 2
Rhizophoraceae (Tr)	2	5
Rosaceae (C,T)	5 & 1	9 & 6
Rubiaceae (C,Tr)	41 & 2	144 & 5
Rutaceae (Tr)	6 & 3	22 & 10
Salicaceae	0 & 1	0 & 1
Samydaceae (Tr)	6	16
Sapindaceae (Tr)	10 & 2	19 & 2
Sapotaceae (Tr)	7	26
Serophulariaceae (C)	16 & 3	26 & 7
Simarubaceae (Tr)	6	7
Solanaceae (C)	13 & 1	48 & 1
Staphyleaceae	1	1
Stereuliaceae (Tr)	7 & 1	16 & 2
Symplocaceae	1	2
Theaceae (Tr)	4 & 1	9 & 1
Thymeliaceae (C,T)	2	3
Tiliaceae (C,T)	4	12 & 1
Tovariaceae	1	1
Turneraceae (Tr)	2	4
Ulmaceae (C,T)	2	5
Umbelliferae (C,T)	3	6
Urticaceae (C,T)	7	55
Vacciniaceae	1	1
Verbenaceae (C,Tr)	16	40
Violaceae (C,T)	0 & 1	0 & 2
Zygophyllaceae	3	3

*Introductions are shown as additions.

The largest family is Rubiaceae with 149 species (Cuba 296), the genus *Psychotria* alone accounting for some 41. The largest order is Leguminosae with 203 species. Other large families are: Compositae 136, Euphorbiaceae 119 and Papilionaceae 117.

Families with 50 species or over are: Malvaceae, Melastomaceae, Solanaceae, Myrtaceae, Urticaceae, Convolvulaceae, Piperaceae.

Families with 35 species or over are: Caesalpini-

ceae, Boraginaceae, Verbenaceae, Mimosaceae, Solanaceae, Acanthaceae.

Almost half of the orders, 26, are represented by but a single family while of the 129 families in Table 2, 42 have two or less species. These figures reflect the great number of small families and genera in the flora. There are several reasons that can be suggested for this feature. Some dozen or more families are naturally small and often of restricted distribution, eg. Cyrillaceae, Clethraceae, Quiinaceae, Garryaceae, Lacistemaee, Tovariaceae, Batidaceae and Brunelliaceae. A few, although belonging to a large family, are represented by virtue of a single wide-ranging genus (Cunoniaceae, Linaceae). Others are temperate families represented in the tropics in montane regions (Plantaginaceae Geraniaceae, Primulaceae) : others, such as Staphylaceae are disjunct. Nevertheless, there still remain large tropical families of wide distribution that are barely represented in Jamaica eg. Iacacinaceae, Leethidaceae, Loasaceae, Vacciniaceae.

Notable pan-tropic families which are not represented at all are Monimiaceae, Marantaceae, Myristicaceae and Proteaceae. The last named is found in South America and extends up to Mexico but to none of the West Indies.

Table 2 shows that 48 families are considered to be cosmopolitan. Eight of these have a predominantly tropical and 19 a predominantly temperate distribution. The overall picture shows 64 tropical families out of the 129 present. Seven families have their greater affinities in the southern hemisphere—these are Cunoniaceae, Myrtaceae, Oxalidaceae, Thymeliaceae, Goodeniaceae, Canellaceae and Rutaceae while there are nine families having predominantly northern hemisphere affinities—Ulmaceae, Garryaceae, Nymphaeaceae, Papaveraceae Staphyleaceae, Geraniaceae, Rosaceae, Primulaceae and Aquifoliaceae.

PHYTogeOGRAPHY

A list of plants with authorities mentioned in the paper, is given in the Appendix. This list does not include the marine algae or the mosses since, for these, authorities are given in the text.

The flora of Jamaica may be grouped as follows:

- i. A cosmopolitan element
- ii. A West Indian element
- iii. An endemic element
- iv. A continental element

COSMOPOLITAN ELEMENT

These are mainly pan-tropical and pan-Caribbean plants, the highest percentage of which are strand plants such as:

- Batis maritima*
- Borrichia arborescens*
- Casuarina bonducella*
- Canavalia obtusifolia*
- Coccoloba uvifera*
- Euphorbia buxifolia*
- Heliotropium curassavicum*
- Ipomea pes-capri*
- Sesuvium portulacastrum*

Sporobolus virginicus
Tournefortia gnaphaloides

Common lowland pasture and roadside weeds reflect the 400 years of European communication in the Caribbean with such plants as:

Abrus precatorius
Achyranthes indica
Amaranthus spinosus
Argemone mexicana
Asclepias curassavicum
Boerhavia scandens
Bryophyllum pinnatum
Bidens pilosa
Borreria laevis
Calotropis procera
Capsicum frutescens
Cleome spinosa
Crotalaria retusa
Desmodium supinum
Gomphrena globosa
Gynandropsis pentaphylla
Hyptis pectinata
Jatropha curcas
Lantana involucrata
Mirabilis jalapa
Mimosa pudica
Petiveria alliacea
Pisonia aculeata
Psidium guajava
Ricinus communis
Sida rhombifolia
Stachytarpheta indica
Tribulus cistoides
Turnera ulmifolia
Urena lobata
Waltheria americana

There are others too numerous to mention. Also in this category are the many introduced trees which have become thoroughly naturalized in the lowlands, i.e., Bamboo, Mango and many Leguminosae.

WEST INDIAN ELEMENT

Although this may be regarded as an element common to the Caribbean islands, there are two aspects. Firstly, there are those plants common to all the West Indies and, secondly, a Greater Antillean element more or less distinct from the Lesser Antilles. Investigation of these two groups must await fuller botanical knowledge of these islands. The West Indian element, as a whole, includes many trees of the seasonal formations:

Acacia farnesiana
Amyris balsamifera
A. elemifera
Brosimum alicastrum
Bocconia frutescens
Brya ebenus
Bucida buceras
Bursera simaruba
Canella winterana
Ceiba pentandra
Clusia rosea
Chlorophora tinctoria
Crescentia cujete
Diphollis salicifolia
Exostema caribaeum

Guaiacum officinale
Guazuma ulmifolia
Krugiodendron ferreum
Petitia domingensis
Spondias monbin
Swietenia mahogoni

There are many common genera in the wet uplands, although the species may differ in the various regions.

Brunellia comocladifolia
Clethra occidentalis
Cyrilla racemiflora
Calophyllum spp.
Guarea glabra
Homalium spp.
Hedyosmum arboreum
Ilex spp.
Juniperus spp.
Laplacea spp.
Nectandra patens
Ocotea spp.
Prunus occidentalis
Weinmannia pinnata.

ENDEMIC ELEMENT

Endemism has been over-rated in the Caribbean and, as knowledge of the flora increases, there will be a modification of the past estimates. Complete figures are as yet unavailable for Jamaica; however, Burns (1947) confining his studies to indigenous trees and shrubs, found 440 endemics out of a total of 969 species. Fawcett & Rendle record 73 endemic orchids out of a total of 194 species. If we combine these figures and assume, for the present purpose, that there are few other endemics in the total angiosperm population of 2,500 species, a figure of 20.5% is obtained which represents the lowest possible percentage of endemism in the Jamaica flora. This figure compares with Trinidad (belonging to the Venezuelan Botanical district) of 7%; the Lesser Antilles 12% and Porto Rico 13%. Data for Cuba are unavailable at present but, judging from the large number of endemics in the flora as published so far, the overall figure will probably be the highest in the Greater Antilles.

Of the Pteridophytes, which have received recent attention by Proctor (1953), there are 90 endemic species (14%) as compared with 4% in Porto Rico. The ferns do not bear out the oft-expressed view that Jamaica represents a centre of high endemism.

Local endemism will be an interesting and fruitful field of study in Jamaica as it has proved in Cuba where no less than 16 endemic genera are localized in the north-west mountains (Carabia 1943, 1945a).

CONTINENTAL ELEMENT

The flora of the Greater Antilles is predominantly Central American in origin and its analysis sketches the history of past migrations. Biogeographers have, in the past, generally agreed that migration has taken place from Central America to the Greater Antilles. This aspect will be discussed later, however, the immediate purpose being not to prove migration but to indicate the position of Jamaica in the present-day plant distribution of the area.

The Continental element can be divided into three groups having Central, North and South American affinities respectively. It seems to be agreed that the more limited northern element was already present in parts of Central America when the Greater Antilles were linked by a land connection to the mainland. The distribution of these groups shows that, in many cases, Jamaica stands in sharp contrast to the other islands of the Caribbean. Table 3 gives a preliminary list of genera and species which illustrates this point.

TABLE 3. Distribution of the Continental elements of the Flora of the Caribbean.

	Cuba	Hispaniola	Porto Rico	L. Ant.	Jamaica
NORTHERN					
<i>Magnolia</i>	X	X	X	0	0
<i>Juglans</i>	X	—	X	0	0
<i>Fraxinus</i>	X	—	X	0	0
<i>Pinus caribaei</i>	X	0	0	0	0
<i>Pinus occidentale</i>	X	X	0	0	0
<i>Phyllostylon</i>	X	X	0	0	0
<i>Salix</i>	X	—	0	0	0
<i>Quercus</i>	X	—	0	0	0
CENTRAL					
<i>Didymopanax</i>	X	X	X	X	0
<i>Bombax</i>	X	X	X	0	0
<i>Lysiloma</i>	X	X	X	0	0
<i>Brosimum alicastrum</i>	X	X	X	X	X
SOUTHERN					
<i>Podocarpus urbani</i>	0	0	0	X	X
<i>Ocotea martinicensis</i>	0	0	0	X	X
<i>Protium attenuatum</i>	0	0	0	X	X
<i>Turpinia occidentale</i>	0	0	0	X	X
<i>Zygia latifolia</i>	0	0	0	X	X
<i>Sterculia caribea</i>	0	0	0	X	X
<i>Crataeva tapia</i>	0	0	0	X	X
<i>Sympomia globulifera</i>	0	X	0	X	X
<i>Dacryodes excelsa</i>	0	0	X	X	0
<i>Epidendrum</i> (3 spp.).....	0	0	0	0	X
<i>Brassavola nodosa</i>	0	0	0	0	X
<i>Maxillaria rufescens</i>	0	0	0	0	X
<i>Cryptarrhena lunata</i>	0	0	0	0	X
<i>Euterpe globosa</i>	X	X	X	X	0
<i>Carapa guianensis</i>	X	X	X	X	0

The northern group is illustrated by seven genera which are found in Cuba, by-pass Jamaica, yet may reach Porto Rico but not the Lesser Antilles. Proceeding eastwards, genera may be represented by fewer species. For example, there are 3 species of *Pinus* in Cuba but only one in Hispaniola. As far as is known with certainty, Fagaceae and Saliaceae reach their southern insular limit in Cuba but *Phyllostylon* is represented in Hispaniola. *Phyllostylon* is also found in South America, however, and the family Ulmaceae is represented in Jamaica by *Celtis*. The genus *Magnolia* which extends to Porto Rico is absent from Jamaica and the Lesser Antilles. From the ecological standpoint too, the northern element, as a whole, becomes less important as one proceeds eastwards.

The group with Central American affinities includes several species of *Didymopanax* that are common in

the Greater and Lesser Antilles but the genus is not found in Jamaica. *Bombax* and *Lysiloma* are similarly situated. On the other hand, there are species such as *Brosimum alicastrum* that have a wide distribution, being present throughout the islands and also occurring in South America.

Analysis of the southern element is more complex. The distribution pattern, as illustrated, shows that there are many species which are either common to the Lesser Antilles and Jamaica or to the entire Antilles but not Jamaica. It will be convenient here to discuss this distribution from a migrational aspect.

Much of the Southern element appears to have reached the Greater Antilles via the Lesser Antilles rather than Central America although this latter route cannot be ruled out. This view is supported by the fact that floristic representation and ecological importance is centred in the Lesser Antilles and diminishes as one proceeds northwards to Jamaica. There are many examples of such plants reaching the other Greater Antilles but missing out Jamaica. Table 3, however, is more concerned with showing examples where Jamaica is alone represented. The species of *Ocotea*, *Protium* and *Turpinia* may be regarded as Lesser Antillean indigenous species which have reached Jamaica as waifs by sea and wind currents. *Licania*, common in the Lesser Antilles, does not reach the Greater Antilles. *Dacryodes excelsa*, an important dominant of Lesser Antillean rain forest associations, goes no further than Porto Rico. In a similar category is *Sloanea*, represented by at least 6 species in the Lesser Antilles but by one infrequent species in Jamaica. *Pterocarpus officinalis* is present in Jamaica but never forms the large swamp forests found in the Lesser Antilles. The palm, *Euterpe globosa*, and *Carapa guianensis* are examples of southern species, widespread in the Caribbean, which by-pass Jamaica but reach Central America. There are 6 South American orchid species which occur only in Jamaica of all the West Indies. Five of these have probably reached Jamaica via Central America according to the present distribution pattern. The species of *Zygia* and *Sterculia* are South American species found in the Lesser Antilles and Jamaica. Together with *Crataeva tapia*, they are disjunct species of very problematical significance.

Some southern elements, e.g. *Weinmannia pinnata*, are of wide distribution throughout tropical America and could have equally well migrated by various routes. There are others, however, which by their distribution pattern in the West Indian islands, definitely indicate an isthmus route. The genus *Clethra* is found in South America and Trinidad but is not recorded for the Lesser Antilles or Porto Rico. However, it extends up to Mexico via Panama and extends westwards to Cuba and Jamaica. *Podocarpus urbani*, also present in Central America, is confined to Jamaica and the Lesser Antilles.

Fawcett & Rendle (1910) analysing the Orchidaceae of Jamaica, found that the greatest affinity is with Cuba where 82 of the 121 cosmopolitan species are found. Fourteen species are restricted to the

two islands. Hispaniola has 29 common species and only 2 restricted to the two islands. Porto Rico, with a better-known flora, has some 40 species in common with Jamaica. Further afield, the species of orchids in common with Jamaica are:

Bahamas	10
Florida	15
Lesser Antilles	38
Trinidad	40
Central America	44
South America	48

Many of these are cosmopolitan throughout Tropical America and have little significance in geographical affinities. However, it will be seen that there is almost equal affinity with South America and Central America and this would suggest two broad migration routes—one from the south and one from the west. Before this evidence can be fully evaluated, however, it will be necessary to relate the precise numerical affinities between each of the Greater Antilles in turn.

The fern flora of Jamaica shows a relationship with both the Greater Antilles and Central America. Affinities within the Greater Antilles are, however, not so strong with Jamaica as between the other members of this group. This follows the pattern seen in the higher plants.

It will be evident then that Jamaica holds a unique floristic position in the Caribbean either by having or lacking certain species. In this connection an observation by Woodson (1940) is of interest. Dealing with the present records of distribution of the Apocynaceae, he states "not a single instance is found of a distribution common to Hispaniola and Jamaica, except in the case of species of wide distribution throughout the Greater Antilles as a whole. On the other hand, numerous instances of distributions involving Cuba and Hispaniola are found." This conclusion is further borne out when considering the distribution of *Pinus occidentalis* and *Podocarpus purdeanus*. The former only occurs in Cuba and Hispaniola and the latter only in south-east Cuba and Jamaica.




FIG. 7. Bio-geographical relations within the Caribbean. Migration to the Greater Antilles has taken place from the west by former land connections to Honduras, and from the Lesser Antilles by long-distance migration.

Until more is known of the plant ecological and floristic affinities in the Caribbean area, the examples given here must merely suggest the many problems awaiting solution. The main lines of approach must consist in careful consideration of migration methods together with the ecology of the species concerned. Broadly speaking, it would seem that the explanation of the geographical affinities of the flora of the Greater Antilles is to be found in migration from Central America by way of land connections and from the east and south by long-distance migration (Fig. 7). This will now be discussed with particular reference to the relationship of the Jamaican flora.

LAND BRIDGE HYPOTHESES

Biogeographers have postulated Central American-Antillean land bridges in at least two places, one from the Honduran peninsula to the Greater Antilles and one from the peninsula of Yucatan to Cuba. Schuchert (1935), who is the most recent authority on this subject, sees no evidence for the latter and does not believe that it ever existed. A series of diagrams representing the ideas of Schuchert on the geological history of this region are given by Woodson (1940) and are summarized in Table 4.

The important features that can be drawn from the data available are that submergences following the wide land connections during both Cretaceous and Lower Oligocene times were incomplete as far as Cuba, and possibly Hispaniola and Porto Rico, were concerned, some land remaining above the sea. After the second submergence, in the Middle Oligocene, Cuba was separated from the rest of the greater Antilles. During Upper Eocene and again in Middle Oligocene times, the southeastern tip of Cuba alone was directly connected to Honduras. In Upper Miocene times, a land connection existed between Honduras, Jamaica, Hispaniola and Porto Rico. With the exception of the centre and west of Cuba, there appears to have been no further major submergence as far as these islands were concerned, but the land bridge to Honduras was finally broken in Upper Pliocene times.

It is clear that migrations by the Antillean-Honduran land bridge will explain the broad general affinities between the floras of the Greater Antilles and Central America. Furthermore, the conjectural geographical history pointing to early isolation of Cuba, is also borne out by the presence there of such relicts as the endemic *Microcystis* and *Zamia* spp. and the high number of endemics generally, as shown in the recent flora. An explanation can also be offered of the affinity between Cuba, Hispaniola and Porto Rico, and the anomalous position of Jamaica as far as some members of the northern group of Central American plants are concerned. This could have occurred not necessarily by inter-island migration but by the continued presence in these three islands of refugia when the rest of the land, and the whole of Jamaica, was submerged. Such refugia would provide secondary centres of distribution in the islands concerned when the land, as we know it to-day,

TABLE 4. The Conjectural Land Connections of the Greater Antilles (after Schuchert 1935).

Geological Period	Land Connections	Remarks
Upper Pliocene—Pleistocene	None	Cuba submerged except for S. E.
Upper Miocene—Lower Pliocene	Honduras—Jamaica—Hispaniola—Porto Rico	Narrow separation from Cuba to N. W. Hispaniola
Lower Middle Miocene	Honduras—Jamaica—Hispaniola	Much of Cuba & Hispaniola submerged
Upper Oligocene	No connections to mainland	Cuba submerged except for south-east corner; Jamaica completely submerged
Middle Oligocene	Honduras—S. E. Cuba; Jamaica—Hispaniola; Hispaniola—Porto Rico	Cuba finally isolated from rest of Greater Antilles
Lower Oligocene	Honduras—G. Antilles—Bahamas	
Upper Eocene	Honduras—S. E. Cuba; Jamaica—Hispaniola; Hispaniola—Porto Rico	All low parts of islands submerged
Upper Cretaceous	S. America—C. America—G. Antilles—Bahamas—S. Florida	

subsequently emerged. Such a scheme would account for distribution of northern species following the Magnolia pattern and central species following the Bombax, Lysiloma pattern. It does, of course, assume that such plants had already migrated from Central America by these times. As to whether this is a reasonable assumption or not, we have no evidence. However, if they were present, the question might well be asked why did migration not take place to Jamaica from Hispaniola during the land connection of Upper Miocene times?

The only other alternative would seem to be that there may have been inter-island migration of these species which arrived in Cuba during those periods when Cuba alone was connected to Honduras. It is difficult to find any adequate reasons why Jamaica should have been left out of such migrations.

Species distributed according to the pattern of *Pinus caribaea* and the oaks which appear to be confined to Cuba, are in a different category from the above and may well be taken as an example of the danger of putting too much emphasis on land bridges alone without considering the ecological implications of the species concerned. The distribution of *Pinus caribaea* in the Caribbean area is indicated in Fig. 7. Ecologically, it is confined to a habitat where the soil consists of flat siliceous sands (which may be at sea level or up to 2,000 ft.), supporting a Pine savanna of sedge and grasses dotted with pine trees, oaks and a few characteristic associated shrubs. It is absent from Jamaica and possibly the other islands of the Antilles, because there are no habitats present with the edaphic conditions required. It should, furthermore, be noted that, at the present time, *Pinus caribaea* is confined to the west of Cuba where such soils occur. This area, according to Schuchert, was submerged as late as Upper Pliocene-Pleistocene times and it is extremely unlikely that any land pres-

ent in south-east Cuba would provide a refuge for the members of this specialized community. It must be assumed, then, that *Pinus caribaea* arrived in Cuba at a later date and it is tempting to suggest that this was via a land-bridge extending from Yucatan to Cuba, 160 mi away.

It is generally agreed that no land-bridge ever existed between Cuba and Florida and North America. Southward, a continuous land-bridge has been postulated through the Lesser Antilles to South America. This is very doubtful owing to the differing age and volcanic origin of these smaller islands. Beard (1948) has shown that the past geologic history of the Lesser Antilles is reflected in the flora which supports the view that these islands were never linked by a land-bridge.

LONG DISTANCE MIGRATION

Within the Caribbean as a whole, emphasis has been upon continuous land connections. Long distance migration has been relegated to a minor role. Nevertheless, there is much evidence that migration of plants with wind-borne disseminules, has occurred. Strong winds blow down from Cuba to Jamaica in the months of November and December. This may account for the affinity between these two islands of the Orchidaceae and the Pteridophytes. Furthermore, sea and wind currents, often of hurricane force, pass upwards from the Lesser Antilles to the Greater Antilles and the Caribbean shores of Central America for most of the year. Dealing with the distribution of the Apocynaceae, Woodson gives evidence supporting the assumption that wind-borne seeds account for the migration of Greater Antillean species such as *Echites lutea*, *Echites umbellata* and *Rhabdadenia biflora* to the mainland of Central America. In the absence of any evidence for a land-bridge connecting the Greater and Lesser Antilles to South America, it must be assumed that many of the species in Jamaica

that have affinities with the Lesser Antilles, have arrived by long distance migration. How this has occurred should be worked out for each species in detail, bearing in mind that there are many examples, such as those of southern species given in Table 3, which are not common to the Greater Antilles as a whole. Chapman (1947) presented figures showing the geographical relationship of the seaweeds of Jamaica to Florida, the Bahamas and the Virgin Islands. These affinities are undoubtedly due to the prevailing sea currents.

The importance of birds and bats as agents for the spread of plants in the Caribbean has been stressed by Beard (1949). He estimates that at least two-thirds of the rain forest trees of the Lesser Antilles produce fruit attractive to these animals and thinks that this may account for their migration from South America.

Paleobotanical evidence for the Caribbean is fragmentary. Genera and families have been traced back to the Cretaceous and Tertiary, and there appears to have been a sudden flush of new species since that time which, no doubt, has some bearing upon the high endemism claimed for the West Indies (Hollieck 1924, 1928). The Jamaican record has not been studied. A well-preserved palm type has been recently found. However, all the fossils date back to a pre-emergence period and so belong to a lost flora.

COMMUNITY STUDIES

INTRODUCTION

The vegetation of Jamaica, with the exception of the montane regions, is tropical. Typical of Caribbean islands, the major ecological boundaries are determined by a wet northern coast, a central montane region and a dry southern plain. The determining factor is rainfall which, in turn is dependent upon the prevailing wind and the mountain barriers in its path. Within these boundaries, however, many local factors operate to bring about a very diverse and varied vegetational pattern.

The vegetation as a whole has never been described. The sparse literature which exists on the subject comprises, for the most part, descriptive essays written by visiting botanists, or more intensive investigations of a single aspect. The Caribbean Commission (1947) on forest research for Jamaica reported that "no appreciable data are available on plant ecology or other aspects of forest biology" and stated that there was an urgent need for a detailed ecological survey of the forest types and successional groups.

In 1846, the Danish botanist, Ørsted, visited Jamaica and later published a map and description of the vegetation, outlining the broad plant communities. There are also some brief generalized descriptions of historical interest by Sir Hans Sloane, Browne, Swartz and other early workers. Harshberger (1903) and Shreve (1910) describe some of the coastal vegetation, while Shreve (1914) gives an account of the higher montane forest together with some physio-

logical investigations. A lengthy paper by H. & M. Brockmann-Jerosch (1925) is remarkable if only for the fact that it was based upon a three-week visit in 1913. A cross section of the eastern (Blue Mountain) end of the island is well described from the route taken and perhaps remains as one of the best general reports to date. In 1939, Cambridge University sent a small expedition to study mangrove communities. The results were published by Chapman (1940, 1944).

Although initially the sole objective was to describe the local vegetation, it soon became evident that account must be taken of the ecological correlations necessary to fit the vegetation of Jamaica into the Caribbean area as a whole. Beard (1944a) has specifically discussed the problems of ecological classification within the Caribbean and proposed a system for general use throughout Tropical America. He has already used his system to advantage in his excellent monographs on Trinidad (1946) and the Lesser Antilles (1949). An attempt to correlate the vegetation of Jamaica with the vegetation types recognized by Beard is made in the discussion (p. 403).

There are various ways of presenting the vegetation of Jamaica. Swabey (1942) proposed an outline classification which was based on that of Burtt-Davy (1938). While somewhat unwieldy, the main points may be summarized. In Jamaica, he recognized climatic and edaphic climax formations. The former included moist woodland with 70 in. of rain, dry woodland with 50 in. and montane formations. Edaphic climax formations included wet limestone woodlands, dry limestone woodlands and the maritime swamp and aquatic formations. Finally, there were a number of seral "formations". It is interesting to note that the "limestone formations" were regarded as edaphic climaxes. The simplest approach is that adopted by Gleason & Cook (1927) for the vegetation of Porto Rico where they described separately a wet northern region, a central montane and a dry southern region.

In Jamaica, however, the major headings fall more naturally under coastal, lowland and montane.

The coastal communities include the marine vegetation, communities on sand beaches, sand spits and coral rock. Under this heading too, special consideration is given to the small islands known as Cays that occur off the coast. The lowland communities are discussed under four main headings. These are limestone, alluvium, shale, and the swamps and marshes. The montane vegetation is conveniently divided into lower montane rain forest, montane sclerophyll, and montane mist forest with its elfin woodland aspect.

The method of classification is mainly one of convenience. It is based largely upon the edaphic factor. While the recognition of an edaphic climax is not advocated, it cannot be denied that soil type is an important factor in Jamaica where limestone rock and derived soils, "shale" hills and alluvial plains form the major soil patterns. It has already been emphasized how closely the soil-boundaries are correlated with the geological map. It is further remarkable how con-

veniently the rainfall and topographic maps correspond in Jamaica. Only in the John Crow Mountains is there any major deviation. Edaphic factors are also important elsewhere in the Caribbean, for example the volcanic soils of the Lesser Antilles and the serpentine and siliceous savannas of Cuba.

COASTAL COMMUNITIES

The coastline of Jamaica consists, for the most part, of limestone rock or low-fringing coral shelves. High sea cliffs and headlands are infrequent. Variation is given by the occurrence of bays with sandy spits, beaches and by alluvial deposits at river estuaries. The larger bays and sheltered estuaries are mainly on the south coast.

The south coast is drier than the north and the average rainfall is often below 30 in. (Fig. 5, 6). On the north coast, the only area which approaches this figure is that between Falmouth and Montego Bay, where the average is about 40 in.

Since the bulk of the rain falls in the two rainy seasons embracing the months of May and October, long periods of drought occur. The arid nature of this halophytic habitat is further intensified by the fierce insolation, porosity of the substratum and salt-laden winds which may reach hurricane force.

Apart from restricting the height of the vegetation, wind is an important factor in the development of marine currents which bring about deposition of rock, gravel and sand, forming the beaches, spits and cays which are eventually colonised by the strand communities.

At the river mouths and in sheltered areas, a sediment of fine silt is built up to form mud banks. These are colonised by mangroves which assist the processes of accretion and land formation by retaining the silt.

Tidal range in Jamaica is very restricted, being less than 16 in. and averaging 10 in. This is not only important to the marine vegetation but also affects the sand communities. These are confined to a narrow fringe rarely elevated more than a few feet above the sea, so that the substratum is open to modification by accretion or erosion during heavy storms and hurricanes. Strand vegetation is generally in a state of flux and the beaches and spits may show both accretion and erosion occurring simultaneously.

MARINE VEGETATION

Marine algae are not abundant around the coast of Jamaica in comparison with temperate zones. This is, no doubt, due to the small tidal fluctuation and the steeply shelving shore line. Much of the substratum is also unsuitable for the establishment of algal communities.

In shallow waters of sheltered and silty bays, the marine grasses *Thalassia testudinum* and *Cymodocea manatoria* are often abundant.

Green algae are well represented, especially the lime-encrusting members of the Siphonales such as *Halimeda*, *Acetabularia* and *Cymophyllum* but there is a general paucity of brown seaweeds. The red algae

are chiefly small, insignificant and epiphytic. Chapman (1947) gives the following figures:

Chlorophyceae	105
Cyanophyceae	51
Phaeophyceae	36
Rhodophyceae	135
Total	327

Zonation, which is usually associated with large tidal fluctuations together with a sloping shoreline, is virtually absent in Jamaica. The communities are seldom, if ever, uncovered by water, and the habitat groups that may be recognized are based upon protection, substratum and, to a lesser extent, depth of water.

Flat coral shelves extending from the shore line and covered by a foot or so of water together with reefs up to 10 ft. under the sea and several hundred yards off shore, provide the most frequent habitats.

In deeper water, on the seaward and exposed edges of the rocks buffeted by the waves, the following may be found:

<i>Sargassum vulgare</i> C. Ag.
<i>S. lendigerum</i> (L.) Kuetz
<i>S. polyceratum</i> Mont. var. <i>ovatum</i> Collins
<i>Dictyota dentata</i> Lamx.
<i>Cutleria</i> sp.
<i>Turbinaria turbinata</i> (L.) Kuntze
<i>Dictyopteris delicatula</i> Lamx.
<i>Acanthophora spicifera</i> (Vahl) Börgs.
<i>Bryothamnion triquetrum</i> (Gmel.) Howe
<i>Gelidiopsis rigida</i> (Vahl) Weber-van Bosse
<i>Gratelouphia filicina</i> (Wulf) C. Ag.
<i>Avrainvillea nigricans</i> Decne.
<i>Gracilaria lacinulata</i> (Vahl) Howe

Most of the algae are to be found on the shallow flat surface of the coral shelves. This has several aspects; the rock may be bare or covered with a layer of silt; it may be exposed, sheltered or have deeper parts where there are indentations or pools.

Caulerpa racemosa (Forsk.) Weber-van Bosse var. *uvifera* (Turner) Weber-van Bosse, grows best on the silt where it forms long rhizomes. Here, also, are to be found *Udotea flabellum* (Ellis & Solander) Howe, *Acetabularia crenulata* Lamx., *Penicillus capitatus* Lamarek and *P. pyriformis* A. & E. S. Gepp.

In quieter waters the following may occur:

<i>Caulerpa taxifolia</i> (Vahl) Ag.
<i>C. crenulata</i> (Vahl) Ag.
<i>Codium tomentosum</i> (Huds.) Stackh.
<i>Cymoplia barbata</i> (L.) Haw.
<i>Halimeda tuna</i> (Ellis et Solander) Lamx.
<i>H. opuntia</i> (L.) Lamx.
<i>H. incrassata</i> (Ellis et Solander) Lamx.
<i>Zonaria zonalis</i> (Lamx.) Howe
<i>Padina vickeriae</i> Hoyt
<i>Rhipocephalus phoenix</i> (Ellis et Solander) Kütz.
<i>Laurencia obtusa</i> (Huds.) Lamx.
<i>L. intricata</i> Lamx.

The tidal fringe provides another habitat where the component species are just covered by water and usually of low moss-like life form. Species common here are:

- Ectocarpus* sp.
Chaetomorpha sp.
Microdictyon boergesenii Setchell
Valonia ventricosa J.Ag.
V. aegagrophila C.Ag.
Cladophoropsis membranaceae (Ag.) Børgs.
Polysiphonia sp.
Dictyosphaeria cavernosa (Forskul) Børgs. *favulosa*
 (Ag.) Deesne.
Cladophora sp.
Enteromorpha sp.
Ulva sp.

We are indebted to Prof. V. J. Chapman of Auckland University College, New Zealand for many of the determinations.

STRAND COMMUNITIES

In Jamaica, coastal communities are found on three types of substratum, viz. sand, limestone and coral rock, and mud (See Fig. 12). The last named is occupied by mangrove swamps which are described later when dealing with the swamp formations. The mangroves, although found on shallow reefs and sand banks, reach their full development along sheltered, muddy estuaries. They are subject to tidal inundation and escape the drought conditions of the land plants growing on porous sands and rocks. It is, thus, not surprising that the mangrove association is so clearly distinct from the other coastal communities. However, all share the common feature of having successional patterns involving topographical changes.

In describing the vegetation on sand and coral rock, a simple classification upon a seral basis is the most satisfactory. Davis (1942) suggested the following for the Caribbean:

- i. a strand-beach associes
- ii. a strand-dune associes
- iii. a strand-scrub associes
- iv. a strand-woodland association

He recognized these communities on the Florida Keys where they corresponded to a pioneer open community on mobile sand, an herbaceous community of the fixed dunes, a scrub community and a climax of woodland. It is under these general headings that the strand communities of Jamaica will be described.

THE STRAND-BEACH ASSOCIES

The beach sands of Jamaica are highly calcareous, containing high proportions of weathered and pulverised limestone and coral rock, together with sea shells and calcareous algae. In a few localities, the sand is predominantly of *Halimeda* remains. There are no coastal dunes such as seen in Porto Rico, the nearest approach being on the Palisadoes where sand has been built up over a shingle bank to a height of 20 ft.

Coastal beaches are to be distinguished from those found on the seaward side of spits. True beaches are few in number, the best examples being on the north coast where they are popular resorts and, consequently, the natural vegetation has suffered. A few are bounded by cliffs and restricted in width. Modifications have been brought about by the plant-

ing of coconuts and the proximity of the coastal roads. It is on these beaches that the best development of the pioneer phases is to be seen. On the spits, continual erosion and modification restricts this community.

This pioneer associes begins above the tidal limits on initially, and potentially, mobile sand. Salinity is high but there is adequate moisture immediately below the surface. Wind and storms may bring about local or major alterations in topography due to sand removal or accretion.

The important plants are halophytes and psammophytes with quick vegetative propagation by runners. *Ipomea pes-capri* flourishes on the mobile sand front and, together with *Sporobolus virginicus*, an important sand-binder, is invariably present in the pioneer associes. The long, trailing runners of *Ipomea* spread for a length of 40 ft. or more across the beach, whilst the tough, wiry rhizomes of *Sporobolus* bind the sand.

Other plants, abundant to frequent, are: *Euphorbia buxifolia*, *Sesuvium portulacastrum*, *Cakile lanceolata* and the grasses *Cenchrus tribuloides*, *Chloris petraea*, *Spartina patens* var. *juncea* and *Paspalum vaginatum*.

The extent and development of the pioneer zone depends much upon the width and elevation of the beach as well as interference by man. On beaches where active erosion is taking place, the pioneer zone is marked only by a sparse drift-line flora fronting a hedge of *Coccoloba* and *Thespesia* (Fig. 11). Even where well developed, it is essentially a very open, successional community.

Plants of the strand-beach associes:

Sand runners and binders:

- Ipomea pes-capri*
Sporobolus virginicus
Cenchrus tribuloides
Paspalum vaginatum

Herbs:

- Euphorbia blodgettii*
E. buxifolia
E. hypericifolia
Sesuvium portulacastrum
Cakile lanceolata
Heliotropium curassavicum
Erigeron canadensis
Lippia reptans
Philoxyerus vermicularis
Atriplex cristata
Manisurus altissima
Portulaca pilosa

Grasses and Sedges:

- Chloris petraea*
Spartina patens
Cyperus brunneus
Cenchrus pauciflorus
Uniola virgata

THE STRAND-DUNE ASSOCIES

This is the second phase in the succession and is characterized by being a closer and predominantly herbaceous community on the fixed dunes. It may emerge from the pioneer zone or, where this has been destroyed by wave action, begin sharply from the eroded shelf. This community is well represented.

The seaward fringe is usually dominated by small tussocks of *Sporobolus* in close formation. Intermingled with this, or forming an inner zone is *Spartina patens*, a typical psammophytic grass with rolled leaves.

Other common plants are *Cenchrus*, *Chloris*, *Cyperus brunneus*, *Euphorbia buxifolia*, *Sesuvium portulacastrum*, *Alternanthera ficoidea*, *Opuntia tuna* and, on the Palisadoes, *Tribulus cistoides*.

Occasional adventives from the scrub are found including *Scaevola plumerii*, *Tournefortia gnaphalodes*, *Suriana maritima*, *Borrichia arborescens*, *Caesalpinia bonducella*.

Wind-stunted Acacia and *Coccoloba* may also be present.

Sharp definition between this zone and the next is rare. More usually there is a gradual merging shown by an increase in the shrub population. Plants of the strand-dune associoes:

Herbs:

Sporobolus virginicus
Spartina patens
Cenchrus tribuloides
Chloris petraea
Cyperus brunneus
C. laevigatus
C. ligularis
Hymenocallis sp.
Euphorbia buxifolia
Fimbristylis ferrugineus
F. spadicea
F. glomerata
Sesuvium portulacastrum
Phloxeris vermicularis
Alternanthera ficoidea
Canavalia obtusifolia
Opuntia tuna
Stemodia maritima
Erigeron canadensis
Eustoma exaltatum
Talinum paniculatum
Isocarpha oppositifolia
Egletes prostrata
Eragrotis ciliaris

Occasional shrubs:

Scaevola plumerii
Tournefortia gnaphalodes
Borrichia arborescens
Suriana maritima
Caesalpinia bonducella
Coccoloba uvifera
Acacia tortuosa
Spilanthes urens
Melochia crenata
Turnera ulmifolia

THE STRAND-SCRUB ASSOCIES

This associoes is seldom well developed on the sand beaches due to their limited width and human interference. It is best seen on sand overlying raised coral rock shelves.

Beaches of this type are found on the north coast and here the scrub is dominated by *Suriana*, *Tournefortia*, *Borrichia* and *Scaevola*. Other frequent species are:

Caesalpinia bonducella
Ernodia littoralis
Erithralis fruticosa
Solanum havanense
Morinda royoc
M. citrifolia
Colubrina asiatica
Coccoloba uvifera
Phyllanthus epiphyllanthus

The open spits of the south coast have a different succession, the climax often being replaced by a caetus thorn-scrub related to that of the alluvial plains. Here, the strand scrub stage is most frequently made up of low, sparse, bushes of *Acacia tortuosa*, *Prosopis juliflora*, *Boerhavia scandens*, *Capparis ferruginea*, *Coccoloba uvifera*, *Pithecellobium unguis-cati*, *Jatropha gossypifolia* and cacti together with the usual grasses, sedges and halophytic herbs. The introduced *Calotropis procera* is present on the Palisadoes.

THE STRAND WOODLAND ASSOCIES

The most frequent woodland on the strand is the *Coccoloba*-*Thespesia* association which may be seen on many beaches around the coast. Although usually exclusive, the following may also be present: *Conocarpus erecta*, *Dalbergia ecastaphyllum*, *Colubrina asiatica*, *Morinda citrifolia*, *Piscidia piscipula*, and *Sophora tomentosa*.

On the north and east coasts, particularly where there is a raised coral shelf, a characteristic strand woodland develops. This is a low (10-20 ft. tall) scrubby, open forest which may include palms and a mixture of trees and shrubs, many of which are from the arid limestone regions. Morant Point in the southeast, exposed to the full force of the easterly trade winds, supports a stunted woodland, more aptly described as "scrub" but rendered conspicuous by the presence of the palm *Thrinax parviflora* occasionally reaching a height of 25 ft. (Fig. 8). The important constituents here are:



FIG. 8. Stunted palm woodland at Morant Point. In the foreground are to be seen *Cordia sebestana* on the left and *Morinda citrifolia* in the centre. The palm is *Thrinax parviflora* and the wind-swept trees in the background are *Conocarpus*, *Capparis* and *Hippomane*. *Croton flavens* forms the ground layer.

Hippomane mancinella
Coccoloba uvifera
Cordia sebestana
Conocarpus erecta
Capparis ferruginea
Eugenia buxifolia
Jacquinia armillaris
Plumiera alba
Colubrina asiatica
Erithralis fruticosa

Also recorded are:

Thespesia populnea
Opuntia tuna
Machaerina rotundata
Croton linearis
C. flavens
Melochia lupulina
Ficus morантensis
Helicteres jamaicensis
Suriana maritima
Morinda royoc
M. citrifolia

On the coastal ledge near Falmouth, which may be up to 200 yd. wide and is relatively protected, the woodland is much more strongly developed. Some of the larger species such as *Thespesia*, *Coccoloba* and *Lonchocarpus* are up to 10 in. in trunk diameter and at least 30 ft. tall, whilst the average height of the canopy is 20-25 ft. There are no palms here but a much greater variety of species than at Morant Point. Nevertheless, it will be seen from the species lists that many are common to both these regions.

Important species occurring here are as follows:

Trees:

Pithecellobium unguis-cati
Hippomane mancinella
Coccoloba uvifera
Conocarpus erecta
Thespesia populnea
Capparis ferruginea
C. cynophallophora
Eugenia sp.
Tecoma leucoxylon
Comocladia sp.
Leucaena glauca
Rhacoma crossoptatum
Gyminda latifolia
Amyris elemifera
Lonchocarpus latifolius
Peltophorum brasiliense

Shrubs:

Jacquinia armillaris
Erithralis fruticosa
Randia aculeata
Helicteres jamaicensis
Melochia tomentosa
Lantana involucrata
Yucca aloifolia
Abutilon giganteum
Croton flavens
Citharexylum fruticosum

Herbs and Climbers:

Tournefortia volubilis
Tragia volubilis
Solanum havanense

Aegiphila elata
Echites umbellata
Pisonia aculeata
Opuntia tuna
Aloe sp.

Also recorded from other areas are: *Ficus aurea*, *Coccoloba littoralis*, *Clusia rosea*, *Clusia flava*, the scrambling shrub *Hyperbaea domingensis*, and the palms *Sabal jamaicensis* and *Coccothrinax fragrans*.

THE SAND SPITS

Spit formation is of frequent occurrence in Jamaica and, since the subject has an important bearing upon the formation of lagoons, salt ponds, salinas and beaches, all of which determine the coastal vegetation in general, a fuller account is not out-of-place here. Spits also bear some relation to the Cays to be described in the next section.

The history of spit formation and its subsequent development is easily worked out and a series of diagrams illustrating this process has been constructed from the various examples seen around the coast (Fig. 9). Starting with an open bay, the first stage is the development of a narrow tongue stretching towards the opposite shore (Stage 1). A sheltered lagoon is produced in which mangroves appear and silt is deposited. Next, the spit is thrown completely across the bay to form a sand bar and an inner salt pond (Stage 2). Subsequently, dry land is produced by evaporation from the shallow areas silted up during the lagoon phase (Stage 3). Finally, all that remains is a beach with a fringing background of mangroves (Stage 4). Dry land formation may be accelerated by riverine deposits of silt or seismic elevation of the land proceeding at the same time.




FIG. 9. Diagrams illustrating sand spit formation (1), and the subsequent development of a sand bar and a salt pond (2), followed by a salina (3). In the final stage (4), the usual strand communities are developed on a beach backed by mangrove woodland (X) and a salina that is being invaded by cactus thorn scrub.

All stages of these processes can be seen around the coast of Jamaica from the embryo spit through sand bars backed by shallow lagoons and salinas to popular bathing beaches. The only evidence of this sequence in the final beach stage is the presence of the last few remaining mangroves, marking the inner edge of the former spit.

The vegetation on the spits shows distinct zonation. On the seaward side, a pioneer zone, which may only be represented by a drift line flora containing typical pioneer species, is followed closely by a narrow dune stage. This zone leads directly into a cactus-thorn scrub in the centre of the spit. On the inner edge, mangroves are developed and may show typical zonation of species from *Rhizophora* in shallow water, through *Avicennia* and *Laguncularia* to *Conocarpus* on dry land. This is illustrated by the profile diagram in Fig. 10. That these types of zonation depend entirely upon the age, width and elevation of the spit is clearly demonstrated by the Palisadoes and the Hellshire spits. A diagrammatic representation of the Hellshire spit is also given in Fig. 10. On the narrowest and youngest region, only the pioneer stage is seen. As the spit broadens, the fixed dune stage is included and, on the inner shore, *Rhizophora* appears. Further widening and elevation leads to the appearance of cactus-thorn scrub which may be the final stage, as on the Palisadoes. It should be noted, however, that, at the widest and most mature end of the Hellshire spit, a zonation typical of a beach is shown and the climax is strand woodland of *Coccoloba*




FIG. 10. Diagrammatic representation of zonation on the Hellshire sand spit. The profile is taken across the spit from A-B. Zonation depends upon the age, width and elevation of the spit.



FIG. 11. Strand woodland of *Coccoloba uvifera* and *Thespesia populnea* fringing the Hellshire sand spit. The strand dune zone has been removed by erosion although a remnant may be seen in the sand cliff near the figures. In the foreground is the tidal drift-line zone above which is a zone of mobile sand colonized by *Sporobolus virginicus*.

uvifera. Erosion may cause modification by entirely removing the pioneer, dune and scrub stages so that the *Coccoloba* woodland fringes the shore directly (Fig. 11).

A perfect example showing the sequence of communities on sand postulated by Davis for the sand keys of Florida has yet to be seen in Jamaica. Many of the more extensive sand beaches have become popular holiday resorts and the natural vegetation destroyed or markedly altered. On the south coast, the succession is modified by the invasion of thorn scrub vegetation similar to that on the alluvial plains. On the north coast, the raised coral rock shelves introduce a further complication. It is, therefore, premature at this stage to enter into the details of successional relationships. Nevertheless, the following pattern suggests itself and is illustrated in Fig. 12.

The pioneer strand beach associes is followed by the more stable strand dune associes. Further topographical and edaphic changes lead either to a climax of *Coccoloba*-*Thespesia* strand woodland or, on the sand spits of the south coast, to a climax of cactus thorn scrub. However, where the sand is built up




FIG. 12. Scheme illustrating the relationships between the marine, coastal and adjacent inland communities.

over coastal limestone and coral rock shelves, a different zonation appears. Proceeding inland, a poorly developed strand-dune associes gives way to the Tournefortia-Suriana strand scrub associes which, in turn, is succeeded by Palm-Hippomane strand woodland. The last two coastal communities have, so far, only been seen in Jamaica where there is a substratum of limestone on coral rock. It is interesting to note, however, that Davis found no significant difference between the rock hammocks (woodland) and the sand hammocks of the Florida keys. Whether the strand scrub and Palm-Hippomane woodland are part of the psammosere in Jamaica is doubtful. The latter community would seem to have closer affinity with the coastal fasciation of Dry Limestone woodland. Mangrove woodland develops in protected areas where silt is deposited.




FIG. 13. Diagrammatic representation of the vegetation type on Lime Cay.

THE CAYS

Discussion of the vegetation of the Cays will be limited mainly to some of the Port Royal Cays visited by the Cambridge Expedition in 1939 and the authors in 1952.

A cay (or key) is a small island of sand or coral rock. The cays of Jamaica have been described by Steers, et al. (1940). Those nearest to the coast occur off the Palisadoes and in the Portland Bight. The former are called the Port Royal Cays. In addition, there are the Morant Cays, 30 mi. south-east of Morant Point, and Pedro Cays, 60 mi. south-west of Kingston. The cays are built upon exposed coral reefs in the open sea. Reef exposure is stated by Colman to be due to two causes: a fall in sea level and the formation of beach rock. Beach rock develops as a result of the precipitation of calcium carbonate from sea water at the site of turbulent wave action. Sea water, rich in carbon dioxide, can dissolve calcium carbonate much more readily than normal sea water. The excess is precipitated when the carbon dioxide is removed by violent aeration. The calcium carbonate so deposited binds together the sand and rock particles immediately underneath the surface. Cay formation is initiated by the accumulation of sand in the hol-

lows of flat reefs or behind boulders and rock fragments thrown up by storm action. The sand itself contains a high proportion of the pulverised remains of calcareous marine animals and plants, amongst which *Halimeda opuntia* is conspicuous.

Lime Cay

Lime Cay is the largest of the Port Royal group, being about 30,000 sq. yd. in area. It supports the most advanced type of vegetation. Unfortunately, it has not been surveyed but the sketch shown in Fig. 13 is a reasonably accurate representation of its structure. The cay consists of sand, containing a high proportion of the remains of the lime-encrusting *Halimeda opuntia*, which has been deposited on the leeward side of an arc-shaped reef. Members of the strand pioneer and strand dune associes are represented on the shore-line whilst the elevated central area shows the initiation of cactus-thorn scrub. Two areas, one in the north and one in the south, support *Rhizophora* and *Avicennia*. They were originally lagoons, they later became salt ponds and now are drying up with the result that *Rhizophora* is disappearing. There are also some plants, scattered and few in number, which are representative of strand woodland. The accompanying list shows the species found on this cay arranged according to the communities they represent.

Strand Pioneer and Strand Dune.

- Calonyction tuba*
- Euphorbia buxifolia*
- E. blodgettii*
- Heliotropium curassavicum*
- Boerhavia scandens*
- Cakile lanceolata*
- Philoxerus vermicularis*
- Sesuvium portulacastrum*
- Canavalia obtusifolia*
- Sporobolus virginicus*
- Scaevola plumieri*
- Caesalpinia bonducella*
- Tribulus cistoides*

Cactus-thorn Scrub:

- Pithecellobium unguis cati*
- Acacia tortuosa*
- Capparis ferruginea*
- C. cynophallophora*
- Morinda royoc*
- Opuntia tuna*

Strand Woodland:

- Cordia sebestana*
- Coccoloba uvifera*
- Thespesia populnea*
- Cassia emarginata*
- Piscidia piscipula*
- Morinda citrifolia*

Mangrove Woodland:

- Rhizophora mangle*
- Avicennia nitida*
- Laguncularia racemosa*
- Conocarpus erecta*
- Batis maritima*

Also found were:

- Indigofera tinctoria*
- Passiflora suberosa*




FIG. 14. Vegetation maps of the Port Royal Cays in 1939 (after Chapman) and 1952.

Maiden Cay

Maiden Cay is the smallest of the group and occupies an area of approximately 2,000 sq. yd. It consists entirely of sand, mostly under tidal influence, so that only about 200 sq. yd. is available for colonization. The highest point is only 4 ft. above the high tide mark. Fig. 14 shows that, from the period 1939 to 1952, there has been some slight change in the shore outline of the Cay due to sand movement but that the area occupied by vegetation has remained substantially the same. Only the pioneer and dune associates are represented and the latter is more in evidence at the moment indicating recent erosion. Since 1939, Sesuvium and Sporobolus have increased slightly in area whilst Euphorbia has spread into the Sporobolus zone. Two plants of Canavalia and one of *Paspalum vaginatum* have appeared and there are 4 specimens of Ipomea, none of which is flourishing extensively. Tribulus and Cakile, found in 1939, were not seen in 1952.

The contrast with the neighbouring Lime Cay is marked. Only 6 species of plants are to be found. Nevertheless, this small island, little more than a sand shoal, has remained more or less static for 13 years, during which time it has withstood the force of the major hurricane of 1951.

Gun Cay

Gun Cay was not visited by the authors but Chapman's vegetation map is reproduced (Fig. 14). Having an estimated area of 2,500 sq. yd., it is almost completely colonized so that the vegetation covers over ten times the area of that on Maiden Cay. The sand communities are represented only by Sesuvium and Sporobolus but all the mangroves are present as well as Batis and Thespesia. Chapman states that Tribulus was also recorded but he does not show it on his map.

Strand Pioneer and Strand Dune:

Sesuvium portulacastrum
Sporobolus virginicus
Tribulus cistoides

Strand Woodland:

Thespesia populnea

Mangrove Woodland:

Avicennia nitida
Laguncularia racemosa
Batis maritima
Rhizophora mangle

Drunkenmans Cay

Drunkenmans Cay, having 9,000 sq. yd. of land area, is characterized by having an outer "J" shaped rim of rock fragments in the shelter of which a lagoon has developed. The diagrams (Fig. 14) show that, since 1939, there has been considerable sand accumulation at the sea entrance to the lagoon. This may have been assisted by the protection afforded by a wreck which beached close to the windward east shore in 1948. Consequently, the main changes are an increase in the area covered by the sand communities and some encroachment of Rhizophora in the

lagoon. Representatives of the maritime communities are shown below:

Strand Pioneer and Strand Dune:

Sporobolus virginicus
Sesuvium portulacastrum
Heliotropium curassavicum
Philoxerus vermicularis

Strand Woodland:

Thespesia populnea

Mangrove Woodland:

Rhizophora mangle
Avicennia nitida
Laguncularia racemosa
Conocarpus erecta

Chapman also records one plant each of *Tournefortia gnaphalodes* and *Euphorbia buxifolia*.

South East Cay

South East Cay has an area of 6,000 sq. yd. and has been built up in the shelter of an extensive rampart of rock fragments over 6 ft. high. It has undergone one minor change since 1939. The north shore has lost most of the Rhizophora and Avicennia and a small piece of coast due to storm action. There is also evidence that Avicennia is spreading on the muddy south shore where considerable protection is afforded by the rampart. Batis, too, has spread in the rocky centre of the island occupied by Laguncularia. *Sesuvium portulacastrum* now occupies a small patch of sand on the rampart. The communities present and their representatives are:

Strand Pioneer and Strand Dune:

Sesuvium portulacastrum
Sporobolus virginicus
Philoxerus vermicularis
Heliotropium curassavicum

Strand Woodland:

Thespesia populnea (one plant).

Mangrove Woodland:

Rhizophora mangle
Avicennia nitida
Laguncularia racemosa
Conocarpus erecta
Batis maritima

Morant Cays

The Morant Cays were not visited by the authors but Chapman has a vegetation map of Middle Cay, which has an area of over 14,000 sq. yd. This Cay may be taken as typical of those which are characterized by having a preponderance of Sesuvium and an absence of Batis and mangroves. The mere presence of such species as Boerhavia, Tournefortia and Turnera, occasional members of the tension belt between the dune and scrub associates on the mainland, is noteworthy although they are sparsely distributed.

Strand Pioneer and Strand Dune:

Cakile lanceolata
Ipomea pes-capri
Sesuvium portulacastrum
Portulaca oleracea

- Phloxeris vermicularis*
Sporobolus virginicus
Turnera ulmifolia
Boerhavia scandens

Strand Scrub:

Tournefortia gnaphalodes (one plant).

The cays are very reminiscent of the sand spits, both as regards their vegetation and methods of topographic development. Thus, sand cays such as Maiden Cay and Lime Cay in which the area, elevation and degree of protection depend entirely on sand accumulation, find their counterpart in sandy spits such as the Hellshire Spit. Rocky cays such as Drunkenmans Cay and South East Cay are similar to the Palisadoes where, in the early stages of development, considerable protection is afforded by the piling up of rock fragments. The suggestion that the Palisadoes has been formed by a fusion of isolated cays may appear to be a reasonable one but the possibility of its development by sand accumulation on the lee side of an extensive rampart should be borne in mind. In both types of cay, of course, sand accumulation is a major factor in the development of both topography and vegetation.

The zonation of plant communities also follows the same general pattern on both the cays and the spits. On the small Maiden Cay, as on the narrow and youngest part of the spits, only the Strand Pioneer and Strand Dune associates are represented. Increase in area and elevation, as on Lime Cay and the wider parts of the spit, leads to the appearance of cactus-thorn scrub and members of the *Coccoloba*-*Thespesia* association. Moreover, at any stage where there is sufficient protection (which may be relatively late on the sand cays and early on the rocky cays), mangrove woodland is to be found.

Facts concerning the vegetation of the cays in relation to area and distance from the mainland have been assembled in Table 5. There appears to be direct relationship between number of species and size in the case of the Port Royal Cays. Lime Cay, the largest, yields the greatest number of species (33) representing 5 different communities and it is the only one in which the cactus-thorn scrub is represented. At the other extreme, Maiden Cay only supports 6 species representing two communities. The other members of the Port Royal group are intermediate in size and number of communities present. It would be unwise, however, to conclude that size is the only factor. It has already been pointed out that mangroves will appear at any time providing there is sufficient protection. Such protection is not governed by area but by topography so that the configuration of the land necessary to provide sufficient shelter may be present very early in the life of the rocky cays even when they are small; in the case of the sand cays, this must await increase in elevation by sand accretion which is related to area. Such considerations might equally well be applied to the development of strand woodland. Furthermore, the area-vegetation concept again breaks down in the case of the Morant Cays. Middle Cay is next in size to

TABLE 5. The plant communities on the cays in relation to area and distance from mainland.

Cay	Distance from mainland in miles	Area in sq. yds.	COMMUNITIES REPRESENTED						Total number of spp.
			Pioneer	Dune	Strand woodland	Mangrove	Cactus-thorn scrub	Total	
Lime.....	1.2	30,000	X	X	X	X	X	5	33
Drunkenmans.....	1.8	9,000	X	X	X	X	—	4	11
South East.....	2.8	6,000	X	X	X	X	—	4	10
Gun.....	0.25	2,000	X	X	X	X	—	4	9
Maiden.....	1.8	200	X	X	—	—	—	2	6
Middle.....	30.0	14,000	X	X	—	—	—	2	10
Pedro.....	60.0	X	X	—	—	—	2	5

Lime Cay and would appear to be of the same type, yet only ten species are present. These represent three communities, and mangroves are not included. It may well be that distance from the mainland is a factor here, a hypothesis receiving further support when it is noted that, according to Lewis (1947), the Pedro Cays, 60 mi. from the mainland, have only five species of land plants, *Sesuvium portulacastrum*, *Portulaca oleracea*, *Cyperus brunnescens*, *Tournefortia gnaphaloides* and *Suriana maritima*. In conclusion, therefore, the major factors to be co-related when considering the vegetation of the cays, are area, degree of protection and distance from the mainland.

THE LOWLAND COMMUNITIES

LIMESTONE

Two distinct vegetational units on limestone rock are recognized in this paper, i.e. dry limestone scrub forest and wet limestone forest.

Some two-thirds of the island is a limestone plateau and this is composed of hard limestone rock and derived soils which are about equal in area. The soil is predominantly of the bauxitic "terra rossa." Most of this is under cultivation and is dealt with separately.

On the rocky areas, rainfall is the controlling factor. The coastal limestone ranges of the south are arid to an extreme, weathering is slow and humus and organic matter almost absent. The area is greatly influenced by the seasonal rainfall and has more than six months when rainfall is under 4 in. Inland, the rainfall is high (75-100 in.) and, although the drainage is extremely rapid, the drought factor is moderate (See Fig. 6).

Aspect is related to wind exposure, humidity and drainage, but altitudinal changes, limited to about 3,000 ft., have little effect on the climax vegetation although they may influence secondary growth types.

Geologically, there are six different types of limestone in Jamaica ranging from cretaceous to recent coralline types. By far the most common is the white limestone of the Upper Eocene and Lower Miocene. This is hard and massive and, under certain climatic conditions, gives a characteristic clinker honeycomb rock common to the arid coastal hills and much of the karst area of the Cockpits. The most important

distinction for ecology is, however, between hard and soft (chalky) limestones which give rise to different soils.

Swabey (1942) considered the "limestone formations" as edaphic climaxes. Dry limestone scrub forest is developed where the rainfall is not more than 40 in. per year, with some months having below 2 in. rainfall. It is particularly evident on the south coast and includes an arid coastal faciation. Wet limestone forest with its moist faciation is typical of the more elevated inland areas such as the Cockpit country and Mt. Diablo plateau. In the dry regions, the rock is almost bare, such soil as exists being confined to the pockets and fissures of the rock. In the wet regions, erosion has proceeded more rapidly to give small hollow basins (Cockpits) in which bauxite soil has accumulated to a depth of from 5-50 ft. The wet limestone areas, then, necessarily have two aspects—the rocky slopes of the rim surrounding the cockpit and floor of the basin. Many of the latter are now under cultivation.

DRY LIMESTONE SCRUB FOREST AND THE ARID COASTAL FACIATION

Along the southern coast of Jamaica there is a series of rocky limestone hills and ranges, for the most part under 2,000 ft. elevation. These areas have a rainfall of less than 40 in. during the year, distributed in two rainy seasons. There is a drought period of more than six months. The north coast has few such areas, an example being the north coast of Trelawny (Fig. 6). Never very far from the sea and often extending through to the coast, these arid hills are of fissured, honey-comb, white limestone. They support a xeric scrub-forest of particular interest to the ecologist.

The main limestone areas of the south coast from east to west are found at Morant Bay, Yallahs, Dallas and Long Mountain, and, west of Kingston harbour, at the Hellshire Hills and Portland Ridge. Further inland and running at right angles to the coast, are the Don Figueroa and Santa Cruz Mts., in St. Elizabeth Parish. Around Negril at the extreme western part of the island, is an elevated coral shelf. The dry limestone hills of Trelawny on the north coast are low-lying and some three miles from the sea. Where the limestone reaches the sea, there is a narrow zonal aspect which is here considered as an arid coastal faciation.

DRY LIMESTONE SCRUB FOREST

Dry limestone scrub forest is a sparse, vegetational cover of low forest and tall scrub growing on bare limestone rock. No soil is present except for that deposited in small crevices or washed down to level areas. Leaf litter is almost nil and the floor is either a jumble of broken stone or a more or less continuous mass of jagged honeycomb rock.

The plants find support and sustenance by developing long, branched root systems which sprawl over and around the rocks to enter every crevice and canny. Species make-up may change rapidly from station to station, depending on aspect, slope and drainage as well as on soil deposits.

Red birch (*Bursera simaruba*) is ubiquitous as a scattered emergent tree, prominent for its copper red, flaky bark and complete deciduousness in the dry season. It has been suggested that its prevalence is due to fire resistance and selective cutting as the wood is regarded as useless for lumber.

In sheltered ravines, the tall cotton tree (*Ceiba pentandra*) may invade from the dry coastal plains and savanna. *Tecoma stans*, also deciduous, appears out of place here with its soft green pinnate leaves and yellow trumpet flowers. A host of small xerophytic trees and shrubs make up the mass of the vegetation. On a typical ridge such as Long Mountain, three narrow altitudinal zones may be correlated with soil accumulation and drainage. There is a lower zone where species from the adjoining alluvial plains form a tension belt on the bauxite soil. These include *Acacia*, *Haematoxylon*, *Guaiacum*, *Tamarindus*, *Albizia*, *Piscidia*, *Leucaena* and *Guazuma*.

A middle belt is dominated by low shrubs (*Croton*, *Tecoma*, *Lantana*, *Morinda*, *Eugenia*, *Vernonia*). The summit is often a thicket of *Rhus*, *Dipholis*, *Gymnanthes*, *Hypelate* and others. Larger trees are found in the moist ravines and may include a few of the trees from the moist inland limestone (*Brosimum*, *Nectandra*).

There is no distinct stratification, and structure varies from low scrub to a thin forest with trees 40 ft. and over. The canopy is always thin, even when continuous. Many of the species are semi-deciduous during the dry season. The trees are thin-boled, spindly and branch low to the ground.

Climbing and scrambling plants are well represented, although true lianes are absent. Epiphytes are restricted to xerophytic bromeliads, orchids and cacti. While some of the plants are thorny (*Randia*), this is not characteristic. Leaf size and texture is predominantly microphyllous and coriaceous. Many of the plants (*Croton linearis*) show a prolonged daily wilting at the height of the dry season. Compound leaves are seen in *Bursera*, *Rhus*, *Piscidia*, *Tecoma*, *Hypelate*, *Simaruba*, *Spathelia*, *Amyris*, *Comoeladria*, and *Pieramnia*.

A ground layer is frequently absent and, when present, is made up of a few ferns, *Peperomia* sp., cacti or woody perennial herbs. Xerophytic pasture weeds readily colonize open areas where depth of soil is sufficient.

Small palms and agaves give character when present.

FLORISTIC COMPOSITION—Trees (from 15-40 ft. according to station) arranged in approximate order of frequency are:

- Bursera simaruba*
- Tecoma stans*
- T. leucozyalon*
- Hypelate trifoliata*
- Bauhinia divaricata*
- Rhus metopium*
- Amyris balsamifera*
- A. elemifera*
- Gymnanthes lucida*
- Dipholis salicifolia*

Spathelia sorbifolia
Clusia rosea
Capparis ferruginea
Gymnanthes elliptica
Comocladia pinnatifolia
Picramnia antidesma
Simarauba glauca
Capparis cynophallophora
Diospyros tetrasperma
Krugiodendron ferreum
Guettarda argentea
Sarcophalus laurinus
Piper amalago
Sabal jamaicensis, the broadthatch palm, may be locally dominant.

Under Shrubs

Small shrubs of 10 ft. and more: (Order given approximates to frequency except that all members of a genus are grouped together).

Eugenia axillaris
E. monticola
E. rhombea
Allophylus pachyphyllus
A. jamaicensis
A. cominia
Psidium albescens
Brya ebenus
Capparis flexuosa
Pisonia aculeata
Tournefortia hirsutissima
T. astrotrichia
Guettarda elliptica
Canella winterana
Morinda royoc
Portlandia grandiflora
P. latifolia
Erythroxylon rotundifolium
Bernardia caprinifolia
Adelia ricinella
Senecio discolor
Casearia hirsuta
C. nitida
Cestrum diurnum
Ayenia laevigata
Phyllanthus angustifolius
Bocconia frutescens
Psychotria balbisiana
Eupatorium dalea
Rondeletia trifolia
R. hirta
Caesalpinia sepiaria (naturalized)
Daphanopsis occidentalis

Small bushes up to 5 ft. or more include:

Croton linearis
C. humilis
C. glabellus
C. flavens
C. grisebachianus
Lantana crocea
L. involucrata
Randia aculealata
R. jamaicensis
Malpighia glabra
M. puniceifolia
Schaefferia frutescens
Abutilon giganteum
Vernonia divaricata

Rondeletia tomentosa
Melochia tomentosa
Castela macrophylla
Acalypha scabrosa
Helicteres jamaicensis
Argythamnia candicans

Climbers:

Abrus precatorius
Stigmaphyllon emarginatum
Cissampelos pareira
Clematis dioica
Smilax balbisiana
Passiflora perfoliata
P. rubra
P. suberosa
Paulinia barbardense
Serjania laevigata
Chiococca alba
Echites suberecta
E. umbellata
Thunbergia fragrans
Cucumis anguria
Ipomea quinquefolia
I. umbellata
I. sidaefolia
I. jamaicensis
Cardiospermum grandiflorum
Gouania lupuloides
Centroserma virginianum
Mucuna pruriens
Galactia pendula
Bidens reptans
Durantia plumieri
Lasciacis divaricata (rambling bamboo grass)
Aristolochia odoratissima
Metastelma sp.
Mikania sp.

Epiphytes:

Tillandsia recurvata
Broughtonia sanguinea
Cereus flagelliformis

Parasites:

Phorandendron spp.

Herbs:

Stachytarpheta indica
Corchorus siliquosus
Heliotropium parviflorum
Asclepias curassavica
Andrographis paniculata
Blechum brownei
Tournefortia volubilis
Cordia globosa
C. cylindrostachya
Erigeron canadensis
Eupatorium odoratum
Sonchus oleraceus
Verbesina pinnatifida
Vernonia divaricata
Amaranthus viridis
A. spinosus
Euphorbia hirta
Tragia volubilis
Priva echinata
Spigelia anthelmia
Sida procumbens
S. rhombifolia
Argemone mexicana

Rivina humilis
Crotalaria retusa
C. verrucosa
Desmodium supinum
Waltheria americana
Turnera ulmifolia
Croton ovalifolius
Salvia occidentalis
Hyptis capitata

The ferns *Pteris longifolia*, *Adiantum melanoleucum*, *A. tenerum*, *Cheilanthes microphylla* are widespread but not abundant.

The above type of forest which is typical of the dry coastal hills at the present time, has been subject to much human interference by periodic burning. The larger trees have been removed for timber and firewood so that little remains of former canopy and sub-canopy trees. Fortunately, there is still one relatively undisturbed area at Portland Ridge on the south coast. Some of the more valuable timber trees here have been removed but, nevertheless, sufficient remain to give a good representation of the type of forest that formerly existed in the dry limestone regions. The identity of many of the trees from this area forms the subject of a valuable report by Lewis (1947) who recognized over 80 species. Since the lower layers of the forest are essentially similar in structure and species to that already described, although less well-developed in this denser woodland, remarks are confined to the tree layers. The canopy is open and varies in height according to station from 40-60 ft. with sporadic emergents up to 80 ft. or more. The sub-canopy is more closed, reaches a height of 20-35 ft. and may include the palms *Coccothrinax fragrans* and *Thrinax parviflora* (Fig. 15). The maximum development is reached in the flatter areas of the ridge where leaf litter and humus cover the limestone. Only four species are completely deciduous. *Swietenia mahogoni*, *Bursera simaruba*, *Spondias monbin* and *Plumiera* sp. However, many of the species classed as evergreen lose some of their leaves during the prolonged dry season.

Floristic Composition

Occasionally emergent (70-80 ft.)

Chlorophora tinctoria
Pisonia fragrans
Rhamnidium jamaicense
Sapota sideroxylon
Swietenia mahogoni
Tecoma leucoxylon

Canopy (40-60 ft.)

Adelia ricinella
Allophylus jamaicensis
Amyris elemifera
Brosimum alicastrum
Bumelia retusa
Bursera simaruba
B. simplicifolia
Caesalpinia vesicaria
Calyptrothecia chytracaulis
C. pallens
C. zuzygium
Citharexylum fruticosum



FIG. 15. An open aspect of Dry Limestone Forest at Portland Ridge. The palm is *Thrinax parviflora*.

Colubrina reclinata
Cordia gerascanthoides
C. nitida
Diospyros tetrasperma
Erythroxylon areolatum
Eugenia axillaris
Exothea paniculata
Exostema caribaeum
Guazuma ulmifolia
Hippomane mancinella
Krugiodendron ferreum
Matayba apetala
Melieocca bijuva
Picrodendron baccatum
Pisonia obtusata
Rhus metopium
Sideroxylon foetidissimum
Tamarindus indica

Sub-Canopy (20-30 ft.)

Albizia berteriana
Anona squamosa
Bauhinia divaricata
Bourreria succulenta
Bumelia rotundifolia
Brya ebenus
Canella winterana
Capparis cynophallophora
C. ferruginea
C. flexuosa
Cascaria quianensis
Cassia emarginata
Cereus peruvianus
Clusia flava
Coccoloba spp.
Coccoloba krugii
Coccothrinax fragrans
Comoeladria velutina
Cordia sebestana
Crescentia cujete
Croton glabellus
Erythroxylon rotundifolium
Eugenia monticola
Guicum officinale
Gymnanthes lucida
Haematoxylon campechianum
Hypelate trifoliata
Linociera ligustrina
Ocotea jamaicensis

- Oxandra lanceolata*
Peltophorum brasiliense
Phyllanthus acuminata
Piscidia piscipula
Plumeria sp.
Sarcomphalus laurinus
Thrinax parviflora
Ximenia americana
Zanthoxylum fagara
Z. spinosum

It will be evident then that the forest here has a very rich flora. It provides an important remnant that would well repay detailed study.

ARID COASTAL FACIATION OF DRY LIMESTONE SCRUB FOREST

The environment of the arid coastal regions of the Caribbean has been graphically described by Seifriz (1943): "The heat is intense, the light blinding, every plant armed, no water, no shade, no trail leading anywhere, as awe-inspiring, as fearful, as superb a picture of the eternal persistence of life under the most adverse conditions that nature can produce."

Perhaps the most notable aspect is the mass of bare, jagged, honey-comb limestone over which one must clamber. The coastal limestone has weathered into a sharp clinker-like, honey-comb rock which has a metallic ring to hob-nailed boots. Sharp-pointed and knife-like edges surround the many small crevices and pockets where the sparse soil and organic debris may accumulate. Every crack and fissure is utilized for the tenacious, if precarious, root-hold of tree and shrub, many of which appear to be growing out of solid rock.

For a distance of over 50 mi. along the south coast from Morant Bay to Portland Point, there is an almost unbroken rim of low limestone hills where this aspect of the limestone vegetation is well developed. The area is arid to an extreme, total rainfall is below 30 in. and there are 6 to 10 months having less than 4 in. with some under 1 in.

The xerophytic vegetation is tolerant of salt spray and sweeping winds. The air has a high evaporating power. It is a littoral woodland on coastal limestone made up of hard-leaved, dry limestone shrubs together with cacti, halophytes and salt-resistant trees not encountered inland. The flora is relatively rich for such an area and merges imperceptibly into typical dry limestone scrub-forest species.

A feature of the faciation is the presence of cacti of which the dildo (*Cereus*), forming columns up to 20 ft. high, is abundant. Such cacti occur only here and in the thorn-scrub in Jamaica.

In sheltered bays, the littoral fringe of the limestone may be occupied by *Rhizophora* although Chapman regarded this as an unstable community. Where mud has been deposited, other mangroves may be present in front of the limestone rock. Small bays in the limestone coast are frequently enclosed by spit formation, first forming a lagoon then a saline flat with typical halophyte communities. In the same way, sand deposited to seaward gives typical strand trees (*Coccoloba*, *Thespesia*) within limestone areas. Again,

where alluvial plains reach the coast, *Prosopis* and *Acacia* may invade soil pockets on the limestone.

Of the limestone scrub-forest itself, typical coastal trees are:

- Hippomane mancinella*
Cordia sebestana
Plumiera alba
Piscidia piscipula
Comocladia velutina
Chrysobalanus icaco
Parkinsonia aculeata (naturalized).
Jacquinia armillaris
Citharexylum fruticosum
Capparis ferruginea
Caesalpinia vesicaria
Pithecellobium unguis-cati

Other common trees and shrubs in approximate order of frequency are:

- Rhus metopium*
Bauhinia divaricata
Hypelate trifoliata
Bursera simaruba
Pisonia obtusata
P. fragrans
Gymnanthes lucida
Eugenia monticola
Amyris elemifera
Matayba apetala
Calyptrotheces chytraculia
C. pallens
Capparis flexuosa
C. cynophallophora
Krugiodendron ferreum
Peltophorum brasiliense
Clusia flava
Canella winterana
Sapota sideroxylon
Guaiacum officinale
Oxandra lanceolata
O. laurifolia
Beurera succulenta
Cordia gerascanthoides
C. nitida
Bursera simplicifolia
Coccoloba spp.
Tecoma leucoxylon
Ficus populnea

Occasional trees and shrubs:

- Exostema caribaeum*
Erythroxylon aveolatum
E. rotundifolioum
Melochia tomentosa
Laetia thamnia
Diospyros tetrasperma
Eugenia axillaris
Sarcomphalus laurinus
Zanthoxylum fagara
Z. flavum
Exothea paniculata
Meliococca bijuga
Bumelia retusa
B. rotundifolia
Lasiocroton macrophyllus
Securinega acidoton
Picradendron baccatum
Adelia ricinella
Phyllanthus acuminatus

P. linearis
Sideroxylon foetidissima
Gymnidia latifolia
Casuarina guianensis
Swietenia mahogoni
Catesbeia parviflora
Agava americana
Croton linearis

Palms:

Coccothrinax fragrans
Thrinax excelsa
T. parviflora

Herbs:

Heliotropium curassavicum
Melochia tomentosa
Schaefferia frutescens
Ambrosia hispida
Helicteres jamaicensis
Jatropha gossypifolia

Climbers:

Chiococca alba
Cissampelos pareira
Cissus cucurbitacea
Metastelma harrisi

Cacti:

Cereus peruvianus
Opuntia tuna
O. jamaicensis
O. spinosissima
Melocactus communis

WET LIMESTONE FOREST

This community is developed in areas of limestone rock where the rainfall is over 75 in. and may range up to 150 in. For the most part, it is at elevations from 1,000 ft.-2,500 ft. and situated inland. No doubt, there are different fasciations within this type which will be recognized with further study.

The largest area of wet limestone forest is in the Cockpit country, although the type is found on certain of the limestone peaks (Mt. Diablo and Dolphin Head) along the central axis of the island. It is also present on some coastal ridges of the wet northwest coast.

Mt. Diablo, 3,300 ft. is a precipitous, calcareous plateau, very similar to the Cockpit area further west. Dolphin Head (1,780 ft.) is a limestone peak in the centre of calcareous shales.

The John Crow Mountains of limestone in the extreme eastern end of Jamaica do not fall into this unit. Due to the very high rainfall of some 300 in. per year, and their height of almost 4,000 ft., they are clothed with lower montane rain forest on the lower slopes and an elfin woodland on the summit.

The Cockpit country is so-called because the whole area is made up of circular depressions (dolinas) up to 500 ft. deep filled by bauxitic soils with accumulated humus from the surrounding rim of limestone rock. The area is typical karst country where underground drainage, subterranean rivers, sinkholes and caves are common. Many of the depressions are the result of sinkholes from underground streams.

The reduction of the Cockpit plateau has not yet proceeded to the level seen in comparable areas in

Cuba and Porto Rico where the "mogotes" or haystack hills represent blocks of rock partly weathered down and having their bases engulfed by the rising plain of soil about them. With time, many of the smaller valleys and depressions in the Cockpit country will join up until only the highest hills will be left.

Burning and cultivation have affected many areas, both in the valleys and on slopes with the production of "ruinate"; however, much still remains relatively undisturbed.

The forest is of two types—that on the bare limestone rock and that on the valley floor. The soil of the latter, which may be anything from 5 to 50 ft. in depth, is bauxitic and has very free drainage, a neutral to alkaline pH and a general deficiency in phosphates. There is, of course, much overlapping of species between the two types of forest and it is mainly the general physiognomy that is different on rock and soil (Fig. 16). As previously mentioned, most of the valleys have been cleared and planted with bananas and other crops such as sugar, yams, coco (Colocasia) and cassava. Fig. 17 shows a clearing in the forest that has been planted with Colocasia and Yams.




FIG. 16. Diagrammatic profile of typical Cockpit country supporting Wet Limestone Forest. Different aspects can be related to depth of soil which is greatest in the valley bottom and grades to bare rock on the slopes. The larger valleys are now under cultivation or pasture.



FIG. 17. Cultivation of Colocasia and Yam in interior of Cockpit country.

The wet limestone forest (Fig. 18) is far more mesophytic and luxuriant than the dry type, having more forest trees, epiphytes and lianes, aroids, bromeliads and orchids. Undergrowth is often sparse due either to the rocky substratum or the dense shade. However, there are more open aspects where *Aechmea paniculigera*, *Polypodium* spp., and aroids are prevalent (Fig. 19).



FIG. 18. View of Wet Limestone Forest looking down into one of the cockpits.



FIG. 19. An open aspect of Wet Limestone Forest showing climbing aroids, *Aechmea paniculigera* (bottom right) and *Polypodium crassifolium* (bottom left).

STRUCTURE

The canopy of the forest is more or less uniform at 50-60 ft., except that in the Cockpits *Terminalia latifolia* and *Cedrela odorata* are emergent trees, 80-100 ft. high. The canopy, although typically closed, is never dense and is made up of tall thin-boled trees 1-2 ft. in girth and branching at 40 ft., into a wide spreading crown with little depth.

A sub-storey of small trees averaging 40 ft. in height may be distinguished. This is often quite dense and made up of its own characteristic species.

The shrub and field layers are sparser than the strata above and often merge together. In fact, undergrowth is generally lacking due to the rocky substratum. Small buttresses may be seen but are not a feature. *Clusia* and *Lucuma* develop stilt roots.

The larger trees show scattered distribution but frequently the forest is composed of closely spaced, small trees and shrubs of similar life form, forming a type of tall thicket.

Much local variation occurs due to sharp changes in aspect and drainage. The vegetation of valleys is consistent but the slopes may reflect wet and dry aspects by the presence and luxuriance of the epiphytes and ground herbs, whilst tree species differ. Soil depth and leaf litter are noticeably less in the drier aspects. The profile diagram (Fig. 20) is of a typical moist slope forest.




FIG. 20. Profile diagram of Wet Limestone Forest measured in the Cockpit country. The trees are included from a strip 100 ft. long and 25 ft. wide.

KEY TO SYMBOLS: Ai. *Andira inermis* (Wormwood); Ba *Brosimum alicastrum* (Breadnut); Bb *Bucida buceras* (Olive bark tree); Bc. *Buchenavia capitata* (Mountain wild olive); Bl. (Bloodwood); C. *Coccoloba* sp. (Grape); Cc. *Cinnamomum corticosum* (Mountain cinnamon); Cs. *Comocladia* sp. (Maiden Plum); D. *Dipholis* sp. (Bulletwood); Dn. *Dipholis nigra* (Galimenta, Red Bullet); Ds. *Dipholis salicifolia* (Black bullet); E. *Evodia* sp. (Rodwood); Ep. *Exotheca paniculata* (Wild Ginep); Fo. *Faramea occidentalis* (Wild Coffee); Gg. (Alligator, Wild akee); H. *Drypetes lateriflora* (*G. area* glabra (Alligator, Wild akee); Kf. *Krugiodendron ferreum* (Black (Whitewood, Guiana plum); Lf. *Lasioscrotum fawcettii*; N. *Nectandra* sp. (Sweet-ironwood); Ol. *Oxandra lanceolata* (Black lancewood); Op. *Ochroma pyramidalis* (Balsamwood); Pla. *Plumeria alba* (Frangipani); Si. g. *Simaruba glauca* (Bitter damson); Sj. *Sapium jamaicense* (Blind-eye); Vu. *Vitex umbrosa* (Fiddlewood, Boxwood); U.K. Unidentified.

Lianes are represented by climbers, some of which may also be classed as epiphytes. Climbing aroids are abundant throughout and everywhere their long, dry, thin roots hang down from the trees. *Polypodium* is a common climbing fern. Other climbers are: the prickly Smilax, *Dioscorea*, *Rajania cordata* (found only in the Cockpit area), and leguminous vines.

Of the bromeliads, species of *Tillandsia* are trunk epiphytes. *Hohenbergia* is a true perching epiphyte confined to the crowns of the larger trees at 40 ft. *Aechmea paniculigera* is common on the floor in rock crevices, together with ferns and mosses.

Small shrubs are represented by Piperaceae, Melastomaceae, Rubiaceae and the short-thatch palm *Thrinax tesselata*. Tree ferns are localized although many other ground ferns reach considerable size.

The trees are evergreen but some are semi-deciduous in other habitats. Leaves, for the most part, are

simple, but compound leaves occur in *Pithecellobium*, *Simaruba*, *Mesquitylon*, *Picraena* and *Zanthoxylum*.

FLORISTIC COMPOSITION

The true climax in the small deep valleys is a *Terminalia*-*Cedrela* association. Both of these however, are valuable timber trees and the fact that they are found growing in fertile valleys has led to intensive exploitation.

A floristic name cannot be so easily applied to the forests of the rocky limestone slopes surrounding these valleys.

Large trees which are common enough throughout are: *Terminalia latifolia*, *Cedrela odorata*, *Calophyllum jacquinii* and *Pithecellobium alexandri*.

Frequent dominants of the canopy layer are the breadnut (*Brosimum alicastrum*), the sweetwoods (*Nectandra* spp.), and the bulletwoods (*Dipholis* spp.).

A list of the trees in approximate order of frequency is:

Brosimum alicastrum
Nectandra antillana
N. sanguinea
Dipholis nigra
Pithecellobium arboreum
Zizyphus chloroxylon
Ficus sp.
Zanthoxylum martinicense
Sloanea jamaicensis
Buchenavia capitata
Lucuma mammosa
Mimusops excisa
Podocarpus purdieanus
Cecropia peltata
Prunus occidentalis
Psidium montanum

The sub-canopy layer of small trees reaches 40 ft. and contains many species, not all of which may be present in any one aspect. This stratum often merges into the canopy above and species may be common to both.

Trophis racemosa
Antirrhoea jamaicensis
Zanthoxylum flavum
Simaruba glauca
Spathelia glabrescens
Mesquitylon jamaicense
Picraena excelsa
Eugenia axillaris
Eugenia spp.
Colubrina ferruginea
Matayba apetala
Prunus myrtifolia
Lagetta lagetto
Miscentea triandra
Gucrea glabra
Exothea paniculata
Sapium jamaicense
Amyris balsamifera
Bauhinia divaricata
Rondeletia spp.
Hufelandia pendula
Ocotea staminea
Xylopia muricata
Oxandra lanceolata

Comocladia pinnatifolia
Andira inermis

The families Sapotaceae and Lauraceae are well represented in the above lists.

The shrub layer includes a few smaller trees and bushes such as *Piper nigrinodum*, many Melastomaceae, *Bocconia arborescens*, *Carica jamaicensis*, *Coccoloba longifolia*, *C. diversifolia*, *Acidotom urens* and *Clusia rosea*.

Climbers and epiphytes include the climbing shrub, *Rourea paucifoliata*, aroids, *Begonia glabra*, *Dioscorea polygonoides*, the wild yam, and abundant *Polypodium heterophyllum*.

Bromeliads are frequent, many species of *Hohenbergia* being present such as *H. distans*, *H. eriostachya* etc.

The field layer has a few herbs such as *Gyrotaenia spicata*, *Peperomia amplexicaulis*, *P. cordifolia*, *Pilea ciliata*, *P. crassifolia* and *P. reticulata*, *Boehmeria jamaicensis* and, on the edges of clearings, *Pachystachys coccinea*.

Ferns are prominent and frequent members of this layer. The commonest dryopteroid is *Thelypteris serrulata*. Other characteristic species are *T. oligophylla*, *T. patens*, *T. venusta*, *T. sagittata*, *Ctenitis ampla* and *C. effusa*.

On limestone ledges throughout is to be found *Thelypteris asterothrix* especially where the rocks overhang above. One can assume that this species thrives best where not exposed to direct wetting.

Other ferns of the wet limestone forest include *Lomariopsis underwoodii*, found sterile except where scendent on tree-trunks, *Campyloneurum augustifolium*, *Polystichum christiana*, *Polypodium dissimile*, *P. heterophyllum*, the filmy ferns *Trichomanes* spp. and the ferns of moist glades such as *Dennstaedtia bipinnata*, with fronds up to 10 ft. long, *D. cicutaria* and *Pteris quadriaurita*.

Mosses are sparse but include:

Thuidium involvens (Hedw.) Mitt.
Leucobryum antillarum Schp.
Entodon macropus (Hedw.) C. M.
Fissidens donnellii Aust.
Hookeriopsis fissidentoides (H. & W.) Jaeg.
Orthostichopsis tetragona (Hedw.) Broth. (pendant from trees).
Isopterygium tencrum (Sw.) Mitt. (on trunks).

Wet limestone forest also includes a moist fasciation seen in such areas as the Red Hills, west of Kingston. Rainfall is the controlling factor and there is a mingling of the dry and wet limestone forest species with some modification in structure. This fasciation has not been fully studied.

VEGETATION OF THE BAUXITE PLATEAUS

This cannot be regarded as an ecological formation but it is a convenient heading for the areas of extensively cultivated limestone soils of Jamaica. The original vegetation was forest which merged into the present limestone forest types.

It has been observed that Jamaica is comprised of some three-quarters limestone and its derived soils.

Much of the limestone is still seen as rocky hills and ranges but, in karst areas, there are frequent depressions and sinkholes in which soil has accumulated. The parishes of Manchester and St. Ann constitute upland plateaus with a covering of residual bauxitic soils, varying much in depth, over the parent limestone. Such bauxitic soils are known locally as "terra rossas" and are now being actively exploited for bauxite, the ore of aluminium.

"Terra rossa" soils vary according to development and there are many "degraded" types where impurities give impeded drainage, or fertility values are lowered. Phosphate deficiency is general. The soils of such large inland valleys as St. Thomas in the Vale are regarded as lacustrine deposits from ancient lakes. The soil here is heavy clay with poor drainage and is generally acid (pH 4-5).

Rainfall over the "terra rossa" areas is, with one notable exception, over 60 in. per annum and there are few months with less than 4 in. The exception (Fig. 6) is along the north coast of Trelawny where the total annual rainfall is below 40 in. and there are at least six "drought" months with less than 4 in. "Terra rossa" here forms a narrow coastal strip under sugar cane or cattle pasture in which the tree members are essentially those migrating from the low, inland limestone ridges. *Swietenia mahogoni* and *Peltophorum brasiliense* are abundant, together with *Cordia gerascanthoides*, *Guazuma ulmifolia*, *Zanthoxylum flavum*, *Bauhinia divaricata*, *Esenbeckia pentaphylla*, *Pimenta officinalis*, *Bursera simaruba*, *Diphlois salicifolia* and *Haematoxylum campechianum*.

The plateaus of St. Ann and Manchester have a higher rainfall and are areas of intensive agriculture and grazing. The induced savannas of the cattle pens have many of the larger forest trees from the surrounding wet limestone forests (Fig. 21).

Magnificent specimens of the cotton tree (*Ceiba pentandra*) and the West Indian Cedar (*Cedrela odorata*) are seen. Common, too, are large fig and guango trees (*Samanea saman* syn *Enterolobium saman*). These old trees are laden with epiphytes,




FIG. 21. Remnant limestone forest and pasture on the Manchester plateau.

climbers, stranglers such as *Solandra grandiflora* with its showy, pale yellow, trumpet blooms, *Schlegelia parasitica*, figs and Clusiads, bromeliads, orchids, ferns and aroids.

Smaller trees which are also frequent as isolated specimens are:

Guazuma ulmifolia
Terminalia latifolia
Pithecellobium arboreum
Cecropia peltata
Nectandra patens
Hibiscus elatus
Hura crepitans
Pimenta officinalis

Widespread shrubs include:

Pisonia aculeata
Psidium guajava
Morinda royoc
Solanum verbascifolium

Roadside and pasture weeds are:

Lantana involucrata
Sida spp.
Bidens pilosa
Borreria laevis
Asclepias curassavica
Mimosa pudica
Crotalaria retusa
Rivina humilis
Bryophyllum pinnatum
Eupatorium odoratum
Desmodium supinum
Stachytarpheta indica
Cordia globosa
Waltheria americana
Achyranthes indica
Tragia volubilis
Bromelia pinguin
Barleria prionitis
Anemone adiantifolia

Climbers:

Echites umbellata
E. suberecta
Stigmaphylloem marginatum
Cissampelos pareira
Cissus sicyoides
Abrus precatorius
Clematis dioica
Momordica charantia
Tournefortia volubilis

In St. Ann, is found the largest area of rendzina soil developed over soft limestone. The hills are rolling, fertile, mainly under pasture and similar in aspect to the wetter shale regions.

While most of the "terra rossas" are free draining and neutral to alkaline depending upon the depth over the parent rock, the lacustrine basins are of greater age and the soil is termed "degraded terra rossa." Drainage is impeded, acidity high, whilst the texture is a heavy clay. These areas are extensively cultivated and grow breadfruit, citrus, mangoes, bananas, pineapples, coconuts, sugar and naseberry (*Achras sapota*).

Amongst the trees, many of the genera common to the plains and shale valleys are seen, such as:

Tamarindus, Chrysophyllum, Ceiba, Cecropia, Catalpa, Bauhinia, Samanea, Blighia, Poinciana, Terminalia, Erythrina, Spondias, Crescentia and Pithecellobium alexandrii.

The savannas of epiphyte-laden mango and guango trees, clumps of bamboo and *Haematoxylon* thickets show a distinct relation with the alluvial plains and lower shale hills, not shown by other communities. *Acacia* species are, however, altogether absent since they fall out rapidly once the alluvium is left.

ALLUVIUM

VEGETATION OF THE ALLUVIAL PLAINS

Large low-lying coastal plains are located along the dry southern coast. These have been formed as the broad alluvial deltas of meandering rivers. The rich alluvium of mixed sand, gravel and loam is spread out as a thick cover over the down-faulted limestone. On the coastline where ancient shallow bays were silted up, the alluvium is interbedded with layers of heavy marine clay. In the Ferry area, the history of such a plain can be easily visualised. The Rio Cobre has changed its course four times within colonial history, each time broadening out the plain. Alluvium is some 15 ft. in depth overlying 60 ft. of marine clay. In 1838, the river was diverted to a new mouth by a cut of some 7 mi. and, since that time, 200 acres of silt has been deposited at the mouth. Much of the saline area is at present being brought under cultivation from mangrove vegetation. Drainage first lowers the water table which kills the mangrove pneumatophores. Further flooding, drainage and cultivation prepares the land for para grass (*Panicum purpurascens*). Leaching continues and, within three years, sugar cane may be planted, being tolerant of salt in a concentration of 15 p.p.m.

The plains were fully exploited during the boom in sugar before the Napoleonic wars. With the decline in sugar after 1840, the cane areas decreased, large portions reverting to degraded, secondary types or turned over to cattle raising. For the most part, the alluvial plains are arable, fertile and suitable for irrigation, hence they are areas of high population and well-developed agriculture.

Starting from the east, are the small plains and river terraces of the Plantain Garden river in St. Thomas and the small deltas at the mouths of the Morant and Yallahs rivers.

Behind Kingston lies the dry Liguanea plain rising to 700 ft. a few miles inland and extending westward to join the Plains in Lower Clarendon parish. There are two important rivers here, the Rio Cobre and the Rio Minho. Much of the area is of irrigated cane fields. Nearer to the coast, there is a mosaic of highly saline and alkaline marine clays with Rio Minho alluvium and degraded "terra rossa" soils. The coastal areas are, therefore, unproductive and infertile (Croucher 1938).

The broad Manchester limestone plateau and limestone ranges separate the Clarendon Plains from the basin of the Black River which forms a low-lying flood plain reaching inland some 20 mi. along its course.

The whole area of some 100 sq. mi. is an extensive sedge marsh, with the exception of the small dry Plains of Pedro skirting the Santa Cruz ranges.

In Westmoreland Parish, are the small George plains traversed by the Cabaritta river. They are low-lying and swampy.

On the north coast, the descent from the central mountains is more abrupt with shorter and swifter rivers. Alluvium is found as river terraces where the soil is more silty and free from saline influences.

The plains may vary from a few to 15 mi. in width and may be continuous for 20 mi. Generally they are hot, dry, low-lying and usually bounded by limestone hills although the Liguanea Plain merges into shale hills. Rainfall is often as low as 30-35 in., though most of the plains have from 40-60 in. Rainfall is uneven and the drought shadow of less than 4 in. of rain, for six or more months, falls over the coastal fringe. There is a coastal aspect marked from the inland aspect by soil, altitude and rainfall. Differences in soils, which may be the fertile free-draining alluvium, heavy saline and acid or alkaline clays with impeded drainage, degraded bauxitic soils or mixture of all, have profound influence upon the vegetation over small areas. Some of the marine clays have a salinity of 500 p.p.m. for the upper 6 in. and a pH of 5.2. At 2 ft. depth, the figures are 3,000 p.p.m. and a pH of 4.5. The plant communities are all induced, modified, secondary types and will be described under the following headings:

- a) Cultivated areas and induced savanna.
- b) Secondary communities.
- c) Thorn scrub.

CULTIVATED AREAS AND INDUCED SAVANNA

In 1938, bananas provided over 50% of Jamaica's export trade and much of this banana land was on the plains. Coconuts are not extensive on alluvial soils. Important minor crops are tobacco, pineapples, tomatoes, maize, cashew nuts and, more recently, rice which will grow on irrigated marine clays.

The drier regions, where irrigation is so far impracticable, are given over to grazing without cultivation. Much is induced savanna of guinea grass (*Panicum maximum*) and the guango tree (*Samanea saman*) (Fig. 22). Guinea grass, now a common fodder grass, was unwittingly introduced from Africa with a consignment of birds in 1744. Other introduced grasses of the pastures are: *Andropogon pertusus*, *Cynodon dactylon*, *Sporobolus indicus*, *Paspalum fimbriatum* and *P. saccharoides*.

Many of the characteristic trees of the southern plains were introduced in early colonial times and have become naturalised. It would be difficult to reconstruct the original vegetation but, no doubt, it was a sparse forest composed of such trees as *Guiaicum officinale* ("lignum vitae"), a small tree with a bushy crown; *Catalpa longissima*, the handsome yokewood reaching up to 80 ft.; the deciduous cotton tree, *Ceiba pentandra*, of 120 ft.; *Hymenaea courbaril*; the gru gru palm, *Acrocomia aculeata*; and *Chlorophora tinctoria*. Smaller trees would have included



FIG. 22. Induced savanna of Guinea grass (*Panicum maximum*) and Guango trees (*Samanea saman*) on the alluvial plains.

Guazuma ulmifolia, *Cordia gerascanthoides*, *Melicocca bijuga*, *Ficus mamillifera*, *Enterolobium mangense*, *Crescentia cujete*, *Sapindus saponaria*, *Buchenavia capitata*, *Cassipourea elliptica*, *Trichilia hirta*, *Quiinia jamaicensis*, *Crataeva gynandra*, *Piscidia piscipula*, and *Cordia alba*. The shrubs *Sarcomphalus laurinus*, *Erythroxylon areolatum*, *Brya ebenus*, *Anacardium occidentale* and *Cassia emarginata* would also have been present.

It is sometimes doubtful whether a tree is naturalized or native since there are more introductions here than anywhere else on the island. *Samanea*, with its sweet, mucilaginous pods relished by stock, was introduced with cattle from Central America. Together with the ubiquitous mango tree, it forms a characteristic part of the landscape. Both are laden down with epiphytic bromeliads and cacti.

Albizia lebbeck, from the Far East, is the now familiar "woman's tongue" which reached Jamaica along with mango in 1782. The name refers to the chattering of the dry pods with the slightest breeze. Introduced *Cassia* species are common, together with such trees as: *Terminalia catappa*, *Adenanthera pavonina*, *Poinciana regia*, *Spathodia campanulata*, *Tamarindus indica*, *Casuarina equisetifolia*, *Moringa oleifera*, *Haematoxylon campechianum*, *Spondias mombin*, *S. purpurea*, *Zizyphus jujuba*, Coconut palms and several *Acacia* species.

Common roadside plants are: *Ricinus communis*, *Jatropha gossypifolia* and *Datura stramonium*. Also frequent are the twining, parasitic *Cuscuta americana* and polygonaceous *Antigon leptopus*.

SECONDARY COMMUNITIES

Ruinate is the delightfully apt term given to all marginal and abandoned lands in Jamaica. It is applied to the infinitely varied, secondary growth types of vegetation developing after burning, catch cropping and abandonment. On the alluvial plains, the common form that occurs is a thorn thicket or, under moderate cutting and grazing, a thorn-savanna (Fig. 23).

The type of ruinate found indicates the previous history of the area, whether, for instance, it is abandoned arable or pasture land, or marginal land



FIG. 23. Thorn savanna, a second growth type, on the alluvial plains. The trees are *Acacia lutea*, *Pithecellobium dulce* and *Acacia farnesiana*. The grass is *Andropogon pertusus*.

after catch cropping. The details of succession require further study.

Frequent pioneer weeds on denuded ground are: *Amaranthus spinosus*, *A. viridis*, *Cyperus rotundus*, *Heliotropium parviflorum*, *Portulaca oleracea*, *Cleome serrata*, *C. spinosa*, *Achyranthes indica*, *Pectis febrifuga*, *P. ciliaris*, *Phyllanthus niruri* and other Euphorbiaceae. These species are common following the rains as ephemeral adventives emphasizing the fact that succession in the dry lowland tropics is complicated by seasonal factors.

The initial stages merge into a more stable community of secondary invaders, a few pioneers remaining to take their place in the final stages of the succession. Such secondary invaders include: *Cynodon dactylon* and *Andropogon pertusus* with herb weeds such as *Crotalaria retusa*, *C. verrucosa*, *Waltheria americana*, *Sida rhombifolia*, *Mimosa pudica* and other semi-woody perennials.

At this stage, shrub and tree seedlings of such plants as *Psidium guajava*, *Haematoxylon campechianum*, *Capsicum frutescens*, *Cestrum diurnum*, *Lantana crocea*, *Pisonia aculeata*, *Acacia lutea* and *Solanum verbascifolium* make their appearance.

Such ruinate may pass through to a thorn thicket or thorn savanna dominated by *Acacia lutea*, with common pasture weeds such as *Borreria laevis*, *Desmodium supinum*, *Sida* spp., *Boerhavia scandens*, *Euphorbia heterophylla*, *E. hypericifolia*, *Alternanthera repens*, *A. parvifolia*, *Celosia argentea*, *Amaranthus* spp., *Leonurus sibiricus*, *Bidens pilosa*, *Gomphrena globosa*, *Plumbago scandens* and, dominating the shade circle of trees, *Ruellia tuberosa* and *Petiveria alliacea*.

THORN SCRUB

Described by Beard (1944) as thorn woodland, the type is scrubby and usually open, or if closed, thin and sparse. The total flora is poor, mesquite (*Prosopis juliflora*) making up 75% of the trees, the remainder being other members of the Mimosaceae and Caesalpiniaceae (Fig. 24). The slender, umbrella-like acacia trees are microphyllous, sparsely leaved, semi-deciduous and spiny. They may be 10-30 ft.

remains of
l that along
and Cymo-
part in the
n Kingston
that there
This was
little doubt
e aquatics,
rooted and
inks in the

the shallow
1 coast and
ocky Cays.
nts of Rhi-
, trees are
e holes and
; boulders.
the whole
rly defined
the limited
es between
ophora re-
ent of its
developed,
ral feet of
ne seaward
he deeper
d swamps.
wamps be-
hose areas
as in the
rarely or
ria is as-
margin of
m a pure
present on
a member
is regards
community
, and the
from his
the Cono-
Diagrams
es in the
by Chap-

depends
level and
s bringing
elevation
mangrove
the man-
x, usually
nd climax
quently a
mmunities
may also
at tenden-
tion—for



FIG. 24. *Prosopis juliflora* forming open thorn woodland on the alluvial plains.

high or windspread, stunted scrub. The conditions in Jamaica appear to be more unfavourable than those indicated by Beard for his type and most of the species show partial defoliation in the extreme dry period. While he recognized a separate climax in cactus scrub, this is regarded here as a littoral faciation.

Thorn scrub has been much modified by drainage operations, burning and shifting cultivation. At the present time, it is more characteristic of the coastal areas where it covers the infertile soils of mixed alluvium and saline clays. The climate is severe: high constant temperatures, and continuous sea breezes reduce humidity while rainfall is low and unevenly distributed so that, for 6 to 10 months of the year, it is never over 4 in. and often below 1 in.

Brockmann-Jerosch (1925) regarded the thorn scrub of Jamaica as an insular "facies" of the caatinga of tropical America and Brazil, while Harshberger (1911) erroneously likened it to chaparral. There is no doubt that much confusion has arisen because of its proximity to dry and arid limestone types with which it often mingles to form an ecotone with exchange of species, although a few are common to both. Floristic Composition.

Trees: *Prosopis juliflora*, *Acacia tortuosa*, *A. lutea*, *A. farnesiana*, *A. villosa*, *Haematoxylon campechianum*, *Caesalpinia vesicaria*, *C. coriaria*, and *Pithecellobium unguis-cati*. Less frequently: *Pisidia pisicupula*, *Guaiacum officinale* and the introduced *Morinda oleifera*, *Tamarindus indica*, *Albizia lebbeck* and *Leucaena glauca* are in association.

Shrubs are *Capparis ferruginea*, which is often abundant, *Cassia emarginata*, *Brya ebenus*, *Croton linearis*, *Capparis lanceolata* and the spiny ground bromeliad, *Bromelia pinguin*. Masses of xerophytic epiphytes include the bromeliads *Tillandsia recurvata* and *T. balbisiana* and the sinuous climbing cacti *Cereus triangularis* and *C. flagelliformis*.

The coastal faciation is often a low, sparse scrub, dominated by *Prosopis* with *Acacia tortuosa* and *Pithecellobium* and the characteristic cacti *Cereus peruvianus*, *Opuntia tuna*, *O. spinosissima*, and *Melo-*



FIG. 25. Cactus-Thorn Scrub on the fringe of a salina. The cacti are *Cereus peruvianus* and *Opuntia spinosissima*. The thorns are *Acacia tortuosa* and *Prosopis juliflora*. The ground layer is a pure cover of *Batis maritima*.

cactus communis. A ground cover of the halophytic herbs, *Batis maritima* and *Alternanthera ficoidea*, may be present (Fig. 25).

A cactus-thorn scrub, in which desert-like cacti, *Acacia tortuosa*, *Prosopis juliflora* and many species from the thorn woodland of the plains are blended together is found on sand spits of sufficient width and elevation. A good example is to be seen on the Palisadoes. Chapman (1944) regarded the acacia scrub as secondary in seral progression to a cactus scrub. However, although cacti, especially *Cereus peruvianus*, may become locally dominant, the climax is best regarded as an acacia thorn-scrub which includes cacti. The whole question of the relationship of cacti to this community requires further study.

To summarize, then, it may be said that the dry alluvial plains once supported several types of seasonal evergreen or deciduous forests. Man's activities have now produced successional communities which may lead to a disclimax of thorn thicket or thorn savanna dominated by *Acacia lutea*. The areas of mixed alluvium and saline clays have a characteristic thorn-scrub vegetation dominated by *Prosopis juliflora*. This community has a coastal faciation in which cacti predominate.

SWAMPS AND MARSHES

Beard (1944a) regards swamp and marsh formations as edaphic climax and defines marshes as seasonal swamps because of seasonal fluctuation in water level. His classification is a sound one and, applied to Jamaica, shows the following types:

- Herbaceous swamp.
- Mangrove woodland.
- Palm-Sedge.
- Marsh forest.

HERBACEOUS SWAMPS

The freshwater swamps of Jamaica are limited to a few small examples. They usually have an appreciable salt content. The upper reaches of the

short Ferry River, west of Kingston, may be taken as typical herbaceous freshwater swamp. The water here has 250 p.p.m. of salt and a pH of 8.0, and the vegetation is similar to such swamps occurring in Cuba and Porto Rico.

Floating and submerged plants are: *Pistia stratiotes*, *Hydrocotyle umbellata*, *Ceratophyllum demersum*, *Riccia fluitans*, *Potamogeton*, *Myriophyllum*, *Utricularia stellaris*, *Nymphaea alba*, *Eichornia crassipes*, *Azolla caroliniana*, *Lemna minor* and *Sagittaria*.

Reeds and sedges include: *Typha angustifolia*, *Phragmites communis*, *Arundo donax*, *Echinodorus* sp., *Eleocharis* sp., *Mariscus jamaicensis*, *Cyperus giganteus* and *Cyperus* spp.

The more saline examples, which are the nearest tropical equivalent to temperate zone salt marshes, and bear some successional relationship to mangrove woodland are characterized by scattered mangroves, usually Rhizophora, *Acrostichum aureum* (the golden swamp fern), various tall reeds and sedges and a ground cover of fringing halophytes.

MANGROVE WOODLAND

This formation has a stable and uniform physiognomy throughout the Caribbean. The four New World species, *Rhizophora mangle*, *Laguncularia racemosa*, *Avicennia nitida* and *Conocarpus erecta* are all represented. The red mangrove, Rhizophora, is pre-eminently a seaward pioneer and possesses strut roots and the viviparous habit. Laguncularia, the white mangrove, and Avicennia, the black mangrove, have pneumatophores and are characteristic of the mud swamps. The button mangrove, Conocarpus, is found on dry land. It does not possess strut roots or pneumatophores, neither does it exhibit vivipary and is regarded by some as not being a true mangrove.

The authors agree with Chapman (1939, 1944) that the habitats for mangrove woodland in Jamaica are silt, sand, peat and coral reefs.

Silt Mangroves.—Silt provides the most important habitat, and extensive areas of this kind are found associated with the protected river estuaries and the coastal margin of the alluvial plains. Mangroves play an important part in land elevation and seaward extension of the coast in these areas. The type of mud that occurs is variable and may be of fine, calcareous clay or have a coarser, sandy component. The organic content may be high in both cases, due to local conditions favouring peat formation. The marine "grasses" *Thalassia testudinum* (turtle grass) and *Cymodocea manatorum* (manatee grass) assist in the formation of marine deposits of mud and are of importance in preparing a bed for the mangrove seedlings.

Sand Mangroves.—A sandy habitat is found on many of the spits and cays as well as the sand shoals known as the Bogue Islands in Montego Bay.

Peat Mangroves.—Mangroves growing over deposits of peat were described by Chapman on the Palisadoes, the Fort Augusta sand-spit and in the Portland Bight. Davis (1940) found peat deposits in the mangrove swamps of Florida and, upon examination,

the peat was seen to consist of the remains of Rhizophora and Avicennia. He also noted that along the outer edges of the swamps, Thalassia and Cymodocea, together with marine algae, play a part in the formation of peat soils. Recent borings in Kingston harbour near the Palisadoes have shown that there are extensive deposits of marine peat. This was examined by the authors and there is little doubt that it has been formed from the marine aquatics, especially Thalassia which is frequently uprooted and deposited by marine currents in large banks in the sheltered bays around the coast.

Reef Mangroves.—This habitat includes the shallow reefs such as those occurring on the north coast and the boulders forming the shore of the rocky Cays. The vegetation is limited to scattered plants of Rhizophora and occasional Avicennia. The trees are held by their roots which penetrate into the holes and crevices of the coral, and between the boulders. Chapman regards them as adventives and the whole community as being unstable. Zonation within the mangrove community is, on the whole, poorly defined in Jamaica and this may be partly due to the limited tidal range which, at Kingston, fluctuates between 8-10 in. and is of short duration. Rhizophora requires shallow water for the establishment of its seedlings but, later, when strut roots are developed, the plants are able to extend out into several feet of water. Accordingly, this plant occupies the seaward fringe of the coastal mangroves and the deeper waters of the lagoons, salt ponds and inland swamps. Avicennia occupies the major part of the swamps behind the Rhizophora zone. It occurs in those areas periodically inundated by the tide as well as in the stagnant, saline and shallow water-flats rarely or never under tidal influence. Laguncularia is associated with Avicennia on the landward margin of the swamp where it may occasionally form a pure zone. Conocarpus, although invariably present on dry land fringing the mangrove swamp, is a member of other coastal communities as well. Davis regards Conocarpus as belonging to a transitional community or ecotone between the mangrove swamp and the strand communities. Chapman, however, from his observations in Jamaica, prefers to regard the Conocarpus associates as a seral community. Diagrams illustrating the principal successional stages in the mangrove swamps near Kingston are given by Chapman (1944).

Zonation is variable and its definition depends greatly upon the degree of protection, water level and salinity. The necessary edaphic conditions bringing about zonation are, however, effected by land elevation which will result in zonal advance of the mangrove woodland towards the open water. Finally, the mangroves will be superceded by an inland climax, usually regarded as climatic. In Jamaica, the inland climax which follows mangrove woodland is frequently a cactus thorn scrub, although salt flat communities (salinas) and brackish herbaceous swamps may also play a part. Thus, although there is a recent tendency to regard mangroves as a climax formation—for



FIG. 30. Bamboo brake on the shale hills.

Common among the introduced and cultivated trees are: *Mangifera indica*, *Meliococca bijuga*, *Artocarpus integrifolia*, *A. incisa*, *Cola acuminata*, *Theobroma cacao*, *Cocos nucifera*, *Gliricidia sepium*, *Bambusa vulgaris*, *Tamarindus indica*, *Scamanea saman*, *Eugenia jambos*, *Inga vera*, *Anona reticulata*, *Albizia lebbeck*, *Eugenia malaccensis*; the trailing "Spanish moss" *Tillandsia usneoides* is a characteristic epiphyte.

The remnant forests of the lower stream valleys include the following trees:

- Ceiba pentandra*
- Cecropia peltata*
- Cedrela odorata*
- Pithecellobium arboreum*
- Chlorophora tinctoria*
- Zanthoxylum flavum*
- Ochroma pyramidale*
- Chrysophyllum ovaliforme*
- Catalpa longissima*
- Ficus suffocans*
- F. wilsonii*
- Prunus myrtifolia*

On the slopes, the sparser forest of smaller trees contains:

- Andira inermis*
- Hura crepitans*
- Bursera simaruba*
- Guazuma ulmifolia*
- Spondias monbin*
- S. purpurea*
- Comocladia pinnatifolia*
- Spathelia sorbifolia*
- Bauhinia divaricata*
- Crescentia cujete*
- Amyris balsamifera*
- Nectandra sanguinea*
- N. antillana*
- Cordia gerascanthoides*
- Coccoloba laurifolia*
- Celtis trinervia*
- Trema micrantha*
- Byrsonima coriacea*
- Pimenta officinalis*
- Guarea glabra*
- Clusia rosea*
- Drypetes lateriflora*
- Picramnia antidesma*

- Heteropteris laurifolia*
- Matayba apetala*
- Brya ebenus*
- Senecio discolor*

The ground flora includes:

- Bletia purpurea*
- Bidens cynapiifolia*
- B. pilosa*
- Asclepias nivea*
- Vernonia divaricata*
- Iresine paniculata*
- Piper umbellatum*
- P. amalago*
- P. aduncum*
- Peperomia verticillata*
- Pilea microphylla*

and the ferns:

- Asplenium pumilum*
- Aneimia underwoodiana*
- A. hirsuta*
- Dryopteris pedata*
- D. resinifera*
- Blechnum unilaterale*
- Gymnopteris rufa*
- Lygodium volubile*
- Notholaena trichomanoides*
- Trismeria trifoliata*

The open banks have quite a rich and characteristic moss flora with such species as:

- Breutelia jamaicensis* (Mitt) Jaeg.
- Pleuropus bonplandii* (Hook) Broth.
- Brachythecium steropoma* (Spruce) Jaeg.
- Papillaria nigrescens* (Hedw) Jaeg.
- Anoectangium euchloron* Schwaegr.
- Philonotis glaucescens* (Hornsch) Par.
- Bryum crugeri* Hpe.
- B. argenteum* Hedw.
- B. truncorum* Brid.
- Calliergonella cuspidata* (Hedw) Loeske
- Campylopus introflexus* (Hedw) Brid.
- Haplocladium microphyllum* (Hedw) Jaeg.
- Meteoriopsis remotifolia* (Hornsch) Broth.

Others found in the moist forest remnants are:

- Helicophyllum tortuatum* (Hook) Brid.
- Entodon macropodus* (Hedw) C. M.
- Fissidens similiretis* Sull.
- Heliodontium capillara* (Hedw) Jaeg.
- Samatophyllum galipense* C. M.
- Squamidium nigricans* (Hook) Broth.

MONTANE COMMUNITIES

LOWER MONTANE RAIN FOREST

A typical lowland tropical rain forest cannot be recognized in Jamaica to-day. The closest approach is seen in the type of rain forest occupying a narrow zone of the lower northern slopes of the Blue Mountain range and the western slopes of the John Crow Mountains. Annual rainfall in the former regions is well over 100 in. while 275 in. has been recorded at Millbank in the vicinity of the latter. There is no marked dry season. On the Blue Mountain range, much of this original forest has been cleared and that which remains occupies the more inaccessible slopes and gradually merges into Mist Forest at altitudes above 3,500 ft. In the east, Corn Puss Gap at 2,000

ft., and Cuna Cuna Gap at 2,500 ft., represent the lowest points at which this type of forest has been observed on the Blue Mountain range. The western slopes of the John Crow Mountains support rain forest from 1,500 ft.-2,500 ft. Above this, it thins out to a mixed palm brake and mist forest, and finally and abruptly to elfin woodland. The John Crow Mountains form the eastern boundary of the low-lying, humid Rio Grande valley. It is in this valley that the remnant vegetation of tree ferns, Heliconias and aroids clearly indicate a locality where lowland tropical rain forest may have flourished in the past and spread up to the type now to be described.

We have called this community Lower Montane Rain Forest since it resembles closely the type so named by Beard for the lesser Antilles and Trinidad. It may be thought of as a variant or facies of lowland tropical rain forest from which it appears to differ in the following respects. The main canopy is lower, being here at 80 ft. There is a less well-defined stratification, few trees with buttresses, and lianes and epiphytes, although present, are not abundant.

The stands of lower montane rain forest which have been examined are all on sloping ground and




FIG. 31. A profile diagram of Lower Montane Rain Forest measured in the western slopes of the John Crow Mountains at 1,800 ft. The diagram represents the trees present in a belt 100 ft. long and 25 ft. wide.

KEY TO SYMBOLS: Al. *Alchornea latifolia* (Jumba); Cj. *Calophyllum jacquinii* (Santa Maria); Cs. *Calyptrochoma swartzii*; E. *Eugenia* sp. (Rodwood). Ep. *Exothea paniculata* (Wild Ginep); F. *Ficus* sp.; Fo. *Faramea occidentalis* (Wild coffee); Gg. *Guarea glabra* (Alligator, Wild akee); Ma. *Matayba anetala* (Wannika); N. *Nectandra* sp. (Sweetwood); Pa. *Pithecellobium alegandrii* (Shadbank); Pm. *Psidium montanum* (Mountain guava); R. *Rubiaceous* sp.; Sg. *Sympomia globulifera* (Hog gum); Tr. *Trophis racemosa* (Ramoon).

the substratum is limestone in the case of the John Crow Mountains and shale on the Blue Mountains. Structure.

A profile diagram of a typical area at 1,800 ft., on the western slopes of the John Crow Mountains is given in Fig. 31. Here is a mixed forest with occasional huge emergent trees of *Psidium montanum* reaching up to 120 ft. and having a girth of 9-11 ft. The main canopy, which is not dense, is at 60-80 ft., and the girth varies from 4-6 ft. A somewhat ill-defined lower layer occurs from 30-50 ft., and may include palms. Tree ferns are rare.

A profile diagram (Fig. 32) of the forest taken near Cuna Cuna Gap at 2,500 ft. also shows a closed canopy from 60-80 ft. but it consists almost exclusively of *Calophyllum jacquinii* (Santa Maria). There is an ill-defined sub-canopy from 40-55 ft., and a lower layer from 10-30 ft. The relatively steep slope here does not aid the clear demarcation of the lower strata.

It is from such areas as these that the following preliminary description of this type of forest oc-




FIG. 32. Profile diagram of Lower Montane Rain Forest measured at Cuna-Cuna Gap in the Blue Mountain Range. The diagram represents the trees present in a strip 100 ft. long and 25 ft. wide.

KEY TO SYMBOLS: Al. *Alchornea latifolia* (Jumba); Cj. *Calophyllum jacquinii* (Santa Maria); Co. *Clethra occidentalis*; Cy. *Cyathea* sp.; D. *Dipholis* sp. (Bullet wood); E. *Eugenia* sp. (Rodwood); Ep. *Exothea paniculata* (Wild Ginep); Gg. *Guarea glabra* (Alligator, Wild akee); He. *Hernandia* sp. (Womanwood); M. *Melastome* sp. (Soapwood); Ma. *Matayba apetala* (Wannika); Me. *Mimusops excisa* (Wild naseberry); 'Mw' Unidentified ('Milkwood'); N. *Nectandra* sp. (Sweetwood); P. *Psychotria* sp.; Sj. *Sapium jamaicense* (Blindeye); Uk. Unidentified.

example, Beard (1944a) classified Mangrove Woodland as an edaphic climax within his Swamp Formation series—the seral nature of the component zones should not be forgotten.

There are few associates of the mangroves, *Batis maritima*, *Salicornia ambigua*, *Acrostichum aureum* and *Alternanthera ficoidea* are often present as undergrowth in the open woodland. On sand, *Sporobolus virginicus* may occur.

Salinas, or salt flats, frequently occur as a landward fringe of the mangrove woodland on the south coast. They are to be found where the substratum is a heavy saline clay. Such areas are sufficiently low-lying to be inundated by the sea during storms or exceptionally high tides. On the other hand, they are sufficiently elevated to prevent normal tidal inundation. The clay soil tends to retain the surface water but the evaporation rate is so high that there are long periods of the year when the surface is dry. However, due to evaporation, salinity is high. In some cases, salt crystals can be seen on the surface. Such areas are bare of vegetation but, more frequently, the salinity, while high, is not sufficient to prevent the extreme halophyte *Batis maritima* from being present. This plant frequently forms a pure cover. Occasional associates are *Sesuvium putulacastrum* and *Salicornia ambigua*. Where salinity falls, *Heliotropium curassavicum* and *Portulaca oleracea* may be present. The landward side of the larger salinas gradually pass over to cactus thorn scrub (Fig. 26).

The processes concerned with the formation of salinas need further attention. They may be formed from salt ponds by evaporation, as outlined on page 374 and then will never have been occupied by mangroves. On the other hand, they may be produced as a result of drying out following artificial drainage, change in the course of rivers or elevation of the land in relation to sea level. In such areas, the salinas may have formerly supported mangroves. Chapman noted that the salinas in the Hunts Bay region contained old stumps of *Avicennia* and suggested that alterations in the course of the Rio Cobre, which are known to have occurred, have resulted in drying out and subsequent death of the *Avicennia* community, formerly occupying the present salinas.




FIG. 26. A large salina at Port Henderson covered by *Batis maritima*. Occasional stunted bushes of *Laguncularia* are present. Thorn scrub and cactus-thorn scrub are seen in the background.

PALM-SEDGE MARSH

The largest swamp area of Jamaica is really a sedge marsh with scattered palms. This covers almost the whole of the low-lying flood plains of the Black River, the delta being some 100 sq. mi. in extent. During the rainy season, it is inundated with freshwater from the swollen rivers, whilst, in the dry season, much of the higher ground dries out to marsh.

The many tributaries of the Black River find their confluence a mile or so above the mouth, this part being known as Broadwater. Above this there are two main branches, the lesser being of dark, sluggish water and open to boats for some five miles. The main river branch is navigable for some 30 mi., flows more swiftly and is often muddy yellow with silt. The river is tidal and brackish for about five miles from the sea. The vegetation shows two aspects: a marginal zone along the river banks and the marsh area inland. The west bank of Broadwater has a dense border of mangroves, chiefly *Rhizophora mangle*, up to 30-40 ft., in height; *Laguncularia* and *Avicennia* are occasional. *Rhizophora* may continue for 6 mi. up the river, although it is reduced in height and becomes less and less frequent. The mangroves are correlated closely with the current and formation of silt banks, being strictly confined to the inner bends of the river. Where the river takes a sharp turn and silt is deposited on the opposite bank, the mangrove belt also changes sides. Opposite the mangroves, there is a zone of *Typha angustifolia* and *Phragmites communis*. In the absence of mangroves it is usually *Typha* which is dominant and, here, *Typha* has the same distributional pattern in relation to silt deposition as had *Rhizophora*. This marginal fringe has also the following species: *Acrostichum aureum*, *Jussiaea suffruticosa*; and the climbers: *Ipomea fastigiata*, *Ipomea carnea*, *Aristolochia trilobata* and *Mikania micrantha*.

Close to the banks, the floating or submerged flora includes: *Nymphaea*, *Sagittaria*, *Eichornia* and others of the freshwater swamps.

By far the greater portion of the Black River area is a sedge marsh which is spread out between the many branches of the river. This is dominated by an almost pure cover of the large *Cyperus giganteus* although other *Cyperus* and *Carex* species are present. Scattered throughout this area, are groups of the royal palm, *Roystonea princeps*, up to 35 ft., and endemic to this area of Jamaica. Nearer the sea, this is joined by *Calyptrotrona swartzii*.

MARSH FOREST

Beard (1946) described marsh forest on alluvial terraces in Trinidad. The same association is known from British Guiana but does not occur in the Lesser Antilles. Marsh forest (Fig. 27) in Jamaica is limited to a few square miles of the upper reaches of the Black River and may also occur further along the coast near Cave. Typically developed on the higher ground where two tributaries meet, the forest is subject to periodic inundation. The floor remains wet and muddy after rain.



lina.
sissi-
juli-
nari-

tytie
deea,

acti,
eies
ided
idth
the
acia
etus
reus
max
in-
ship

dry
son-
ties
nich
orn
of
stic
uli-
in

na-
ea-
ter
ied

to
ip-
the



FIG. 27. Marsh Forest bordering the Black River. The canopy tree is *Sympetrum globulifera* with a lower storey of *Grias cauliflora* and the palms, *Roystonea princeps* and *Calyptroprona swartzii*.

The canopy is closed at about 30 ft., and made up of palms (*Roystonea princeps*, *Calyptroprona swartzii*) which are also members of the palm marsh. The dominant tree is hog gum (*Sympetrum globulifera*) with aerial roots and an exudate of yellow gum when cut. *Sympetrum* frequently forms the emergent layer. *Grias cauliflora* is a very characteristic small tree with large pendant leaves. Other trees, many of which have invaded from neighbouring communities, are: *Calophyllum jacquinii*, *Terminalia latifolia*, *Piscidia piscipula*, *Ficus* spp., *Haematoxylon campechianum* and *Spondias monbin*. Buttress roots are not characteristic.

Epiphytes are aroids and orchids but bromeliads are rare. *Colocasia* and ferns form a ground layer.

SHALE

VEGETATION ON THE LOWER SHALE HILLS

Some of the most cultivated aspects of Jamaica are the "shale" soils of the lower hills and valleys. There are three shale regions, the Blue Mountains in the east which proceed to high elevations, and the central and western regions, both of which are below 2,500 ft., Bull Head (2,782 ft.) in the central region being an exception.

Geologically, two major series are included: the Carbonaceous shales of the Lower Eocene (black shales, coarse conglomerates, limestone and sandstone) and the Cretaceous shales. These latter are red and purple shales, tuffs, breccias and conglomerates interbedded with fossiliferous limestone of the Upper Cretaceous. These rocks produce a complex of local soil types that is included under the term "shale".

Most of the "shale" soils are very porous but, in Jamaica, the consequent free drainage is offset by the high rainfall over the regions where they occur. The profile is often degraded or truncated, much of the area being of "forest soils" which have been badly eroded following deforestation and poor agricultural practices.

The lower shales were originally covered with a sparse, mesophytic forest but the exact nature of this forest is now a matter of conjecture from the



FIG. 28. Remnant mesophytic forest of the lower shale valleys of St. Andrew. Dominant trees are *Ceiba*, *Cecropia*, *Cedrela*, *Pithecellobium* and the introduced *Samanea saman*. Note the steep slopes in the distance from which the forest has been removed and which now support grassland with scattered shrubs and trees.

remnants still to be found in the lower valleys following the stream beds. On the St. Andrew slopes of the Blue Mountain system (Fig. 28) there is a rather sparse forest of large spreading trees and tall spindly shrubs forming a thin canopy broken by emergent cotton trees. This extends up the ravines but, elsewhere, soon tails off into open grassland with a few low hardy shrubs and scattered trees. Much of the present vegetation of the open slope is due to man's interference. The upper soil layers have been rapidly lost after deforestation, continued burning and shifting patchwork cultivation. This has been greatly accelerated by surface erosion due to the steep topography. In St. Andrew, and best seen on the slopes of the Port Royal ranges, this has led to a characteristic hill savanna of guinea grass and mango trees where once there was continuous cover (Fig. 29). *Bambusa vulgaris* with large feathery branches is common in some areas (Fig. 30).



FIG. 29. A view of the lower shale hills in the Yallahs river valley where erosion is heavy. The vegetation consists of induced hill savanna and scattered mango trees.

curring in Jamaica is built up. Apart from occasional and widely spaced emergents such as *Psidium montanum*, *Sympodia globulifera* and *Ficus suffocans* reaching up to 120 ft., the main canopy extends from 60-80 ft. The lowermost branching of this layer is at 60 ft., and a wide-spreading flat-topped crown is developed. Spacing of the trees may be close but variation occurs due to soil depth and drainage.

The canopy is often dense where there is a single dominant such as Santa Maria, but more open in the mixed forest. The lower tree layer is similarly variable in density between 30-50 ft., and with ill-defined stratification.

The shrub layer from 10-20 ft. is usually sparse, due to the poor light, and is represented by spindly shrubs of the Rubiaceae and Melastomaceae, together with *Eugenia*, *Piper* and *Helicoria* species.

The field layer is predominantly of ferns, tree seedlings and bryophytes.

Among the trees a few, such as *Psidium* and *Ficus*, show buttressing while stilt roots are seen in *Sympodia* which, together with *Ficus* and *Sapium*, has latex. The tree fern, *Cyathea*, is the only thorny member. Tree ferns, and more especially palms, characterize the lower strata of the forest.

Lianes are not abundant but there are many small vines, including *Smilax*, several aroids, *Marcgravia* and other climbing herbs and ferns.

Epiphytes are represented by bromeliads which are here limited to the higher tree branches at 60-70 ft.; ferns, bryophytes and orchids occur at the shrub level.

All the tree species are evergreen with simple leaves, except in the case of Matayba, *Pithecellobium* and Guarea. Leaf size falls into Raunkiaer's mesophyll group.

The forest floor is of moist soil, rich in humus and organic litter, usually with a cover of ground ferns and Santa Maria (*Calophyllum*) seedlings. Bryophyte societies occur on small rocks and fallen branches with *Thuidium* and *Leucobryum* as the dominants.

Floristically, there is some relation with the wet limestone forest.

Emergent trees:

Sympodia globulifera
Psidium montanum
Ficus suffocans

Trees of the main canopy:

Calophyllum jacquinii
Matayba apetala
Nectandra antillana
Clethra occidentalis
Exothea paniculata
Dipholis sp.
Cyrilla racemiflora
Prunus occidentalis

Sub-canopy trees:

Guarea glabra
Pithecellobium arboreum
Sapium jamaicense

Nectandra sanguinea
N. coriacea
Eugenia disticha
E. monticola
Hufelandia pendula
Oxandra laurifolia
Ocotea martinicensis
Hernandia catalpifolia
Alchornea latifolia
Dipholis spp.
Cecropia peltata
Calyptronoma swartzii

Shrub layer: Melastomaceae:

Miconia laevigata
Ossaea sperifolia
Conostegia superba
Blakea trinervia

Rubiaceae:

Psychotria pedunculata
P. nervosa
Faramea odoratissima

Also:

Gilibertia ovalifolia
G. arborea
Rheedia pendula
Hedyosmum arborescens
Columnea hirsuta
Lacistema aggregatum
Rytidophyllum grande
Piper nigrinodum
P. discolor
P. scabrum
Helicoria caribaea adds a distinctive note.

Tree ferns, which are not frequent in this forest, are:

Cyathea tussacii
C. grevilleana
Ctenicis villosa
Alsophila swartziana
A. aspera

The field layer:

Besleria lutea
Pachystachys coccinea
Lobelia acuminata
Gryotaenia microcarpa
Peperomia hispida
P. crassicaulis
Scybalium jamaicensis, the root parasite.

Ground ferns:

Asplenium cuneatum
Diplazium faydenii
D. pectinatum
Camptodium pedatum
Bolbitis pergamantacea
B. aliena
B. nicotianaeifolia
Dryopteris gemmipara
D. deltoidea
D. decussata
Thelypteris serrulata
Pteris hexagona
Tectaria incisa

Polybotrya cervina
Nephrolepis rivularis
Danaea jenmanii
D. nodosa

Ground mosses:

Thuidium acuminatum Mitt.
Leucobryum antillarum Schp.
Epiphygium wrightii (Sull.) Lindb.
Leucoloma serrulatum Brid.
Fissidens elegans Brid.
F. mollis Mitt.
F. polypodioides Hedw.
Lepidopilum polytrichoides (Hedw.) Brid.

Epiphytes and climbers:

Begonia glabra
Philodendron tripartitum
P. lacerum
P. cordatum
Smilax domingensis
Marcgravia brownei
Schradera involucrata
Elleanthus capitatus
Dichaea graminea
Epidendrum fragrans

Epiphytic and climbing ferns:

Trichomanes scandens
T. crispum
T. hymenophylloides
T. rigidum
T. holopterum
T. osmundoides
Hymenophyllum crinitum
H. hirtellum
H. hirstum
Polypodium mollissimum
P. jubaeforme
P. lycopodioides
P. exornans
Elaphoglossum simplex
E. petiolatum
Polytaenium feei

Epiphytic mosses:

Octoblepharum albidum Hedw.
Lepyrodontopsis trichophylla (Sw.) Broth.
Pireela filicina (Hedw.) Card.
Pilotrichum amazonum Mitt.
Meteriopsis remotifolia (Hornsch.) Broth.

MONTANE SCLEROPHYLL

At about 2,500 ft. on the St. Andrew (leeward) slopes of the Blue Mountain ranges, the vegetation changes quite suddenly and abruptly to a sclerophyll zone which continues up to 4,000 ft. This zone would appear to be unique in the West Indies, although such dry belts at higher altitudes are not unknown elsewhere in the tropics. There is a similar aspect called "ceja" on the Andes and such belts are also said to occur in Borneo and Ceylon. The montane sclerophyll has two aspects. At the lower altitudes it consists of a low open shrubby community under the periodic influence of shifting cultivation. Above

this is a less disturbed thicket of small trees and shrubs which extends up to the Mist Forest.

The lower sclerophyll belt is of low, bushy shrubs, sparse in cover and scant of foliage. Xeromorphic features are characteristic, the most common being small, hard, thick leaves.

Two very characteristic shrubs restricted to this community are the mountain broom, *Baccharis scoparia*, with minute, green leaves and green, twiggy stems and *Dodonea viscosa*, a small hardwood shrub up to 12 ft. *Dodonea viscosa* is a common lowland coastal species found throughout the Pacific. In Jamaica, it is restricted to this inland elevated community.

Other small shrubs present are:

Clusia rosea
Lantana involucrata
Eugenia spp.
Senecio discolor
Vernonia divaricata
Solanum verbascifolium
Lyonia jamaicensis
Heterotrichum umbellatum
Ascyrum hypericoides

Hard, stiff-leaved pteridophytes are typical of the ground layer, notably:

Pteridium arachnoidium
Gleichenia jamaicensis
Polypodium thysanolepis
Lycopodium cernuum

At this level, guinea grass (*Panicum maximum*) gives way to molasses grass (*Melinis minutiflora*), also an introduced species which is spreading fast and becoming characteristic of this region.

A number of small trailing woody herbs are very characteristic and include:

Chusquea abietifolia—the climbing bamboo
Relbunium hypocarpium
Manettia lygistum
Coccocypselum herbaceum
Micromeria obovata

Herbs include:

Crotalaria retusa
Bidens pilosa
Cordia cylindrostachya
Pavonia rosea
Bryophyllum pinnatum
Flemingia strobilifera
Leianthus longifolius
Gnaphalium americanum

The lichens *Usnea* and *Cladonia* and terrestrial mosses are also present.

Moist shaded banks support *Pilea microphylla*, *Selaginella* spp., *Achimenes coccinea* and other hylorrhophilous plants.

The low, bushy sclerophyll merges into a dry, evergreen thicket at the higher altitudes (3,000-4,000 ft.). This does not correspond to the montane thicket described by Beard (1949) for the Lesser Antilles, which would seem to be equivalent to our Mist Forest. The environment of the sclerophyllous thicket recog-

nised on the south slopes of the Blue Mountains is much drier than Mist Forest with which it has, nevertheless, some floristic affinity. The canopy is more open and from 20-30 ft. high. Dominant among the small trees and tall, slender shrubs are:

Cyrilla racemiflora
Vaccinium meridionale
Clethra occidentalis
Eugenia spp.
Viburnum villosum
Persea alpigena
Coccoloba laurifolia
Clusia rosea
Guettarda argentea
Nectandra patens
Bocconia frutescens

Many of the members belong to the Ericales and Myrtales.

The sclerophyll belt as a whole is rather difficult to explain. In the lower aspect, a number of lowland representatives reappear, such as Lantana, Bryophyllum and Solanum. These belong to secondary growth "ruinate" areas of the dry limestone and alluvium, but occur here together with the bona fide sclerophylls such as Baccharis, Pteridium and Gleichenia.

Grevillea robusta and Eucalyptus, both native to the drier regions of Australia do well here. Drought, however, is not evident even allowing for the sharp decrease in rainfall when compared to the north slopes, where, at the corresponding level, remnants of lower montane rain forest occur.

Brockman-Jerosch (1925) suggested that the factor of most importance for the sclerophyll belt was that of lowered humidity. It is true that the trade winds are depleted of moisture by the time they reach the southern slopes and that the sclerophyll belt does not enter the Blue Mountain mist level. Shreve (1914) found a continued high humidity of 95% for the north ravines but, at Cinchona on the south side, a fluctuating value from 50% to 90%. Evaporation rates were four to five times greater than those of the north slopes. The relative transpiration rates of Dodonea, a typical sclerophyll shrub, was here double that of Clethra from the mist forest where high humidity retarded transpiration. His data would indicate that humidity and evaporation-transpiration ratios are the controlling factors for the sclerophyll.

Ecological considerations provide another line of attack to this problem. At first it is difficult to see anything of a seral nature in the sclerophyll belt. However, on the wet Portland slopes at 2,500 ft., there are minor areas which, from their species, are very reminiscent of the sclerophyll. Such small bushes such as Gleichenia, Pteris and Baccharis occur with the trees Clethra, Viburnum, Coccoloba and Nectandra. Here, however, there is ample evidence of past clearing and burning and the vegetation is undoubtedly seral. Again, in the central shale region on Bull Head (1,782 ft.), where there was once a lower montane rain forest of Santa Maria, there is now a type of secondary community which includes:

Cyrilla, Clethra, Eugenia, Clusia, Persea, *Myrica microcarpa*, Lyonia, Baccharis, *Miconia albidans* and *Clidonia strigillosa* together with the ground herbs: Gleichenia, Lantana, and Pteridium with the grasses *Andropogon virginicus* and *Melinis minutiflora*.

These facts indicate that the lower sclerophyll belt is a second growth community which, if left undisturbed, might eventually lead to sclerophyllous thickets or, on the wetter northern slopes, to lower montane rain forest. Such development may very well be a long-term process in regions such as this where leaching and erosion have followed repeated burning and clearing practised for shifting cultivation. These processes not only prevent regeneration of the trees but have completely changed the edaphic conditions. The fertile surface layers of the soil have gone and, on the steeper slopes, the whole topsoil has been washed down the mountainside. Colonization is by light-seeded weeds and shrubs. There are large patches in the sclerophyll belt where an almost pure cover of either *Melinis minutiflora* or *Gleichenia pectinata* prevails as a result of repeated burning.

MONTANE MIST FOREST

One of the few remaining retreats of original vegetation in Jamaica is to be seen in the montane mist forest clothing the upper reaches of the Blue Mountains which form the central mountain system in the eastern end of the island.

The mist forest may be recognised from about 4,500 ft. upwards together with an elfin woodland aspect. Reference to this forest is found in most accounts of visiting botanists. Ørsted (1857) thought it poor and monotonous in character; Boergesen (1923) "the most peculiar vegetation imaginable," and Brockman-Jerosch (1925) wrote of the confusing abundance of species and the rich plant life in a muggy, glasshouse atmosphere. The fullest account is in Shreve's monograph (1914).

This type of forest has been described throughout Central America and the Caribbean islands. There is little doubt that mist, which cuts down the light and increases humidity, is a far more important factor than rainfall. Pittier (1939) has described the close association in the Venezuelan Andes between such forest and the mist belt. Carr (1949) alludes to the tear-like condensation of water on the trees of the cloud forests of Honduras and Miranda & Sharp (1950) to the fog forest in Eastern Mexico. Gleason & Cook (1927), describing the mossy forests of the Luquillo forests of Porto Rico, describe the peaks as wrapped in cloud which "raises humidity, brings about daily rainfall and reduces the sunshine to short intervals."

In the Blue Mountains of Jamaica, the upper ridges are enveloped daily in mist from 10 A.M.-4 P.M., that is for a period of at least 6 hours (Fig. 33). The mist persists even during rain but much of the moisture, enabling the hymenophyllous ferns and hygrophilous plants to flourish, is due to condensation.

The soil is extremely porous and a high proportion



FIG. 33. General view showing the Mist Forest of the Blue Mountains. In the lower foreground is a plantation of Blue Mahoe (*Hibiscus elatus*).

of the rainfall is rapidly drained off, due to the steep slopes. The rainfall is distributed in light showers but actual figures are reduced in significance, due to the low evaporating power of the air.

Shreve concluded that sunlight was cut down to one quarter of the normal, the nights being clear but mist forming 5-15 minutes after sunrise each day. He estimated the daily duration of mist as a percentage of the total daylight hours and gives the following figures for the leeward and windward slopes above 4,500 ft.

	North.	South.
Feb., July, August.....	30%	10%
Other months.....	70%	30%

Working on the physiological aspects, Shreve was able to correlate a daily march of transpiration rates with mist formation.

Variation in the structural and floristic composition of the mist forest is wide and almost entirely correlated with degree of exposure. Shreve describes the forest of the ravines, the ravine slopes and the ridges. The chief variation is between that of the deep, sheltered, ravines and that of the exposed ridges. The main structural difference is in the canopy which, with progress towards the exposed ridge, becomes lower, open and more uniform, whilst the species in shrub and ground layers change to the more xerophytic ones. Shreve showed that, on the exposed ridges, air movement lowered the humidity greatly.

STRUCTURE—The mist forest is low-canopied, wet and misty. The evergreen trees with sombre dark-green, broad leaves are associated with an abundance of small, melastomaceous and rubiaceous undershrubs, pteridophytes and bryophytes (Fig. 34). In both structure and floristics, it shows many temperate forest features.

The profile diagram (Fig. 35) is for mist forest at 4,200 ft. near Catherine's Peak on a range subsidiary to the Blue Mountains.



FIG. 34. An open aspect of Montane Mist Forest at St. Catherine's Peak. The canopy trees are Cyrilla, Alchornea and Podocarpus. Tree ferns, shrubs and epiphytes are conspicuous.




FIG. 35. Profile diagram of Mist Forest measured at 4,500 ft. in the Blue Mountain Range. The diagram represents the trees present in a belt 200 ft. long and 25 ft. wide.

KEY TO SYMBOLS: Al. *Alchornea latifolia* (Jumba); Cl. *Clausia* sp. (Wild Fig); Co. *Clethra occidentalis*; Cr. *Cyrilla racemiflora* (Bloodwood, Beetwood); Cy. *Cyathea* sp.; E. *Eugenia* sp. (Rodwood); Ha. *Hedysymum arborescens* Sw.; M. *Melastomaceous* sp. (Soapwood); N. *Nectandra* sp. (Sweetwood); Pu. *Podocarpus urbani* (Yacca); So. *Solanaceous* sp.; Uk. Unidentified.

Emergent trees are few, the canopy at 40-50 ft. being uneven and somewhat open. A sub-canopy of varying density can be recognised at 30 ft. The shrub layer at 10 ft. is far more scattered while the field layer is a close and often luxuriant cover of herbs, ferns and bryophytes.

The dominant trees of the canopy, Podocarpus, Cyrilla and Alchornea are spaced at about 10 ft. and average 2 ft. 6 in. in girth. Most tend to be tall and spindly, with bushy crown branching. Cyrilla is frequently inclined in the direction of the slope. Shreve was of the opinion that this leaning indicated erosion movement of the shallow topsoil. It appears, however, that there are other factors such as wind exposure, growth towards light where trees have been felled by hurricanes etc. Similar features can be seen in Beard's profiles for montane rain forest in Trinidad.

Tree ferns are frequent, adding character, but palms are absent from this community in Jamaica. Compound leaves are seen in Brunellia, Wienmannia and Guarea, but generally the leaves are simple, often coriaceous, evergreen and small. Podocarpus and Cyrilla have flaky bark, but buttress roots and caulinflory (seen to a limited extent in Solanum and Mecranium) are generally absent. Clusia has aerial roots and begins life as an epiphyte. Lianes are rare but are represented by climbers up to 25 ft. such as Smilax, Maregravia and Schradera. Epiphytes do not extend much above 15 ft. and include shrubs, bromeliads, orchids, ferns and bryophytes. Many are to be found again as members of the ground flora on fallen debris.

FLORISTIC COMPOSITION—Floristically, the montane mist forest is simple, the same species occurring in the different aspects with great uniformity.

The dominants are *Podocarpus urbani* and *Cyrilla racemiflora* and this leads logically to naming this montane community the Podocarpus-Cyrilla association.

Associated with the canopy layer are:

Podocarpus urbani
Cyrilla racemiflora
Alchornea latifolia
Clethra occidentalis
Brunellia comocladifolia
Solanum punctulatum
Laplacea haematoxylon
Ficus harrisi
Guarea glabra

The sub-canopy of under trees includes:

Clusia havetioides
Eugenia biflora
E. fragrans
Podocarpus urbani
Clethra occidentalis
Cyrilla racemiflora
Vaccinium meridionale
Gilibertia arborea
G. nutans
Acnistus arborescens
Weinmannia pinnata
Sciadophyllum brownii

The following less frequently occurring trees, are characteristic of differing aspects:

Juniperus barbadensis
Psidium montanum
Myrica microcarpa
Garrya fadyenii
Citharexylum caudatum
Nectandra patens
Dipholis montana
Rapanea ferruginea
Viburnum villosum
Turpinia occidentalis
Lyonia jamaicensis
Rhamnus sphaerosperma
Eugenia marchiana
Eroteum theoides

The shrub layer includes many species. Not all

of these will be found in any one area as this layer shows the greatest variation with aspect. The species are:

Hedyosmum arborescens
Psychotria corymbosa
P. brachiata
P. brownei
Cephaelis elata
Palicourea crocea
Meriania purpurea
M. leucantha
Heterotrichum umbellatum
Miconia rubens
M. quadrangularis
M. dodecandra
Mecranium purpureescens
M. virgatum
M. amygdalinum
Tournefortia cymosa
Ilex montana
Myrica microcarpa
Columnnea hirsuta
Ocotea jamaicensis
Hufelandia pendula
Acalypha virgata
Cestrum hirtum
Datura suaveolens
Bocconia frutescens
Bochmeria caudata
Piper hispidum
P. otophyllum
P. arboreum

The field layer includes the following herbs: *Begonia acutifolia*, many Peperomias and Pileas, *Lobelia assurgens*, *Besleria lutea*, *Panicum palmifolium*, *Uncinia hamata*, and *Rhynchospora eggersiana*. Many introduced, temperate species found on the open paths belong to such genera as: *Plantago*, *Rumex*, *Ranunculus* and *Fragaria*.

Ferns are too numerous to mention in detail. Among the ground ferns are:

Danea jamaicensis
Gleichenia pectinata
Diplazium centripetale
D. brunneo-viride
D. costale
D. heteroclita
Cryopteris denticulata
Dennstaedtia globulifera
Elaphoglossum eggersii
Asplenium pramorum
A. harpeodes
A. alatum
A. radicans
Blechnum polypodioides
B. lineatum
B. jamaicense
Nephrolepis pectinata
Cystopteris fragilis

Tree ferns:

Lophosoria quadrripinnata
Marrattia alata

- Orthepteris domingensis*
Cyathea nigrescens
C. tussacii
C. pubescens
C. armata

Epiphytic ferns: *Polypodium loricatum* is common in mist forest. *Grammitis*, with 22 species occurring as epiphytes on mossy tree trunks and branches, is a characteristic genus of the montane formations. *Grammitis serrulata* is very frequent.

The filmy ferns, *Hymenophyllum sericeum*, *H. polyanthos* and *H. lanatum* are frequent to abundant, as well as many *Trichomanes* spp.

Climbers and scramblers, some of which are epiphytic shrubs:

- Passiflora penduliflora*
Maurandia erubescens
Marcgravia brownei
Cassia viminea
Schradera involucrata
Blakea trinervia
Solandra grandiflora
Metastelma fawcettii
M. atrorubens
Begonia glabra
Manettia lygistrum
Cionosicys pomiformis
Smilax celastroides

True epiphytes are the bromeliads, *Thecophyllum sintensii*, *Tillandsia incurva*, *Hohenbergia* and *Guzmania* spp., and the orchids: *Dichaea graminea*, *Dichaea glauca*, *Lepanthes concinna*, *L. tridentata*, *L. concolor*, *Epidendrum verrucosum*, *Liparis elata* and *Stelis ophioglossoides*.

Mosses, terrestrial and on logs:

- Campylium chrysophyllum* (Brid.) Bryhn.
Pleurozium bonplandii (Hook.) Broth.
Syrrhopodon lycopodioides (Sw.) CM.
S. gaudichaudii Mont.
Pilopogon gracilis (Hook.) Brid.
Ditrichum rufescens (Hpe.) Broth.
Aongstroemia jamaicensis CM.
Campylopus harrisi (CM.) Par.
C. aretocarpus (Hornschr.) Mitt.
Entosthodon bonplandii (Brid.) Mitt.
Hookeriopsis undata (Hedw.) Jaeg.
Hypopterygium tamariscinum (Hedw.) Brid.
Leucobryum antillarum Schp.
L. giganteum CM.
Polytrichum juniperinum Hedw.
Pogonatum robustum Mitt.
Atrichum synoicum (CM.) Par.
Rhizogonium spiniforme (Hedw.) Bruch.
Sphagnum meridense (Hpe.) CM.

On tree trunks or pendant from branches:

- Meteoriopsis patula* (Hedw.) Broth.
Phyllogonium fulgens (Hedw.) Brid.
Pilotrichella flexilis (Hedw.) Jaeg.
Macromitrium longifolium (Hook.) Brid.
M. punctatum (Hook. & Grey) Brid.
Pirella cavifolia (Card & Herz) Card.
Prionodon densus (Hedw.) CM.

Common genera of hepaticas present are:

Anthoceros, *Riccia*, *Pallavicinia*, *Riccardia*, *Plagiochila*, *Dumortiera*, *Moerchia*, *Lepidozia*, *Metzgeria*, *Marchantia*, *Trichocollea*, *Frullania*, *Calypogeia* and *Lejeunea*.

The question of nomenclature for the montane forests of the Caribbean has already received considerable attention. Beard (1942, 1944a) recognised a number of synonyms, preferring "montane rain forest" for the tropics but "temperate rain forest" as a world formation, the formation also occurring in New Zealand, Chile and Formosa. However, in his publication on the vegetation of the Windward and Leeward Isles (1949), he does not mention this term at all but introduces "montane thicket" which, from his description, corresponds to the former "montane rain forest". He also commended the term "cloud forest". Beard does not use the term "montane" as a synonym for "mountain," a wise precaution in the Caribbean where local conditions may bring "montane" formations to low levels.

Barbour (1942) gives details of the nomenclature covering rain forest, cloud forest and so on with equivalents in other languages but does little to clarify the basic concept.

Carabia (1945a) suggests Koppen's climatic system as a possible basis. He considers the term "rain" misleading and suggests "antillean montane forest" as a sub-climax to true rain forest and attempts a broad definition.

The present writers agree that "rain" is misleading. The mist forests of the Caribbean are not controlled by rain but by mist and cloud. The only present solution is to use a suitable term to cover the formation in this region and "montane mist forest" although not ideal, is recommended.

ELFIN WOODLAND

In Jamaica, elfin woodland is found on the exposed summits and northern ridges of the Blue Mountains at 5,000 ft., and over. It is an open woodland of gnarled and twisted trees, often short, windblown and laden with mosses, lichens, ferns and epiphytes. (Fig. 36).



FIG. 36. Elfin woodland on the exposed ridges of the Blue Mountains at 5,000 ft. The pendant moss is *Phyllogonium fulgens*.

The canopy is reduced to 20 ft., the dominants being *Clusia* and *Clethra*. Regarded as climax vegetation by Beard (1944a), it is here placed as a faciation of the montane mist forest.

Structure

There is but a single woody stratum; many of the trees which are tall at lower altitudes here mingle with the shrubs. Branching is rambling and distorted, frequently beginning close to ground level. Crowns are sheared and sloping due to wind exposure. Some of the plants have leaves that are fleshy, e.g. *Clusia* and *Hedyosmum*, tomentose e.g. *Clethra*, or small and coriaceous e.g. *Eugenia alpina* and *Ilex obcordata*. The dominant trees are: *Clethra alexandri* and *Clusia havetiooides* whilst the following are also conspicuous in the canopy layer: *Vaccinium meridionale*, *Podocarpus urbanii*, *Cyrilla racemiflora*, *Eugenia alpina*, *Weinmannia pinnata* and *Ilex montana*.

Common shrubs are: *Hedyosmum arborescens*, *Palicourea crocea*, *Ilex obcordata*, *Sciadophyllum brownii* and *Blakea trinerva*.

The field layer contains species from the mist forest including *Pilea parietaria*, *Peperomia* spp., and *Lycopodium cervicum* is locally present.

Mosses hang in festoons and, with liverworts, form mats on every trunk providing a niche for the small epiphytes, orchids, bromeliads, ferns and lichens.

Elfin woodland is present on the wet limestone slopes of the John Crow Mountains as low as 2,500 ft., and proceeds up to the plateau at 3,500 ft. The latter is worthy of special mention.

Climbing up from the Rio Grande Valley on the western slopes, the attainment of the summit plateau is sudden and definite, so abrupt is the edge of the escarpment. This plateau, tilted eastwards and 2-3 mi. in width, is made up of pinnacled and crevassed limestone simulating the broken surface of glacier ice.

The vegetation is a low, sprawling tangle of mossy trees and shrubs. Every leaf and twig is festooned with dripping bryophytes. Horizontal boughs, sinuous roots and sprawling *Clusias* lie hidden under a wet verdant mass of epiphytes. The whole is laid, like a tangle of ropes and spars, across the hopeless confusion of large and small fissures and projecting pinnacles of broken limestone. Crevasses and small ravines lie at all angles and are from 6-20 ft. in depth. The only possible progress is across the top of the tangled vegetation where every foothold must be tested before relinquishing the last. Large rock castles up to 20 ft. in height are frequent and each must be circumnavigated. It is small wonder that such an area has been little explored.

The tree layer forms a low but uneven canopy at 15-20 ft., due to the broken topography.

Trees and Shrubs:

Clusia clarendonensis is very abundant, *Clethra occidentalis*, *Guarea glabra*, *Eugenia* sp., *Columnnea hirsuta*, *Cyathea gracilis*, *Psychotria nervosa*, *Conostegia balsamians*, *Blakea trinerva*, *Sciadophyllum brownii*, *Rheedia sessiliflora*, *Piper geniculatum*.

Epiphytes include ferns, orchids, bromeliads, herbs and bryophytes:

Begonia glabra, *Pilea* sp., *Besleria lutea*, *Lycopodium wilsonii*, *Cochlidium graminoides*, *C. seminudum*, *Polypodium hartii*, *Rhipidopteris peltata*, *Trichomanes herbaceum*, *Diplazium fadyenii*, *Grammitis trifurcata*, *Elaphoglossum villosum*, *Hymenophyllum fucoides*, and *Ctenitis villosa*.

Bryophytes:

Meteoriopsis remotifolia (Hornsch) Broth.

Lepyrodontopsis trichophylla (Sw.) Broth.

Pilotrichum amasonum Mitt.

Rhizogonium spiniforms (Hadw.) Bruch.

On the exposed ridges and peaks of the Blue Mountains, many of the trees of the elfin woodland are stunted and may only reach shrub height forming a low, open forest (Fig. 37). Some of the open places are occupied by thickets of the climbing bamboo *Chusquea abietefolia* or the ferns *Gleichenia jamaicensis*, *Odontosoria aculeata*, *Histiopteris incisa*, *Paesia viscosa* and *Blechnum lineatum*. In other open areas, the ground is covered with a mat of *Lycopodium clavatum*, *L. cernuum* or *L. fawcettii*.



FIG. 37. An open aspect of Elfin woodland showing dwarfed *Podocarpus* and *Clethra alexandrii*. The floor cover is composed of *Glychenia* and *Lycopodium* sp.

On the exposed north west face of Sir John Peak and Mossmans Peak, the alpine grass *Danthonia shrevei* is found.

Shreve (1914) states that alpine grassland was encountered by Volkens on Kilamanjaro at 7,800 feet only (400 ft. higher than Blue Mountain Peak and 15 degrees lower in latitude). He suggests that, owing to the rapid erosion taking place, *Danthonia* may be a relict from the comparatively recent times when the Blue Mountains were much higher than the tree limit. The suggestion that such stunted, open woodland represents an approach towards *paramo* or Antillean meadow is tempting. Owing to its very limited expression in Jamaica, however, it is best regarded as an aspect faciation of elfin woodland which may be due to exposure and poor soil development. In some cases, as on Blue Mountain Peak where clearing has taken place in connection with

the erection of triangulation points and huts, such vegetation may well represent a stage of second growth.

SUMMARY AND DISCUSSION

The plant formations and secondary communities recognised in the coastal, lowland and montane regions of Jamaica are shown in Fig. 38. The ecological status and inter-relationship of these communities will now be discussed.

On the coastal sands, an open strand-beach associates of pioneers leads to a closed herbaceous strand dune associates and finally to a climax of *Coccocloba-Thespesia* strand woodland. In the vicinity of the southern alluvial plains, the strand sere may be deflected. Here, the stable sand is invaded by cactus-thorn scrub of *Cereus opuntia* and *Acacia tortuosa* which forms a climax community on many of the sand spits. Where raised beaches of coral rock occur, an associates of *Tournefortia-Suriana* strand scrub is to be found and this leads to a different type of strand woodland in which the palm *Thrinax parviflora* and *Hippomane mancinella* are conspicuous.

In protected coastal areas, where silt is deposited, Mangrove Woodland develops in which the four New World species *Rhizophora mangle*, *Avicennia nitida*, *Laguncularia racemosa* and *Conocarpus erecta* are present.

Swamp formations include Mangrove Woodland, Herbaceous Swamp, Palm-Sedge Marsh and Marsh Forest.

Salt flats or salinas frequently fringe the landward margins of Mangrove Woodland in the vicinity of the alluvial plains. They are dominated by an almost exclusive cover of *Batis maritima*. Large salinas show evidence of invasion by adjacent cactus-thorn scrub. These relationships of the coastal communities have been given in Fig. 12.

Freshwater herbaceous swamps are very limited in extent, being confined to low-lying river margins. Typical plants are *Typha*, *Phragmites* and *Arundo donax* together with floating and submerged aquatics. Palm-Sedge Marsh and Marsh Forest are confined to the low-lying flood plains of which the Black River produces the most noteworthy example. The former is dominated by *Cyperus* spp. with the sporadic occurrence of the palms *Roystonea princeps* and *Calyptronoma swartzii*. Marsh Forest occurs in slightly higher ground and these two palms together with *Sympogonia globulifera* form a closed canopy.

Thorn-scrub dominated by *Prosopis juliflora* is found on the alluvial plains. The ecological conditions necessary for the development of thorn-scrub and its cactus faciation need to be determined before their relationship to one another and to the coastal communities can be fully understood.

The coastal limestone hills support a sub-climax community of Dry Limestone Scrub Forest. This is a relatively open stunted woodland with a great number of species and few large trees. It has been subject to human interference and formerly included a much richer tree flora. This is well shown to-day in




FIG. 38. Scheme to show the Formations and Secondary Communities recognized in Jamaica, their ecological status and inter-relationship.

the Dry Limestone Forest on the ridge at Portland Point where over 80 tree species have been identified forming a canopy at 50-60 ft. and a sub-canopy at 20-30 ft. Arid Limestone Scrub Forest is a coastal faciation of this plant community. Palm-Hippomane Strand Woodland and Arid Limestone Scrub Forest have many species in common which are evergreen and have thick, leathery leaves. The latter community differs mainly in having a larger number of species and these are common to the Dry Limestone Scrub

Forest. We have regarded it as a faciation of Dry Limestone Scrub Forest but it may well be an ecotone between this community and Strand Woodland.

The former presence of forest on the alluvial plains is conjectured from the number of remnant scattered native trees that could form woodland were it not for interference of man and his animals. At the present time, it is replaced by induced savanna, with scattered trees of *Acacia lutea* and *Samanea saman* or by Acacia or logwood thicket.

A remnant forest of large spreading trees including *Ceiba* and *Cedrella* occurs in the lower valleys of the shale hills. This was formerly much more widespread on the hills than it is to-day. Shifting cultivation and erosion has led to secondary communities of either induced hill savanna with scattered mango trees or of bamboo brake.

Wet Limestone Forest occurs where the rainfall is over 100 in. per year. It is seen to best advantage at the present time in the elevated Cockpit country. This is a luxuriant evergreen forest which has physiognomic and floristic affinity to Lower Montane Rain Forest. It is considered to be a faciation of Tropical Rain Forest.

No typical Tropical Rain Forest exists in Jamaica to-day but, from the Tree Fern Brake of the Rio Grande Valley, its former presence is a very reasonable conjecture. Tree Fern Brake is regarded as a secondary or sub-climax community to Rain Forest. It does not occur at the present time outside the Rio Grande Valley. Ørsted, who spent six weeks in Jamai-

ca in 1846, made the tree-ferns of Jamaica classical. He wrote "There is probably no other place on earth where the tree-ferns attain such considerable height and where, furthermore, these ferns form such large continuous woods displacing all other plants, as can be found in the Jamaican mountains." He is taken to task rather sharply over these remarks by Brockman-Jerosch who visited Jamaica in 1914 but failed to see tree-fern brake. However, Ørsted's map (Fig. 39) is highly informative. He supports his description by sketching in a wide belt of tree-fern brake below the mist forest on the northern slopes of the Blue Mountains. This is clearly the result of clearing which is also indicated. Man and not the tree-ferns have "displaced" the other plants. It is to be noted that the tree-fern brake is shown for the north slopes where it has apparently followed clearing of lower montane rain forest.

Lower Montane Rain Forest is still to be found in the less accessible wet northern slopes of the Blue Mountains and the western slopes of the John Crow Mountains bounding the Rio Grande Valley. It is a montane faciation of lowland Tropical Rain Forest. Two types have been observed. One is a mixed forest dominated by tall, straight boled trees such as *Psidium montanum*, *Calophyllum jacquinii*, *Matayba apetala* and *Sympodia globulifera*. In the other type, *Calophyllum jacquinii* alone is dominant.

Montane Sclerophyll is typical of the dry, southern slopes of the Blue Mountains between altitudes of 2,500-4,000 ft. It has two aspects—a lower, open




PLATE 3.—AN EARLY ATTEMPT TO ILLUSTRATE THE VEGETATIONAL ZONES OF A TROPICAL ISLAND.—From A. S. ØRSTED, "Jamaica, En Natur-skildring" (1857).—Courtesy Arnold Arboretum of Harvard University.

FIG. 39. Vegetation of Jamaica (From A. S. Ørsted). Courtesy of Arnold Arboretum of Harvard University. Note the coastal vegetation, the dry southern plains and the upland plateau of Manchester, with scattered bottle-shaped cotton trees, rising to the wet cockpit areas. The north-eastern aspect is shown on the right and includes the Lowland Tropical Rain Forest of the Rio Grande valley with the John Crow Mountains on the left. The lower slopes of the Blue Mountains already show coffee plantations, patchwork cultivation and second growth tree fern brake. The higher reaches support Mist Forest.

zone consisting of xerophytic shrubs widely dispersed in a ground flora of grasses. Shifting cultivation prevails and burning here prevents the spread of the shrubs but encourages the spread of the introduced grass *Melinis minutiflora* and the fern *Gleichenia pectinata*. These plants are rapidly dominating large areas. Above this zone is a less disturbed and more closed sclerophyllous thicket of dwarf, spindly trees and shrubs in which *Cyrilla racemiflora*, *Clethra occidentale* and *Vaccinium meridionale* are conspicuous. The lower bushy sclerophyll zone is thought to be a secondary developmental community which, if undisturbed, would lead to sclerophyllous thicket in these seasonally dry areas. There is, however, some indication that, where it occurs in wetter regions, it may bear the same relationship to Lower Montane Rain Forest.

Montane Mist Forest occurs extensively in the Blue Mountain range above 4,500 ft. It is shrouded in mist for most of the day. Low-canopied evergreen trees are associated with an abundance of undershrubs, ferns, bryophytes and lichens. Elfin Woodland which occurs on the exposed ridges and summits, is regarded as an open faciation of the Mist Forest. Fig. 40 shows the distribution of some of the vegetation units as seen in a diagrammatic cross section of the island from the Parish of Portland in the north-east running south-west to the Parish of St. Andrew.




FIG. 40. Diagrammatic representation of the vegetation seen in a cross section of the eastern end of the island passing through the parishes of Portland and St. Andrew.

It will be profitable to relate the vegetation as a whole with the Formations proposed by Beard (1944a) for use throughout Tropical America and the Caribbean. Beard follows the present trend in tropical ecology and recognises physiognomic units, as distinct from floristic units, each with a characteristic essential habitat. He distinguishes an optimal Formation "where every condition for plant growth is as ideally favourable as it can be in the tropics" and Formation Series "within which are structures and life forms expressing every degree of transition from the optimum to extreme adversity." The optimal Formation type is Rain Forest. The Formation Series, each of which is based on a major type of habitat, are as follows:

- i. Seasonal Formations
- ii. Dry Evergreen Formations

- iii. Montane Formations
- iv. Swamp Formations
- v. Marsh (Seasonal Swamp) Formations

The swamp and marsh formations may be considered to be edaphic climax formations—the remainder, including the optimal Formations are climatic climax formations. Within each Formation Series are units arranged in regard to moisture relations but distinguished purely on physiognomic data. In the Seasonal Formations, for example, where the characteristic of the habitat is seasonal drought, there is Evergreen Seasonal Forest ranging through Deciduous Seasonal Forest to Scrub and finally Desert.

With time and practical application, Beard has seen fit to modify and make additions to his system for which reference should be made to his publications. The Pan-Caribbean Forestry Meeting in 1946 adopted Beard's system until "further work should confirm or decry its utility." The following discussion is a contribution towards that end.

In Table 6 may be seen the correlation of the major vegetation types recognised in Jamaica by the authors with the Formations proposed by Beard.

TABLE 6. The Vegetation of Jamaica in relation to the Caribbean Climax Formations of Beard (1944a).

Main Vegetation Types Recognized in Jamaica	Equivalent Vegetation Types Recognized by Beard
Conjectured from presence of Tree Fern Brake	1. OPTIMAL FORMATION Rain Forest
Wet Limestone Forest? Dry Limestone Scrub Forest Thorn Scrub Cactus-Thorn Scrub	2. SEASONAL FORMATIONS Evergreen Seasonal Forest? Semi-Evergreen Seasonal Forest? Thorn Woodland Cactus Scrub
Strand Woodland	3. DRY EVERGREEN FORMATIONS Littoral Woodland
Lower Montane Rain Forest Montane Mist Forest Palm Brake? Elfin Woodland Montane Sclerophyll	4. MONTANE FORMATIONS Lower Montane Rain Forest Montane Rain Forest Palm Brake Elfin Woodland No equivalent
Herbaceous Swamp Mangrove Woodland	5. SWAMP FORMATIONS Herbaceous Swamp Mangrove Woodland
Marsh Forest Palm-Sedge Marsh	6. MARSH (SEASONAL SWAMP) FORMATION Marsh Forest Palm Marsh

The Optimal Formation type, i.e. Tropical Rain Forest, as defined by Beard, permits of no physiognomic variation since the habitat conditions are uniformly favourable all the year round. This is, perhaps, too rigid a definition. It is felt that both Evergreen Seasonal Forest, with which our Wet Limestone Forest corresponds in structure and seasonal moisture relations, and Lower Montane Rain

Forest might well be regarded as faciations of typical Rain Forest.

Although the Dry Limestone Forests undoubtedly belong to the Seasonal Formation Series, more details of their physiognomy are required before they can be correlated with certainty. At the moment, all that can be said is that there would seem to be a greater affinity to Semi-Evergreen Seasonal Forest than Deciduous Seasonal Forest although the severe habitat conditions would lead one to expect the latter formation type. No doubt, in former times, there were present in Jamaica more luxuriant gradations of this type of seasonal forest in areas of decreasing seasonal drought, which culminated in Wet Limestone Forest. Such areas are now mainly under cultivation and pasture. The remnants that are left have not yet been studied.

Cactus-thorn Scrub is considered by the authors to be a faciation of Thorn Scrub.

The habitat conditions for the Dry Evergreen Formations as defined by Beard are well-drained lands where the moisture supply is not seasonal but fairly consistently inadequate all the year round. This may be due to strong winds and excessively porous soil. The typical species are hard-leaved evergreens. Strand Woodland corresponding to Beard's Littoral Woodland is a member of this formation series.

In Jamaica, there is a seasonal drought in all low-lying coastal areas, with the possible exception of the northeast, which is just as severe as that prevailing on the coastal limestone hills supporting deciduous seasonal forest. Yet there is this abrupt change to small, hard-leaved, evergreen woodland. It is suggested that here it is the presence of salt laden winds that are responsible rather than consistent drought. The fact that the coastal faciation of dry limestone forest has many evergreen species in common with strand woodland is significant in this respect.

Montane Mist Forest corresponds to Beard's Montane Rain Forest. Daily mist is the most notable characteristic of the habitat and the authors suggest that Mist Forest is the better term. Elfin Woodland is regarded as an open stunted faciation of Mist Forest, brought about by more exposed conditions, since, for the most part, it is made up of species common in the Mist Forest. Montane Sclerophyll has no counterpart in Beard's classification. Later he uses the term "Montane Thicket" which might well be a suitable one for the upper sclerophyllous thicket of Jamaica, but, from his description of this community, it would seem to be a synonym for his Montane Rain Forest. A community that would seem to correspond to Palm Brake occurs in the western slopes of the John Crow Mountains above Lower Montane Rain Forest. It has not yet been investigated. The swamp and marsh communities of Jamaica fit in very well with Beard's nomenclature and need no further comment.

In conclusion, it may be said that Beard's classification as applied to Jamaica, receives general ratification. Whether it is desirable to separate Ever-

green Seasonal Forest and Lower Montane Rain Forest so markedly from Tropical Rain Forest is doubtful. It is suggested that the controlling habitat factor for Littoral Woodland is the presence of salt-laden winds, not consistent drought. A type of seasonal forest called Dry Limestone Scrub Forest is produced in Jamaica under the latter conditions. The addition of Montane Sclerophyll to the Montane Formation series is desirable.

The present vegetation of Jamaica enables reasonably accurate reconstruction of the original vegetation of the island to be made. Early descriptions of dense jungles with the impression of extensive lowland tropical rain forest must be discounted.

LITERATURE CITED

- Barbour, W. R. 1942. Forest types of tropical America. *Caribbean Forester* **3**: 137-150.
- Beard, J. S. 1942. Montane vegetation in the Antilles. *Caribbean Forester* **3**: 61-74.
- _____. 1944a. Climax vegetation in tropical America. *Ecology* **25**: 127-158.
- _____. 1944b. The natural vegetation of Tobago, B. W. I. *Ecol. Monog.* **14**: 135-163.
- _____. 1945. Progress of succession on the Soufrière of St. Vincent. *Jour. Ecol.* **33**: 1-9.
- _____. 1946. The natural vegetation of Trinidad. *Oxford Forestry Mem.* **20**: 1-152.
- _____. 1949. The natural vegetation of the Windward and Leeward Islands. *Oxford Univ. Press*. 191 pp.
- Brennan, J. F. 1936. Meteorology of Jamaica. *Jamaica: Govt. Print.* 32 pp.
- Britton, N. L. & P. Wilson. 1923-26. The Botany of Porto Rico and the Virgin Islands. In *Scientific Survey of Porto Rico and the Virgin Islands*. Vol. 5 (1923-24) & 6 (1925-26). N. Y. Acad. Sci.
- Brockman-Jerosch, H. & M. 1925. *Jamaika—Vegetationsbilder* **16**: Reihe, Heft 5/6, Jena.
- Brown, E. P. 1756. *A civil and natural history of Jamaica*. London. (Ed. 2. 1789.)
- Borgesen, F. 1923. Et Besög; Jamaica's Bregneskove. In "Naturens Verden" 7, Aarg, Heft 7. Copenhagen.
- Burtt-Davy, J. 1938. The classification of tropical woody vegetation-types. *Imp. Forestry Inst. Pap.* No. 13. Oxford, England.
- Burns, L. V. 1947. A preliminary study of the forest vegetation of Jamaica. (Unpublished.)
- Carabia, J. P. 1943. Endemism in the Cuban Flora. (Abst.) *Suppl. Amer. Jour. Bot.* **29**(10): 4s.
- _____. 1945a. Vegetation of Sierra de Nipe Cuba. *Ecol. Monog.* **15**: 324-341.
- _____. 1945b. A brief review of the Cuban flora. In Verdoorn, F. *Plants and plant science in Latin America*. Chronica Botanica Co. pp. 68-70.
- Caribbean Commission. 1947. *Forest research within the Caribbean area*. Washington. Pp. 78.
- Carr, A. F. 1949. La Montana Llorona. *Sci. Monthly* **68**: 225-234.
- Chapman, V. J. 1940. In Steers, J. A., V. J. Chapman, J. Colman & J. A. Lofthouse. *Sand Cays & Mangroves in Jamaica*. *Geog. Jour.* **96**(5): 305-328.

- . 1944. The 1939 Cambridge University Expedition to Jamaica. Part I. A study in the botanical processes concerned in the development of the Jamaican shore-line. *Linn. Soc. London Jour.* **52**: 407-447.
- . 1947. The geographical elements of the seaweed flora of Jamaica. *Nat. Hist. Soc. Jamaica. Nat. Hist. Notes* **3**: 88-89.
- Charter, C. F.** 1941. A reconnaissance survey of the soils of British Honduras. *Trinidad: Govt. Printing Office.*
- Croucher, H. H.** 1938. A soil survey of a portion of the Lower Clarendon Plain. *Jamaica: Govt. Printer.*
- Dansereau, P.** 1951. Description and recording of vegetation upon a structural basis. *Ecology* **32**: 172-229.
- Davis, J. H.** 1940. The ecology and geologic role of mangroves in Florida. *Carnegie Inst. Wash. Pub.* **517**: 303-412.
- . 1942. The ecology of the vegetation and topography of the sand keys of Florida. *Carnegie Inst. Wash. Pub.* **524**: 115-195.
- Fawcett, W. & A. B. Rendle.** 1910-1936. Flora of Jamaica. *Brit. Museum Nat. Hist. London. Vol. I, III, IV, V, VII.*
- Grisebach, A. H. R.** 1864. Flora of the British West Indian Islands. *London.*
- Gleason, H. A. & M. I. Cook.** 1927. The plant ecology of Porto Rico. *Scientific survey of Porto Rico and the Virgin Islands. Vol. 7. N. Y. Acad. Sci.*
- Gill, T.** 1931. Tropical forests of the Caribbean. *Trop. Plant Res. Found. Washington, D. C.*
- Gooding, E. G. B.** 1947. Observations on the sand dunes of Barbados. *Jour. Ecol.* **34**: 111-125.
- Good, R.** 1947. The geography of the flowering plants. *London. Pp. 13-403.*
- Hardy, F.** 1951. Soil productivity of the British Caribbean. *Imp. Coll. Trop. Agr. Jour.* **28**: 3-31.
- Harshberger, J. W.** 1911. Phytogeographic survey of North America. *Veget. der Erde. Vol. 13*: 790 pp.
- . 1903. Notes on the strand flora of great Inagua, Haiti and Jamaica. *Torreya* **3**: 67-70.
- Hitchcock, A. S.** 1936. Manual of the grasses of the West Indies. *U. S. Dept. Agr. Misc. Pub.* **243**: 1-439.
- Hollick, A.** 1924. A review of the fossil flora of the West Indies. *N. Y. Bot. Gard. Bull.* **12**: 259-323.
- . 1929. Palaeobotany of Porto Rico. *Scientific Survey of Porto Rico and the Virgin Islands. Vol. 7. N. Y. Acad. Sci.*
- Holdridge, L. R.** 1945. A brief sketch of the flora of Hispaniola. In Verdoorn, F. Plants and plant science in Latin America. Pp. 67-68. *Waltham, Mass.: Chronica Botanica Co.*
- Howard, R. A.** 1952. The vegetation of the Grenadines, Windward Isles, British West Indies. *Gray Herbarium, Harvard Univ. Contrib. No. 174.*
- Hutchinson**, 1926 & 1934. The families of flowering plants. (2 vols.) *London.*
- Jenman, G. S.** 1890-93. Synoptical list with descriptions of the ferns and fern allies of Jamaica. *Bot. Dept. of Jamaica Bull.*
- Lanjouw, J.** 1945. On the location of botanical collections from Central and South America. In Verdoorn, F. Plants and plant science in Latin America. Pp. 224-235. *Waltham, Mass.: Chronica Botanica Co.*
- Leon, H.** 1946. *Flora de Cuba. Vol 1. Habana.*
- Leon, H. & H. Alain.** 1951. *Flora de Cuba. Vol. 2. Habana.*
- Lewis, C. B.** 1943. Notes on the trees of Portland Ridge. *Nat. Hist. Soc. Jamaica* **14-17**.
- . 1947. A trip to the Morant and Pedro Cays. *Nat. Hist. Soc. Jamaica* **3**: 105-108.
- Long, E.** 1774. The history of Jamaica. *London. 3 vols.*
- Lunan, J.** 1814. *Hortus Jamaicensis. Jamaica.*
- Marie-Victorin, Frere & Frere, Leon.** 1942. *Itinéraires Botaniques dans l'Île de Cuba. Vol. 1. Montreal.*
- Macfadyen, J.** 1837. *Flora of Jamaica. London. Vols. 1 & 2 [1850].*
- Miranda, F. & A. J. Sharp.** 1950. Characteristics of the vegetation in certain temperate regions of Eastern Mexico. *Ecology* **31**: 313-333.
- Ørsted, A. S.** 1857. *Jamaica. En Naturskildring. Copenhagen.*
- Pittier, H.** 1939. *Supplements a las plantas usuales de Venezuela. Editorial Elite. Caracas.*
- Proctor, G. R.** 1953. A preliminary check-list of the Jamaican pteridophytes. *Inst. Jamaica. Sci. Bull. Ser. 5*: 1-89.
- Raunkiaer, C. F.** 1934. The life forms of plants and statistical plant geography. *Oxford, England.*
- Richards, P. W., A. G. Tansley & A. S. Watt.** 1939. The recording of structure, life-form and flora of tropical forest communities as a basis for their classification. *Imp. Forestry Inst. Pap. No. 19. Oxford, England.*
- Sawkins, J. G.** 1869. Memoirs of the geological survey. Reports on the geology of Jamaica or Part II of the West Indian Survey. *London: H. M. S. O. 340 pp.*
- Schuchert, C.** 1935. Historical geology of the Antillean-Caribbean region. *New York: Wiley & Son.*
- Seifriz, W.** 1943. *Plant Life of Cuba. Ecol. Monog.* **13**: 375-426.
- Shreve, F.** 1910. The coastal deserts of Jamaica. *Plant World* **13**: 129-134.
- . 1914. A montane rain forest. *Carnegie Inst. Washington Pub.* **119**: 1-110.
- . 1945. The vegetation of Jamaica. In Verdoorn, F. Plants and plant science in Latin America. Pp. 287-289. *Waltham, Mass.: Chronica Botanica Co.*
- Sloane, H.** 1696. Catalogus plantarum quae insula Jamaica sponte proveniunt. *London.*
- . 1707-1725. Voyage to . . . Jamaica, with the natural history. *London. 2 vols.*
- Standley, P. C.** 1941. The forests of Guatemala. *Tropical Woods.* **67**: 1-18.
- Stehlé, H.** 1935. *Essai d'éologie et de géographie botanique. Flore de Guadeloupe. Basse-terre, Guadeloupe.*
- . 1945. Les conditions écologique la vegetation et les ressources agricoles de l'Archipel des Petits Antilles. In Verdoorn, F. Plants and plant science in Latin America. Pp. 85-99. *Waltham, Mass.: Chronica Botanica Co.*
- . 1945. Forest types of the Caribbean islands. *Caribbean Forester* **6** (Suppl.): 273-408.

- Steers, J. A., V. J. Chapman, J. Colman & J. A. Loft-house. 1940. Sand Cays and mangroves in Jamaica. *Geog. Jour.* 96(5): 305-328.
- Stevenson, N. S. 1942. Forest associations of British Honduras. *Caribbean Forester* 3: 164-171.
- Steyermark, J. A. 1950. Flora of Guatemala. *Ecology* 31: 368-372.
- Swabey, C. 1942. The classification of vegetation in Jamaica. *Nat. Hist. Soc. Jamaica. Nat. Hist. Notes* 1(7): 12-17.
- . 1941. The principal timbers of Jamaica. *Dept. Sci. & Agr. Jamaica. Bull.* 29.
- Urban, I. 1898. *Symbolae Antillanae*. Berlin.
- Verdoorn, F. 1945. Plants and plant science in Latin America. *Chronica Botanica Co.* XXXVII. 381 pp.
- Woodson, R. E. 1940. The Apocynaceous Flora of the Yucatan Peninsula. *Carnegie Inst. Wash. Pub.* 522.

APPENDIX

The nomenclature adopted by Fawcett & Rendle has been followed wherever possible. For plants not covered by this Flora, reference has been made to the Jamaica Herbarium and the following works:

Flora of the British West Indian Islands by A. H. R. Grisebach (1864); The Botany of Porto Rico and the Virgin Islands by N. L. Britton (1923-26); Flora de Cuba by Leon (1946) and Leon & Alain (1951); Manual of the grasses of the West Indies by A. S. Hitchcock (1936); A preliminary check-list of Jamaican pteridophytes by G. R. Proctor (1953).

The mosses have been checked by E. Bartram, and the marine algae by Prof. V. J. Chapman. They are not included in the appendix since authorities are given in the text.

A check-list of plants mentioned in the text (excluding the bryophytes and marine algae) follows.

DICOTYLEDONS

Acanthaceae

Andrographis paniculata Ns.*Barleria prionitis* L.*Blechum brownei* Juss.*Pachystachys coccinea* Ns.*Ruellia tuberosa* L.*Thunbergia fragrans* Roxb.

Amarantaceae

Achyranthes indica Mill.*Alternanthera fieoidea* Roem. & Schult.*A. parvifolia* Fawc. & Rendle*A. repens* Kuntze.*Amarantus viridis* L.*A. spinosus* L.*Celosia argentea* L.*Gomphrena globosa* L.*Iresine paniculata* Kuntze*Philoxerus vermicularis* Beauv.

Anacardiaceae

Anacardium occidentale L.*Comocladia pinnatifolia* L.*C. velutina* Britton*Mangifera indica* L.*Mosquitoxylon jamaicense* Kr. & Urb.*Rhus metopium* L.*Spondias monbin* L.*S. purpurea* L.

Anonaceae

Anona glabra L.*A. palustris* L.*A. reticulata* L.*A. squamosa* L.*Oxandra lanceolata* Braill.*Oxandra laurifolia* A. Rich.*Xylopia muricata* L.

Apocynaceae

Echites suberecta Jacq.*E. umbellata* Jacq.*Plumeria alba* L.

Araliaceae

Ilex montana Griseb.*I. obcordata* Sw.

Gilibertia arborea March.

G. nutans March.*G. ovalifolia* Fawc. & Rendle*Sciadophyllum brownei* Spreng.

Aristolochiaceae

Aristolochia odoratissima L.*A. trilobata* L.

Asclepiadaceae

*Asclepias curassavica**A. nivea* L.*Metastelma atrorubens* Schlecht.*M. fawcettii* Schlecht.*M. harrisii* Schlecht.

Balanophoraceae

Seybalium jamaicense Schott & Endl.

Batidaceae

Batis maritima L.

Begoniaceae

Begonia acutifolia Sw.*B. glabra* Aubl.

Bignoniaceae

Catalpa longissima Sims.*Crescentia cujete* L.*Schlegelia parasitica* Mrs.*Spathodea campanulata* Beauv.*Tecoma leucoxylon* Mart.*T. stans* Juss.

Bombacaceae

Ceiba pentandra Gaertn.*Ochroma pyramidale* Urb.

Boraginaceae

Beureria succulenta Jacq.*Cordia alba* R. & Sch.*C. collococca* L.*C. cylindristachya* R. Sch.*C. gerascanthoides* H. B. K.*C. nitida* Vahl*C. sebestana* Jacq.*Heliotropium curassavicum* L.*H. parviflorum* L.*Tournefortia astrotrichia* DC.*T. cymosa* (L.) DC.*T. gnaphalodes* R. Br.*T. hirsutissima* L.*T. volubilis* L.

Brunelliaceae

Brunellia comocladifolia Humb. & Bonpl.

Burseraceae

Bursera simaruba Sarg.*B. simplicifolia* DC.

Caetaceae

Cereus flagelliformis Miller*C. peruvianus* Miller*C. triangularis* Haw.*Melocactus communis* Link & Otto*Opuntia jamaicensis* Britt. & Harris*O. spinosissima* Miller*O. tuna* Miller

Caesalpiniaceae

Bauhinia divaricata L.*Caesalpinia bonducella* Fleming*C. coriaria* Willd.*C. vesicaria* L.*Cassia emarginata* L.*C. viminea* L.*Haematoxylon campechianum* L.*Hymenaea courbaril* L.*Parkinsonia aculeata* L.*Peltophorum brasiliense* Urb.*Poinciana regia* Boj.*Tamarindus indica* L.

Campanulaceae

Lobelia acuminata Sw.*L. assurgens* L.

Canellaceae

Canella winterana Gaertn.*Cinnamodendron corticosum* Miers.

Capparidaceae

Capparis cynophallophora L.*C. ferruginea* L.*C. flexuosa* L.*Cleome serrata* Jacq.*C. spinosa* Jacq.*Crataeva gynandra* L.*C. tapia* L.

Caprifoliaceae

Viburnum villosum Sw.

Caricaceae

Carica jamaicensis Urb.

Casuarinaceae	Cunoniaceae	Goodeniaceae
<i>Casuarina equisetifolia</i> Forst.	<i>Weinmannia pinnata</i> L.	<i>Scaevola plumieri</i> (L.) Vahl
Celastraceae	<i>Cyrillaceae</i>	Guttiferae
<i>Gyminda latifolia</i> Urb.	<i>Cyrilla racemiflora</i> L.	<i>Calophyllum jacquinii</i> Fawc. & Rendle
<i>Rhacomia crossopetalum</i> L.	Ebenaceae	<i>Clusia clarendonensis</i> Britton
<i>Schaefferia frutescens</i> Jacq.	<i>Diospyros tetrasperma</i> Sw.	<i>C. flava</i> Jacq.
Ceratophyllaceae	Ericaceae	<i>C. havetoides</i> Planch.
<i>Ceratophyllum demersum</i> L.	<i>Lyonia jamaicensis</i> Don	<i>C. rosea</i> Jacq.
Chenopodiaceae	Erythroxylaceae	<i>Mammea americana</i> L.
<i>Atriplex cristata</i> Humb. & Bonpl.	<i>Erythroxylon areolatum</i> L.	<i>Rheedia pendula</i> Urb.
<i>Salicornia ambigua</i> Michx.	<i>E. rotundifolium</i> Lun.	<i>R. sessiliflora</i> Planch.
Chloranthaceae	Euphorbiaceae	<i>Sympmania globulifera</i> L. f.
<i>Hedosymnum arborescens</i> Sw.	<i>Acalypha scabrosa</i> Sw.	Hernandiaceae
<i>H. nutans</i> Sw.	<i>A. virgata</i> L.	<i>Hernandia catalpifolia</i> Britt. & Harris
Clethraceae	<i>Acidotoma urens</i> Sw.	Hypericaceae
<i>Clethra alexandri</i> Griseb.	<i>Adelia ricinella</i> L.	<i>Ascyrum hypericoides</i> L.
<i>C. occidentalis</i> (L.) Steud.	<i>Alchornea latifolia</i> Sw.	Labiatae
Combretaceae	<i>Argythamnia candicans</i> Sw.	<i>Hyptis capitata</i> Jacq.
<i>Bucida buceras</i> L.	<i>Bernardia carpinifolia</i> Griseb.	<i>Leonurus sibiricus</i> L.
<i>Buchenavia capitata</i> Eichl.	<i>Croton flavens</i> L.	<i>Micromeria obovata</i> Benth.
<i>Conocarpus erecta</i> L.	<i>C. glabellus</i> L.	<i>Salvia occidentalis</i> Sw.
<i>Laguncularia racemosa</i> Gaertn.	<i>C. grisebachianus</i> Muell.	Lacistemaceae
<i>Terminalia catappa</i> L.	<i>C. humilis</i> L.	<i>Lacistema aggregatum</i> Fawc. & Rendle
<i>T. latifolia</i> Sw.	<i>C. linearis</i> Jacq.	Lauraceae
Compositae	<i>C. ovalifolius</i> Vahl.	<i>Hufelandia pendula</i> Nees
<i>Ambrosia hispida</i> Pursh	<i>Drypetes lateriflora</i> Kr. & Urb.	<i>Misanthea triandra</i> Mez.
<i>Baccharis scoparia</i> (L.) Sw.	<i>Euphorbia buxifolia</i> Lam.	<i>Nectandra antillana</i> Meisn.
<i>Bidens cynapiifolia</i> H. B. K.	<i>E. blodgettii</i> Engelm.	<i>N. coriacea</i> Griseb.
<i>B. pilosa</i> L.	<i>E. heterophylla</i> L.	<i>N. patens</i> Griseb.
<i>B. reptans</i> (L.) G. Don.	<i>E. hirta</i> L.	<i>N. sanguinea</i> Roland ex. Rottb.
<i>Borreria arborescens</i> (L.) DC.	<i>E. hypericifolia</i> L.	<i>Ocotea jamaicensis</i> Mez.
<i>Egletes prostrata</i> (Sw.) Kuntze	<i>Gymnanthes elliptica</i> Sw.	<i>O. martinicensis</i> Mez.
<i>Erigeron canadensis</i> L.	<i>G. lucida</i> Sw.	<i>O. staminea</i> Mez.
<i>Eupatorium dalea</i> L.	<i>Hippomane mancinella</i> L.	<i>Persea alpigena</i> Spreng.
<i>E. odoratum</i> L.	<i>Hura crepitans</i> L.	Lentibulariaceae
<i>Gnaphalium americanum</i> Mill.	<i>Jatropha gossypifolia</i> L.	<i>Utricularia stellaris</i> L.
<i>Isocharpha oppositifolia</i> (L.) R. Br.	<i>Lasiocroton macrophyllus</i> Griseb.	Loganiaceae
<i>Pectis ciliaris</i> L.	<i>Phyllanthus acuminatus</i> Vahl	<i>Spigelia anthelmia</i> L.
<i>P. febrifuga</i> van Hall	<i>P. angustifolius</i> Sw.	Malpighiaceae
<i>Senecio discolor</i> (Sw.) DC.	<i>P. epiphyllanthus</i> L.	<i>Byrsonima coriacea</i> DC.
<i>Sonchus oleraceus</i> L.	<i>P. linearis</i> Sw.	<i>Heteropteris laurifolia</i> A. Juss.
<i>Spilanthes urens</i> Jacq.	<i>P. niruri</i> L.	<i>Malpighia glabra</i> L.
<i>Verbesina pinnatifida</i> Sw.	<i>Picroidendron baccatum</i> Kr. & Urb.	<i>M. puniceifolia</i> L.
<i>Vernonia divaricata</i> Sw.	<i>Ricinus communis</i> L.	<i>Stigmaphyllon emarginatum</i> A. Juss.
Connaraceae	<i>Sapium jamaicense</i> Sw.	Malvaceae
<i>Rourea paucifoliolata</i> Planch.	<i>Securinega acidoton</i> Fawc. & Rendle	<i>Abutilon giganteum</i> Sweet.
Convolvulaceae	<i>Tragia volubilis</i> L.	<i>Hibiscus elatus</i> Sw.
<i>Calonyction tuba</i> Schlecht.	Ficoideaceae	<i>Pavonia rosea</i> Schlecht.
<i>Cuscuta americana</i> L.	<i>Sesuvium portulacastrum</i> L.	<i>Sida procumbens</i> Sw.
<i>Ipomea carnea</i> Jacq.	Flacourtiaceae	<i>S. rhombifolia</i> L.
<i>I. fastigiata</i> Swt.	<i>Casearia guianensis</i> Urb.	<i>Thespesia populnea</i> Solander
<i>I. jamaicensis</i> G. Don.	<i>C. hirsuta</i> Sw.	Maregraviaceae
<i>I. pes-caprae</i> Sw.	<i>C. nitida</i> Jacq.	<i>Marcgravia brownei</i> Kr. & Urb.
<i>I. quinquefolia</i> L.	<i>Laetia thamnia</i> L.	Melastomaceae
<i>I. sidifolia</i> Choisy.	Garryaceae	<i>Blakea trinervia</i> L.
<i>I. umbellata</i> Mey.	<i>Garrya fadyenii</i> Hook.	<i>Conostegia balbisiana</i> Ser.
Crassulaceae	Gentianaceae	<i>C. superba</i> Naud.
<i>Bryophyllum pinnatum</i> Kurz.	<i>Eustoma exaltatum</i> Griseb.	<i>Heterotrichum umbellatum</i> Urb.
Cruciferae	<i>Leianthus longifolius</i> Griseb.	<i>Mecranium amygdalinum</i> Triana.
<i>Cakile lanceolata</i> O. E. Schulz.	Gesneriaceae	<i>M. purpurascens</i> Triana.
Cucurbitaceae	<i>Achimenes coccinea</i> Pers.	<i>M. virgatum</i> Triana.
<i>Cionosicyos pomiformis</i> Griseb.	<i>Besteria lutea</i> L.	<i>M. leucantha</i> Sw.
<i>Cucumis anguria</i> L.	<i>Columnea hirsuta</i> Sw.	<i>M. purpurea</i> Sw.
<i>Momordica charantia</i> L.	<i>Rytidophyllum grande</i> Mart.	<i>Miconia dodecandra</i> Cogn.

<i>M. laevigata</i> DC.	<i>P. guajava</i> L.	Polygonaceae
<i>M. quadrangularis</i> Naud.	<i>P. montanum</i> Sw.	
<i>M. rubens</i> Naud.		<i>Antigon leptopus</i> Hook.
<i>Ossaea asperifolia</i> Triana.		<i>Coccoloba diversifolia</i> Jaeq.
Meliaceae		<i>C. krugii</i> Lindau.
<i>Cedrela odorata</i> L.	<i>Boerhavia scandens</i> L.	<i>C. laurifolia</i> Jaeq.
<i>Guarea glabra</i> Vahl	<i>Mirabilis jalapa</i> L.	<i>C. littoralis</i> Urb.
<i>Swietenia mahogani</i> Jaeq.	<i>Pisonia aculeata</i> L.	<i>C. longifolia</i> Fisch.
<i>Trichilia hirta</i> L.	<i>P. fragrans</i> Dumont.	<i>C. uvifera</i> L.
Menispermaceae	<i>P. obtusata</i> Jaeq.	
<i>Cissampelos pareira</i> L.		Portulacaceae
<i>Hyperbaena domingense</i> Benth.		<i>Portulaca oleracea</i> L.
Mimosaceae		<i>Talinum paniculatum</i> Gaertn.
<i>Acacia lutea</i> Hitchc.		Quiinaceae
<i>A. tortuosa</i> Willd.	<i>Nymphaea alba</i> L.	<i>Quiinia jamaicensis</i> Griseb.
<i>A. villosa</i> Willd.		Ranunculaceae
<i>Adenanthera pavonina</i> L.	<i>Linociera ligustrina</i> Sw.	<i>Clematis dioica</i> L.
<i>Albizia lebbeck</i> Benth.	<i>Ximenia americana</i> L.	Rhamnaceae
<i>A. berteriana</i> (Balbis) Maza.		<i>Colubrina asiatica</i> Brongn.
<i>Enterolobium mangense</i> Fawc.		<i>C. ferruginosa</i> Brongn.
Rendle		<i>C. reclinata</i> Brongn.
<i>E. saman</i> Prain.		<i>Gouania lupuloides</i> Urb.
<i>Inga vera</i> Willd.		<i>Krugiodendron ferreum</i> Urb.
<i>Leucaena glauca</i> Benth.		<i>Rhamnus sphaerosperma</i> Sw.
<i>Mimosa pudica</i> L.		<i>Rhamnidium jamaicense</i> Urb.
<i>Pithecellobium alexandri</i> Urb.		<i>Sarcophalus laurinus</i> Griseb.
<i>P. arboreum</i> Urb.		<i>Zizyphus chloroxylon</i> Oliv.
<i>P. unguis-cati</i> Benth.		<i>Z. jujuba</i> Lam.
<i>P. dulce</i> Benth.		Rhizophoraceae
<i>Prosopis juliflora</i> DC.		<i>Cassipourea elliptica</i> Poir.
Moraceae		<i>Rhizophora mangle</i> L.
<i>Artocarpus incisa</i> L.		Rosaceae
<i>Artocarpus integrifolia</i> L.		<i>Chrysobalanus icaco</i> L.
<i>Brosimum alicastrum</i> Sw.		<i>Prunus myrtifolia</i> Urb.
<i>Cecropia peltata</i> L.		<i>P. occidentalis</i> Sw.
<i>Chlorophora tinctoria</i> Gaudich.		Rubiaceae
<i>Ficus aurea</i> Nutt.		<i>Antirrhoea jamaicensis</i> (Griseb.) Urb.
<i>F. harrisi</i> Warb.		<i>Borreria laevis</i> (Lam.) Griseb.
<i>F. mammillifera</i> Warb.		<i>Catesbeia parviflora</i> Sw.
<i>F. morantensis</i> Britton		<i>Cephaelis elata</i> Sw.
<i>F. populnea</i> Willd.		<i>Chiococca alba</i> (L.) Hitchc.
<i>F. suffocans</i> Griseb.		<i>Coccocypselum herbaceum</i> Aubl.
<i>F. wilsonii</i> Warb.		<i>Erthalis fruticosa</i> L.
<i>Trophis racemosa</i> Urb.		<i>Ernodea littoralis</i> Sw.
Moringaceae		<i>Exostema caribaeum</i> (Jacq.) Roem.
<i>Moringa oleifera</i> Lam.		& Schult.
Myricaceae		<i>Faramea occidentalis</i> (Jacq.) A. Rich.
<i>Myrica microcarpa</i> Benth.		<i>Guettarda argentea</i> Lam.
Myrsinaceae		<i>G. elliptica</i> Sw.
<i>Rapanea ferruginea</i> R. & P.		<i>G. longiflora</i> Griseb.
Myrtaceae		<i>Machaonia rotundata</i> Griseb.
<i>Calyptanthes chytraculia</i> Sw.		<i>Manettia lygistum</i> (L.) Sw.
<i>Calyptanthes pallens</i> Griseb.		<i>Morinda citrifolia</i> L.
<i>Eugenia alpina</i> Willd.		<i>M. roya</i> L.
<i>E. axillaris</i> Willd.		<i>Palicourea crocea</i> (Sw.) Roem. &
<i>E. biflora</i> DC.		Schult.
<i>E. buxifolia</i> Willd.		<i>Portlandia grandiflora</i> L.
<i>E. disticha</i> DC.		<i>P. latifolia</i> Britt. & Harris.
<i>E. fragrans</i> Willd.		<i>Psychotria balbisiana</i> DC.
<i>E. jambos</i> L.		<i>P. brachiata</i> Sw.
<i>E. malaccensis</i> L.		<i>P. brownei</i> Spreng.
<i>E. marchiana</i> Griseb.		<i>P. corymbosa</i> Sw.
<i>E. monticola</i> DC.		<i>P. nervosa</i> Sw.
<i>E. rhombea</i> Kr. & Urb.		<i>P. pedunculata</i> Sw.
<i>Pimenta officinalis</i> Lindl.		<i>Randia aculeata</i> L.
<i>Psidium albescens</i> Urb.		<i>Relbunium hypocarpium</i> (L.) Hemsl.
		<i>Rondeletia hirta</i> Sw.
		<i>R. tomentosa</i> Sw.

- R. trifolia* Jacq.
Schraderia involucrata (Sw.) Schum
 Rutaceae
Amyris balsamifera L.
Amyris elemifera L.
Esenbeckia pentaphylla Griseb.
Spathelia glabrescens Planch.
Spathelia sorbifolia L.
Zanthoxylum fagara Sarg.
Zanthoxylum flavum Vahl
Zanthoxylum martinicense DC.
Zanthoxylum spinosum Sw.
 Sapindaceae
Allophylus cominia Sw.
A. jamaicensis Radlk.
A. pachyphyllus Radlk.
Cardiospermum grandiflorum Sw.
Dodonaea viscosa Jacq.
Exothea paniculata Radlk.
Hypelate trifoliata Sw.
Malayba apetala Radlk.
Melicocca bijuga L.
Paullinia barbadensis Jacq.
Sapindus saponaria L.
Serjania laevigata Radlk.
 Sapotaceae
Achras sapota L. (= *Sapota sideroxyton* Griseb.)
Bumelia retusa Sw.
B. rotundifolia Sw.
Chrysophyllum cainito L.
C. oliviforme Lam.
Dipholis montana Griseb.
D. nigra Griseb.
D. salicifolia A. DC.
Lucuma mammosa Gaertn.
Mimusops excisa Urb.
Sideroxylon foetidissimum Jacq.
 Serophulariaceae
Maurandia erubescens (Don) A. Gray
 Simarubaceae
Castela macrophylla Urb.
Picraena excelsa Lindl.
Picramnia antidesma Sw.
Simaruba glauca DC.
Suriana maritima L.
 Solanaceae
Aenistus arborescens Schlecht.
Capiscum frutescens L.
Cestrum diurnum L.
C. hirtum Sw.
Datura stramonium L.
D. suaveolens Humb. & Bonpl.
Solandra grandiflora Sw.
Solanum havanense Jacq.
S. punctulatum Dun.
S. torvum Sw.
S. verbascifolium L.
 Staphylaceae
Turpinia occidentalis G. Don
 Sterculiaceae
Ayenia laevigata Sw.
Cola acuminata Schott & Endl.
Guazuma ulmifolia Lam.
Helicteres jamaicensis Jacq.
Melochia crenata Vahl
 K. *M. lupulina* Sw.
M. tormentosa L.
Theobroma cacao L.
Waltheria americana L.
 Ternstroemiacae
Eroteum theoides Sw.
Laplacea haematoxylon G. Don
 Thymelaeaceae
Daphnopsis occidentalis Kr. & Urb.
Lagetta lagetto Nash
 Tiliaceae
Corchorus siliquosus L.
Sloanea jamaicensis Hook.
 Turneraceae
Turnera ulmifolia L.
 Ulmaceae
Celtis trinervia Lam.
Trema micrantha Bl.
 Umbelliferae
Hydrocotyle umbellata L.
 Urticaceae
Boehmeria caudata Sw.
B. jamaicensis Urb.
Gyrotienia microcarpa Fawc. & Rendle
G. spicata Wedd.
Pilea ciliata Blume
P. crassifolia Blume
P. reticulata Wedd.
P. microphylla Liebm.
P. parietaria Griseb.
 Vacciniaceae
Vaccinium meridionale Sw.
 Verbenaceae
Aegiphila elata Sw.
Avicennia nitida Jacq.
Citharexylum caudatum L.
C. fruticosum L.
Duranta plumieri Jacq.
Lantana crocea Jacq.
L. involucrata L.
Lippia reptans H. B. K.
Priva echinata Juss.
Stachytarpheta indica Vahl
Vitex umbrosa Sw.
 Vitaceae
Cissus cucurbitacea Britt.
C. sicyoides L.
 Zygophyllaceae
Guaicum officinale L.
Tribulus cistoides L.
 MONOCOTYLEDONS
 Amaryllidaceae
Agave americana L.
Hymerocallis speciosa Salisb.
 Araceae
Philodendron spp.
Pistia stratiotes L.
 Bromeliaceae
Aechmea paniculigera Griseb.
Bromelia pinguin L.
Hohenbergia distans Griseb.
H. erythrostachya Brongn.
Thecophyllum sintenisii (Baker) Mez.
Tillandsia balbisiana Schult.
T. incurva Griseb.
T. recurvata L.
T. usneoides L.
 Cyperaceae
Cyperus brunneus Sw.
C. giganteus Vahl
C. retundus L.
Timbristylis ferruginea (L.) Vahl
Mariscus jamaicensis Britton
Rhynchospora eggersiana Boekl.
Uncinia hamata (Sw.) Urb.
 Dioscoraceae
Dioscorea polygonoides H. & B.
Rajania caudata L.
 Gramineae
Andropogon pertusus (L.) Willd.
A. virginicus L.
Arundo donax L.
Cenchrus pauciflorus Benth.
C. tribuloides L.
C. echinatus L.
Chloris petraea Swartz
Chusquea abietifolia Griseb.
Cynodon dactylon (L.) Pers.
Danthonia schrebei Britton
Eragrostis ciliaris (L.) Link
Lasiacis divaricata (L.) Hitchc.
Manisuris altissima (Poir) Hitchc.
Melinis minutiflora Beauv.
Panicum maximum Jacq.
P. palmifolium Willd.
P. purpurascens Raddi.
Paspalum fimbriatum H. B. K.
P. saccharoides Nees
P. vaginatum Swartz
Phragmites communis Trin.
Spartina patens var. *junccea* H.
 Hitchc.
Sporobolus indicus (L.) R. Br.
S. virginicus (L.) Kunth
Uniola virgata (Poir) Griseb.
 Hydrocharidaceae
Thalassia testudinum Konig.
 Lemnaceae
Lemna minor L.
 Liliaceae
Smilax balbisiana Kunth
S. domingensis Willd.
 Musaceae
Heliconia caribaea Lam.
 Orchidaceae
Bletia purpurea (Lam.) DC.
Broughtonia sanguinea (Sw.) R. Br.
Dichaea glauca Lindl.
D. graminea Griseb.
Elleanthus capitatus Rehb.
Epidendrum fragrans Sw.
E. verrucosum Sw.
Lepanthes concinna Sw.
L. concolor Fawc. & Rendle
L. tridentata Sw.
Liparis elata Lindl.
Stelis ophioglossoides Sw.

Palmaceae	<i>Cochlidium graminoides</i> (Sw.) Kaulf (<i>C. seminudatum</i> (Willd.) Maxon)	<i>Lomariopsis underwoodii</i> Holttum
<i>Acrocomia aculeata</i> (Jacq.) Lodd.	<i>Ctenitis ampla</i> (H. B. W.) Copel.	<i>Lophosoria quadripinnata</i> (Gmel) C. Chr.
<i>Calyptronoma swartzii</i> Griseb.	<i>C. effusa</i> (Sw.) Copel.	<i>Lycopodium cernuum</i> L.
<i>Coccothrinax fragrans</i> Burret.	<i>Ctenis villosa</i> (L.) Copel.	<i>L. fawcettii</i> Lloyd & Underw.
<i>Cocos nucifera</i> L.	<i>Cyathea armata</i> (Sw.) Domin.	<i>L. wilsonii</i> Underw. & Lloyd
<i>Roystonea princeps</i> (Becc.) Burret.	<i>C. aspera</i> (L.) Sw.	<i>L. clavatum</i> L.
<i>Sabal jamaicensis</i> Becc.	<i>C. gracilis</i> Griseb.	<i>Lygodium volvabile</i> Sw.
<i>Thrinax excelsa</i> Lodd.	<i>C. tussacii</i> Desv.	<i>Marattia alata</i> Sw.
<i>T. parviflora</i> Sw.	<i>C. nigrescens</i> (Hook) J. Sm.	<i>Nephrolepis pectinata</i> (Willd.) Shott.
<i>T. tesselata</i> Becc.	<i>C. pubescens</i> Mett. ex. Kuhn.	<i>N. rivularis</i> (Vahl) Mett.
Pontederiaceae	<i>C. grevilleana</i> Mart.	<i>Notholaena trichomanoides</i> (syn. <i>Cheilanthes trichomanoides</i> (L.) Mett.)
<i>Eichhornia crassipes</i> Solms.	<i>Cystopteris fragilis</i> (L.) Bernh.	<i>Orthopteris domingensis</i> Spreng.
Potamogetonaceae	<i>Danaea jamaicensis</i> Underw.	<i>Paesia viscosa</i> St. Hil.
<i>Cymodocea manatorum</i> Aschers	<i>D. jenmani</i> Underw.	<i>Polybotrya cervina</i> (L.) Kaulf.
Typhaceae	<i>D. nodosa</i> (L.) J. E. Smith.	<i>Polypodium angustifolium</i> Sw.
<i>Typha angustifolia</i> L.	<i>Dennstaedtia bipinnata</i> (Cav.) Maxon	<i>P. dissimile</i> L.
GYMNOSPERMAE	<i>D. cicutaria</i> (Sw.) Moore	<i>P. exornans</i> Maxon
Pinaceae	<i>D. globulifera</i> (Poir.) Hieron	<i>P. hartii</i> (syn. <i>Grammitis hartii</i> (Jenn.) Proctor)
<i>Juniperus barbadensis</i> L.	<i>Diplazium brunneo-viride</i> (Jenm.) C. Chr.	<i>P. heterophyllum</i> L.
Taxaceae	<i>D. centripetale</i> (Baker) Maxon	<i>P. jubaeforme</i> Kaulf.
<i>Podocarpus purdianus</i> Hook.	<i>D. costale</i> (Sw.) Presl	<i>P. loriceum</i> L.
<i>P. urbani</i> Pilger (<i>P. coriaceus</i> Rich.)	<i>D. fadyeni</i> (Hook) Proctor	<i>P. lycopodioides</i> L.
PTERIDOPHYTA	<i>D. pectinatum</i> (Fee) C. Chr.	<i>P. mollissimum</i> Fee
<i>Acrostichum aureum</i> L.	<i>Doryopteris pedata</i> (L.) Fee	<i>P. thyrsanolepis</i> A. Br. ex. Klotzsch.
<i>Adiantum melanoleucum</i> Willd.	<i>D. decussata</i> (L.) Proctor	<i>Polystichum christiana</i> (Jenm.) Underw. & Maxon
<i>A. tenerum</i> Sw.	<i>D. deltoidea</i> (syn. <i>Thelypteris deltoidea</i> (Sw.) Proctor)	<i>Pteridium arachnoideum</i> (Kaulf.) Maxon
<i>Alsophila aspera</i> R. Br. (<i>Cyathea aspera</i> Sw.)	<i>D. denticulata</i> (Sw.) Ktze.	<i>Pteris hexagona</i> (L.) Proctor
<i>A. swartziana</i> Mart. (<i>C. armata</i> Sw.)	<i>D. gemmipara</i> (C. Chr.) Maxon	<i>P. longifolia</i> Proctor
<i>Anemia hirsuta</i> (L.) Sw.	<i>D. heteroclita</i> (syn. <i>Thelypteris heteroclita</i> (Desv.) Proctor)	<i>P. quadriaurita</i> Retz.
<i>A. underwoodiana</i> Maxon	<i>D. decussata</i> (L.) Proctor	<i>Rhipidopteris peltata</i> (Sw.) Schott.
<i>A. adiantifolia</i> (L.) Sw.	<i>D. dryopteris</i> (syn. <i>Thelypteris dryopteris</i> resinifera (Desv.) Proctor)	<i>Tectaria incisa</i> Cav.
<i>Asplenium alatum</i> Humb. & Bonpl.	<i>D. resinifera</i> (Desv.) Proctor	<i>Thelypteris asterothrix</i> (Fee) Proctor
<i>A. cuneatum</i> Lam.	<i>Elaphoglossum eggersii</i> (Baker) Christ.	<i>T. decussata</i> (L.) Proctor
<i>A. harpeodes</i> Kunze	<i>E. petiolatum</i> (Sw.) Urban	<i>T. deltoidea</i> (Sw.) Proctor
<i>A. praemorsum</i> Sw.	<i>E. simplex</i> (Sw.) Schott.	<i>T. heteroclita</i> (Desv.) Proctor
<i>A. pumilum</i> Sw.	<i>E. villosum</i> (Sw.) J. Sm.	<i>T. oligophylla</i> (Maxon) Proctor
<i>A. radicans</i> L.	<i>Gleichenia jamaicensis</i> (Underw.) Proctor	<i>T. patens</i> (Sw.) Small
<i>Azolla caroliniana</i> Willd.	<i>G. pectinata</i> (Willd.) Presl	<i>T. sagittata</i> (Sw.) Proctor
<i>Blechnum jamaicense</i> (Broadb.) C. Chr.	<i>Grammitis hartii</i> (Jenm.) Proctor	<i>T. serrulata</i> (Sw.) Proctor
<i>B. lineatum</i> (Sw.) C. Chr.	<i>G. serrulata</i> (Sw.) Sw.	<i>T. venusta</i> (Hew) Proctor
<i>B. polypodioides</i> (Sw.) Kuhn	<i>G. trifurcata</i> (L.) Copel.	<i>Trichomanes crinitum</i> Sw.
<i>B. unilaterale</i> Sw.	<i>Gymnopteris rufa</i> (L.) Bernh.	<i>T. crispum</i> L.
<i>Bolbitis aliena</i> (Sw.) Alston	<i>Histiopteris incisa</i> (Thunb.) J. Sm.	<i>T. holopterum</i> Kunze
<i>B. nicotianaefolia</i> (Sw.) Alston	<i>Hymenophyllum fucoides</i> Sw.	<i>T. hymenophylloides</i> v. d. B.
<i>B. pergamantacea</i> (Maxon) Ching.	<i>H. hirsutum</i> (L.) Sw.	<i>T. osmundoides</i> DC. ex. Poir.
<i>Camptodium pedatum</i> (Desv.) Fee	<i>H. hirtellum</i> Sw.	<i>T. rigidum</i> Sw.
<i>Campyloneuron angustifolium</i> (syn. <i>Polypodium angustifolium</i> Sw.)	<i>H. lanatum</i> Fee	<i>T. scandens</i> L.
<i>Cheilanthes microphylla</i> Sw.	<i>H. polyanthos</i> (Sw.) Sw.	<i>Trismeria trifoliata</i> (L.) Diels
<i>C. trichomanoides</i> (L.) Mett.	<i>H. sericeum</i> (Sw.) Sw.	

