

LETTER • OPEN ACCESS

## Feedback between drought and deforestation in the Amazon

To cite this article: Arie Staal *et al* 2020 *Environ. Res. Lett.* **15** 044024

View the [article online](#) for updates and enhancements.

### Recent citations

- [Tracking the global flows of atmospheric moisture and associated uncertainties](#)  
Obbe A. Tuinenburg and Arie Staal



## LETTER

## Feedback between drought and deforestation in the Amazon

## OPEN ACCESS

RECEIVED  
24 June 2019

REVISED  
17 December 2019

ACCEPTED FOR PUBLICATION  
5 February 2020

PUBLISHED  
2 April 2020

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Arie Staal<sup>1,2,7</sup> , Bernardo M Flores<sup>3</sup> , Ana Paula D Aguiar<sup>1,4</sup>, Joyce H C Bosmans<sup>5</sup>, Ingo Fetzer<sup>1</sup> and Obbe A Tuinenburg<sup>6</sup>

<sup>1</sup> Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden

<sup>2</sup> Aquatic Ecology and Water Quality Management Group, Wageningen University, Wageningen, The Netherlands

<sup>3</sup> Department of Plant Biology, University of Campinas, Campinas, Brazil

<sup>4</sup> Earth System Science Centre, National Institute for Space Research (INPE), São José dos Campos, Brazil

<sup>5</sup> Department of Environmental Science, Radboud University, Nijmegen, The Netherlands

<sup>6</sup> Department of Environmental Science, Copernicus Institute for Sustainable Development, Utrecht University, Utrecht, The Netherlands

<sup>7</sup> Author to whom any correspondence should be addressed.

E-mail: [ariestaal@gmail.com](mailto:ariestaal@gmail.com)

**Keywords:** land use change, remote sensing, moisture recycling, fire, modeling, forest clearing, agriculture

Supplementary material for this article is available [online](#)


### Abstract

Deforestation and drought are among the greatest environmental pressures on the Amazon rainforest, possibly destabilizing the forest-climate system. Deforestation in the Amazon reduces rainfall regionally, while this deforestation itself has been reported to be facilitated by droughts. Here we quantify the interactions between drought and deforestation spatially across the Amazon during the early 21st century. First, we relate observed fluctuations in deforestation rates to dry-season intensity; second, we determine the effect of conversion of forest to cropland on evapotranspiration; and third, we simulate the subsequent downwind reductions in rainfall due to decreased atmospheric water input. We find large variability in the response of deforestation to dry-season intensity, with a significant but small average increase in deforestation rates with a more intense dry season: with every mm of water deficit, deforestation tends to increase by 0.13% per year. Deforestation, in turn, has caused an estimated 4% of the recent observed drying, with the south-western part of the Amazon being most strongly affected. Combining both effects, we quantify a reinforcing drought-deforestation feedback that is currently small, but becomes gradually stronger with cumulative deforestation. Our results suggest that global climate change, not deforestation, is the main driver of recent drying in the Amazon. However, a feedback between drought and deforestation implies that increases in either of them will impede efforts to curb both.

Deforestation, the human-driven cropping of tree cover, has been gradually decreasing the extent of the Amazon rainforest over the last decades, mainly for the expansion of pastures and soybean plantations [1–3]. The spatial and temporal distribution of deforestation is not random, as a range of socioeconomic and institutional factors affect deforestation. For example, population density and accessibility by road contribute to deforestation [4, 5]. In contrast, designating forest as protected area successfully inhibits deforestation [6–8]. Sometimes, cause-and-effect relations are more complex, as when positive feedbacks are in play. An example of such an amplifying causal loop is a two-way interaction between road density

and deforestation: accessibility of the forest increases deforestation, but that deforestation is in turn also used as justification for constructing more roads [9]. Besides being a product of socioeconomic complexity, deforestation also interacts with climatic and ecological processes in the Amazon, e.g. [10–14]. However, not all causal pathways have been studied so far.

In the Amazon, deforestation is often accompanied by fire [15, 16]. Because a more intense dry season makes the Amazon forest more flammable [4, 17–21], it seems reasonable to hypothesize that increasing dry-season intensity facilitates clear-cut deforestation in several ways: (1) traditional slash-and-burn agriculture depends on a sufficiently dry



season [15, 17]; (2) pastures are often repeatedly cleared of encroaching woody vegetation and weeds with the use of fire [15, 22]; and (3) deforestation using mechanical methods is often followed by the combustion of the remaining vegetation, which is easier when the material is drier [15, 17, 23, 24]. Furthermore, dry seasons facilitate the escape of fires into neighboring forest areas, especially in fragmented landscapes [25], potentially making them more attractive for subsequent deforestation. While dry seasons may facilitate the deforestation process, deforestation in turn affects the regional water cycle [26–29]. Evapotranspiration from forests is higher than from other land covers, in part due to enhanced evaporation directly from leaves (interception evaporation) and in part due to transpiration, in which trees pump water from the soil and release it through their leaves during photosynthesis [30]. Thus, forests maintain high evapotranspiration in the dry season by transpiring water that is stored in deeper soil layers, which act as a buffer during droughts. Part of this water subsequently precipitates over the forest, alleviating the intensity of the dry season by disproportionately contributing to rainfall under drier conditions [29]. Taking a viewpoint at the Amazonian scale, we see a positive feedback emerging: as deforestation reduces forest area, less water can be recycled and dry seasons intensify regionally; the more intense dry seasons become, the more deforestation tends to occur. These dynamics may amplify regional-scale deforestation and in principle contribute to a self-propagating loop of forest loss [31]. However, the extent to which dry seasons and deforestation affect one another remains unclear [32] and a hypothetical drought-deforestation feedback has never been explicitly addressed. Therefore, we here integrate several state-of-the-art approaches to disentangle the causal interactions between deforestation and dry-season intensity, and analyze the ‘drought-deforestation feedback’ in the Amazon rainforest.

To estimate how deforestation has interacted with regional rainfall patterns, we relate remotely sensed time series of forest cover change [33] to changing

dry-season intensity during the early 21st century. We combine these results with: (1) a hydrological model that estimates forest evapotranspiration relative to that of cropland on a monthly basis [34, 35]; and (2) a high-resolution atmospheric moisture tracking algorithm to determine the fate of that evapotranspiration [29, 36]. In addition, we account for multiple evapotranspiration-rainfall cycles of water (‘cascading moisture recycling’). This allows us to quantify the strength of the drought-deforestation feedback, analyze how it has changed over the course of more than a decade, and discuss its implications for the stability of the Amazon rainforest.

## Methods

We quantify two causal effects: the effect of drought on deforestation and the effect of deforestation on drought. Below we outline how each was estimated, followed by a description of how we used those results to calculate the feedback strength in the Amazon forest. A diagram of the feedback loop, of which we quantify all steps for the Amazon, is given in figure 1.

### The effect of drought on deforestation

We define drought as the intensity of a dry season in a given calendar year as given by the Maximum Climatological Water Deficit (MCWD in mm). The independent calculation of MCWD for each calendar year means we do not account for possible continuation of droughts into a new calendar year. MCWD is a common measure of drought in the Amazon [37, 38], capturing the cumulative difference between precipitation  $P$  and evapotranspiration  $E$  in a certain year. MCWD values are negative, so a more negative value of MCWD means a more intense dry season. Monthly precipitation and actual evapotranspiration data were taken from the GLDAS2 dataset [39].

Deforestation estimates were obtained from the Landsat satellite dataset of [33]. This dataset provides forest loss and forest gain data on  $0.00025^\circ$  ( $\sim 25$  m) resolution. Forest loss is provided on an annual basis,

whereas forest gain is provided only once for the entire time period of 2000 through 2017. If a cell has become forested during this period, we assumed that the increase in tree cover is linearly distributed over the years. Forest degradation by understory fires or logging, for instance, is not detected by the algorithm behind the dataset, so our analysis accounts for clear-cut deforestation only. All forest loss and gain data were aggregated to net deforestation (forest loss minus gain) on a  $0.25^\circ$  basis to match the resolution of the hydrological simulations (see section *The effect of deforestation on drought*).

We related dry-season intensity to deforestation to better understand how the former affects the latter. However, correlation analysis can be problematic, because the relation between deforestation and dry-season intensity may be confounded by other spatially non-random factors. For example, deforestation is mostly concentrated in the drier southern and eastern parts of the Amazon, along the ‘arc of deforestation’. This region is not only naturally drier than the central-western parts of the Amazon, but it also holds more infrastructure. To correct for such geographical factors, we determined the effect of dry-season intensity (MCWD in mm) on deforestation on a per-cell basis over time: for this we spatially averaged the climatic and deforestation data to  $1^\circ$  and linearly regressed the annual local deforestation anomaly (in % of the local multi-year average) to MCWD. For each cell, this resulted in the slope of the effect of MCWD on deforestation, or  $\Delta\text{Deforestation}/\Delta\text{MCWD}$  in  $\% \text{ yr}^{-1} \text{ mm}^{-1}$ . Because we found that the effect of MCWD on  $\Delta\text{Deforestation}/\Delta\text{MCWD}$  is small (figure S5), we simply took its average across MCWD levels as the effect of dry-season intensity on deforestation.

To check for robustness of our results we repeated the above analysis using the *Projeto de Monitoramento do Desmatamento na Amazônia Legal por Satélite* (PRODES) dataset from the Brazilian *Instituto Nacional de Pesquisas Espaciais* (INPE). However, because this dataset is only available for the Brazilian Amazon, it underestimates total deforestation. Therefore, we present the data from Landsat for our main analyses and a comparison with PRODES in the Supplementary Information (available online at [stacks.iop.org/ERL/15/044024/mmedia](https://stacks.iop.org/ERL/15/044024/mmedia)).

### The effect of deforestation on drought

To estimate the effect of deforestation on drought we combined output of the PC Raster Global Water Balance hydrological model (PCR-GLOBWB) [34, 35] with an atmospheric moisture tracking algorithm [36] forced with atmospheric and surface flux data. Below we outline how the effects of deforestation on rainfall were calculated. We provide a simple schematic of the methodological steps as figure S7, and for the equations of the moisture tracking algorithm we refer to [29].

We estimated the monthly forest contributions to evapotranspiration during 2003–2014 using the PCR-GLOBWB model at  $0.5^\circ$  resolution. The model estimates the partitioning of evapotranspiration into transpiration, interception evaporation, and remaining bare-soil evaporation for a number of possible land cover types, with a per-cell and monthly output [34, 40]. We calculated the actual forest contribution to evapotranspiration by multiplying its transpiration and interception-evaporation estimates for full forest cover by the fractions of the cells that are actually covered by forest [33]. We then replaced these forest evapotranspiration fluxes by those for rainfed cropland (such as soybean plantations) [34], maximized at the original fluxes. The output of PCR-GLOBWB shows good agreement with independent dry-season evapotranspiration estimates [29] and discharge data [41] from the Amazon river. We show the differences in evapotranspiration between forest and rainfed cropland on a monthly basis in the supplementary material (figure S6).

To estimate the regional effects of evapotranspiration reductions on rainfall due to deforestation, we used a Lagrangian atmospheric moisture tracking model [36]. This model tracks precipitation backwards in time by separating the precipitation into a large number of moisture parcels. These are initially released within cells of  $0.25^\circ$  at random heights in the atmospheric column with a probability scaled with the humidity profile. The trajectories of the parcels are simulated back in time using three-dimensional estimates of wind speed and direction using linearly interpolated ERA-Interim data at  $0.75^\circ$  and 6 h resolution [42]. This is done for each time step of fifteen minutes throughout the study period of 2003–2014. Each time step, water in the parcels that are tracked are updated using interpolated three-hourly evapotranspiration (ET) and precipitation (P) estimates from the GLDAS2 dataset [39] and precipitable water (PW) from ERA-Interim. Every time step, the fraction ET/PW of the moisture in the parcel is assumed to have originated from evapotranspiration at that location. That amount of moisture is allocated there and the moisture present in the parcel is updated. Thus, the amount of tracked moisture in the parcels decreases along their trajectories backward in time from precipitation to evaporation. Parcels of water were tracked either until more than 95% of it had been allocated, 30 d have passed since tracking started, or it has left the study domain of tropical South America ( $81.5^\circ \text{ W}$ – $34^\circ \text{ W}$ ;  $13^\circ \text{ N}$ – $35^\circ \text{ S}$ ). For precipitation in each cell in tropical South America, we determined the corresponding upwind evaporation location on a monthly basis. Thus, we obtained the monthly moisture flows between each pair of  $0.25^\circ$  cells in tropical South America, from which we used those located in the Amazon. By multiplying these moisture flows with the fractional change in evapotranspiration due to deforestation, we estimated the effects of deforestation on

monthly precipitation throughout the Amazon. Implicitly, we assumed that deforestation does not affect wind patterns.

Upon precipitating, moisture can re-evaporate and precipitate again in what has been called ‘cascading moisture recycling’ [43]. Thus, evapotranspiration reduction by deforestation could affect precipitation multiple times. Reference [29] determined that in the Amazon forest this happens up to six times. This cascading moisture recycling accounts for about half of the forest’s contribution to rainfall. Therefore, we tracked the atmospheric trajectories of re-evapotranspired moisture for seven re-evaporation times according to the method of [43]. Although we only present moisture circulation within the Amazon, the calculations were done for the entire tropical South America. We updated our estimates of the deforestation effects on rainfall with this cascading moisture recycling. We show the deforestation-induced changes in MCWD by 2014 and smoothed them spatially using a Gaussian filter with a standard deviation of  $0.5^\circ$ . We also mapped the trend in MCWD across 2000–2014 and subtracted from that the annual contribution of deforestation (the total by 2014, divided by 14) and smoothed it likewise.

### Quantifying the feedback

With the above described method we estimated how deforestation during 2003–2014 has affected the frequency distribution of MCWD. By multiplying the estimated frequency distribution of MCWD in the absence of deforestation by the observed effect of MCWD on deforestation we could estimate deforestation in the absence of a drought-deforestation feedback. The feedback strength, measured as the ‘drought-deforestation feedback factor’  $F_{DD}$ , represents the factor by which deforestation is multiplied due to the feedback. Hence, it is the ratio of observed deforestation  $D_{obs}$  over estimated deforestation in the absence of the drought-deforestation feedback  $D_{est}$ :

$$F_{DD} = \frac{D_{obs}}{D_{est}}.$$



As in [29], due to limited availability of data that are consistent across the different models that we used, we only analyze the period between 2003 and 2014: atmospheric data pre-2003 have discrepancies with those from 2003 onward due to differences in the data sources [42], and the input data for PCR-GLOBWB are available until 2014 [34]. However, we could estimate how the feedback strength has increased with cumulative deforestation during the early 21st century to obtain the effect of historical deforestation on the feedback strength. We took the estimate that goes furthest back in time [44], which is 1960, although it only accounts for the Brazilian Amazon.

## Results

Our hydrological simulations show that deforestation, which has predominantly occurred in the south-eastern part of the Amazon (figure 2(A)), has made dry seasons more intense over the early 21st century (figure 2(B)). By 2014, the mean decrease in Maximum Climatological Water Deficit (MCWD in mm) across the Amazon in response to 21st century deforestation was *c.* 1.6 mm (figure S8), or a decrease of  $0.11 \text{ mm yr}^{-1}$  (linear  $R^2 = 0.87$ ). This corresponds to 3.8% of the average decrease in MCWD. We postulate that the remainder of the decrease (figure 2(C)) is due to global climate change, although natural fluctuations such as the El Niño Southern Oscillation [45], or other atmospheric effects of deforestation such as changed circulation [46] may have contributed as well. Although the largest drying trends have occurred in the central and western parts of the Amazon (figures 2(C) and S1), deforestation has intensified dry seasons especially in the south-western Amazon ( $\sim 4 \text{ mm}$ ; figure 2(B)). This demonstrates the international effects of deforestation on rainfall [47]: deforestation in the Brazilian Amazon causes part of the drying in Bolivia and Peru (figure S10).


Looking on a per-cell basis, we find that, on average, deforestation increases in years with a more intense dry season: the sensitivity of deforestation to MCWD is significantly below zero with a mean of  $-0.13\% \text{ yr}^{-1} \text{ mm}^{-1}$  (95% CI  $[-0.14, -0.12]$ ), which is the annual deforestation change for each mm change of water deficit ( $n = 14$  for each cell). Note that, because MCWD is negative, a negative sensitivity implies that deforestation increases as conditions become drier. There are, however, large spatial differences in the sensitivity of deforestation to drought ( $\sigma = 0.30$ ). Especially in the central Amazon we find locations with increases in deforestation with drought (figure 2(D)). In 69% of the Amazon (regardless of *p*-value), deforestation increases with a more intense dry season (see figure S2 for examples). When areas with a non-significant ( $\alpha = 0.05$ ) effect of drought are excluded, this proportion rises to 80% (figure S3). In other words, in four out of five significant effects, deforestation increased with drought. Among the remaining areas, we find a number of cells with significant effects of drought on deforestation near major roads in a new deforestation frontier in the western Amazonia of Brazil (figure S4).

Although our study period covers only a part of the historical deforestation, we are able to estimate the increase in feedback strength within that period, showing that the feedback has gained strength by an average of  $5.1 \times 10^{-4}$  per year compared to the feedback strength in 2000 (linear  $R^2 = 0.75$ ,  $p \approx 0$ ; figure S9). Put otherwise, for every 1000  $\text{km}^2$  that was deforested, the feedback increased on average by  $2.7 \times 10^{-5}$  (linear  $R^2 = 0.69$ ,  $p \approx 0$ ; figure 3).



Aside from analyzing drought-deforestation interaction during the recent past, we can assess where potential future deforestation would intensify dry

seasons. We therefore calculate the effects of conversion from forest to cropland on the amount of evapotranspiration that directly precipitates in the Amazon



within a dry season. On average, converting one hectare of forest to cropland would cause a reduction of 0.5 million L per year of evaporated water that subsequently precipitates within the Amazon during a dry season (i.e. during months when evapotranspiration exceeds precipitation). Locally, this amount ranges between 0 and 2 million  $\text{L ha}^{-1} \text{ yr}^{-1}$  (figure 4).

## Discussion

### Drought-deforestation feedback: weak, but getting stronger

We analyzed the interactions between deforestation and dry-season intensity in the Amazon, suggesting a drought-deforestation feedback. The strength of this feedback is a function of the cumulative historical forest loss (deforestation minus reforestation). Although the feedback is still small and the effect of drought on deforestation is highly variable, the drought-deforestation feedback may be a previously unidentified hidden driver of deforestation which is becoming stronger as progressively more forest is lost every year. Historical deforestation has a perpetual legacy effect in increasing dry-season intensity, particularly in the south-western Amazon, presumably because using fire to clear the landscape has become easier [17, 19, 49]. Taking the relation between feedback strength and cumulative deforestation, we can estimate to what extent historical deforestation amplifies current deforestation. It is estimated that between 1960 and 2014 in the Brazilian Amazon alone, cumulative deforestation amounts to  $649\,000 \text{ km}^2$  [44]. If the correlation between deforestation and drought represents a causal effect, then by 2014, in the order of 1% of annual deforestation may have taken

place due to the drought-deforestation feedback (best estimate is 1.7%, or around  $300 \text{ km}^2$  for 2014). It should be noted that this estimate depends on uncertain effects of drought on deforestation and a short period of twelve years on which our feedback analysis is based. Also, in assuming a linear increase in feedback strength with cumulative deforestation we adopted a conservative approach. Longer time series and improved satellite monitoring can reduce these uncertainties.

### The roles of climate change and fire

Climate change intervenes with the drought-deforestation feedback by increasing the duration, intensity, and frequency of droughts [32, 50–52]. Given our finding that no more than 4% of recent drying in the Amazon has resulted from deforestation, it appears that climate change is the main driver of drought-facilitated forest loss. Intensified dry seasons in the Amazon increase the opportunities for wildfires to spread from human-ignited sources into standing forest. Wildfires not only degrade a forest, but may destabilize the remaining forest cover through flammability and erosion feedbacks favouring open vegetation [53–56]. There could be a critical threshold of drying above which forest loss self-perpetuates even in the absence of further deforestation [31, 57].

### The role of governance

In 2019, the rates of deforestation and fire occurrence in Brazil were unusually high for a year with normal rainfall levels [15], whereas during most of our study period, those rates were gradually declining due to strong forest governance [58]. This illustrates how the effects of drought on deforestation can be

overwhelmed by changes in governance. Raising societal awareness of the drought-deforestation feedback that we have identified could avoid runaway forest loss. Despite the legacy effect of past deforestation, the feedback decreases in importance with reduced deforestation and ceases to affect the system when deforestation is halted altogether. Governance interventions, involving (inter)national political [59] and financial actors [60], as well as local populations, could realize this [61]. Special attention should be given to Amazonian regions where the forest contributes the most to dry-season rainfall recycling, such as the Brazilian states of Acre, Amapá and Pará, as well as parts of Peru and French Guiana (figure 4).

### Limitations

Our results are limited to the observed spatial distribution of dry season intensity across the Amazon. In areas where droughts do not occur yet, and where the absence of infrastructure limits deforestation, we do not know the sensitivity to more intense droughts, increasing the uncertainty of the drought-deforestation feedback in the relatively remote wetter regions. Furthermore, if, due to global climate change, natural fluctuations or deforestation itself, wind patterns change, then the effects of deforestation may change as well. Nevertheless, it seems very likely that the western Amazon will remain a main destination of forest-generated atmospheric moisture, given the large-scale easterly trade winds and the blockage of these winds by the Andes [62]. On smaller scales, however, presumed changes in winds are more complex [63]. We did not account for sub-grid convective cloud transport [64] while models suggest that small-scale deforestation may enhance precipitation locally [65]. This occurs especially in the dry season when the system is less dominated by strong easterly winds [63]. Thus, the effects of deforestation on drought could be scale-dependent, which is not accounted for in our simulations.

The complexity of deforestation dynamics derives from multiple nonlinear interactions between nature and society, including climatic conditions and societal rules that are continuously changing. This means that our results do not warrant prediction of future forest loss. However, they do highlight one social-ecological feedback that constitutes these dynamics. The drought-deforestation feedback has been overlooked so far, and its explicit incorporation in scenarios of future deforestation could improve our projections. As a step in the direction of disentangling the complexities of the Amazon forest, we here unveil a feedback mechanism between drought and deforestation that has been increasingly impacting the dynamics of this important ecosystem.

## Conclusion

We presented a previously unrecognized feedback between drought and deforestation in the Amazon. We analyzed the spatial patterns of the causal effects between drought and deforestation by analyzing remotely sensed forest loss data in response to dry-season intensity, and by simulating the atmospheric trajectories of forest-induced evapotranspiration. On average, deforestation becomes higher with a more intense dry season; this deforestation intensifies dry seasons in the south-western Amazon in particular. We conclude that the drought-deforestation feedback has a small, but increasing, effect on deforestation.

## Acknowledgments

We thank the two anonymous reviewers for their helpful comments. AS and IF acknowledge support from the European Research Council project Earth Resilience in the Anthropocene (743080 ERA). BMF is funded by São Paulo Research Foundation Grant FAPESP 2016/25086-3. OAT acknowledges support from the research program Innovational Research Incentives Scheme 016. veni.171.019, funded by the Netherlands Organization for Scientific Research (NWO).

## Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## ORCID iDs

Arie Staal  <https://orcid.org/0000-0001-5409-1436>

Bernardo M Flores  <https://orcid.org/0000-0003-4555-5598>

Ingo Fetzer  <https://orcid.org/0000-0001-7335-5679>

Obbe A Tuinenburg  <https://orcid.org/0000-0001-6895-0094>

## References

- [1] Curtis P G, Slay C M, Harris N L, Tyukavina A and Hansen M C 2018 Classifying drivers of global forest loss *Science* **361** 1108–11
- [2] Macedo M N, DeFries R S, Morton D C, Sticker C M, Galford G L and Shimabukuro Y E 2012 Decoupling of deforestation and soy production in the southern Amazon during the late 2000s *Proc. Natl Acad. Sci.* **109** 1341–6
- [3] Tyukavina A, Hansen M C, Potapov P V, Stehman S V, Smith-Rodriguez K, Okpa C and Aguilar R 2017 Types and rates of forest disturbance in Brazilian Legal Amazon, 2000–2013 *Sci. Adv.* **3** e1601047
- [4] Laurance W F, Albernaz A K M, Schroth G, Fearnside P M, Bergen S, Venticinque E M and Da Costa C 2002 Predictors of deforestation in the Brazilian Amazon *J. Biogeography* **29** 737–48

- [5] Aguiar A P D, Câmara G and Escada M I S 2007 Spatial statistical analysis of land-use determinants in the Brazilian Amazonia: exploring intra-regional heterogeneity *Ecol. Modell.* **209** 169–88
- [6] Nepstad D *et al* 2006 Inhibition of Amazon deforestation and fire by parks and indigenous lands *Conserv. Biol.* **20** 65–73
- [7] Oliveira P J C, Asner G P, Knapp D E, Almeyda A, Galván-Gildemeister R, Keene S, Raybin R F and Smith R C 2007 Land-use allocation protects the Peruvian Amazon *Science* **317** 1233–6
- [8] Soares-Filho B *et al* 2010 Role of Brazilian Amazon protected areas in climate change mitigation *Proc. Natl Acad. Sci.* **107** 10821–6
- [9] Fearnside P M 1987 Deforestation and international economic development projects in Brazilian Amazonia *Conserv. Biol.* **1** 214–21
- [10] Coe M T *et al* 2013 Deforestation and climate feedbacks threaten the ecological integrity of south-southeastern Amazonia *Phil. Trans. R. Soc. B* **368** 20120155
- [11] Malhi Y, Roberts J T, Betts R A, Killeen T J, Li W and Nobre C A 2008 Climate change, deforestation, and the fate of the Amazon *Science* **319** 169–72
- [12] Davidson E A *et al* 2012 The Amazon basin in transition *Nature* **481** 321–8
- [13] Laurance W F and Williamson G B 2001 Positive feedbacks among forest fragmentation, drought, and climate change in the Amazon *Conserv. Biol.* **15** 1529–35
- [14] Nepstad D C, Stickler C M, Soares-Filho B and Merry F 2008 Interactions among Amazon land use, forests and climate: prospects for a near-term forest tipping point *Phil. Trans. R. Soc. B* **363** 1737–46
- [15] Barlow J, Berenguer E, Carmenta R and França F 2020 Clarifying Amazonia's burning crisis *Glob. Change Biol.* **26** 319–21
- [16] Qin Y *et al* 2019 Improved estimates of forest cover and loss in the Brazilian Amazon in 2000–2017 *Nat. Sustain.* **2** 764–72
- [17] Aragão L E O C, Malhi Y, Barbier N, Lima A, Shimabukuro Y, Anderson L and Saatchi S 2008 Interactions between rainfall, deforestation and fires during recent years in the Brazilian Amazonia *Phil. Trans. R. Soc. B* **363** 1779–85
- [18] Alencar A A, Brando P M, Asner G P and Putz F E 2015 Landscape fragmentation, severe drought, and the new Amazon forest fire regime *Ecological Appl.* **25** 1493–505
- [19] Morton D C, Le Page Y, DeFries R, Collatz G J and Hurr G C 2013 Understorey fire frequency and the fate of burned forests in southern Amazonia *Phil. Trans. R. Soc. B* **368** 20120163
- [20] Aragão L E O C *et al* 2018 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions *Nat. Commun.* **9** 536
- [21] Chen Y, Velicogna I, Famiglietti J S and Randerson J T 2013 Satellite observations of terrestrial water storage provide early warning information about drought and fire season severity in the Amazon *J. Geophys. Res.: Biogeosciences* **118** 495–504
- [22] Nepstad D, Carvalho G, Cristina Barros A, Alencar A, Paulo Capobianco J, Bishop J, Moutinho P, Lefebvre P, Lopes Silva U and Prins E 2001 Road paving, fire regime feedbacks, and the future of Amazon forests *Forest Ecology Manage.* **154** 395–407
- [23] Lima A, Silva T S F, de Feitas R M, Adami M, Formaggio A R and Shimabukuro Y E 2012 Land use and land cover changes determine the spatial relationship between fire and deforestation in the Brazilian Amazon *Appl. Geogr.* **34** 239–46
- [24] Aguiar A P D *et al* 2012 Modeling the spatial and temporal heterogeneity of deforestation-driven carbon emissions: the INPE-EM framework applied to the Brazilian Amazon *Glob. Change Biol.* **18** 3346–66
- [25] Hissa L, de B V, Müller H, Aguiar A P D, Hostert P and Lakes T 2018 Historical carbon fluxes in the expanding deforestation frontier of southern Brazilian Amazonia (1985–2012) *Reg. Environ. Change* **18** 77–89
- [26] Spracklen D V and Garcia-Carreras L 2015 The impact of Amazonian deforestation on Amazon basin rainfall *Geophys. Res. Lett.* **42** 9546–52
- [27] Spracklen D V, Baker J C A, Garcia-Carreras L and Marsham J 2018 The effects of tropical vegetation on rainfall *Annu. Rev. Environ. Resour.* **43** 193–218
- [28] Zemp D C, Schleussner C F, Barbosa H M J and Rammig A 2017 Deforestation effects on Amazon forest resilience *Geophys. Res. Lett.* **44** 6182–90
- [29] Staal A, Tuinenburg O A, Bosmans J H C, Holmgren M, van Nes E H, Scheffer M, Zemp D C and Dekker S C 2018 Forest-rainfall cascades buffer against drought across the Amazon *Nat. Clim. Change* **8** 539–43
- [30] Van der Ent R J, Wang-Erlandsson L, Keys P W and Savenije H H G 2014 Contrasting roles of interception and transpiration in the hydrological cycle: II. Moisture recycling *Earth Syst. Dyn.* **5** 471–89
- [31] Zemp D C, Schleussner C F, Barbosa H M J, Hirota M, Montade V, Sampaio G, Staal A, Wang-Erlandsson L and Rammig A 2017 Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks *Nat. Commun.* **8** 14681
- [32] Marengo J A, Souza C A, Thonicke K, Burton C, Halladay K, Betts R A, Alves L M and Soares W R 2018 Changes in climate and land use over the Amazon region: current and future variability and trends *Frontiers Earth Sci.* **6** 228
- [33] Hansen M C *et al* 2013 High-resolution global maps of 21st-century forest cover change *Science* **342** 850–3
- [34] Bosmans J H C, van Beek L P H, Sutanudjaja E H and Bierkens M F P 2017 Hydrological impacts of global land cover change and human water use *Hydrol. Earth Syst. Sci.* **21** 5603–26
- [35] Sutanudjaja E H *et al* 2018 PCR-GLOBWB 2: a 5 arcmin global hydrological and water resources model *Geoscientific Model Dev.* **11** 2429–53
- [36] Tuinenburg O A, Hutjes R W A and Kabat P 2012 The fate of evaporated water from the Ganges basin *J. Geophys. Res.: Atmos.* **117** D01107
- [37] Lewis S L, Brando P M, Phillips O L, van der Heijden G M F and Nepstad D 2011 The 2010 Amazon drought *Science* **331** 554
- [38] Malhi Y, Aragão L E, Galbraith D, Huntingford C, Fisher R, Zelazowski P, Sitch S, McSweeney C and Meir P 2009 Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest *Proc. Natl Acad. Sci.* **106** 20610–5
- [39] Rodell M *et al* 2004 The global land data assimilation system *Bull. Am. Meteorol. Soc.* **85** 381–94
- [40] Van Beek L P H, Wada Y and Bierkens M F P 2011 Global monthly water stress: 1. Water balance and water availability *Water Resour. Res.* **47** W07517
- [41] Van Schaik E, Killaars L, Smith N E, Koren G, van Beek L P H, Peters W and van der Laan-Luijkx I T 2018 Changes in surface hydrology, soil moisture and gross primary production in the Amazon during the 2015/2016 El Niño *Phil. Trans. R. Soc. B* **373** 20180084
- [42] Dee D P *et al* 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q. J. R. Meteorol. Soc.* **137** 553–97
- [43] Zemp D C, Schleussner C F, Barbosa H M J, van der Ent R J, Donges J F, Heinke J, Sampaio G and Rammig A 2014 On the importance of cascading moisture recycling in South America *Atmos. Chem. Phys.* **14** 13337–59
- [44] Aguiar A P D *et al* 2016 Land use change emission scenarios: anticipating a forest transition process in the Brazilian Amazon *Glob. Change Biol.* **22** 1821–40
- [45] Marengo J A 2004 Interdecadal variability and trends of rainfall across the Amazon basin *Theor. Appl. Climatol.* **78** 79–96
- [46] Boers N, Marwan N, Barbosa H M J and Kurths J 2017 A deforestation-induced tipping point for the South American monsoon system *Sci. Rep.* **7** 41489
- [47] Wang-Erlandsson L, Fetzer I, Keys P W, van der Ent R J, Savenije H H G and Gordon L J 2018 Remote land use impacts on river flows through atmospheric teleconnections *Hydrol. Earth Syst. Sci.* **22** 4311–28
- [48] Amazonia Socioambiental (<https://amazoniasocioambiental.org/en/maps/#download>)

- [49] Brando P M *et al* 2014 Abrupt increases in Amazonian tree mortality due to drought-fire interactions *Proc. Natl Acad. Sci.* **111** 6347–52
- [50] Boisier J P, Ciais P, Ducharme A and Guimberteau M 2015 Projected strengthening of Amazonian dry season by constrained climate model simulations *Nat. Clim. Change* **5** 656–60
- [51] Fu R *et al* 2013 Increased dry-season length over southern Amazonia in recent decades and its implication for future climate projection *Proc. Natl Acad. Sci.* **110** 18110–5
- [52] Sena E T, Dias M A F S, Carvalho L M V and Dias P L S 2018 Reduced wet-season length detected by satellite retrievals of cloudiness over Brazilian Amazonia: a new methodology *J. Clim.* **31** 9941–64
- [53] Staver A C, Archibald S and Levin S A 2011 The global extent and determinants of savanna and forest as alternative biome states *Science* **334** 230–2
- [54] Wuyts B, Champneys A R and House J I 2017 Amazonian forest-savanna bistability and human impact *Nat. Commun.* **8** 15519
- [55] Van Nes E H, Staal A, Hantson S, Holmgren M, Pueyo S, Bernardi R E, Flores B M, Xu C and Scheffer M 2018 Fire forbids fifty-fifty forest *PLoS One* **18** e0191027
- [56] Flores B M, Staal A, Jakovac C C, Hirota M, Holmgren M and Oliveira R S 2019 Soil erosion as a resilience drain in disturbed tropical forests *Plant Soil* Accepted (<https://doi.org/10.1007/s11104-019-04097-8>)
- [57] Lovejoy T E and Nobre C 2018 Amazon tipping point *Sci. Adv.* **4** eaat2340
- [58] Nepstad D *et al* 2014 Slowing Amazon deforestation through public policy and interventions in beef and soy supply chains *Science* **344** 1118–23
- [59] Nepstad D *et al* 2009 The end of deforestation in the Brazilian Amazon *Science* **326** 1350–1
- [60] Galaz V, Crona B, Dauriach A, Scholtens B and Steffen W 2018 Finance and the Earth system—Exploring the links between financial actors and nonlinear changes in the climate system *Glob. Environ. Change* **53** 296–302
- [61] Levis C, Flores B M, Mazzochini G G, Manhães A P, Campos-Silva J V, Amorim P B, Peroni N, Hirota M and Clement C R 2020 Help restore Brazil's governance of globally important ecosystem services *Nat. Ecology Evol.* **4** 172–3 Accepted
- [62] Jones C and Carvalho L M V 2013 Climate change in the South American monsoon system: present climate and CMIP5 projections *J. Clim.* **26** 6660–78
- [63] Lawrence D and Vandecar K 2015 Effects of tropical deforestation on climate and agriculture *Nat. Clim. Change* **5** 27–36
- [64] D'Almeida C, Vörösmarty C J, Hurtt G C, Marengo J A, Dingman S L and Keim B D 2007 The effects of deforestation on the hydrological cycle in Amazonia: a review on scale and resolution *Int. J. Climatol.* **27** 633–47
- [65] Avissar R and Liu Y 1996 Three-dimensional numerical study of shallow convective clouds and precipitation induced by land surface forcing *J. Geophys. Res.: Atmos.* **101** 7499–518