

RESEARCH ARTICLE

Journal of Ecology



Environmental drivers of *Sphagnum* growth in peatlands across the Holarctic region

Fia Bengtsson¹  | Håkan Rydin¹  | Jennifer L. Baltzer²  | Luca Bragazza^{3,4,5} | Zhao-Jun Bu^{6,7}  | Simon J. M. Caporn⁸ | Ellen Dorrepaal⁹  | Kjell Ivar Flatberg¹⁰ | Olga Galanina^{11,12}  | Mariusz Gałka¹³  | Anna Ganeva¹⁴ | Irina Goia¹⁵ | Nadezhda Goncharova¹⁶ | Michal Hájek¹⁷  | Akira Haraguchi¹⁸  | Lorna I. Harris¹⁹  | Elyn Humphreys²⁰  | Martin Jiroušek^{17,21}  | Katarzyna Kajukato²² | Edgar Karofeld²³  | Natalia G. Koronatova²⁴  | Natalia P. Kosykh²⁴ | Anna M. Laine^{25,26}  | Mariusz Lamentowicz²²  | Elena Lapshina²⁷  | Juul Limpens²⁸  | Maiju Linkosalmi²⁹ | Jin-Ze Ma^{6,7} | Marguerite Mauritz³⁰ | Edward A. D. Mitchell^{31,32}  | Tariq M. Munir^{33,34}  | Susan M. Natali³⁵  | Rayna Natcheva¹⁴ | Richard J. Payne^{36,37*} | Dmitriy A. Philippov³⁸  | Steven K. Rice³⁹ | Sean Robinson⁴⁰  | Bjorn J. M. Robroek⁴¹  | Line Rochefort⁴²  | David Singer^{31,43}  | Hans K. Stenøien¹⁰  | Eeva-Stiina Tuittila²⁵  | Kai Vellak²³ | James Michael Waddington⁴⁴  | Gustaf Granath¹ 

¹Department Ecology and Genetics, Uppsala University, Uppsala, Sweden; ²Biology Department, Wilfrid Laurier University, Waterloo, ON, Canada; ³Department of Life Science and Biotechnologies, University of Ferrara, Ferrara, Italy; ⁴Swiss Federal Institute for Forest, Snow and Landscape Research, WSL Site Lausanne, Lausanne, Switzerland; ⁵Ecole Polytechnique Fédérale de Lausanne EPFL, School of Architecture, Civil and Environmental Engineering ENAC, Laboratory of Ecological Systems ECOS, Lausanne, Switzerland; ⁶Institute for Peat and Mire Research, State Environmental Protection Key Laboratory of Wetland Ecology and Vegetation Restoration, Northeast Normal University, Changchun, China; ⁷Key Laboratory of Geographical Processes and Ecological Security in Changbai Mountains, Ministry of Education, School of Geographical Sciences, Northeast Normal University, Changchun, China; ⁸Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK; ⁹Climate Impacts Research Centre, Department of Ecology and Environmental Science, Umeå University, Abisko, Sweden; ¹⁰NTNU University Museum, Norwegian University of Science and Technology, Trondheim, Norway; ¹¹Institute of Earth Sciences, St. Petersburg State University, St. Petersburg, Russia; ¹²Komarov Botanical Institute Russian Academy of Sciences, St. Petersburg, Russia; ¹³Department of Geobotany and Plant Ecology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland; ¹⁴Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia, Bulgaria; ¹⁵Department of Taxonomy and Ecology, Faculty of Biology and Geology, Babeş-Bolyai University, Cluj Napoca, Romania; ¹⁶Institute of Biology of Komi Scientific Centre of the Ural Branch of the Russian Academy of Science, Syktyvkar, Russia; ¹⁷Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic; ¹⁸Department of Biology, The University of Kitakyushu, Kitakyushu, Japan; ¹⁹Department of Geography, McGill University, Montreal, QC, Canada; ²⁰Department of Geography and Environmental Studies, Carleton University, Ottawa, ON, Canada; ²¹Department of Plant Biology, Faculty of AgriSciences, Mendel University in Brno, Brno, Czech Republic; ²²Climate Change Ecology Research Unit, Adam Mickiewicz University in Poznań, Poznań, Poland; ²³Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia; ²⁴Laboratory of Biogeocenology, Institute of Soil Science and Agrochemistry, Siberian Branch of Russian Academy of Sciences, Novosibirsk, Russia; ²⁵Peatland and Soil Ecology Group, School of Forest Sciences, University of Eastern Finland, Joensuu, Finland; ²⁶Department of Ecology and Genetics, University of Oulu, Oulu, Finland; ²⁷Yugra State University, Khanty-Mansiysk, Russia; ²⁸Plant Ecology and Nature Conservation Group, Wageningen University, Wageningen, The Netherlands; ²⁹Finnish Meteorological Institute, Helsinki, Finland; ³⁰Center for Ecosystem Science and Society (Ecos), Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ, USA; ³¹Laboratory of Soil Biodiversity, Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland; ³²Jardin botanique de Neuchâtel, Neuchâtel, Switzerland; ³³Department of Geography, University of Calgary, Calgary, AB, Canada; ³⁴Department of Geography and Planning, University of Saskatchewan, Saskatoon, SK, Canada; ³⁵Woods Hole Research Center, Falmouth, MA, USA; ³⁶Environment and Geography, University of York, York, UK; ³⁷Lomonosov Moscow State University, Moscow, Russian Federation; ³⁸Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok, Russia; ³⁹Department of Biological Sciences, Union College, Schenectady, NY, USA; ⁴⁰Department of Biology, SUNY-Oneonta, Oneonta, NY, USA; ⁴¹Aquatic Ecology & Environmental Biology, Institute for Water and Wetland Research, Radboud University, Nijmegen, The Netherlands; ⁴²Department of Plant Science and Center for Northern Studies, Laval University, Québec, QC, Canada; ⁴³Department of Zoology, Institute of Biosciences, University of São Paulo, São Paulo, Brazil and ⁴⁴School of Earth, Environment and Society, McMaster University, Hamilton, ON, Canada

Richard J. Payne: deceased.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society

Correspondence

Fia Bengtsson

Email: fia.bengtsson@ebc.uu.se

Funding information

W. Garfield Weston Foundation Fellowship for Northern Conservation; Extensius; Università degli Studi di Ferrara, Grant/Award Number: FAR 2013 and FAR 2014; Vetenskapsrådet, Grant/Award Number: 2015-05174; Academy of Finland, Grant/Award Number: Project code 287039; National Science Foundation, Grant/Award Number: NSF-1312402; Jilin Provincial Science and Technology Development Project, Grant/Award Number: 20190101025JH; The Russian Science Foundation, Grant/Award Number: 19-14-00102; The Russian Foundation for Basic Research, Grant/Award Number: 14-05-00775, 15-44-00091, 19-05-00830, 18-04-00988 and 18-44-860017; The Polish National Centre for Research and Development, Grant/Award Number: 203258; The National Science Centre, Poland, Grant/Award Number: 2015/17/B/ST10/01656; Institutional Research Funds from the Estonian Ministry of Education and Research, Grant/Award Number: IUT34-7; The Natural Sciences and Engineering Research Council of Canada; The Czech Science Foundation, Grant/Award Number: 19-28491X; The National Natural Science Foundation of China, Grant/Award Number: 41471043 and 41871046; The Ministry of Science and Higher Education of the Russian Federation; The Swiss National Science Foundation, Grant/Award Number: P2NEP3_178543

Handling Editor: Rien Aerts**Abstract**

1. The relative importance of global versus local environmental factors for growth and thus carbon uptake of the bryophyte genus *Sphagnum*—the main peat-former and ecosystem engineer in northern peatlands—remains unclear.
2. We measured length growth and net primary production (NPP) of two abundant *Sphagnum* species across 99 Holarctic peatlands. We tested the importance of previously proposed abiotic and biotic drivers for peatland carbon uptake (climate, N deposition, water table depth and vascular plant cover) on these two responses. Employing structural equation models (SEMs), we explored both indirect and direct effects of drivers on *Sphagnum* growth.
3. Variation in growth was large, but similar within and between peatlands. Length growth showed a stronger response to predictors than NPP. Moreover, the smaller and denser *Sphagnum fuscum* growing on hummocks had weaker responses to climatic variation than the larger and looser *Sphagnum magellanicum* growing in the wetter conditions. Growth decreased with increasing vascular plant cover within a site. Between sites, precipitation and temperature increased growth for *S. magellanicum*. The SEMs indicate that indirect effects are important. For example, vascular plant cover increased with a deeper water table, increased nitrogen deposition, precipitation and temperature. These factors also influenced *Sphagnum* growth indirectly by affecting moss shoot density.
4. **Synthesis.** Our results imply that in a warmer climate, *S. magellanicum* will increase length growth as long as precipitation is not reduced, while *S. fuscum* is more resistant to decreased precipitation, but also less able to take advantage of increased precipitation and temperature. Such species-specific sensitivity to climate may affect competitive outcomes in a changing environment, and potentially the future carbon sink function of peatlands.

KEYWORDS

climate, global change, net primary production, nitrogen deposition, PAR, peat mosses, plant–climate interactions, structural equation model

1 | INTRODUCTION

Net primary production in peatlands is relatively low, but because, in general production exceeds decomposition, peatlands have remained important carbon (C) sinks throughout the Holocene (Yu, 2012). As a result, northern peatlands store c. 500 Gt C (Loisel et al., 2014; Yu, Loisel, Brosseau, Beilman, & Hunt, 2010). A recent analysis suggests that the stock may even be >1,000 Gt (Nichols & Peteet, 2019), which is more than the C in the atmosphere today (829 Gt; IPCC, 2013). In northern peatlands, the growth of peat mosses (genus *Sphagnum*) is vital for C sequestration and storage, as they effectively engineer a wet and acidic environment that inhibits decomposition (Rydin & Jeglum, 2013). However, despite their crucial role in peatland C cycling, we still lack a clear understanding of the factors controlling *Sphagnum* growth at global and local scales.

The important role of *Sphagnum* in C sequestration suggests a need to include *Sphagnum* growth in terrestrial ecosystem models. For example, Charman et al. (2013) suggested that biomass input (i.e. NPP) has driven the peat accumulation rate over the last millennium, while variation in decomposition has mattered less. Peatland models have recently included *Sphagnum*-specific growth functions (Turetsky et al., 2012), but peatland modules are generally lacking in Earth system models (e.g. ORCHIDEE, Qiu et al., 2018), but see Bechtold et al. (2019). A literature survey revealed that attempts to link variation in *Sphagnum* NPP to important environmental drivers have come to different conclusions (Table 1). Gunnarsson (2005) found that temperature, precipitation, altitude and latitude explained 40% of the variation in productivity. Moore (1989) identified annual mean temperature as a key driver of *Sphagnum* production, while Krebs, Gaudig, and Joosten (2016) reported that growth of *S. papillosum* was primarily influenced by water availability (precipitation frequency

TABLE 1 Previous studies relating *Sphagnum* growth to climatic variables in temperate and boreal peatlands

Reference	Data	Modelling approach	Response(s)	Explained variance
Asada et al. (2003)	1 peatland, 9 bryophyte species, 6 <i>Sphagnum</i>	Correlations between Growth and Climate Index (CI, based on mean daily prec and mean, max, min temp), or each predictor separately	NPP 1 year, within season variation	For <i>S. fuscum</i> , daily prec. $r = 0.6$ (ns), temp. $r = -0.4$ – 0.05 (ns), CI $r = 0.7$ (*)
Gunnarsson (2005)	Metastudy: 55 publications, 35 species	Multiple regression with predictors temp, prec, alt, lat, distance to sea	NPP	Temp and prec most important: Model explains 40%; including species as a factor, 36%
Krebs et al. (2016)	1) New data, 2 species, 3 sites for <i>S. papillosum</i> , 2 sites for <i>S. palustre</i> , mostly 2 years. 2) Metastudy of <i>S. papillosum</i> from 18 publications	1) Multiple regression with climate, lat, 'main site variables'. 2) Boosted regression tree	LI, NPP	1) Model explains 22% of NPP, 52% of LI. 2) Microhabitat explains 27%, number of rain-free days: 15%, prec/temp: 10%, temp: 7%
Limpens et al. (2011)	Metastudy of 107 nitrogen fertilization field experiments	Meta-regressions	NPP	Higher summer temperature and increased annual precipitation intensified the negative effects of nitrogen deposition
Loisel et al. (2012)	Metastudy with 52 sites, <i>S. fuscum</i> and <i>S. magellanicum</i>	Simple and multiple regressions	LI	$R^2 = 0.23$ for PAR, P/Eq had barely an effect. Multiple regression only for 'continental' sites, does not improve simple regression ($R^2 = 0.31$ for cont. sites)
Moore (1989)	1) New data, 6 species, of which 2 measured at 2 sites, 2 years. 2) Metastudy of 14 publications, and one new site	1) Correlations and regressions. 2) Regression	NPP	1) Mean daily temp, daily prec, bright sunshine hrs. 2) Mean annual temp

and precipitation/temperature quotient). Loisel, Gallego-Sala, and Yu (2012) found that cumulative photosynthetically active radiation (PAR) for days over 0°C was the most important driver of growth. Globally, terrestrial NPP is largely determined by precipitation and water availability, and in northern regions (above 50°N) temperature and solar radiation become increasingly important (Gallego-Sala et al., 2018; Schloss, Kicklighter, Kaduk, Wittenberg, & The Participants of the Potsdam NPP Model Intercomparison, 2001).

The contrasting findings of these previous studies may result from inconsistencies in methodology. First, meta-analyses use data collected using different methods, in different habitats (e.g. pH and degree of ground-water influence) and over different time periods (Gunnarsson, 2005; Krebs et al., 2016; Limpens et al., 2011; Loisel et al., 2012). Second, many of the previous studies are geographically limited (Asada, Warner, & Banner, 2003; Moore, 1989; Walker et al., 2017) and consequently cover a narrow range of climatic conditions. Third, many studies include different species for different areas and species can covary with environmental conditions, which may add bias and uncertainty to the results. To overcome these problems we performed a Holarctic, coordinated sampling effort, focusing on common but ecologically divergent species.

In addition to climatic drivers, local conditions and air pollutants such as nitrogen (N) and sulphur (S) deposition affect *Sphagnum* growth. Limpens et al. (2011) examined the effects of increased N deposition in a meta-analysis comprising 29 studies across Europe

and North America. They showed that small N additions to sites with low background N deposition stimulated growth, while adding higher levels of N depressed growth. Interestingly, increased vascular plant cover caused the shift from a positive to a negative growth response to N to occur at lower N addition levels (Limpens et al., 2011). Experiments have shown that N addition leads to increased cover of vascular plants that will reduce light availability, increase litter and lower surface temperature, thereby reducing *Sphagnum* productivity (Berendse et al., 2001; Chong, Humphreys, & Moore, 2012). Moreover, Hayward and Clymo (1983) reported reduced moss mass growth when PAR was lowered. In addition to vascular plant cover, the position above the water table, which is a proxy for water availability, has been correlated with *Sphagnum* NPP. Production was lower for *Sphagnum* growing higher above the water table (e.g. Gunnarsson, 2005; Krebs et al., 2016). These local factors and N availability are known to modify *Sphagnum* NPP, but to what extent relative to climatic drivers has not been investigated.

Here, we set out to test the importance of global and local environmental drivers of length growth and C sequestration potential (i.e. NPP) of two common peat-forming *Sphagnum* species in northern peatlands. Length (height) increment is essential for the shoot to keep up with vascular plants and to compete for light with neighbouring *Sphagnum* shoots. NPP is more directly related to peatland C sequestration, and depends on the combination of length increment (LI) and the bulk density of the material produced. The two

Sphagnum species, *S. fuscum* and *S. magellanicum* were selected to represent different niche preferences within peatlands, i.e. lawns and hummocks. Our aim was to explain and predict variation in *Sphagnum* productivity within sites and across the Holarctic using statistical modelling—including structural equation models (SEMs, Shipley, 2009). By sampling in a consistent and standardized manner across sites and years, we tested the relative contributions of previously identified environmental drivers (e.g. water table position, temperature, precipitation, PAR) to growth. We did this by exploring both direct and indirect effects (e.g. impact on vascular plant cover, shoot density) of these drivers. This allows us to test to what extent climate-change effects on *Sphagnum* will be modified and/or mitigated by local conditions and if the response is species-specific. More broadly, improving our understanding of the drivers of *Sphagnum* NPP will support forecasting of changes in peatland carbon dynamics under ongoing climate change.

2 | MATERIALS AND METHODS

2.1 | Study species and sampling design

Two *Sphagnum* species belonging to different subgenera (sections) were selected for this study: *Sphagnum fuscum* (Schimp.) H. Klinggr. and *S. magellanicum* Brid. *Sphagnum fuscum* in the subgenus *Acutifolia* is characterized by its brown colour and its small and densely packed capitula (Figure 1). It typically forms hummocks in bogs, but also in minerotrophic peatlands where its hummocks often are high enough

to be ombrotrophic. Recently, some individuals of *S. fuscum* in Europe have been identified as conspecific to the North American *S. beothuk* R. E. Andrus (Kyrkjeeide et al., 2015). Genetic analyses performed on one sample from each of 26 of our 102 sites indicated a small proportion (two samples) of *S. beothuk* (N. Yousefi, K. Hassel, & H.K. Stenøien, unpubl.). *Sphagnum magellanicum* belongs to the subgenus *Sphagnum*, which is a group of large and stout species with cucullate leaves. *Sphagnum magellanicum* stands out in its subgenus due to its colour that varies from deep red to almost completely green (Figure 1). Also, recently, *S. magellanicum* in northern peatlands has been split into two species: *S. medium* Limpr. that commonly occupies the bog expanse, where it grows close to the water table and at relatively intense light conditions, and *S. divinum* Flatberg & Hassel that occupies bog margins and poor fens, growing further from the water table and at lower light intensity (Hassel et al., 2018). Genetic analysis (N. Yousefi, K. Hassel, & H.K. Stenøien, unpubl.) on one sample each from 62% of our sites indicates that both species were included in approximately equal proportions. Given that *S. beothuk*/*S. fuscum* and *S. medium*/*S. divinum* are sister species and were treated collectively in the past, and that they had not been differentiated at the time of data collection, we here follow Granath et al. (2018) and treat our study species as *S. fuscum* coll. and *S. magellanicum* coll.

In 2013 and 2014 we sampled throughout the species' Holarctic distributions to cover the climatic variation of sites where the species co-occur. In total, we sampled at 102 sites, of which three sites were excluded from the analysis due to inconsistent sampling (Hani and Mangui, China) or lack of sample tissue N concentration data (Verh-Tarka, Russia). Of the remaining sites, 76 had both species,




FIGURE 1 *Sphagnum magellanicum* coll. (red, left) and *S. fuscum* coll. (brown, right). Illustration: F. Bengtsson

and 85 had data from both 2013 and 2014 (Figure 2; Table S1). The sample sites covered a large proportion of the boreal and temperate biomes where northern peatlands are present (Figure 3).

Our aim was to sample from similar habitats across the sites, and we therefore sampled species in their typical open, nutrient-poor habitats, ranging from predominantly bogs to poor fens. At each site, four mono-specific, homogeneous non-treed patches (1 m²) were chosen for each species, with each patch at least 10 m apart.

2.2 | Length growth and production

Sphagnum growth is easy to measure as they have no roots or rhizoids (Shaw et al., 2010) and new growth is confined to the top of the shoot. We assessed *Sphagnum* growth using two different approaches; LI (mm), and biomass production (NPP; g/m²). To measure LI we inserted a minimum of three 'brush wires' in each patch at the start of the growing season (Figure S1). The brush wires resemble bottlebrushes and were constructed by attaching paintbrush bristles to c. 10–15 cm long wires. This is an improved version of the cranked wire method described in Clymo (1970), and is now the most commonly applied method (Rydin & Jeglum, 2013). The brushes were inserted c. 5–10 cm into the *Sphagnum* carpet without disturbing the moss surface by initially pressing the bristles into a narrow tube, which was then pulled out while the brush was left in the *Sphagnum* carpet allowing the bristles to spread and 'anchor' in the vegetation. The height of the part of the brush wire protruding from the *Sphagnum* carpet was measured each year at the beginning of the season, and then remeasured at the end of the season. The


difference between spring and autumn measurements gave us the LI (mm). Measurements with large negative values (<–5 mm growth) indicated physical disturbance and were removed from analyses.

To estimate stem bulk density and shoot density, we collected one moss core (5–10 cm diameter) from each sampling patch at the end of the season at the same time as the final wire length measurements. The capitula (i.e. the moss top with a height of c. 0.5 cm in *S. fuscum* and 1 cm in *S. magellanicum*) were removed and dried, and the 3-cm stem sections immediately below the capitula were dried (24 hr at 60–65°C) and weighed to obtain stem bulk density (BD; kg/m³). The area-based biomass growth (NPP; g/m²) was calculated as BD × LI. In addition to measurements of LI and production for the growing season, we calculated LI and NPP per day during the sampling periods. We also counted the number of capitula in the core to obtain numerical shoot density (cm^{–2}).

2.3 | Local predictors: Water table and vascular plant cover

For each patch and year, we recorded two variables that we expected would relate to growth: height of the moss surface above the water table (HWT; cm) and vascular plant cover (vertical projection on 1 m² surrounding the patch, %) at the start and end of the season. For a few sites, these measurements were only taken once per year at either the start or end of the season. For sites where these metrics were made both early and late in the season, the correlation between start and end values of HWT was $r \approx 0.80$ ($df = 83$ in 2013, and 85 in 2014) and between start and end values of vascular plant

FIGURE 2 The sampling design included 102 sites distributed across the Holarctic region where either *Sphagnum fuscum* or *S. magellanicum*, or both were sampled. At each site and for each year, we aimed to sample four patches (red circles) of each species if they were both present at the site, and at each patch we took three measurements of length increment, and one measurement of bulk density to calculate biomass accumulation. Illustration: F. Bengtsson






FIGURE 3 Mean annual temperature (°C) and annual precipitation (mm) extracted from WorldClim (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005) for the study sites (black rings) superimposed on the Whittaker biome chart (Biome chart adopted from Ricklefs (2008) plotted with the R package PLOTBIOMES, <https://github.com/valentininelav/plotbiomes>)

cover $r \approx 0.90$ (df 76 in 2013 and 64 in 2014) respectively. To include all sites in the analyses, we therefore used the average for each year for all sites.

Dried capitula were sent for elemental analysis (C, N, P, K, Mg, Mn, Ca, Cd, Cr, Cu, Al, B, Fe, Na, Ni, Zn) at Forest Research, Alice Holt Lodge, UK. For total C and N, around 10 mg (6 decimal balance) of milled and dried (70°C) sample was analysed using the combustion method with a Carlo Erba CN analyser (Flash1112 series). For cations and trace metals, 0.6 grams of plant material was digested with a microwave system (Anton Paar Multiwave 3000) using 7.5 ml of HNO_3 at 185°C for 30 min (20 min ramp from ambient to 185°C). The extract was then adjusted to 25 ml and analysed with an ICP-OES dual view (Thermo ICap 6500). Most samples were from 2013 and in general two patches per species and site were analysed. Total element concentrations were expressed as percentages or mg/kg. In this study we use mean N concentration per site, but include all data in our publication for completeness.

2.4 | Global predictors: Weather and nitrogen deposition

Data on meteorological variables for each site and year were extracted from the NASA GESDISC (Global Modeling and Assimilation Office (GMAO), 2015) land surface and flux diagnostic products (M2T1NXLND, M2TMNXFLX). We used temperature (T2M; K), precipitation (PRECTOTLAND; $\text{kg m}^{-2} \text{s}^{-1}$), evaporation (EVLAND; $\text{kg m}^{-2} \text{s}^{-1}$), PAR (diffuse flux: PARDF; W/m^2 and beam flux PARDR; W/m^2), time during an hour with no precipitation (HOURNORAIN; s)

as predictors. HOURNORAIN was recalculated as the mean rain free period (average time between rain events; days). These data are derived from satellite and meteorological station observations and are gridded with a grid cell resolution of longitude 0.667° and latitude 0.5° . We extracted data for the periods of growth measurements at each site, and calculated the average temperature (°C), total precipitation (kg/m^2), evaporation (kg/m^2) and PAR (PARDF+PARDR, W/m^2) and the average number of consecutive days without rain (discarding days with <1 mm precipitation; d; Table S2). Data on nitrogen deposition were extracted from the model synthesis of Lamarque et al. (2013). They modelled global total N deposition for the time period 1995–2005 (model output given as an estimate for the year 2000) with a resolution of 0.5° longitude by 0.5° latitude. Elevation and coordinates were extracted from Google Maps (Table S1).

2.5 | Data analysis

To test the importance of the controlling factors for *Sphagnum* growth, we selected variables based on previous studies (see Table 1). Temperature and precipitation have been used in most studies and were the main predictors in the meta-analysis by Gunnarsson (2005). He also included the geographical variables latitude, elevation and distance from the sea, but we did not test these variables as they are not mechanistically related to growth, and their direct effects on growth should be included in the climatic data. Loisel et al. (2012) compiled global growth data for *S. fuscum* and *S. magellanicum*, and modelled seasonal LI as a response to growing season

length, PAR and P/E_q (index of moisture balance: the quotient between annual precipitation [P] and the annually integrated equilibrium evapotranspiration [E]). We included PAR as a predictor and tested if P/E or $P-E$ would perform as better predictors compared to P and E individually. An additional climatic variable was average length of rain free periods (Krebs et al., 2016). Nitrogen availability has been related to *Sphagnum* growth (Granath, Strengbom, & Rydin, 2012; Limpens et al., 2011) and our model included N deposition (site level) and tissue N concentration (site level but species-specific). Within-site predictors were vascular plant cover (Gavazov et al., 2018), shoot numerical density and height above the water table (Table 1). To test for species-specific responses we added interaction terms with species for all predictors.

For estimation and testing of predictor effects, we used linear mixed models (LMMs) to set up a piecewise structural equation model (pSEM, Lefcheck, 2016) in R (R Core Team, 2017) in the package `LME4` (ver `lme4_1.1-20`; Bates, Maechler, Bolker, & Walker, 2015). Since we sampled approximately four patches per species and site, we used 'site' as a random factor in our models, and the site effect was allowed to vary between years (random intercept-slope model). As a first step, we fitted LMMs (classic multivariable regression) including the global (temperature, precipitation, evaporation, PAR, average length of rain free periods, N dep) and local (HWT, vascular plant cover, N tissue) predictors, species and year as fixed factors, including interaction effects between global variables and species. We ran the models without interaction effects to explore main effects and their R^2 values. p values for interaction and main effects were extracted from type II ANOVAs using the `CAR` package (Fox & Weisberg, 2011). R^2 values can quantitatively evaluate the fit of the model and were produced for site (R^2_{site} , explained between site variation) and within-site level (residual error) by calculating the proportional reduction in these variance components under the fitted model when compared to the null model (following Johnson, 2014; Nakagawa & Schielzeth, 2013 and using code from the R package `MuMIn`, Bartoń, 2018). To give a measurement of overall model fit we also estimated the amount of total variance explained by fixed effects (Marginal R^2). A parametric bootstrap procedure was employed to produce approximate confidence intervals (CIs) of R^2 s and we present the 90% percentile CI.

As a second step we extended the above analyses to a pSEM to also examine indirect effects. A piecewise SEM is flexible as it allows multilevel structures and that each response (endogenous variables; here NPP or LI, shoot density, tissue N concentration, vascular plant cover and HWT) is modelled individually (Shipley, 2009). We performed SEMs for LI and NPP, and due to important species interactions found in initial multivariable regressions, the SEMs were done for each species. The paths included in the SEM (i.e. model specification) were the same for each species and LI and NPP. In principle, all paths were included, but we removed paths for which there were no clear mechanisms. For example, many of the variables cannot affect HWT, such as tissue N concentration, and tissue N concentration is expected to be modified by vascular plant cover, but vascular plant cover is not affected by tissue N. The

complete model specifications can be found in Tables S7 and S8. To simplify the figure illustrations of models we removed links that had minor impact on the response ($p \geq 0.1$). To be able to compare direct and indirect effects we standardized both endogenous and exogenous variables by dividing by one standard deviation.

Standard residual analyses were performed to check homogeneity and normality of errors, distribution of random effects and influence of individual data points. Residuals showed increasing variance and Box-Cox transformations were performed using the function `powerTransform` in the `CAR` package (ver 3.0-0, Fox & Weisberg, 2011) to achieve homogeneous errors. We used correlation analysis to examine potential issues with multicollinearity.

3 | RESULTS

3.1 | Descriptive statistics

Across species and years, growth measurements showed as much variation within sites as between sites. The proportion of within-site variance to total variation was 47% for LI and 57% for NPP measured on an annual basis (year^{-1} ; Table S3). The larger species, *S. magellanicum*, had on average greater LI than *S. fuscum*, while *S. fuscum* had somewhat higher NPP than *S. magellanicum* due to its higher bulk density (Figure 4; Table S5).

On average, LI was slightly higher in the second year (mean \pm SE; 2013: 16.9 ± 1.1 , 2014: 18.2 ± 1.2 mm), while NPP was lower in the second year (2013: 194.6 ± 12.0 ; 2014: 182.7 ± 11.2 g/m²; Figure 4; Table S5). Correlations between the years at the site level were $r = 0.68$ ($df = 68$) and $r = 0.57$ ($df = 65$) for *S. fuscum* LI and NPP respectively ($p < 0.0001$). For *S. magellanicum* correlations between years were $r = 0.69$ ($df = 77$) and $r = 0.48$ ($df = 73$), respectively ($p < 0.0001$). Between year differences were only detected for NPP (Table S4), indicating a lower NPP in 2014, but the effect was small (12.6 g/m² difference).

The correlation (at site level) between N deposition and N tissue concentration was $r = 0.35$ ($df = 97$), and N deposition and vascular plant cover was $r = 0.25$ ($df = 95$).

3.2 | Multivariable regression models

Our full models with LI as response explained 53% of the growth calculated per day (day^{-1} ; CI 40%–64%) and 52% on an annual basis (year^{-1} ; CI 38%–63%) of variance between sites (Table S4). The models explained less variation in NPP between sites than in LI, and models showed similar R^2_{site} values on a daily basis (31%; CI 11%–46%) and annual basis (32%; CI 12%–47%). The correlations between LI and NPP were $r = 0.51$ and 0.55 for *S. fuscum* in 2013 and 2014, respectively ($p < 0.0001$, $df = 74$ and 73), and $r = 0.56$ and 0.57 for *S. magellanicum* in 2013 and 2014, respectively ($p < 0.0001$, $df = 79$ and 82).

We detected species-specific responses to precipitation (LI and NPP), temperature (LI), evaporation (NPP) and numerical density




FIGURE 4 Growth and bulk density for *Sphagnum fuscum* and *S. magellanicum*, for each year (2013 and 2014 respectively). (a) Season length increment (LI; mm/year), (b) bulk density (BD; kg/m³) and (c) seasonal production (NPP; g m⁻² year⁻¹). Boxplots show median (horizontal lines), 25th and 75th percentiles (boxes), values within a 1.5 interquartile range (vertical lines) and outside values (points). In white: mean \pm SE

(NPP; Table S4). These interactions suggest different effect sizes and similar directions of the response (e.g. the positive effect of precipitation was weaker and often not distinguished from zero in *S. fuscum* compared to a clear positive effect on *S. magellanicum* in all models).

Using precipitation minus evaporation (precipitation surplus, $P-E$), as a predictor instead of using the two variables as separate predictors gave similar results (Table S6).

For comparison with Loisel et al. (2012) we ran the LI model using PAR, species and year as predictors, which gave an R^2_{site} of 22% and an R^2_{marginal} of 12% respectively.

TABLE 2 Path coefficients for direct and indirect effect on length increment (LI) extracted and calculated from the piecewise SEM presented in Figure 5. Only coefficients with $p < 0.15$ were included in the calculations. No eff. = no path coefficients $p < 0.15$

Variable	Effect type on LI	<i>Sphagnum fuscum</i> Path coefficient	<i>Sphagnum magellanicum</i> Path coefficient
Temp	Direct	0.249	0.475
	Indirect	-0.017	No eff.
Precipitation	Direct	No eff.	0.330
	Indirect	-0.009	-0.011
Evaporation	No effects		
PAR	No effects		
HWT	Indirect	-0.069	-0.035
Vascular plant cover	Direct	-0.069	-0.102
No rain days	Direct	No eff.	-0.123
N deposition	Indirect	0.042	0.014
N concentration	Direct	0.157	0.106
Numerical density	No effects		

3.3 | Structural equation modelling

Using structural equation modelling built from mixed effects models, we investigated direct and indirect effects of studied variables on length (LI; Table 2; Figure 5) and biomass growth (NPP; Table 3; Figure 6). The full output from SEM and effects on responses other than growth (i.e. shoot density, tissue N concentration and vascular plant cover) in the piecewise SEM are shown in Tables S7 and S8.

3.3.1 | Direct and indirect effects of global predictors

Regardless of species, temperature had positive direct effects on growth (Tables 2 and 3), and the effect was strongest for LI in *S. magellanicum* (path coefficient = 0.475, $p < 0.0001$). Precipitation had a similar direct effect on LI for *S. magellanicum* (path coefficient = 0.330, $p < 0.001$), but no direct effect on LI for *S. fuscum* nor on NPP. Both these climatic variables had indirect effects on growth through numerical density and/or vascular plant cover (although there were no indirect effects for temperature in *S. magellanicum* growth). For example, precipitation had no direct effects on NPP, but its effects on vascular plant cover and numerical density led indirectly to negative effects on growth (*S. fuscum*: path coefficient = -0.089, $p < 0.03$, *S. magellanicum*: path coefficient = -0.041, $p < 0.08$; Table 3). Evaporation had virtually no direct effect on growth, but had a positive effect on moss shoot density (*S. fuscum*: path coefficient = 0.32, $p < 0.0001$, *S. magellanicum*: path




FIGURE 5 Graphical representation of piecewise structural equation models using mixed effects models to describe effects of the environmental variables from the models in Table S4 on annual length increment. Num density = numerical density of *Sphagnum* shoots, PAR = photosynthetic active radiation, No rain = mean length of rain free periods, HWT = height above the water table. White boxes are exclusively predictor variables, grey boxes are response (length increment), or predictor and response. Arrowheads show the direction of effects and line thickness indicates the effect size. For simplicity, arrows associated with $p \geq 0.1$ have been removed. See Table S7 for complete model results

coefficient = 0.09, $p < 0.001$; Tables S7 and S8). The number of rain free days had consistently direct negative effects on growth, although the effect was not significant for LI in *S. fuscum*. Only a direct effect of PAR on growth was detected (only significant for NPP in *S. magellanicum*), while nitrogen deposition had indirect effects by increasing N content in the mosses and vascular plant cover.

TABLE 3 Path coefficients for direct and indirect effect on net primary productivity (NPP) extracted and calculated from the piecewise SEM presented in Figure 6. Only coefficients with $p \leq 0.1$ were included in the calculations. No eff. = no path coefficients $p < 0.11$

Variable	Effect type on NPP	<i>Sphagnum fuscum</i> Path coefficient	<i>Sphagnum magellanicum</i> Path coefficient
Temp	Direct	0.225	0.249
	Indirect	-0.191	No eff.
Precipitation	Indirect	-0.089	-0.041
Evaporation	Indirect	0.097	0.060
PAR	Direct	0.209	0.264
HWT	Indirect	0.057	-0.042
Vascular plant cover	Direct	-0.100	-0.124
	Indirect	-0.043	-0.016
No rain days	Direct	-0.185	-0.213
N deposition	Indirect	No eff.	-0.028
N concentration	No effects		
Numerical density	Direct	0.306	0.714

3.3.2 | Direct and indirect effects of local predictors

Among predictors varying within site, shoot density (i.e. *Sphagnum* capitulum density) had strong positive effects on NPP in both species. Vascular plant cover had negative effects on growth in all models regardless of species, both directly and via negative effects on shoot density. Similarly, increasing HWT decreased growth through effects on vascular plant cover. However, the positive effect HWT had on shoot density for *S. fuscum* compensated for the negative effect through increased vascular plant cover. Moss tissue N concentration had a weak positive effect on LI but not on NPP.

4 | DISCUSSION

4.1 | Climatic predictors

Precipitation was expected to increase growth (Asada et al., 2003; Backéus, 1988; Bragazza et al., 2016; Gunnarsson, 2005; Moore, 1989) as it helps the poikilohydric *Sphagnum* maintain a sufficient moisture content to remain active and photosynthesise (Schipperges & Rydin, 1998). In our study the effects of precipitation were species-specific and a positive effect on growth was only observed for the wetter-growing species, *S. magellanicum*. However, more general support for the importance of precipitation was indicated in our data by a consistent negative effect on growth of *mean length of rain free periods* (cf. Krebs et al., 2016). Interception by vascular plants means that a minimum amount of precipitation per rain event is required to add moisture to *Sphagnum*; to account for this we counted days




FIGURE 6 Graphical representation of piecewise structural equation models using mixed effects models to describe effects of the environmental variables from the models in Table S4 on annual net primary productivity. Num density = numerical density of *Sphagnum* shoots, PAR = photosynthetic active radiation, No rain = mean length of rain free periods, HWT = height above the water table. White boxes are exclusively predictor variables, grey boxes are response (net primary production), or predictor and response. Arrowheads show the direction of effects and line thickness indicates the effect size. For simplicity, arrows associated with $p \geq 0.1$ have been removed. See Table S8 for complete model results

with precipitation <1 mm as rain free (Farrick & Price, 2009; Price, Rochefort, & Quinty, 1998). Hence, our results imply that extreme dry-periods affect *Sphagnum* growth but that the amount of precipitation is a better predictor of growth in non-hummock species.

Evaporation is often incorporated with precipitation into a moisture index to reflect water availability. We calculated the amount

of available water ($P-E$), but our results corroborate those of Loisel et al. (2012) who found no effect of this differential on growth. Keeping precipitation and evaporation as separate predictors resulted in a better fitting model.

Temperature is a major climatic driver of plant production. There is a mechanistic relationship between temperature and photosynthesis (Farquhar, von Caemmerer, & Berry, 1980), exhibiting a unimodal pattern where temperature optimum seems to be rather high in *Sphagnum* (Harley, Tenhunen, Murray, & Beyers, 1989; Skre & Oechel, 1981). As expected, our results consistently show that warmer temperatures result in increased growth, but this effect was weaker in *S. fuscum* than in *S. magellanicum*. In fact, the summed effect of temperature on NPP in *S. fuscum* was zero due to the indirect effect of reduced density. The strong temperature effect on LI in *S. magellanicum* illustrates that this species can grow well in length even with little increase in biomass accumulation (Mazziotta, Granath, Rydin, Bengtsson, & Norberg, 2019), the result of which would be patches with low bulk density. For *S. fuscum* this would not be a viable strategy since maintaining high BD is crucial to maintaining a high water content (McCarter & Price, 2014; Nijp et al., 2014).

PAR is strongly related to temperature and directly controls NPP (Chapin III, Matson, & Vitousek, 2011) because PAR is a key driver of carbon fixation in plants. Using PAR as the only predictor aside from year and species, we found some support for Loisel, Gallego-Sala, and Yu (2012) results that PAR is an important driver of NPP in *Sphagnum*. When controlling for other factors (i.e. our full multivariable models), there was no or little effect of PAR on length growth, but still a positive effect on annual NPP. This indicates that other variables, most notably temperature, with which PAR was correlated and which also control evaporation, were the main drivers of growth.

4.2 | Effects related to vascular plant cover

Low levels of *N deposition* can increase *Sphagnum* growth in N-limited systems (Limpens et al., 2011), but evidence of large-scale negative effects of greater N deposition on *Sphagnum* growth are lacking. Experimental evidence shows that high levels of N deposition will promote vascular plant growth, and thereby intensify the effects of competition for light from vascular plants (Berendse et al., 2001; Bubier, Moore, & Bledzki, 2007; Limpens et al., 2011). Indeed, one of our most consistent results was the negative effect of vascular plant cover on all growth responses and N deposition was associated with higher vascular plant cover in the *S. magellanicum* patches. However, the summed impact of N deposition on growth showed a positive effect on length growth due to higher tissue N concentration. The positive effect of N deposition is consistent with the positive low-dose response reported from experiments (Juutinen et al., 2015; Limpens et al., 2011), but we could detect no such effect on NPP. It is possible that length growth is promoted under elevated N availability to compete with vascular plants at the cost of less dense *Sphagnum* mats.

Water table position has frequently been associated with *Sphagnum* growth, suggesting that a more surficial water table promotes

Sphagnum growth (Gunnarsson, 2005; Li, Glime, & Liao, 1992). Our models did indicate a weak negative effect on growth with deep water tables through indirect effects (vascular plant cover and shoot density). Interestingly, for *S. fuscum* there was a weak positive effect of HWT on NPP due to higher shoot density. Likely, the observed weak effect of water table on growth cannot be easily disentangled from the species effect, as the HWT effect appears stronger between species than within species (Bengtsson, Granath, & Rydin, 2016). These findings suggest that the effects reported in experimental studies (e.g. Li et al., 1992) are weaker in the field. It is possible that strong effects are only apparent during a short period of the season (Walker et al., 2017), or in dry years when the water table is significantly lowered for an extended period (Bragazza, 2008). Unless the drought is severe, hydrological feedbacks will maintain sufficiently high moisture content for photosynthesis across the water table gradient within a species niche (Waddington et al., 2015).

4.3 | Comparing species, years and growth responses

The two species show large differences in LI and bulk density, but less so in NPP. In fact, NPP was higher in *S. fuscum* even though this species had lower LI. This has been reported previously (Lindholm & Vasander, 1990) and is a result of the high BD in *S. fuscum*, and NPP being the product of LI and BD. The underlying mechanism behind the rather similar NPP is a trade-off between maximizing length growth and maintaining high water content (Laing, Granath, Belyea, Allton, & Rydin, 2014; Mazziotta et al., 2019). Generally, *S. fuscum* grows in higher hummocks, where its higher bulk density and shoot density increases capillary water uptake and reduces evaporative loss. *Sphagnum magellanicum* on the other hand, mainly grows in lawn habitats and relies on being closer to the water table to maintain a high water content. Hence, *S. fuscum* is less dependent on a continuous wet climate than other peat mosses, and can remain photosynthetically active during drought events and during dry years (Rydin, 1993). This can explain the stronger effect of precipitation and temperature on *S. magellanicum* and it follows that *S. fuscum* has a more stable growth and is more resistant to changes in climate, while *S. magellanicum* can take advantage of higher amounts of rain and grow faster when rain events occur (Mazziotta et al., 2019). This is in line with a paleoecological study showing that wetter climate caused a vegetation dominance shift from *S. fuscum* to *S. magellanicum* with subsequent reduction of carbon sequestration (Belyea & Malmer, 2004). The results may be affected by ecological differences between the newly described taxa within *S. magellanicum*. The two, *S. medium* and *S. divinum*, have been described to occur in contrasting habitats (Yousefi et al., 2017). However, we have observed both species in open as well as shaded habitats (Bengtsson, Granath, Cronberg, & Rydin, 2020). The ecological boundaries between these species need to be further studied, since it seems that both could be plastic enough to occur in either environment.

Our models for LI explained more variance than those for NPP. LI varied less between the two years than NPP, which depended

on stem BD. Hence NPP is also somewhat affected by the previous year's growth. Often, in multi-year studies on *Sphagnum* NPP, a constant BD is assumed, even though it may differ between years. We collected data on BD for each year, which resulted in a higher between year NPP correlation (0.48–0.58) than often reported (e.g. Lindholm & Vasander, 1990). It is clear that annual NPP can be challenging to measure accurately in *Sphagnum* and perhaps a time-lag model is required to better capture the environmental effects (e.g. Back us, 1988 included previous year's weather for predictions).

4.4 | Growth models

When *Sphagnum* moss growth is modelled, HWT is regularly used as the main predictor, and climatic factors are reflected indirectly as changes in HWT (e.g. The Holocene Peat Model, Frothingham et al., 2010). Process-based modules, similar to those used for vascular plants, are rare for mosses in large-scale models but there are some attempts for *Sphagnum* (e.g. Walker et al., 2017). In our models we focus on predictors with documented mechanistic roles. This is also the approach used in previous attempts to understand broad patterns of *Sphagnum* productivity (Table 1). Comparisons are difficult as most of these studies report only one main predictor that fitted the data best. If and how other variables were tested is often not mentioned in these papers. Because we sampled *Sphagnum* growth in the same two species from a broad geographical range, within narrow habitat definitions and with a common methodology, our study is better suited to identify the strongest climatic and environmental predictors for growth. Our dataset and analyses constitute an important step to inform current and future ecosystem models. Indeed, we are able to highlight the important difference between length growth and NPP, and provide direction for what variables to include in ecosystem models. In addition, our study enables comparison and parameterization of local drivers, such as HWT and vascular plant cover, which are two central factors in peatland models (Hayward & Clymo, 1983).

Our models for daily productivity showed the same significant predictors as for the entire growing season, except for the effect of PAR. Using daily growth we could investigate if effects were solely driven by conditions during the growing season, and the observed effect of PAR (summed over the season) supports the previous studies that have highlighted growing season length as an important factor for terrestrial ecosystems in general (Michaletz, Kerkhoff, & Enquist, 2018), and specifically for peatlands (Charman et al., 2013; Koebsch et al., 2020; Loisel et al., 2012). Still, our results suggest that climatic variables do affect growth directly and indirectly, and that growing season length may not be the main driver.

By measuring across the Holarctic we covered a wide range of environmental conditions. However, only 2 years growth limits our ability to study within-site variability and more consecutive years of measurements is needed to understand this variation, and to test if proposed mechanisms hold within a site or region.

4.5 | Implications for peatlands

The scale of our study and the consistent design provide a unique dataset to explore the drivers of growth of two common *Sphagnum* species across Holarctic peatlands. We conclude that variation in growth often is as large within as between peatlands, highlighting the importance of both global climatic factors (precipitation and mean temperature) and local factors (here vascular plant cover). However, it was challenging to determine which climatic or other environmental variables influenced the growth of *Sphagnum*, since the most important variables covaried. Surprisingly, water table depth was not a good predictor of growth within a species.

Undoubtedly, the growth of *Sphagnum* and hence peatland function will be affected by environmental change. *Sphagnum magellanicum* generally has a wider niche in relation to water table level and nutrients, and showed a stronger direct response in LI to precipitation and temperature, which can give it a competitive advantage in warmer and wetter environments, than *S. fuscum*, which engineers an environment that is more resistant to desiccation (Bengtsson, Granath, et al., 2020). However, due to indirect effects and differences in bulk density the responses of the two species in productivity (NPP) were more similar, suggesting that the response of peatland moss NPP to climate change may be robust to species turnover (cf. Jassey & Signarbieux, 2019; Robroek et al., 2017).

ACKNOWLEDGEMENTS

We dedicate this work to the memory of Maria Noskova and Richard Payne, collaborators on this project who tragically passed away before the submission of this paper. The project was supported by the Swedish Research Council (2015-05174), the Russian Science Foundation (grant 19-14-00102), the Russian Foundation for Basic Research (research projects nos. 14-05-00775, 15-44-00091, 19-05-00830, 18-04-00988 and 18-44-860017), University of Ferrara (FAR 2013 and 2014), the Polish National Centre for Research and Development (within the Polish-Norwegian Research Programme: the project WETMAN (Central European Wetland Ecosystem Feedbacks to Changing Climate Field Scale Manipulation, Project ID: 203258), the National Science Centre, Poland (ID: 2015/17/B/ST10/01656), institutional research funds from the Estonian Ministry of Education and Research (grant IUT34-7), the Natural Sciences and Engineering Research Council of Canada, the Czech Science Foundation (Michal Hájek; Centre for European Vegetation Syntheses CEVS—project of Czech Science Foundation no. 19-28491X), the National Natural Science Foundation of China (No. 41471043 and No. 41871046), Jilin Provincial Science and Technology Development Project (20190101025JH), Academy of Finland (Project code 287039), the Ministry of Science and Higher Education of the Russian Federation (according to the state assignment of ISSA SB RAS), the Swiss National Science Foundation projects no. P2NEP3_178543 and with generous support awarded to Lorna I. Harris from the W. Garfield Weston Foundation Fellowship for Northern Conservation, administered by Wildlife

Conservation Society (WCS) Canada, National Science Foundation (NSF-1312402) to S.M.N., and Extensius to F.B. We acknowledge the Adirondack and Maine offices of The Nature Conservancy, the Autonomous Province of Bolzano (Italy), Staatsbosbeheer and Landschap Overijssel (the Netherlands), the Greenwoods Conservancy, NY and the Orono Bog Boardwalk, University of Maine for access to field sites.

AUTHORS' CONTRIBUTIONS

G.G. and H.R. initiated and designed the study; All authors collected data, and F.B. and G.G. performed the analyses; F.B. wrote the first draft and all authors commented on the manuscript.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/1365-2745.13499>.

DATA AVAILABILITY STATEMENT

Data and R-script available from the Dryad Digital Repository <https://doi.org/10.5061/dryad.1ns1rn8rm> (Bengtsson, Rydin, et al., 2020).

ORCID

Fia Bengtsson  <https://orcid.org/0000-0003-0497-8316>
 Håkan Rydin  <https://orcid.org/0000-0002-7582-3998>
 Jennifer L. Baltzer  <https://orcid.org/0000-0001-7476-5928>
 Zhao-Jun Bu  <https://orcid.org/0000-0001-9710-6343>
 Ellen Dorrepaal  <https://orcid.org/0000-0002-0523-2471>
 Olga Galanina  <https://orcid.org/0000-0001-5723-309X>
 Mariusz Gałka  <https://orcid.org/0000-0001-8906-944X>
 Michal Hájek  <https://orcid.org/0000-0002-5201-2682>
 Akira Haraguchi  <https://orcid.org/0000-0002-5010-521X>
 Lorna I. Harris  <https://orcid.org/0000-0002-2637-4030>
 Elyn Humphreys  <https://orcid.org/0000-0002-5397-2802>
 Martin Jiroušek  <https://orcid.org/0000-0002-4293-478X>
 Edgar Karofeld  <https://orcid.org/0000-0001-6533-8473>
 Natalia G. Koronátova  <https://orcid.org/0000-0002-0557-0083>
 Anna M. Laine  <https://orcid.org/0000-0003-2989-1591>
 Mariusz Lamentowicz  <https://orcid.org/0000-0003-0429-1530>
 Elena Lapshina  <https://orcid.org/0000-0001-5571-7787>
 Juul Limpens  <https://orcid.org/0000-0001-5779-0304>
 Edward A. D. Mitchell  <https://orcid.org/0000-0003-0358-506X>
 Tariq M. Munir  <https://orcid.org/0000-0002-4591-0978>
 Susan M. Natali  <https://orcid.org/0000-0002-3010-2994>
 Dmitriy A. Philippov  <https://orcid.org/0000-0003-3075-1959>
 Sean Robinson  <https://orcid.org/0000-0003-0961-4968>
 Bjorn J. M. Robroek  <https://orcid.org/0000-0002-6714-0652>
 Line Rochefort  <https://orcid.org/0000-0001-8450-0376>
 David Singer  <https://orcid.org/0000-0002-4116-033X>
 Hans K. Stenøien  <https://orcid.org/0000-0003-2191-332X>
 Eeva-Stiina Tuittila  <https://orcid.org/0000-0001-8861-3167>
 James Michael Waddington <https://orcid.org/0000-0002-0317-7894>
 Gustaf Granath <https://orcid.org/0000-0002-3632-9102>

REFERENCES

- Asada, T., Warner, B. G., & Banner, A. (2003). Growth of mosses in relation to climate factors in a hypermaritime coastal peatland in British Columbia, Canada. *The Bryologist*, 106(4), 516–527. [https://doi.org/10.1639/0007-2745\(2003\)106\[516:GOMIRT\]2.0.CO;2](https://doi.org/10.1639/0007-2745(2003)106[516:GOMIRT]2.0.CO;2)
- Backéus, I. (1988). Weather variables as predictors of *Sphagnum* growth on a bog. *Holarctic Ecology*, 11, 146–150. <https://doi.org/10.1111/j.1600-0587.1988.tb00793.x>
- Bartoň, K. (2018). *MuMIn: Multi-model inference*. R package version 1.42.1. Retrieved from <https://CRAN.R-project.org/package=MuMIn>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bechtold, M., De Lannoy, G. J. M., Koster, R. D., Reichle, R. H., Mahanama, S. P., Bleuten, W., ... Devito, K. (2019). PEAT-CLSM: A specific treatment of peatland hydrology in the NASA catchment land surface model. *Journal of Advances in Modeling Earth Systems*, 11, 2130–2162.
- Belyea, L. R., & Malmer, N. (2004). Carbon sequestration in peatland: Patterns and mechanisms of response to climate change. *Global Change Biology*, 10(7), 1043–1052. <https://doi.org/10.1111/j.1529-8817.2003.00783.x>
- Bengtsson, F., Granath, G., Cronberg, N., & Rydin, H. (2020). Mechanisms behind species-specific water economy responses to water level drawdown in peat mosses. *Annals of Botany*, 126(2), 219–230. <https://doi.org/10.1093/aob/mcaa033>
- Bengtsson, F., Granath, G., & Rydin, H. (2016). Photosynthesis, growth, and decay traits in *Sphagnum* – A multispecies comparison. *Ecology and Evolution*, 6, 3325–3341. <https://doi.org/10.1002/ece3.2119>
- Bengtsson, F., Rydin, H., Baltzer, J. L., Bragazza, L., Bu, Z.-J., Caporn, S. J. M., ... Granath, G. (2020). Data from: Environmental drivers of *Sphagnum* growth in peatlands across the Holarctic region. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.1ns1rn8rm>
- Berendse, F., Van Breemen, N., Rydin, H. A., Buttler, A., Heijmans, M., Hoosbeek, M. R., ... Wallen, B. O. (2001). Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production in *Sphagnum* bogs. *Global Change Biology*, 7, 591–598. <https://doi.org/10.1046/j.1365-2486.2001.00433.x>
- Bragazza, L. (2008). A climatic threshold triggers the die-off of peat mosses during an extreme heat wave. *Global Change Biology*, 14, 2688–2695. <https://doi.org/10.1111/j.1365-2486.2008.01699.x>
- Bragazza, L., Buttler, A., Robroek, B. J. M., Albrecht, R., Zaccane, C., Jassey, V. E. J., & Signarbieux, C. (2016). Persistent high temperature and low precipitation reduce peat carbon accumulation. *Global Change Biology*, 22, 4114–4123. <https://doi.org/10.1111/gcb.13319>
- Bubier, J. L., Moore, T. R., & Bledzki, L. A. (2007). Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog. *Global Change Biology*, 13, 1168–1186. <https://doi.org/10.1111/j.1365-2486.2007.01346.x>
- Chapin III, F. S., Matson, P. A., & Vitousek, P. M. (2011). *Principles of terrestrial ecosystem ecology* (2nd ed.). New York: Dordrecht, Heidelberg, London: Springer.
- Charman, D. J., Beilman, D. W., Blaauw, M., Booth, R. K., Brewer, S., Chambers, F. M., ... Zhao, Y. (2013). Climate-related changes in peatland carbon accumulation during the last millennium. *Biogeosciences*, 10, 929–944. <https://doi.org/10.5194/bg-10-929-2013>
- Chong, M., Humphreys, E., & Moore, T. R. (2012). Microclimatic response to increasing shrub cover and its effect on *Sphagnum* CO₂ exchange in a bog. *Ecoscience*, 19, 89–97. <https://doi.org/10.2980/19-1-3489>
- Clymo, R. S. (1970). The growth of *Sphagnum*: Methods of measurement. *Journal of Ecology*, 58, 13–49. <https://doi.org/10.2307/2258168>
- Farquhar, G. D., von Caemmerer, S., & Berry, J. A. (1980). A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta*, 149, 78–90. <https://doi.org/10.1007/BF00386231>
- Farrick, K. K., & Price, J. S. (2009). Ericaceous shrubs on abandoned block-cut peatlands: Implications for soil water availability and *Sphagnum* restoration. *Ecohydrology*, 2, 530–540. <https://doi.org/10.1002/eco.77>
- Fox, J., & Weisberg, S. (2011). *An R companion to applied regression* (2nd ed.). Thousand Oaks, CA: Sage. Retrieved from <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>
- Frolking, S., Roulet, N. T., Tuittila, E.-S., Bubier, J. L., Quillet, A., Talbot, J., & Richard, P. J. H. (2010). A new model of Holocene peatland net primary production, decomposition, water balance, and peat accumulation. *Earth System Dynamics*, 1, 1–21. <https://doi.org/10.5194/esd-1-1-2010>
- Gallego-Sala, A. V., Charman, D. J., Brewer, S., Page, S. E., Prentice, I. C., Friedlingstein, P., ... Zhao, Y. (2018). Latitudinal limits to the predicted increase of the peatland carbon sink with warming. *Nature Climate Change*, 8, 907–913. <https://doi.org/10.1038/s41558-018-0271-1>
- Gavazov, K., Albrecht, R., Buttler, A., Dorrepaal, E., Garnett, M. H., Gogo, S., ... Bragazza, L. (2018). Vascular plant-mediated controls on atmospheric carbon assimilation and peat carbon decomposition under climate change. *Global Change Biology*, 24, 3911–3921. <https://doi.org/10.1111/gcb.14140>
- Global Modeling and Assimilation Office (GMAO). (2015). *MERRA-2 tavgM_2d_csp_Nx: 2d, Monthly mean, Time-averaged, Single-level, Assimilation, COSP satellite simulator V5.12.4*, Greenbelt, MD: Goddard Earth Sciences Data and Information Services Center (GES DISC). <https://doi.org/10.5067/BZPOTGJOQKLU>
- Granath, G., Rydin, H., Baltzer, J. L., Bengtsson, F., Boncek, N., Bragazza, L., ... Rice, S. K. (2018). Environmental and taxonomic controls of carbon and oxygen stable isotope composition in *Sphagnum* across broad climatic and geographic ranges. *Biogeosciences*, 15, 5189–5202. <https://doi.org/10.5194/bg-15-5189-2018>
- Granath, G., Strengbom, J., & Rydin, H. (2012). Direct physiological effects of nitrogen on *Sphagnum*: A greenhouse experiment. *Functional Ecology*, 26, 353–364. <https://doi.org/10.1111/j.1365-2435.2011.01948.x>
- Gunnarsson, U. (2005). Global patterns of *Sphagnum* productivity. *Journal of Bryology*, 27, 269–279. <https://doi.org/10.1179/174328205x70029>
- Harley, P. C., Tenhunen, J. D., Murray, K. J., & Beyers, J. (1989). Irradiance and temperature effects on photosynthesis of tussock tundra *Sphagnum* mosses from the foothills of the Philip Smith Mountains, Alaska. *Oecologia*, 79, 251–259. <https://doi.org/10.1007/BF00388485>
- Hassel, K., Kyrkjeeide, M. O., Yousefi, N., Prestø, T., Stenøien, H. K., Shaw, J. A., & Flatberg, K. I. (2018). *Sphagnum divinum* (sp. nov.) and *S. medium* Limpr. and their relationship to *S. magellanicum* Brid. *Journal of Bryology*, 40, 197–222. <https://doi.org/10.1080/03736687.2018.1474424>
- Hayward, P. M., & Clymo, R. S. (1983). The growth of *Sphagnum*: Experiments on, and simulation of, some effects of light flux and water-table depth. *The Journal of Ecology*, 71(3), 845–863. <https://doi.org/10.2307/2259597>
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978. <https://doi.org/10.1002/joc.1276>
- IPCC. (2013). *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge University Press.
- Jassey, V. E. J., & Signarbieux, C. (2019). Effects of climate warming on *Sphagnum* photosynthesis in peatlands depend on peat moisture and species-specific anatomical traits. *Global Change Biology*, 25, 3859–3870. <https://doi.org/10.1111/gcb.14788>
- Johnson, P. C. (2014). Extension of Nakagawa & Schielzeth's R²GLMM to random slopes models. *Methods in Ecology and Evolution*, 5(9), 944–946. <https://doi.org/10.1111/2041-210X.12225>

- Juutinen, S., Moore, T. R., Laine, A. M., Bubier, J. L., Tuittila, E.-S., De Young, A., & Chong, M. (2015). Responses of the mosses *Sphagnum capillifolium* and *Polytrichum strictum* to nitrogen deposition in a bog: Growth, ground cover, and CO₂ exchange. *Botany-Botanique*, 94, 127–138. <https://doi.org/10.1139/cjb-2015-0183>
- Koebisch, F., Sonnentag, O., Järveoja, J., Peltoniemi, M., Alekseychik, P., Aurela, M., ... Peichl, M. (2020). Refining the role of phenology in regulating gross ecosystem productivity across European peatlands. *Global Change Biology*, 26, 876–887. <https://doi.org/10.1111/gcb.14905>
- Krebs, M., Gaudig, G., & Joosten, H. (2016). Record growth of *Sphagnum papillosum* in Georgia (Transcaucasus): Rain frequency, temperature and microhabitat as key drivers in natural bogs. *Mires and Peat*, 18(04), 1–16. <https://doi.org/10.19189/MaP.2015.OMB.190>
- Kyrkjede, M. O., Hassel, K., Stenøien, H. K., Prestø, T., Boström, E., Shaw, A. J., & Flatberg, K. I. (2015). The dark morph of *Sphagnum fuscum* (Schimp.) H. Klinggr. in Europe is conspecific with the North American *S. beothuk*. *Journal of Bryology*, 37, 251–266. <https://doi.org/10.1179/1743282015Y.00000000020>
- Laing, C. G., Granath, G., Belyea, L. R., Allton, K. E., & Rydin, H. (2014). Tradeoffs and scaling of functional traits in *Sphagnum* as drivers of carbon cycling in peatlands. *Oikos*, 123, 817–828. <https://doi.org/10.1111/oik.01061>
- Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C.-U., Shaw, M., Vet, R., ... Nolan, M. (2013). Multi-model mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): Evaluation of historical and projected future changes. *Atmospheric Chemistry and Physics*, 13, 7997–8018. <https://doi.org/10.5194/acp-13-7997-2013>
- Lefcheck, J. S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution*, 7, 573–579. <https://doi.org/10.1111/2041-210X.12512>
- Li, Y., Glime, J. M., & Liao, C. (1992). Responses of two interacting *Sphagnum* species to water level. *Journal of Bryology*, 17, 59–70. <https://doi.org/10.1179/jbr.1992.17.1.59>
- Limpens, J., Granath, G., Gunnarsson, U., Aerts, R., Bayley, S., Bragazza, L., ... Xu, B. (2011). Climatic modifiers of the response to nitrogen deposition in peat-forming *Sphagnum* mosses: A meta-analysis. *New Phytologist*, 191, 496–507. <https://doi.org/10.1111/j.1469-8137.2011.03680.x>
- Lindholm, T., & Vasander, H. (1990). Production of eight species of *Sphagnum* at Suurisuo mire, southern Finland. *Annales Botanici Fennici*, 27, 145–157.
- Loisel, J., Gallego-Sala, A. V., & Yu, Z. (2012). Global-scale pattern of peatland *Sphagnum* growth driven by photosynthetically active radiation and growing season length. *Biogeosciences*, 9, 2737–2746. <https://doi.org/10.5194/bg-9-2737-2012>
- Loisel, J., Yu, Z., Beilman, D. W., Camill, P., Alm, J., Amesbury, M. J., ... Zhou, W. (2014). A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation. *The Holocene*, 24, 1028–1042. <https://doi.org/10.1177/0959683614538073>
- Mazziotta, A., Granath, G., Rydin, H., Bengtsson, F., & Norberg, J. (2019). Scaling functional traits to ecosystem processes: Towards a mechanistic understanding in peat mosses. *Journal of Ecology*, 107, 843–859. <https://doi.org/10.1111/1365-2745.13110>
- McCarter, C. P. R., & Price, J. S. (2014). Ecohydrology of *Sphagnum* moss hummocks: Mechanisms of capitula water supply and simulated effects of evaporation. *Ecohydrology*, 7, 33–44. <https://doi.org/10.1002/eco.1313>
- Michaletz, S. T., Kerkhoff, A. J., & Enquist, B. J. (2018). Drivers of terrestrial plant production across broad geographical gradients. *Global Ecology and Biogeography*, 27, 166–174. <https://doi.org/10.1111/geb.12685>
- Moore, T. R. (1989). Growth and net production of *Sphagnum* at five fen sites, subarctic eastern Canada. *Canadian Journal of Botany*, 67, 1203–1207. <https://doi.org/10.1139/b89-156>
- Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2), 133–142. <https://doi.org/10.1111/j.2041-210x.2012.00261.x>
- Nichols, J. E., & Peteet, D. M. (2019). Rapid expansion of northern peatlands and doubled estimate of carbon storage. *Nature Geoscience*, 12, 917–921. <https://doi.org/10.1038/s41561-019-0454-z>
- Nijp, J. J., Limpens, J., Metselaar, K., van der Zee, S. E. A. T. M., Berendse, F., & Robroek, B. J. M. (2014). Can frequent precipitation moderate the impact of drought on peatmoss carbon uptake in northern peatlands? *New Phytologist*, 203, 70–80. <https://doi.org/10.1111/nph.12792>
- Price, J., Rochefort, L., & Quinty, F. (1998). Energy and moisture considerations on cutover peatlands: Surface microtopography, mulch cover and *Sphagnum* regeneration. *Ecological Engineering*, 10, 293–312. [https://doi.org/10.1016/S0925-8574\(98\)00046-9](https://doi.org/10.1016/S0925-8574(98)00046-9)
- Qiu, C., Zhu, D., Ciais, P., Guenet, B., Krinner, G., Peng, S., ... Ziemblinska, K. (2018). ORCHIDEE-PEAT (revision 4596), a model for northern peatland CO₂, water, and energy fluxes on daily to annual scales. *Geoscientific Model Development*, 11, 497–519. <https://doi.org/10.5194/gmd-11-497-2018>
- R Core Team. (2017). *R: A language and environment for statistical computing*. R version 3.4.0. Vienna, Austria: Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Ricklefs, R. E. (2008). *The economy of nature* (6th ed.). New York, NY: W. H. Freeman and Company.
- Robroek, B. J. M., Jassey, V. E. J., Payne, R. J., Martí, M., Bragazza, L., Bleeker, A., ... Verhoeven, J. T. A. (2017). Taxonomic and functional turnover are decoupled in European peat bogs. *Nature Communications*, 8, 1161. <https://doi.org/10.1038/s41467-017-01350-5>
- Rydin, H. (1993). Mechanisms of interactions among *Sphagnum* species along water-level gradients. *Advances in Bryology*, 5, 153–185.
- Rydin, H., & Jeglum, J. K. (2013). *The biology of peatlands* (2nd ed.). Oxford, UK: Oxford University Press.
- Schipperges, B., & Rydin, H. (1998). Response of photosynthesis of *Sphagnum* species from contrasting microhabitats to tissue water content and repeated desiccation. *New Phytologist*, 140(4), 677–684. <https://doi.org/10.1046/j.1469-8137.1998.00311.x>
- Schloss, A. I., Kicklighter, D. W., Kaduk, J., Wittenberg, U., & The Participants of the Potsdam NPP Model Intercomparison. (2001). Comparing global models of terrestrial net primary productivity (NPP): Comparison of NPP to climate and the Normalized Difference Vegetation Index (NDVI). *Global Change Biology*, 5, 25–34. <https://doi.org/10.1046/j.1365-2486.1999.00004.x>
- Shaw, A. J., Cox, C. J., Buck, R. W., Devos, N., Buchanan, A. M., Cave, L., ... Temsch, E. M. (2010). Newly resolved relationships in an early land plant lineage: Bryophyta class Sphagnopsida (peat mosses). *American Journal of Botany*, 97(9), 1511–1531. <https://doi.org/10.3732/ajb.1000055>
- Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*, 90, 363–368. <https://doi.org/10.1890/08-1034.1>
- Skre, O., & Oechel, W. C. (1981). Moss functioning in different taiga ecosystems in interior Alaska. *Oecologia*, 48, 50–59. <https://doi.org/10.1007/BF00346987>
- Turetsky, M. R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Froliking, S., McGuire, A. D., & Tuittila, E.-S. (2012). The resilience and functional role of moss in boreal and arctic ecosystems. *New Phytologist*, 196, 49–67. <https://doi.org/10.1111/j.1469-8137.2012.04254.x>
- Waddington, J. M., Morris, P. J., Kettridge, N., Granath, G., Thompson, D. K., & Moore, P. A. (2015). Hydrological feedbacks in northern peatlands. *Ecohydrology*, 8, 113–127. <https://doi.org/10.1002/eco.1493>
- Walker, A. P., Carter, K. R., Gu, L., Hanson, P. J., Malhotra, A., Norby, R. J., ... Weston, D. J. (2017). Biophysical drivers of seasonal variability in *Sphagnum* gross primary production in a northern temperate

- bog. *Journal of Geophysical Research Biogeosciences*. <https://doi.org/10.1002/2016JG003711>
- Yousefi, N., Hassel, K., Flatberg, K. I., Kemppainen, P., & Trucchi, E., Shaw, A. J., ... Stenøien, H. K. (2017). Divergent evolution and niche differentiation within the common peatmoss *Sphagnum magellanicum*. *American Journal of Botany*, 104(7), 1060–1072. <https://doi.org/10.3732/ajb.1700163>
- Yu, Z. (2012). Northern peatland carbon stocks and dynamics: A review. *Biogeosciences*, 9, 4071–4085. <https://doi.org/10.5194/bg-9-4071-2012>
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., & Hunt, S. J. (2010). Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL043584>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Bengtsson F, Rydin H, Baltzer JL, et al. Environmental drivers of *Sphagnum* growth in peatlands across the Holarctic region. *J Ecol.* 2020;00:1–15. <https://doi.org/10.1111/1365-2745.13499>