

See corresponding editorial on page 833.

Insects are a viable protein source for human consumption: from insect protein digestion to postprandial muscle protein synthesis in vivo in humans: a double-blind randomized trial

Wesley JH Hermans,¹ Joan M Senden,¹ Tyler A Churchward-Venne,¹ Kevin JM Paulussen,¹ Cas J Fuchs,¹ Joey SJ Smeets,¹ Joop JA van Loon,² Lex B Verdijk,¹ and Luc JC van Loon¹

¹Department of Human Biology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre+, Maastricht, The Netherlands; and ²Laboratory of Entomology, Plant Sciences Group, Wageningen University, Wageningen, The Netherlands

ABSTRACT

Background: Insects have recently been identified as a more sustainable protein-dense food source and may represent a viable alternative to conventional animal-derived proteins.

Objectives: We aimed to compare the impacts of ingesting lesser mealworm- and milk-derived protein on protein digestion and amino acid absorption kinetics, postprandial skeletal muscle protein synthesis rates, and the incorporation of dietary protein-derived amino acids into de novo muscle protein at rest and during recovery from exercise in vivo in humans.

Methods: In this double-blind randomized controlled trial, 24 healthy, young men ingested 30 g specifically produced, intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm- or milk-derived protein after a unilateral bout of resistance-type exercise. Primed continuous L-[ring-²H₅]-phenylalanine, L-[ring-3,5-²H₂]-tyrosine, and L-[1-¹³C]-leucine infusions were applied, with frequent collection of blood and muscle tissue samples.

Results: A total of 73% ± 7% and 77% ± 7% of the lesser mealworm and milk protein-derived phenylalanine was released into the circulation during the 5 h postprandial period, respectively, with no significant differences between groups (*P* < 0.05). Muscle protein synthesis rates increased after both lesser mealworm and milk protein concentrate ingestion from 0.025 ± 0.008%/h to 0.045 ± 0.017%/h and 0.028 ± 0.010%/h to 0.056 ± 0.012%/h at rest and from 0.025 ± 0.012%/h to 0.059 ± 0.015%/h and 0.026 ± 0.009%/h to 0.073 ± 0.020%/h after exercise, respectively (all *P* < 0.05), with no differences between groups (both *P* > 0.05). Incorporation of mealworm and milk protein-derived L-[1-¹³C]-phenylalanine into de novo muscle protein was greater after exercise than at rest (*P* < 0.05), with no differences between groups (*P* > 0.05).

Conclusions: Ingestion of a meal-like amount of lesser mealworm-derived protein is followed by rapid protein digestion and amino acid absorption and increases muscle protein synthesis rates both at rest and during recovery from exercise. The postprandial protein handling

of lesser mealworm does not differ from ingesting an equivalent amount of milk protein concentrate in vivo in humans. This trial was registered at www.trialregister.nl as NL6897. *Am J Clin Nutr* 2021;114:934–944.

Keywords: milk protein, FSR, stable isotopes, protein metabolism, intrinsically labeled protein

Introduction

It has been well established that ingestion of protein increases muscle protein synthesis rates (1). The anabolic properties of various protein sources (2–5) are largely defined by their protein digestion and amino acid absorption kinetics (3, 6–8) and amino acid composition (6, 9, 10). Physical activity and

Supported by an in-kind contribution in the form of stable isotope tracers from Cambridge Isotope Laboratories for the production of intrinsically labeled lesser mealworm.

JJAvL is on the Board of Advisors for Protifarm. Protifarm produces large quantities of sustainable, high-quality insect ingredients for the food and pharmaceutical industry and is based in Ermelo, Netherlands.

Supplemental Methods are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/ajcn/>.

Address correspondence to LJCvL (e-mail: l.vanloon@maastrichtuniversity.nl).

Abbreviations used: EAA, essential amino acid; En%, energy percent; FSR, fractional synthesis rate; iAUC, incremental AUC; MILK, 30 g milk protein concentrate; MPE, mole percentage excess; NEAA, nonessential amino acid; *R*_a, rate of appearance; *R*_d, rate of disappearance; TAA, total amino acid; WORM, 30 g lesser mealworm-derived protein; 1-RM, 1 repetition maximum.

Received October 20, 2020. Accepted for publication March 16, 2021.

First published online May 21, 2021; doi: <https://doi.org/10.1093/ajcn/nqab115>.

exercise performed before protein ingestion can further augment the muscle protein synthetic response to feeding (5, 11–13).

Dietary protein sources can be broadly classified as either animal- or plant-based proteins. Animal-based proteins are generally considered to have stronger anabolic properties than isolated plant-derived proteins (14). This has been attributed to the fact that isolated plant-derived proteins often show an incomplete amino acid profile with low amounts of leucine and/or insufficient lysine, histidine, or methionine (14). The few studies that have compared postprandial muscle protein synthesis rates after ingestion of isolated plant- and animal-derived proteins have reported a lesser increase in muscle protein synthesis rates after ingestion of wheat (15) and soy (10, 12, 16, 17) protein than after dairy protein. However, with regards to sustainability, the production of sufficient amounts of conventional animal-based proteins to meet future global food demands represents a challenge. Consequently, there is a growing interest in the definition of other, (more) sustainable protein sources. Edible insects have gained much interest as an alternative source of dietary protein that may be produced on a more viable and sustainable commercial scale and, as such, may contribute to ensuring global food security (18, 19).

Insects have a high protein content and their amino acid composition indicates that insects may provide a high-quality protein source that can stimulate postprandial muscle protein accretion similarly to high-quality animal-based protein sources (18, 20, 21). Lesser mealworm (*Alphitobius diaperinus* Panzer; Coleoptera: Tenebrionidae)-derived protein has an amino acid composition that closely resembles protein sources such as milk or meat (18, 22). So far, to our knowledge there are no data available on lesser mealworm protein digestion and amino acid absorption kinetics or on the postprandial anabolic response to lesser mealworm protein ingestion. We hypothesize that lesser mealworm protein is properly digested and absorbed, and has the capacity to increase muscle protein synthesis rates in vivo in humans both at rest and during recovery from exercise.

To test our hypotheses, we first fed lesser mealworm larvae with free L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine to produce intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm protein. Combining the ingestion of intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm or milk protein concentrate (23–25) with the intravenous infusion of L-[ring-²H₅]-phenylalanine and L-[1-¹³C]-leucine allowed us to quantify protein digestion and amino acid absorption kinetics, postprandial muscle protein synthesis rates, as well as the incorporation of dietary protein-derived amino acids into de novo muscle protein in vivo in humans (26).

Methods

Production of intrinsically labeled milk- and lesser mealworm-derived protein

Previously, we (25, 27, 28) and others (29–31) have produced different types of intrinsically labeled protein. Intrinsically labeled milk protein was produced by infusing lactating dairy cows with large quantities of L-[1-¹³C]-leucine and L-[1-¹³C]-phenylalanine. Throughout the stable isotope infusion milk

was subsequently collected, processed, and fractionated as described previously (25, 32, 33), providing intrinsically L-[1-¹³C]-phenylalanine [38.3 mole percentage excess (MPE)] and L-[1-¹³C]-leucine (10.8 MPE) labeled milk protein.

For the current study, we applied the same principle to produce intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm. A 2.4-kg batch of 21-d-old lesser mealworm larvae (*A. diaperinus*; Protifarm) were fed 2.4 kg chicken feed (mainly wheat, barley, sorghum, carrot pulp, and apple core) mixed with 16 g L-[1-¹³C]-leucine and 103 g L-[1-¹³C]-phenylalanine. After 7 d, when the larvae had significantly increased in size, a sample was taken for microbiological testing to confirm that the larvae met all chemical and bacteriological specifications for human consumption (Eurofins KBBL). The larvae were sieved from the feed substrate, deprived of food for 3 h to empty their gut content, and then sieved again. Subsequently, the larvae were blanched and freeze-dried for 48 h to provide ~800 g dry lesser mealworm. The freeze-dried lesser mealworm larvae were ground into a fine powder in an experimental kitchen with food-grade status (Human Nutrition Research Unit, Wageningen University). Subsequently, nitrogen content (Dumas method; 34) and amino acid composition (ultra-performance LC-MS; ACQUITY UPLC H-Class with QDa; Waters) of the freeze-dried lesser mealworm powder were measured. The L-[1-¹³C]-leucine, L-[1-¹³C]-phenylalanine, and L-[1-¹³C]-tyrosine enrichments of the freeze-dried lesser mealworm powder were measured in triplicate by GC-MS (Agilent 7890A GC/5975C; MSD) after overnight hydrolysis (9.3, 50.8, and 33.4 MPE, respectively).

Subjects

Twenty-four healthy, young men [mean \pm SD age: 23 \pm 3 y; BMI (in kg/m²): 23.1 \pm 2.7] were selected to participate in this study. **Table 1** details the characteristics of the subjects. The trial was conducted between February 2018 and May 2019 at Maastricht University Medical Centre+, in Maastricht, Netherlands. All subjects were informed of the nature and possible risks of the experimental procedures, before providing written informed consent. This study was approved by the Medical Ethical Committee of the Maastricht University Medical Centre+ and performed in conformance with the principles outlined in the Declaration of Helsinki for use of human subjects and tissue.

Pretesting

Volunteers aged between 18 and 35 y with a BMI between 18.5 and 30.0 kg/m² underwent medical screening to assess body mass, height, blood pressure, and body composition (via DXA; Discovery A, Hologic). The participants were deemed healthy and eligible to participate based upon their response to a medical questionnaire and screening results. Participants were excluded if they were smoking, using medication that affected protein metabolism, intolerant to the investigated proteins, performing structural resistance training, or exercising >5 times/wk. After initial screening participants were familiarized with the exercise testing protocol and exercise equipment. Participants underwent estimates of unilateral 1 repetition maximum (1-RM) strength on the supine leg press and seated leg extension (Technogym BV).

TABLE 1 Subjects' characteristics¹

| | WORM | MILK | <i>P</i> value |
|---------------------------------|-------------|-------------|----------------|
| Age, y | 24 ± 3 | 22 ± 3 | 0.156 |
| Body mass, kg | 77.6 ± 11.1 | 72.0 ± 8.6 | 0.179 |
| Lean body mass, kg | 60.4 ± 7.0 | 56.6 ± 5.6 | 0.395 |
| Appendicular lean mass, kg | 28.2 ± 3.3 | 26.4 ± 3.4 | 0.414 |
| Body fat, % | 20.1 ± 5.4 | 17.9 ± 3.9 | 0.693 |
| Height, m | 1.70 ± 0.07 | 1.80 ± 0.10 | 0.712 |
| BMI, kg/m ² | 24.2 ± 2.8 | 22.1 ± 2.2 | 0.061 |
| Systolic blood pressure, mm Hg | 123 ± 10 | 122 ± 7 | 0.827 |
| Diastolic blood pressure, mm Hg | 70 ± 8 | 69 ± 4 | 0.743 |
| Resting heart rate, bpm | 61 ± 8 | 58 ± 7 | 0.397 |
| 1-RM Leg press, kg | 143 ± 29 | 139 ± 28 | 0.694 |
| 1-RM Leg extension, kg | 62 ± 9 | 60 ± 11 | 0.410 |

¹Values are means ± SDs. *n* = 12/group; data for lean body mass, appendicular lean mass, and body fat are displayed for *n* = 10 in the WORM group. Data were analyzed using an independent-samples *t* test. No differences were detected between groups. MILK, 30 g milk protein concentrate; WORM, 30 g lesser mealworm-derived protein; 1-RM, 1-repetition maximum.

After a 1-legged warm-up on a cycle ergometer (Lode BV) at 75 W for 5 min with the assigned exercise leg, 10 submaximal repetitions were performed on the exercise machine. Next, the subject performed single repetitions with incremental weights for every repetition until failure. Between every attempt, a rest period of ≥2 min was taken and the 1-RM was typically reached between 5 and 7 attempts.

Study design

In this double-blind, randomized, parallel-group trial, subjects were randomly assigned to consume either 30 g intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled milk protein (MILK; *n* = 12) or 30 g intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm-derived protein (WORM; *n* = 12). Randomization was performed using a computerized random-number generator. An independent person was responsible for blinding and drink preparation. Within each treatment group the exercise leg was randomly assigned. In each group 6 subjects exercised with their nondominant and 6 with their dominant leg. All presented muscle data for the WORM group are based on *n* = 11, because of missing muscle biopsies in 1 of the participants. Data derived from the DXA scan were only available for *n* = 10 in the WORM group, because of logistical issues.

Diet and physical activity

All subjects refrained from any sort of strenuous physical activity or exercise for 3 d before the trial and kept their diet as consistent as possible for 2 d before the experiment. Mean ± SD habitual protein intake was 1.5 ± 0.6 g protein · kg body mass⁻¹ · d⁻¹. On the evening before the experimental trial at 20:00, all subjects consumed the same standardized meal (1.71 MJ/405 kcal) providing 22 energy percent (En%) protein, 53 En% carbohydrate, and 26 En% fat.

Intrinsically labeled protein

Subjects ingested 30 g intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm- or milk-derived

protein. To allow ingestion of 30 g protein, subjects ingested either 64 g powdered lesser mealworm or 40 g dried milk protein concentrate dissolved into 300 mL water. The nitrogen content of the ingested lesser mealworm and milk protein concentrate was 10% and 12%, respectively, and conversion factors of 4.76 and 6.25 were used to calculate the protein contents of the dried lesser mealworm (47%) and milk protein concentrate (75%), respectively (35). The dried lesser mealworm provided 312 kcal, with 46En%, 6En%, and 46 En% provided by protein, carbohydrate, and fat, respectively (Kreca). Milk protein concentrate provided 142 kcal, with 91En%, 6En%, and 5 En% provided by protein, carbohydrate, and fat, respectively (Friesland Campina). **Table 2** shows the amino acid composition of the lesser mealworm- and milk-derived protein. Drinks were prepared and provided to the participants in a nontransparent shaker bottle by an independent person. After ingestion, bottles were rinsed with 50 mL water and the remaining fluid was ingested to ensure that all protein was consumed.

Study protocol

At 07:30, after an overnight fast, subjects arrived at the laboratory by car or public transport. A catheter was inserted into an antecubital vein for stable isotope-labeled amino acid infusion. A second catheter was inserted into a dorsal hand vein of the contralateral arm and placed in a hot box (60°C) for arterialized blood sampling (36). After taking a baseline blood sample, the plasma amino acid pools were primed with a single dose of L-[ring-²H₅]-phenylalanine (2.0 μmol/kg), L-[ring-3,5-²H₂]-tyrosine (0.613 μmol/kg), and L-[1-¹³C]-leucine (3.99 μmol/kg), after which a continuous intravenous L-[ring-²H₅]-phenylalanine (0.05 μmol · kg⁻¹ · min⁻¹), L-[ring-3,5-²H₂]-tyrosine (0.015 μmol · kg⁻¹ · min⁻¹), and L-[1-¹³C]-leucine (0.1 μmol · kg⁻¹ · min⁻¹) infusion was initiated (*t* = -180 min). Subsequently, the subjects rested in a supine position for 120 min during which 2 additional arterialized blood samples were drawn (*t* = -120 and -60 min).

At *t* = -50 min, the exercise session was started. The exercise session consisted of 1-legged cycling at 75 W for 5 min followed by 5 sets of leg press and leg extension. For the first set, the workload was set at 50% of the predetermined 1-RM for 10

TABLE 2 Milk- and lesser mealworm-derived protein characteristics¹

| Component | WORM | MILK |
|---|------|------|
| Alanine, g | 2.3 | 1.0 |
| Arginine, g | 1.6 | 0.9 |
| Aspartic acid, g | 2.1 | 1.7 |
| Glutamic acid, g | 3.6 | 6.1 |
| Glycine, g | 1.6 | 0.5 |
| Histidine, g | 0.9 | 0.7 |
| Isoleucine, g | 0.9 | 1.0 |
| Leucine, g | 2.5 | 2.8 |
| Lysine, g | 1.8 | 2.2 |
| Methionine, g | 0.4 | 0.8 |
| Phenylalanine, g | 1.2 | 1.2 |
| Proline, g | 1.8 | 3.4 |
| Serine, g | 1.4 | 1.3 |
| Threonine, g | 1.2 | 1.1 |
| Tyrosine, g | 2.4 | 1.4 |
| Valine, g | 1.3 | 1.2 |
| EAA, g | 10.2 | 11.0 |
| NEAAs, g | 16.9 | 16.3 |
| TAA, g | 27.2 | 27.2 |
| L-[1- ¹³ C]-leucine, MPE | 9.3 | 10.9 |
| L-[1- ¹³ C]-phenylalanine, MPE | 50.8 | 38.3 |
| L-[1- ¹³ C]-tyrosine, MPE | 33.4 | 8.6 |

¹Asparagine, cysteine, glutamine, and tryptophan were not measured. EAA, essential amino acid; MILK, 30 g milk protein concentrate; MPE, mole percentage excess; NEAA, nonessential amino acid; TAA, total amino acid; WORM, 30 g lesser mealworm-derived protein.

repetitions. The following 3 sets were executed at 80% 1-RM for 8–10 repetitions, with the fifth set being performed until volitional failure. The subjects rested for 2 min between all sets and exercises and were verbally encouraged to complete the protocol.

Immediately after the exercise, another arterialized blood sample was collected together with a muscle biopsy sample from the *M. vastus lateralis* of both the exercised (EXERCISE) as well as the rested (REST) leg. Immediately thereafter subjects consumed 30 g intrinsically L-[1-¹³C]-phenylalanine and L-[1-¹³C]-leucine labeled lesser mealworm- or milk-derived protein ($t = 0$ min). Arterialized blood samples were collected at $t = 20, 40, 60, 90, 120, 150, 180, 240,$ and 300 min. Additional muscle biopsy samples were obtained at $t = 120$ and 300 min from both legs to determine mixed muscle protein synthesis rates both at rest and after exercise.

Blood samples were collected in EDTA-containing tubes and centrifuged at $1000 \times g$ at 4°C for 10 min. Aliquots of plasma were frozen in liquid nitrogen and stored at -80°C until further analysis. Muscle biopsy samples were taken from the middle region of the *M. vastus lateralis*, 15 cm above the patella and 3 cm below entry through the fascia, using a modified Bergström needle (37). Muscle samples were dissected carefully, freed from any visible fat, immediately frozen in liquid nitrogen, and stored at -80°C until further analysis.

Plasma and muscle tissue analyses

The **Supplemental Methods** present details of analyses related to the determination of plasma (glucose, insulin, amino acids, L-[1-¹³C]-leucine enrichments, L-[1-¹³C]- and L-[ring-²H₅]-phenylalanine enrichments, L-[1-¹³C]-, L-[ring-3,5-²H₂]-, and L-[ring-²H₄]- tyrosine enrichments, and mixed plasma

proteins) as well as muscle (mixed muscle protein L-[1-¹³C]-leucine and L-[1-¹³C]- and L-[ring-²H₅]-phenylalanine enrichments) data.

Calculations

Ingestion of L-[1-¹³C]-phenylalanine-labeled protein, intravenous infusion of L-[ring-²H₅]-phenylalanine and L-[ring-3,5-²H₂]-tyrosine, and blood sample enrichment values were used to assess whole-body amino acid kinetics in non-steady state conditions. Total, exogenous, and endogenous phenylalanine rate of appearance (R_a) and plasma availability of dietary protein-derived phenylalanine that appeared in the systemic circulation as a fraction of the total amount of phenylalanine that was ingested ($\text{Phe}_{\text{plasma}}$) were calculated using modified Steele's equations (38, 39). For complete details, see the Supplemental Methods.

The primary outcome, fractional synthesis rate (FSR) of mixed muscle protein, was calculated by dividing the increment in enrichment of the product, i.e., protein-bound L-[ring-²H₅]-phenylalanine and L-[1-¹³C]-leucine, by the enrichment of the precursor. The muscle FSR (%/h) was calculated as follows:

$$\text{FSR} = \frac{dE_p}{E_{\text{precursor}}t} \cdot 100 \quad (1)$$

Where dE_p is the Δ increment of protein-bound L-[ring-²H₅]-phenylalanine and L-[1-¹³C]-leucine during incorporation periods, $E_{\text{precursor}}$ is the average (weighed means) plasma L-[ring-²H₅]-phenylalanine and L-[1-¹³C]-leucine enrichment during the period for the determination of amino acid incorporation, and t reflects the time interval (h) between sampling time points. Basal muscle protein synthesis rates were assessed using the plasma proteins ($t = -180$ min) and the single biopsy approach.

To account for the differences in the L-[1-¹³C]-phenylalanine enrichments in the lesser mealworm- and milk-derived protein (50.8 and 38.3 MPE, respectively), incorporation of the protein-derived L-[1-¹³C]-phenylalanine into de novo muscle protein was corrected for the differences in the dietary protein L-[1-¹³C]-phenylalanine enrichments. Individual L-[1-¹³C]-phenylalanine enrichment values in the WORM group were divided by 50.8 and multiplied by 38.3, to allow direct comparison between the groups.

Statistical analysis

All data are expressed as means \pm SDs. Data were checked and normality and sphericity were confirmed. Baseline characteristics, peak plasma concentrations, incremental AUC (iAUC), and dietary protein-derived amino acid availability were compared between groups using an independent-samples t test. For time-dependent plasma variables, repeated-measures ANOVAs with group (MILK or WORM) as a between-subjects factor and time as a within-subjects factor were used. For FSR variables, repeated-measures ANOVAs with group (MILK or WORM) as a between-subjects factor and leg (exercise compared with rest) and time (basal compared with postprandial) as within-subjects factors were used. In case of significant interactions, separate analyses were performed as appropriate (i.e., within each group or within legs, as well as between groups or between legs for every time period separately). In the case of significant time effects, Bonferroni post hoc analyses were performed to locate

the differences for plasma variables. Significance was set at $P < 0.05$. Calculations were performed using SPSS version 25.0 (IBM Corp.).

Sample size was calculated with differences in postprandial muscle FSR as the primary outcome measure. Based on studies comparing muscle protein turnover between groups at rest and after exercise (4, 6), we used a difference between groups in muscle FSR of 20% (within-group SD: 16%). Using a power of 80%, a significance level of 0.05, and a dropout rate of 10%, the final number of participants was calculated as $n = 12/\text{group}$.

Results

Plasma metabolites

Plasma glucose concentrations increased shortly after protein ingestion, and then decreased (peak values: 5.5 ± 0.4 and 5.6 ± 0.6 mmol/L for MILK and WORM, respectively; Time: $P < 0.001$), with no differences between groups (Time \times Group: $P = 0.945$; data not shown). Plasma insulin concentrations increased after protein ingestion in both groups (peak values: 20.9 ± 11.6 and 21.2 ± 9.3 mU/L for MILK and WORM, respectively; Time: $P < 0.001$) and returned back to baseline values within 5 h. No significant differences were observed between groups (Time \times Group: $P = 0.223$; data not shown).

Plasma leucine (Figure 1A), phenylalanine (Figure 1B), and tyrosine (Figure 1C) concentrations increased after MILK and WORM protein ingestion (Time: $P < 0.001$). Plasma leucine concentrations were higher for MILK than for WORM at all time points, except $t = 0$ and 120 min (Time \times Group: $P < 0.001$). Peak leucine concentrations were 27% higher for MILK than for WORM ($P < 0.001$). Plasma phenylalanine concentrations were higher for MILK than for WORM, except between $t = 60$ and 150 min (Time \times Group: $P = 0.002$). Peak plasma phenylalanine concentrations were 15% higher for MILK than for WORM ($P = 0.007$). Plasma tyrosine concentrations were higher for WORM than for MILK from $t = 60$ through 300 min (Time \times Group: $P < 0.001$). Peak plasma tyrosine concentrations were 22% higher for WORM than for MILK ($P < 0.001$).

Figure 2 depicts concentrations of plasma essential amino acids (EAAs; Figure 2A), nonessential amino acids (NEAAs; Figure 2B), and total amino acids (TAAs; Figure 2C). Plasma EAA concentrations increased after protein ingestion (Time: $P < 0.001$) and were higher for MILK than for WORM at $t = 20, 40,$ and 180 min (Time \times Group: $P < 0.001$). Peak plasma EAA concentrations were 12% higher for MILK than for WORM ($P = 0.005$) with no significant differences for postprandial EAA iAUC between groups ($P = 0.231$). Plasma NEAA concentrations increased after protein ingestion (Time: $P < 0.001$) and were higher for MILK than for WORM at $t = 20$ min (Time \times Group: $P = 0.003$). However, the peak plasma NEAA concentrations and the postprandial NEAA iAUC did not differ between groups ($P = 0.466$ and $P = 0.397$, respectively). Plasma TAA concentrations increased after protein ingestion (Time: $P < 0.001$) and were higher for MILK than for WORM at $t = 20$ and 300 min (Time \times Group: $P = 0.001$). However, the peak plasma TAA concentrations and the postprandial TAA iAUC did not differ between groups ($P = 0.181$ and $P = 0.812$, respectively).




FIGURE 1 Plasma leucine (A), phenylalanine (B), and tyrosine (C) concentrations after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 12$) derived protein during 5 h of recovery from a single bout of unilateral exercise in healthy, young men. The dotted line represents the time of protein ingestion. Values represent means \pm SDs. Data were analyzed using repeated-measures (Time \times Group) ANOVA and separate analyses were performed when a significant interaction was detected. Bonferroni post hoc testing was used to detect differences between groups. Time \times Group interactions were observed for plasma leucine, phenylalanine, and tyrosine concentrations (all $P < 0.05$). *MILK significantly different from WORM ($P < 0.05$). MILK, 30 g milk protein concentrate; WORM, 30 g lesser mealworm-derived protein.

Whole-body amino acid kinetics and metabolism

Figure 3 shows exogenous (Figure 3A), endogenous (Figure 3B), and total plasma phenylalanine R_a (Figure 3C) and total plasma phenylalanine rates of disappearance (R_d) (Figure 3D). Exogenous phenylalanine R_a increased after protein ingestion and remained elevated over the entire postprandial period (Time: $P < 0.001$) with the exogenous phenylalanine




FIGURE 2 Plasma EAA (A), NEAA (B), and TAA (C) concentrations and iAUCs after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 12$) derived protein during 5 h of recovery from a single bout of unilateral exercise in healthy, young men. The dotted line represents the time of protein ingestion. Values represent means \pm SDs. Data were analyzed using repeated-measures (Time \times Group) ANOVA and separate analyses were performed when a significant interaction was detected. Bonferroni post hoc testing was used to detect differences between groups. Time \times Group interactions were observed for plasma EAAs, NEAAs, and TAAs (all $P < 0.05$). *MILK significantly different from WORM ($P < 0.05$). EAA, essential amino acid; MILK, 30 g milk protein concentrate; NEAA, nonessential amino acid; TAA, total amino acid; WORM, 30 g lesser mealworm-derived protein.




FIGURE 3 Exogenous phenylalanine R_a (A), endogenous phenylalanine R_a (B), total phenylalanine R_a (C), and total plasma phenylalanine R_d (D) after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 12$) derived protein during 5 h of recovery from a single bout of unilateral exercise in healthy, young men. The dotted line represents the time of protein ingestion. Values represent means \pm SDs. Data were analyzed using repeated-measures (Time \times Group) ANOVA and separate analyses were performed when a significant interaction was detected. Bonferroni post hoc testing was used to detect differences between groups. Time \times Group interactions were observed for exogenous phenylalanine R_a , total phenylalanine R_a , and total phenylalanine R_d (all $P < 0.05$). *MILK significantly different from WORM ($P < 0.05$). MILK, 30 g milk protein concentrate; R_a , rate of appearance; R_d , rate of disappearance; WORM, 30 g lesser mealworm-derived protein.

R_a being higher for MILK than for WORM at $t = 20$ min (Time \times Group: $P = 0.009$). Endogenous phenylalanine R_a decreased after protein ingestion and remained below baseline values during the entire postprandial period (Time: $P < 0.001$), with no significant differences between groups (Time \times Group: $P = 0.269$). After protein ingestion, total phenylalanine R_a increased and remained elevated for 150 min (Time: $P < 0.001$) and the total phenylalanine R_a was higher for MILK than for WORM at $t = 0, 20, 40,$ and $180\text{--}300$ min (Time \times Group: $P = 0.001$). Total phenylalanine R_d increased after protein ingestion and remained elevated for 180 min (Time: $P < 0.001$), and was higher for MILK than for WORM at $t = 0\text{--}60$ and $t = 180\text{--}300$ min (Time \times Group: $P = 0.022$).

Plasma dietary protein-derived phenylalanine availability over the 5 h postprandial period (calculated as a fraction of the total amount of ingested phenylalanine) was $73\% \pm 7\%$ and $77\% \pm 7\%$ for WORM and MILK, respectively, and did not differ between groups ($P = 0.263$) (Figure 4). The calculated absolute amount of dietary protein-derived amino acids that appeared in the circulation during the 5-h postprandial period was 22.0 ± 0.6 and 23.0 ± 0.6 g for WORM and MILK, respectively. Also, when plasma dietary protein-derived amino acid release was expressed as availability over 0–2 h (12.4 ± 2.7 and 12.2 ± 2.1 g) and 2–5 h (9.6 ± 1.8 and 10.8 ± 1.7 g), WORM and MILK were not significantly different ($P = 0.790$ and $P = 0.084$, respectively).

Whole-body protein synthesis rates (0.77 ± 0.06 compared with 0.68 ± 0.08 $\mu\text{mol Phe} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) were 13% higher and breakdown rates (0.55 ± 0.04 compared with 0.48 ± 0.06 $\mu\text{mol Phe} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) were 16% higher for MILK than for WORM

($P = 0.005$ and $P = 0.002$, respectively). However, whole-body protein oxidation rates (0.10 ± 0.02 compared with 0.09 ± 0.05 $\mu\text{mol Phe} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and net balance (0.21 ± 0.04 compared with 0.20 ± 0.06 $\mu\text{mol Phe} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) did not differ between MILK and WORM ($P = 0.782$ and $P = 0.426$, respectively).

Mixed muscle protein synthesis rates

Figure 5 shows mixed muscle protein synthesis rates calculated based upon L-[ring- $^2\text{H}_5$]-phenylalanine. No Time \times Leg \times Group interactions were observed ($P = 0.114$ and $P = 0.549$ for 0–2 h and 0–5 h, respectively). A significant Leg \times Time interaction showed that the increase in muscle protein synthesis rates from basal to postprandial was greater in the EXERCISE than in the REST leg, for both 0–2 h and 0–5 h (both $P < 0.001$).

In the REST leg, muscle protein synthesis rates based on L-[ring- $^2\text{H}_5$]-phenylalanine (Figure 5) increased from basal to postprandial (0–2 h) after both WORM (from $0.025 \pm 0.008\%/h$ to $0.042 \pm 0.019\%/h$) and MILK (from $0.028 \pm 0.010\%/h$ to $0.056 \pm 0.025\%/h$) protein ingestion (Time: $P < 0.001$), with no significant differences between groups (Time \times Group: $P = 0.218$). In the EXERCISE leg, postexercise muscle protein synthesis rates based on L-[ring- $^2\text{H}_5$]-phenylalanine (Figure 5) increased from basal to postprandial (0–2 h) after both WORM (from $0.025 \pm 0.012\%/h$ to $0.071 \pm 0.021\%/h$) and MILK (from $0.026 \pm 0.009\%/h$ to $0.067 \pm 0.021\%/h$) protein ingestion (Time: $P < 0.001$), with no significant differences between groups (Time




FIGURE 4 Dietary protein-derived phenylalanine (expressed as a percentage of total amount of protein-bound phenylalanine ingested) released in the circulation during 5 h after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 12$) derived protein in healthy, young men. Values represent means \pm SDs. Data were analyzed using an independent t test. MILK, 30 g milk protein concentrate; WORM, 30 g lesser mealworm-derived protein.

\times Group: $P = 0.588$). Similarly, muscle protein synthesis rates based on L-[ring- 2 H $_5$]-phenylalanine increased from basal to postprandial (0–5 h) after both WORM ($0.045 \pm 0.017\%/h$) and MILK ($0.056 \pm 0.012\%/h$) protein ingestion (Time: $P < 0.001$) at REST, as well as after EXERCISE (Time: $P < 0.001$; WORM: $0.059 \pm 0.015\%/h$ and MILK: $0.073 \pm 0.020\%/h$), with no significant differences between groups (Time \times Group: $P = 0.089$ and $P = 0.088$, respectively).




FIGURE 5 Mixed muscle protein FSRs assessed using L-[ring- 2 H $_5$]-phenylalanine after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 11$) derived protein for both REST and EXERCISE in healthy, young men. Values represent means \pm SDs. Data were analyzed using repeated-measures (Time \times Leg \times Group) ANOVA, and separate analyses were performed when a significant interaction was detected. A Time \times Leg \times Group interaction was absent ($P > 0.05$). A Time \times Leg interaction was observed ($P < 0.05$). #Both 0–2 h and 0–5 h postprandial FSR significantly different from basal ($P < 0.05$); \$postprandial FSR in EXERCISE significantly different from REST ($P < 0.05$). EXERCISE, exercised leg; FSR, fractional synthesis rate; MILK, 30 g milk protein concentrate; REST, rested leg; WORM, 30 g lesser mealworm-derived protein.

In line with the data for L-[ring- 2 H $_5$]-phenylalanine, no Time \times Leg \times Group interactions were observed for postprandial muscle protein synthesis rates based on L-[13 C]-leucine ($P = 0.372$ and $P = 0.684$ for 0–2 h and 0–5 h, respectively), but significant Leg \times Time interactions showed that the increase in muscle protein synthesis based on L-[13 C]-leucine from basal to postprandial was greater in the EXERCISE than in the REST leg, for both 0–2 h ($P = 0.022$) and 0–5 h ($P < 0.001$). In the early postprandial period (0–2 h), muscle protein synthesis rates based on L-[13 C]-leucine increased to $0.030 \pm 0.026\%/h$ and $0.048 \pm 0.028\%/h$ after WORM and MILK protein ingestion at REST (Time: $P < 0.001$), respectively, and to $0.060 \pm 0.031\%/h$ and $0.050 \pm 0.041\%/h$ after EXERCISE (Time: $P < 0.001$), respectively, with no significant differences between groups (Time \times Group: $P = 0.433$ and $P = 0.871$, respectively). Similarly, muscle protein synthesis rates based on L-[13 C]-leucine increased from basal to postprandial (0–5 h) after both WORM ($0.042 \pm 0.020\%/h$) and MILK ($0.055 \pm 0.017\%/h$) protein ingestion (Time: $P < 0.001$) at REST, as well as after EXERCISE (Time: $P < 0.001$; WORM: $0.061 \pm 0.013\%/h$ and MILK: $0.073 \pm 0.017\%/h$), with no significant differences between groups (Time \times Group: $P = 0.552$ and $P = 0.394$, respectively).

Incorporation of protein-derived L-[13 C]-phenylalanine into de novo muscle protein during the postprandial period increased after protein ingestion at REST (Time: $P < 0.001$) and after EXERCISE (Time: $P < 0.001$), but did not differ between groups (Time \times Group: $P = 0.623$ and Time \times Group: $P = 0.692$, respectively; data not shown). To account for differences in the L-[13 C]-phenylalanine enrichments in the WORM and MILK protein (50.8 compared with 38.3 MPE, respectively), incorporation of protein-derived L-[13 C]-phenylalanine into de novo muscle protein was corrected for the differences in dietary protein L-[13 C]-phenylalanine enrichments. **Figure 6** shows the relative incorporation of dietary protein-derived L-[13 C]-phenylalanine into de novo muscle protein. Relative muscle protein-bound L-[13 C]-phenylalanine incorporation increased after protein ingestion at REST (Time: $P < 0.001$) and during recovery from EXERCISE (Time: $P < 0.001$), with no significant differences between groups (Time \times Group: $P = 0.302$ and Time \times Group: $P = 0.110$, respectively). At $t = 5$ h relative muscle protein-bound L-[13 C]-phenylalanine enrichments were higher for the EXERCISE than for the REST leg ($P < 0.001$).

Discussion

This study shows that ingestion of lesser mealworm and milk protein concentrate are both followed by rapid protein digestion and amino acid absorption, with $>70\%$ of the ingested protein-bound phenylalanine being released into the circulation during the 5-h postprandial period. Ingestion of both mealworm and milk protein concentrate increased muscle protein synthesis rates both at rest and during recovery from resistance-type exercise, with no differences between protein sources.

Conventional animal-based protein sources such as meat, eggs, and milk are considered high-quality protein sources because they are highly digestible and meet all of the current EAA requirements (40). However, production of sufficient amounts of these animal-based protein sources is not feasible to meet the growing global protein demands (41). Consequently, there




FIGURE 6 Corrected L-[1-¹³C]-phenylalanine enrichment in mixed muscle protein over time after ingestion of milk- (MILK; $n = 12$) or lesser mealworm- (WORM; $n = 11$) derived protein for both REST and EXERCISE in healthy, young men. Values for the WORM group were multiplied by 0.75 to correct for the difference in enrichment in the MILK (38.3 MPE) and WORM (50.8 MPE) protein. Values represent means \pm SDs. Data were analyzed using repeated-measures (Time \times Leg \times Group) ANOVA, and separate analyses were performed when a significant interaction was detected. A Time \times Leg \times Group interaction was absent ($P > 0.05$). A Time \times Leg interaction was observed ($P < 0.05$). ⁺ $t = 2$ h significantly different from $t = 0$ h ($P < 0.05$); [#] $t = 5$ h significantly different from $t = 0$ and $t = 2$ h ($P < 0.05$); ^{\$}REST significantly different from EXERCISE for that time point ($P < 0.05$). EXERCISE, exercised leg; MILK, 30 g milk protein concentrate; MPE, mole percentage excess; REST, rested leg; WORM, 30 g lesser mealworm-derived protein.

is an increasing interest in the production of alternative and more sustainable dietary protein sources. Edible insects have been identified as a protein-dense food source that may be produced more sustainably at a viable commercial scale for human consumption (19, 21). From the wide array of edible insects, lesser mealworms (*A. diaperinus*) have been reported to provide a source of high-quality protein containing high amounts of EAAs (18–21). In agreement with previous literature (18, 20, 42), we show that lesser mealworm protein has a relatively high EAA content and a leucine content comparable with that of milk protein (Table 2). This implies that lesser mealworm may also have anabolic properties that are not distinct from high-quality animal-derived protein sources.

In addition to amino acid composition, protein digestion and amino acid absorption kinetics represent another important factor that determines the capacity of a protein source to stimulate postprandial muscle protein synthesis (3, 6–8). After lesser mealworm-derived protein ingestion, we observed a rapid rise in circulating plasma amino acid concentrations that remained elevated over the subsequent hours (Figures 1, 2), which is in line with previous data (42). Because postprandial plasma amino acid responses do not necessarily reflect protein digestion and amino acid absorption, we combined the ingestion of specifically produced intrinsically L-[1-¹³C]-phenylalanine labeled protein with continuous intravenous L-[ring-²H₅]-phenylalanine infusion. Upon ingestion of the lesser mealworm- and milk-derived protein we observed a rapid rise in the release of exogenous protein-derived phenylalanine into the circulation (Figure 3). The postprandial rise in release of exogenous protein-derived phenylalanine was greater after milk than after lesser mealworm-derived protein during the early postprandial phases. Throughout

the full 5-h postprandial period 73% \pm 7% and 77% \pm 7% of the ingested protein-bound phenylalanine had been released into the circulation after the ingestion of 30 g lesser mealworm- and milk-derived protein, respectively ($P > 0.05$) (Figure 4). These data are the first that we know of to show that insect-derived protein can be rapidly digested and effectively absorbed, with a postprandial rise in protein-derived amino acid availability in the circulation that is not different from a high-quality dairy protein source.

It has been well established that protein ingestion stimulates muscle protein synthesis (43, 44), with the postprandial rise in circulating EAAs (45, 46) and leucine in particular (47, 48) being responsible for activating the muscle protein synthetic machinery. In this study, we measured the incorporation of L-[ring-²H₅]-phenylalanine and L-[1-¹³C]-leucine in muscle protein to assess postabsorptive and postprandial muscle tissue protein FSRs. Using the L-[ring-²H₅]-phenylalanine tracer, we observed a substantial increase in muscle protein synthesis rates after both mealworm and milk protein concentrate ingestion (Figure 5). Exercise before protein intake sensitizes skeletal muscle tissue to the anabolic properties of amino acids, resulting in an even greater postprandial increase in muscle protein synthesis rates (11, 49). In the present study, participants performed a single bout of 1-legged resistance-type exercise before protein ingestion, allowing us to assess muscle protein synthesis rates both at rest and during recovery from exercise. In agreement, exercise stimulated muscle protein synthesis rates resulting in an even greater increase in postprandial muscle protein synthesis rates after ingestion of both protein sources (Figure 5). No differences were observed between treatments, implying that the lesser mealworms represent a readily accessible protein source with strong anabolic properties that do not seem to differ from a high-quality protein source like milk.

In the current study, participants ingested specifically produced, intrinsically labeled lesser mealworm- and milk-derived protein with very high L-[1-¹³C]-phenylalanine enrichments (50.8 and 38.3 MPE, respectively). We observed no significant differences in the relative incorporation of dietary protein-derived L-[1-¹³C]-phenylalanine into skeletal muscle protein (Figure 6). Of course, prior exercise further increased the uptake of dietary protein-derived L-[1-¹³C]-phenylalanine into muscle protein, confirming the impact of muscle contraction on stimulating amino acid uptake and subsequent de novo incorporation into muscle protein. These data demonstrate that both the mealworm and the milk protein concentrate provide us with required amino acids as precursors for de novo muscle protein synthesis, confirming that we are what we just ate (26), whether that is milk or mealworm.

In the present study we provided subjects with a meal-sized serving of milk- or lesser mealworm-derived protein. Previous work has reported that ingestion of 20 g high-quality protein is sufficient to maximize postprandial muscle protein synthesis rates in healthy, young adults (4, 50). Therefore, it could be speculated that ingestion of smaller doses of lesser mealworm-derived protein could have revealed (greater) differences in the muscle protein synthetic response to milk- compared with lesser mealworm-derived protein ingestion. However, in the present study we were able to show similar efficacy of both protein digestion and amino acid absorption (Figure 4) and we were unable to detect any measurable differences in postprandial muscle protein synthesis rates (Figure 5). These observations

differ from previous work comparing dairy with plant-based protein isolates, where it was shown that substantially more wheat protein has to be ingested to allow a muscle protein synthetic response equivalent to that seen after the ingestion of 35 g dairy protein (15). Clearly, the high quality of the insect-derived proteins (18) compared with existing commercial plant-based protein concentrates and isolates (51) will be of particular relevance in populations where protein quality is of great(er) importance. The latter includes people that consume less protein and suffer from anabolic resistance (52), such as the older and/or more clinically compromised populations. A more sustainable protein source without the obvious limitations of low digestibility, low EAA content, and/or specific amino acid deficiency would prove to be highly attractive. Together with the additional benefits of insects containing high amounts of micronutrients (53) and antioxidants (54), our data provide a strong incentive for incorporating insects into the Western diet. The consumption of insects is already common for >2 billion people worldwide, mainly in Asia, Africa, and South America (53). It will be interesting to see insects and insect-derived proteins becoming a major protein source in our future food supply (55).

In conclusion, ingestion of a meal-like amount of lesser mealworm is followed by rapid protein digestion and amino acid absorption and a substantial increase in muscle protein synthesis rates both at rest and during recovery from exercise. The postprandial protein handling of lesser mealworm-derived protein does not differ compared with ingestion of the same amount of milk protein in vivo in humans, which shows that insects can provide a viable, high-quality protein source for human consumption.

We acknowledge Maarten Overkamp, Janneau van Kranenburg, Antoine H Zorenc, Joy PB Goessens, Annemie P Gijsen, Wendy Sluijsmans, and Hasibe Aydeniz for their (technical) assistance. We acknowledge Protifarm R&D for their assistance in producing the labeled lesser mealworm.

The authors' responsibilities were as follows—TAC-V and LJCvL: designed the research; WJHH, JMS, TAC-V, KJMP, CJF, JSJS, and JJA vL: conducted the research; WJHH, CJF, JMS, and LJCvL: analyzed the data; WJHH and LBV: performed the statistical analysis; WJHH, TAC-V, and LJCvL: wrote the manuscript; WJHH and LJCvL: hold primary responsibility for the final content; and all authors: read and approved the final manuscript. All other authors report no conflicts of interest.

Data availability

Data described in the article, code book, and analytic code will be made available upon request pending application and approval.

References

- Fuchs CJ, Hermans WJH, Holwerda AM, Smeets JSJ, Senden JM, van Kranenburg J, Gijsen AP, Wodzig W, Schierbeek H, Verdijk LB, et al. Branched-chain amino acid and branched-chain ketoacid ingestion increases muscle protein synthesis rates in vivo in older adults: a double-blind, randomized trial. *Am J Clin Nutr* 2019;110(4):862–72.
- Pennings B, Groen BB, van Dijk JW, de Lange A, Kiskini A, Kuklinski M, Senden JM, van Loon LJ. Minced beef is more rapidly digested and absorbed than beef steak, resulting in greater postprandial protein retention in older men. *Am J Clin Nutr* 2013;98(1):121–8.
- Koopman R, Walrand S, Beelen M, Gijsen AP, Kies AK, Boirie Y, Saris WH, van Loon LJ. Dietary protein digestion and absorption rates and the subsequent postprandial muscle protein synthetic response do not differ between young and elderly men. *J Nutr* 2009;139(9):1707–13.
- Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr* 2014;99(1):86–95.
- Pennings B, Koopman R, Beelen M, Senden JM, Saris WH, van Loon LJ. Exercising before protein intake allows for greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in both young and elderly men. *Am J Clin Nutr* 2011;93(2):322–31.
- Pennings B, Boirie Y, Senden JM, Gijsen AP, Kuipers H, van Loon LJ. Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *Am J Clin Nutr* 2011;93(5):997–1005.
- Boirie Y, Dangin M, Gachon P, Vasson M-P, Maubois J-L, Beaufrère B. Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci U S A* 1997;94(26):14930–5.
- Dangin M, Boirie Y, Garcia-Rodenas C, Gachon P, Fauquant J, Callier P, Ballèvre O, Beaufrère B. The digestion rate of protein is an independent regulating factor of postprandial protein retention. *Am J Physiol Endocrinol Metab* 2001;280(2):E340–8.
- Gorissen SHM, Witard OC. Characterising the muscle anabolic potential of dairy, meat and plant-based protein sources in older adults. *Proc Nutr Soc* 2018;77(1):20–31.
- Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol* 2009;107(3):987–92.
- Trommelen J, Holwerda AM, Kouw IWK, Langer H, Halson SL, Rollo I, Verdijk LB, van Loon LJC. Resistance exercise augments postprandial overnight muscle protein synthesis rates. *Med Sci Sports Exerc* 2016;48(12):2517–25.
- Yang Y, Churchward-Venne TA, Burd NA, Breen L, Tarnopolsky MA, Phillips SM. Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. *Nutr Metab (Lond)* 2012;9(1):57.
- Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, Tarnopolsky M, Phillips SM. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 2012;108(10):1780–8.
- Young VR, Pellett PL. Plant proteins in relation to human protein and amino acid nutrition. *Am J Clin Nutr* 1994;59(5):1203S–12S.
- Gorissen SHM, Horstman AMH, Franssen R, Crombag JJR, Langer H, Bierau J, Respondek F, van Loon LJC. Ingestion of wheat protein increases in vivo muscle protein synthesis rates in healthy older men in a randomized trial. *J Nutr* 2016;146(9):1651–9.
- Wilkinson SB, Tarnopolsky MA, MacDonald MJ, MacDonald JR, Armstrong D, Phillips SM. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am J Clin Nutr* 2007;85(4):1031–40.
- Phillips SM. Nutrient-rich meat proteins in offsetting age-related muscle loss. *Meat Sci* 2012;92(3):174–8.
- Churchward-Venne TA, Pinckaers PJM, van Loon JJA, van Loon LJC. Consideration of insects as a source of dietary protein for human consumption. *Nutr Rev* 2017;75(12):1035–45.
- Van Huis A, Van Itterbeek J, Klunder H, Mertens E, Halloran A, Muir G, Vantomme P. Edible insects: future prospects for food and feed security. Rome, Italy: Food and Agriculture Organization of the United Nations; 2013.
- Rumpold BA, Schluter OK. Nutritional composition and safety aspects of edible insects. *Mol Nutr Food Res* 2013;57(5):802–23.
- van Huis A. Potential of insects as food and feed in assuring food security. *Annu Rev Entomol* 2013;58:563–83.
- Bukkens SGF. The nutritional value of edible insects. *Ecol Food Nutr* 1997;36(2–4):287–319.
- Burd NA, Gorissen SH, van Vliet S, Snijders T, van Loon LJ. Differences in postprandial protein handling after beef compared with milk ingestion during postexercise recovery: a randomized controlled trial. *Am J Clin Nutr* 2015;102(4):828–36.
- Burd NA, Cermak NM, Kouw IWK, Gorissen SH, Gijsen AP, van Loon LJC. The use of doubly labeled milk protein to measure postprandial muscle protein synthesis rates in vivo in humans. *J Appl Physiol* 2014;117(11):1363–70.

25. Van Loon LJC, Boirie Y, Gijsen AP, Fauquant J, De Roos AL, Kies AK, Lemosquet S, Saris WHM, Koopman R. The production of intrinsically labeled milk protein provides a functional tool for human nutrition research. *J Dairy Sci* 2009;92(10):4812–22.
26. Groen BBL, Horstman AM, Hamer HM, de Haan M, van Kranenburg J, Bierau J, Poeze M, Wodzig WKWH, Rasmussen BB, van Loon LJC. Post-prandial protein handling: you are what you just ate. *PLoS One* 2015;10(11):e0141582.
27. Pennings B, Pellikaan WF, Senden JMG, van Vuuren AM, Sikkema J, van Loon LJC. The production of intrinsically labeled milk and meat protein is feasible and provides functional tools for human nutrition research. *J Dairy Sci* 2011;94(9):4366–73.
28. Burd NA, Hamer HM, Pennings B, Pellikaan WF, Senden JMG, Gijsen AP, van Loon LJC. Substantial differences between organ and muscle specific tracer incorporation rates in a lactating dairy cow. *PLoS One* 2013;8(6):e68109.
29. Boirie Y, Fauquant J, Rulquin H, Maubois J-L, Beaufrère B. Production of large amounts of [¹³C]leucine-enriched milk proteins by lactating cows. *J Nutr* 1995;125(1):92–8.
30. Van der Wielen N, Khodorova NV, Gerrits WJJ, Gaudichon C, Calvez J, Tomé D, Mensink M. Blood ¹⁵N:¹³C enrichment ratios are proportional to the ingested quantity of protein with the dual-tracer approach for determining amino acid bioavailability in humans. *J Nutr* 2020;150(9):2346–52.
31. Reitelseder S, Agergaard J, Doessing S, Helmark IC, Lund P, Kristensen NB, Frystyk J, Flyvbjerg A, Schjerling P, van Hall G, et al. Whey and casein labeled with L-[1-¹³C]leucine and muscle protein synthesis: effect of resistance exercise and protein ingestion. *Am J Physiol Endocrinol Metab* 2011;300(1):E231–42.
32. Gorissen SHM, Burd NA, Hamer HM, Gijsen AP, Groen BB, van Loon LJC. Carbohydrate coingestion delays dietary protein digestion and absorption but does not modulate postprandial muscle protein accretion. *J Clin Endocrinol Metab* 2014;99(6):2250–8.
33. Res PT, Groen B, Pennings B, Beelen M, Wallis GA, Gijsen AP, Senden JMG, van Loon LJC. Protein ingestion before sleep improves postexercise overnight recovery. *Med Sci Sports Exerc* 2012;44(8):1560–9.
34. Dumas JBA. Procédés de l'analyse organique. *Ann Chim Phys* 1831;247:198–213.
35. Janssen RH, Vincken J-P, van den Broek LAM, Fogliano V, Lakemond CMM. Nitrogen-to-protein conversion factors for three edible insects: *Tenebrio molitor*, *Alphitobius diaperinus*, and *Hermetia illucens*. *J Agric Food Chem* 2017;65(11):2275–8.
36. Abumrad NN, Rabin D, Diamond MP, Lacy WW. Use of a heated superficial hand vein as an alternative site for the measurement of amino acid concentrations and for the study of glucose and alanine kinetics in man. *Metabolism* 1981;30(9):936–40.
37. Bergström J. Percutaneous needle biopsy of skeletal muscle in physiological and clinical research. *Scand J Clin Lab Invest* 1975;35(7):609–16.
38. Boirie Y, Gachon P, Corny S, Fauquant J, Maubois JL, Beaufrère B. Acute postprandial changes in leucine metabolism as assessed with an intrinsically labeled milk protein. *Am J Physiol* 1996;271(6 Pt 1):E1083–91.
39. Dangin M, Guillet C, Garcia-Rodenas C, Gachon P, Bouteloup-Demange C, Reiffers-Magnani K, Fauquant J, Ballèvre O, Beaufrère B. The rate of protein digestion affects protein gain differently during aging in humans. *J Physiol* 2003;549(2):635–44.
40. van Vliet S, Burd NA, van Loon LJC. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr* 2015;145(9):1981–91.
41. Wu G, Fanzo J, Miller DD, Pingali P, Post M, Steiner JL, Thalacker-Mercer AE. Production and supply of high-quality food protein for human consumption: sustainability, challenges, and innovations. *Ann N Y Acad Sci* 2014;1321(1):1–19.
42. Vangsoe MT, Thogersen R, Bertram HC, Heckmann L-HL, Hansen M. Ingestion of insect protein isolate enhances blood amino acid concentrations similar to soy protein in a human trial. *Nutrients* 2018;10(10):1357.
43. Gorissen SHM, Remond D, van Loon LJC. The muscle protein synthetic response to food ingestion. *Meat Sci* 2015;109:96–100.
44. Wolfe RR. Regulation of muscle protein by amino acids. *J Nutr* 2002;132(10):3219S–24S.
45. Volpi E, Kobayashi H, Sheffield-Moore M, Mittendorfer B, Wolfe RR. Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr* 2003;78(2):250–8.
46. Tipton KD, Gurkin BE, Matin S, Wolfe RR. Nonessential amino acids are not necessary to stimulate net muscle protein synthesis in healthy volunteers. *J Nutr Biochem* 1999;10(2):89–95.
47. Rieu I, Balage M, Sornet C, Giraudet C, Pujos E, Grizard J, Mosoni L, Dardevet D. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol* 2006;575(1):305–15.
48. Churchward-Venne TA, Breen L, Di Donato DM, Hector AJ, Mitchell CJ, Moore DR, Stellingwerff T, Breuille D, Offord EA, Baker SK, et al. Leucine supplementation of a low-protein mixed macronutrient beverage enhances myofibrillar protein synthesis in young men: a double-blind, randomized trial. *Am J Clin Nutr* 2014;99(2):276–86.
49. Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol Endocrinol Metab* 1995;268(3):E514–E20.
50. Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, Prior T, Tarnopolsky MA, Phillips SM. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr* 2009;89(1):161–8.
51. Gorissen SHM, Crombag JJR, Senden JMG, Waterval WAH, Bierau J, Verdijk LB, van Loon LJC. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* 2018;50(12):1685–95.
52. Wall BT, Gorissen SH, Pennings B, Koopman R, Groen BBL, Verdijk LB, van Loon LJC. Aging is accompanied by a blunted muscle protein synthetic response to protein ingestion. *PLoS One* 2015;10(11):e0140903.
53. Rumpold BA, Schlüter O. Insect-based protein sources and their potential for human consumption: nutritional composition and processing. *Anim Front* 2015;5(2):20–4.
54. van Huis A. Nutrition and health of edible insects. *Curr Opin Clin Nutr Metab Care* 2020;23(3):228–31.
55. Aiking H. Future protein supply. *Trends Food Sci Technol* 2011;22(2–3):112–20.