

COMMENT

Open Access



Routine use of indirect calorimetry in critically ill patients: pros and cons

Elisabeth De Waele^{1,2} and Arthur R. H. van Zanten^{3,4*}

Purpose of review

To review the pros and cons of indirect calorimetry (IC) to estimate resting energy expenditure (REE) and define individual nutritional energy targets among critically ill patients. We evaluate pros, (relative) cons and when adjustments of REE are needed (Fig. 1).

Introduction

IC measuring O₂ consumption (VO₂) and CO₂ production (VCO₂) represents real-time energy metabolism [1]. REE assessment by IC to guide nutrition therapy is recommended by international nutrition guidelines for adult ICU patients, although the level of evidence is still low (grade B recommendation) [2].

Indirect calorimetry does not change a patient's outcome—Yes, it does

Indirect calorimetry is not lifesaving but a monitoring tool providing factual information on metabolism. Retrospective data revealed a U-shaped association between 60-day mortality and calorie intake in early critical illness, underlining the importance of targeting individual energy demands and preventing both overfeeding and underfeeding [3]. In a systematic review and meta-analysis in 2020, no improved outcomes were found when IC was used, although prescribed energy targets were more closely met when IC informed energy delivery compared with predictive equations [4]. The more precise intake observed suggests that IC feedback improves feeding

performance, compared with equations, and may prevent underfeeding and overfeeding in individual patients.

In 2021, another meta-analysis showed a 23% reduction in short-term mortality when energy targets were based on IC [5]. In another meta-analysis, lower 28-day mortality was found when isocaloric IC-guided nutrition was compared with (hypocaloric) feeding protocols using predictive equations, however, without effect on nosocomial infections and 90-day mortality [6]. All meta-analyses did not find a shorter length of stay (LOS) or mechanical ventilation duration, possibly due to higher calorie intake leading to higher CO₂ production, potentially prolonging mechanical ventilation duration and LOS.

You can reliably estimate energy expenditure with predictive equations—no, you can't


Personalised nutrition should be based on individual energy targets. However, predictive equations are highly inaccurate, potentially leading to 500–1000 kcal/day nutrition targets higher or lower than individual demands, conferring marked risks of underfeeding and overfeeding [7]. Using only the VCO₂ from mechanical ventilators to estimate REE is inferior to IC and barely better than predictive equations [8]. Responding to varying needs during the patient journey is recommended. IC using a canopy can be expanded to the post-ICU phase [9]. The ICALIC research group helped to develop moderate-cost highly efficient technology to measure REE in ventilated and spontaneous breathing patients [10].

*Correspondence: zantena@zgv.nl

³ Department of Intensive Care Medicine and Research, Gelderse Vallei Hospital, Willy Brandtlaan 10, 6716 RP Ede, The Netherlands
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



You can use it in all patients—no, you can't at any time

In ventilated patients with $FiO_2 > 0.7$ and $PEEP > 12$ cmH_2O , no reliable measurements of VO_2 and VCO_2 are possible. Furthermore, any air leak excludes the use of IC as not all exhaled gas will meet the device's sensors. Typical examples are pneumothorax, subcutaneous emphysema, or tracheal-oesophageal fistula. High-flow nasal oxygen therapy or non-invasive ventilation precludes IC use. As moisture affects the performance of sensors, nebulisation during measurement is not recommended. In non-ventilated patients, measurements can only be taken when oxygen therapy has been stopped [1]. As these limitations are present in many patients, equations are still needed then.

You should give nutritional energy to meet the REE in all patients—no, you shouldn't

REE can be used to set energy targets until a new measurement has been taken. However, endogenous energy production can be marked in the early phase of critical illness (> 1000 kcal/day). Adding full nutrition to this non-inhibitable energy production may induce over-feeding [1]. Gradually advancing to target is recommended. IC should not be followed during this phase. However, over 3–4 days, this effect dissipates in most

patients. Then, IC can be used to set targets. Another reason not to follow IC REE is administering non-nutritional calories (propofol, glucose, and citrate). Although not intended as nutrition, these calories contribute to energy intake and are substantial in individual patients [1]. When after commencement of nutrition, plasma phosphate drops below 0.65 mmol/l, known as refeeding hypophosphatemia (RFH), caloric restriction (< 500 kcal/day) is warranted. Studies have demonstrated increased mortality with high-calorie intake during RFH [1]. Therefore, then REE should not be the target.

You will encounter practical issues—yes, probably

Large tidal volume variations during spontaneous mechanical ventilation can affect reliable measurements. Devices with a mixing chamber have better performance [10]. Early COVID-19 guidelines recommended against IC to reduce infection transmission as ventilator circuit disconnections may enhance aerosol production [11]. However, later studies using strict hygiene protocols show that safe application is feasible and provide crucial insights into metabolic changes of (hyper)metabolism over time [12].

Measuring REE during extracorporeal treatments (like CRRT and ECMO) poses significant challenges. Techniques may disturb REE by impacting temperature and stress levels. Moreover, techniques can also affect

metabolism (adding bicarbonate and/or citrate) or non-pulmonary gas exchange. The effects of these techniques on REE are probably limited but also variable. Performing IC among these patients remains challenging [13]. Using complex equations to adjust for metabolic and gas exchange disturbances is possible but seems less feasible in daily practice [14].

Indirect calorimetry is expensive—no, but there are costs involved

Costs of IC comprise device investment costs, consumables, calibration gas, and service. Additionally, staff time should be added. Typically, a procedure to measure takes 5–10 min. Modern interfaces are intuitive, and training during IC implementation can be limited to 1–2 h. Formal health economic evaluations are not available. Modern devices have acceptable costs per measurement (*personal communication: EdW, AvZ*). The business case is positive when improved outcomes reported in meta-analyses (mortality) translate into daily practice. Optimising nutritional performance may impact readmission rates, long-term functional outcomes (recovery from ICU-acquired weakness), and quality of life. Unfortunately, these endpoints have not been studied yet.

Conclusions

Indirect calorimetry is the gold standard for measuring energy expenditure to set nutrition therapy goals during critical illness. Although recommended, the evidence underlying recommendations is limited. We have summarised when not to follow the REE from IC. Also, technical and patient-related limitations have been addressed. Future developments of IC to monitor REE continuously and during oxygen therapy and non-invasive ventilation are warranted to further enhance its application in daily practice.

Acknowledgements

Not applicable.

Author contributions

EDW and AvZ equally contributed to conceiving and editing the manuscript, tables and figures. Both authors read and approved the final manuscript.

Funding

The authors did not receive any funding for this work.

Availability of data and materials

No datasets were used. Information on datasets and materials from studies reported can be found through the references.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

EDW reported receiving honoraria for advisory board meetings, lectures, and travel expenses from Baxter Healthcare, Danone-Nutricia, Cardinal Health, Fresenius Kabi, GE Healthcare and research support from the Belgian Government and Wetenschappelijk Fonds Willy Gepts. AVZ reported receiving honoraria for advisory board meetings, lectures, research, and travel expenses from AOP Pharma, Baxter, Cardinal Health, Danone-Nutricia, Dim-3, Fresenius Kabi, GE Healthcare, Mermaid, Lyric, and Rousselot.

Author details

¹Department of Clinical Nutrition, Vrije Universiteit Brussel, Laarbeeklaan 101, 1090 Brussels, Belgium. ²Intensive Care Unit, UZ Brussel, Vrije Universiteit Brussel, Laarbeeklaan 101, 1090 Brussels, Belgium. ³Department of Intensive Care Medicine and Research, Gelderse Vallei Hospital, Willy Brandtlaan 10, 6716 RP Ede, The Netherlands. ⁴Division of Human Nutrition and Health, Chair Group Nutritional Biology, Wageningen University and Research, HELIX (Building 124), Stippeneng 4, 6708 WE Wageningen, The Netherlands.

Received: 31 March 2022 Accepted: 25 April 2022

Published online: 05 May 2022

References

1. Moonen HPFX, Beckers KJH, van Zanten ARH. Energy expenditure and indirect calorimetry in critical illness and convalescence: current evidence and practical considerations. *J Intensive Care*. 2021;9(1):8. <https://doi.org/10.1186/s40560-021-00524-0>.
2. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, Hiesmayr M, Mayer K, Montejo JC, Pichard C, Preiser JC, van Zanten ARH, Oczkowski S, Szczeklik W, Bischoff SC. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr*. 2019;38(1):48–79. <https://doi.org/10.1016/j.clnu.2018.08.037>.
3. Zusman O, Theilla M, Cohen J, Kagan I, Bendavid I, Singer P. Resting energy expenditure, calorie and protein consumption in critically ill patients: a retrospective cohort study. *Crit Care*. 2016;20(1):367. <https://doi.org/10.1186/s13054-016-1538-4>.
4. Tatucu-Babet OA, Fetterplace K, Lambell K, Miller E, Deane AM, Ridley EJ. Is energy delivery guided by indirect calorimetry associated with improved clinical outcomes in critically ill patients? A systematic review and meta-analysis. *Nutr Metab Insights*. 2020;13:1178638820903295. <https://doi.org/10.1177/1178638820903295>.
5. Duan JY, Zheng WH, Zhou H, Xu Y, Huang HB. Energy delivery guided by indirect calorimetry in critically ill patients: a systematic review and meta-analysis. *Crit Care*. 2021;25(1):88. <https://doi.org/10.1186/s13054-021-03508-6>.
6. Pertzov B, Bar-Yoseph H, Menndel Y, Bendavid I, Kagan I, Glass YD, Singer P. The effect of indirect calorimetry guided isocaloric nutrition on mortality in critically ill patients—a systematic review and meta-analysis. *Eur J Clin Nutr*. 2022;76(1):5–15. <https://doi.org/10.1038/s41430-021-00919-0>.
7. De Waele E, Opsomer T, Honoré PM, Diltoer M, Mattens S, Huyghens L, Spapen H. Measured versus calculated resting energy expenditure in critically ill adult patients. Do mathematics match the gold standard? *Minerva Anesthesiol*. 2015;81(3):272–82.
8. Koekkoek WAC, Xiaochen G, van Dijk D, van Zanten ARH. Resting energy expenditure by indirect calorimetry versus the ventilator-VCO₂ derived method in critically ill patients: the DREAM-VCO₂ prospective comparative study. *Clin Nutr ESPEN*. 2020;39:137–43. <https://doi.org/10.1016/j.clnesp.2020.07.005>.
9. Van Zanten ARH, De Waele E, Wischmeyer PE. Nutrition therapy and critical illness: practical guidance for the ICU, post-ICU, and long-term convalescence phases. *Crit Care*. 2019;23(1):368.
10. Oshima T, Berger MM, De Waele E, Guttormsen AB, Heidegger CP, Hiesmayr M, Singer P, Wernerman J, Pichard C. Indirect calorimetry in nutritional therapy. A position paper by the ICALIC study group. *Clin Nutr*. 2017;36(3):651–62. <https://doi.org/10.1016/j.clnu.2016.06.010>.

11. Chapple LS, Fetterplace K, Asrani V, Burrell A, Cheng AC, Collins P, Doola R, Ferrie S, Marshall AP, Ridley EJ. Nutrition management for critically and acutely unwell hospitalised patients with coronavirus disease 2019 (COVID-19) in Australia and New Zealand. *Nutr Diet*. 2020;77(4):426–36. <https://doi.org/10.1111/1747-0080.12636>.
12. Whittle J, Molinger J, MacLeod D, Haines K, Wischmeyer PE; LEEP-COVID Study Group. Persistent hypermetabolism and longitudinal energy expenditure in critically ill patients with COVID-19. *Crit Care*. 2020;24(1):581. <https://doi.org/10.1186/s13054-020-03286-7>.
13. Jonckheer J, Demol J, Lanckmans K, Malbrain MLNG, Spapen H, De Waele E. MECCIAS trial: Metabolic consequences of continuous veno-venous hemofiltration on indirect calorimetry. *Clin Nutr*. 2020;39(12):3797–803. <https://doi.org/10.1016/j.clnu.2020.04.017>.
14. De Waele E, Jonckheer J, Pen JJ, Demol J, Staessens K, Puis L, La Meir M, Honoré PM, L N G Malbrain M, Spapen HD. Energy expenditure of patients on ECMO: a prospective pilot study. *Acta Anaesthesiol Scand*. 2019;63(3):360–4. <https://doi.org/10.1111/aas.13287>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

