



Determinants of adherence to recommendations on physical activity, red and processed meat intake, and body weight among lynch syndrome patients

M Hoedjes¹ · A Vrieling² · L de Brauwier¹ · A Visser² · E Gómez García³ · N Hoogerbrugge³ · E Kampman⁴

Received: 4 February 2022 / Accepted: 30 August 2022
© The Author(s) 2022

Abstract

This study aimed to identify determinants of adherence to lifestyle and body weight recommendations for cancer prevention among Lynch Syndrome (LS) patients. Cross-sectional baseline data of LS patients participating in the Lifestyle & Lynch (LiLy) study was used to assess determinants of adherence to the World Cancer Research Fund cancer prevention recommendations on body weight, physical activity, and red and processed meat intake. Adherence and potential determinants of adherence were assessed using questionnaires. Multivariable logistic regression analyses were conducted to identify determinants of adherence. Of the 211 participants, 50.2% adhered to the body weight recommendation, 78.7% adhered to the physical activity recommendation, and 33.6% adhered to the red and processed meat recommendation. Being younger and having a higher level of education were associated with adherence to the recommendation on body weight. Having knowledge about the recommendation was associated with adherence to the recommendations on physical activity and red and processed meat. Results confirm that knowledge about recommendations for cancer prevention is an important determinant for adherence and suggest that strategies to increase knowledge should be included in lifestyle promotion targeted at LS patients, along with behavior change techniques influencing other modifiable determinants.

Keywords Lynch Syndrome · Cancer prevention · Physical activity · Body weight · Red and processed meat intake · Determinants

Introduction

Lynch syndrome (LS) is an inherited cancer syndrome characterized by a high hereditary risk of various cancers, primarily in the colorectum and the endometrium [1].

Worldwide, approximately 28,600 cases of LS are newly diagnosed each year⁽²⁾. LS is caused by a germline mutation in one of the mismatch repair (MMR) genes [2, 3]. LS patients have a risk of developing colorectal cancer (CRC) between 22 and 69% up to age 70, as opposed to 1–5% in the general Western population [4–6]. Significant differences have been reported in cumulative cancer risk and risk of different cancer types according to MMR gene mutation type (*MLH1*, *MSH2*, *MSH6* and *PMS2*) [7, 8]. The clinical phenotype of LS has been shown to vary between families, countries, and continents [8], suggesting the importance of the role of environmental and non-genetic factors, such as lifestyle-related factors [9], in the development of cancer [10, 11]. In addition, low penetrance genetic risk factors may be associated with the observed variety in cancer risk among LS patients [12]. The influence of lifestyle-related factors on CRC among LS patients appears to be comparable or even stronger as compared with the general population [11].

✉ M Hoedjes
m.hoedjes@tilburguniversity.edu

¹ Center of Research on Psychological and Somatic disorders, Department of Medical and Clinical Psychology, Tilburg University, PO Box 90153, 5000 LE Tilburg, The Netherlands

² Department for Health Evidence, Radboud university medical center, Nijmegen, The Netherlands

³ Department of Clinical Genetics, Maastricht University Medical Center, Maastricht, The Netherlands

⁴ Division of Human Nutrition and Health, Wageningen University, Wageningen, the Netherlands

Studies investigating the association between lifestyle-related factors and the occurrence of sporadic cancer have shown that lower levels of physical activity and higher body fatness are associated with an increased risk of different types of cancer, including CRC and endometrium cancer [13]. Also, the intake of red and processed meat has been associated with an increased risk of sporadic CRC [13]. Among LS patients, lifestyle-related factors have also been associated with cancer risk. A recent systematic review has shown that early-adulthood overweight/obese weight status and adulthood weight gain may be associated with increased colorectal cancer risk in LS patients [14]. Moreover, a recent meta-analysis has shown that obesity has been associated with an increased risk for colorectal cancer, but only in men with LS [15]. Furthermore, reviews of the current literature among LS patients have shown that high fruit intake and physical activity have been associated with decreased colorectal cancer risk [14], whereas smoking and alcohol consumption have been associated with an increased colorectal cancer risk in LS patients [16].

Based on a large body of scientific evidence for these observed associations, the World Cancer Research Fund (WCRF) has issued lifestyle and body weight recommendations for cancer prevention [13]. Cancer survivors (i.e. those who have been diagnosed with cancer including those who have recovered) are also recommended to meet these lifestyle and body weight recommendations for cancer prevention. Meeting these recommendations has been associated with favorable health-outcomes, such as a higher health-related quality of life, and a decreased risk for type II diabetes mellitus, cardiovascular disease, second primary cancers, cancer recurrences, and mortality [17–20]. Current guidelines from the European Hereditary Tumour Group (EHTG) and European Society of Coloproctology (ESCP) advise health care providers to inform LS patients about the observed associations between lifestyle, body weight and the risk of cancer [16].

We previously found that adherence to WCRF lifestyle and body weight recommendations in LS patients is low and that providing WCRF health promotion materials increased awareness of and knowledge about WCRF recommendations, without increasing psychological distress. However, this did not affect adherence [21]. Little is known on how adherence to these recommendations can best be promoted. Insight into determinants of health behaviors among LS patients is needed to be able to identify what techniques and strategies can be used to achieve health behavior changes in this specific patient population. Apart from our previous qualitative study on determinants of adherence to lifestyle and body weight recommendations among LS patients [22], to our knowledge, no other study has examined determinants of adherence or health behavior change among LS

patients. Data on non-changeable determinants associated with (non-)adherence (such as sociodemographic and certain health-related determinants specific to LS, including cancer diagnosis in personal history and years since LS diagnosis) provides insight into which LS patients specifically should be targeted to improve adherence. Data on changeable determinants associated with (non-)adherence provides insight into which modifiable determinants should be targeted for change and informs about what type of techniques or strategies can be used to positively influence these changeable determinants. Such changeable determinants relevant for LS patients include psychological determinants, such as cancer worry and symptoms of depression. These psychological determinants have been associated with unfavorable lifestyle behaviors in previous studies [23, 24]. Besides, behavior change concepts that are frequently included in theories and models of health behavior change are knowledge (about the recommendations) and awareness (of the influence of lifestyle-related factors on cancer risk) [25]. Knowledge and awareness have been shown to be determinants of health behavior in other populations [26, 27].

The aim of this cross-sectional study was to explore demographic, health-related, behavior change and psychological determinants for adherence to body weight, physical activity, and red and processed meat intake recommendations among LS patients, as these specific recommendations are relevant for LS-related types of cancer (CRC, endometrium) [13].

Methods

Study design

This study uses cross-sectional, baseline data ($n=218$) from the Lifestyle & Lynch (LiLy) study, a randomized controlled trial to test the effect of providing LS patients with WCRF-NL health promotion materials of the WCRF cancer prevention recommendations [21].

Participants and procedure

The LiLy study recruited participants between April and September 2015 at Radboud University Medical Center and Maastricht University Medical Centre. LS patients aged between 18 and 65 years were eligible for participation if LS diagnoses was confirmed by a germline pathogenic variant in one of the MMR-genes (*MLH1*, *MSH2*, *MSH6* or *PMS2*). LS patients were excluded from participation if they had insufficient understanding of the Dutch language or if they were participating in the GeoLynch study, a prospective

cohort study among LS patients, to prevent interference between both studies [28]. Since only 4% of eligible LS patients participated in the GeoLynch study, this is unlikely to have biased the results of this study. More information on the LiLy study can be found elsewhere [21].

After informed consent was obtained, eligible LS patients were asked to fill in the baseline questionnaire, which took approximately 45 min to complete. The medical ethical research committees of the Radboud University Medical Center and Maastricht University Medical Centre granted permission to perform this study.

Measures

Adherence to the WCRF recommendations

For this study, adherence to the WCRF recommendations on physical activity, body weight, and red and processed meat intake were included. These recommendations were included as these are relevant for LS-related types of cancer (CRC, endometrium) [13] and the smallest group of each of these dichotomous outcome variables (adherence vs. non-adherence to these recommendations) was large enough to be able to be incorporated into the statistical analyses given the sample size ($n = 211$) [29].

Body weight

Self-reported body weight and height were used to calculate Body Mass Index (BMI) (kg/m^2). Participants were categorised into the following BMI categories: <18.5 ; $18.5-24.9$; $25-29.9$; and ≥ 30 kg/m^2 . The category $18.5-24.9$ kg/m^2 was considered as adherent to the body weight recommendation. The other categories were considered not to be adherent to the body weight recommendation.

Physical activity

Adherence to the physical activity recommendation was assessed using the validated Short Questionnaire to Assess Health Enhancing Physical Activity (SQUASH) questionnaire, which contains questions about multiple activities referring to a normal week in the past month. Results were converted to time spent on light, moderate, and vigorous activities, which were then converted to activity scores [30]. When the number of moderate and vigorous exercise activities was at least 30 min a day, for a minimum of 5 days a week, patients were categorized as adherent to the physical activity recommendation.

Red and processed meat

Intake of red and processed meat was measured with an adjusted version of a 40-item, validated Food Frequency Questionnaire (FFQ) specifically developed to assess adherence to the Dutch Guidelines for a healthy diet [31]. Items assessing red meat intake (grams per week) and processed meat intake (grams per week) during the last month were used to determine adherence to the recommendation on red and processed meat intake. When red meat intake was <500 g/w, of which processed meat was <3 g/d, participants were considered to adhere to the recommendation on red and processed meat intake.

Demographic and health-related characteristics

Demographic characteristics were assessed using the baseline questionnaire, which included items on age, gender, marital status, and education. Clinical characteristics were assessed using the same questionnaire by items on personal and family cancer history, colon surgery (colectomy, hemicolectomy, colon resection), time since LS diagnosis, and smoking behaviour.

Behavior change and psychological characteristics

Awareness Awareness of the cancer risk factors as described in the WCRF/AICR recommendations for cancer prevention (referred to as awareness of the WCRF/AICR recommendations) was assessed using a question from the AICR Cancer Risk Awareness Survey: “Do the following factors have a significant influence on whether or not the average person develops cancer?”

From the exposures that were mentioned in the entire Awareness questionnaire reflecting all recommendations, only the exposures related to the recommendations on body weight, insufficient physical activity, and red and processed meat intake were included for the current study. For each exposure, answer options were: “yes, a big influence”; “yes, a small influence”; “no”; and “I do not know”. Participants with correct answers, indicating that the exposures were of influence, were considered to be aware of the specific cancer risk factors while participants with answers “no” and “I do not know” were considered to be unaware.

Knowledge Knowledge of the WCRF recommendations on body weight, physical activity, and red and processed meat intake was assessed using 3 multiple choice questions; 1 for each recommendation. These knowledge questions required more detailed content-specific knowledge about the recommendations. For example, the multiple choice question “What is the minimally recommended amount of time a day you should be spending on physical activity according to the

recommendations for cancer prevention?”, assessed knowledge about the physical activity recommendation. The 5 answer options included: “A recommendation regarding physical activity and cancer risk does not exist”; “A minimum of 30 minutes physical activity per day of moderate intensity (meaning an increased breath and heart rate)”; “A minimum of 60 minutes physical activity per day of moderate intensity”; “A minimum of 90 minutes physical activity per day of moderate intensity”; “I don’t know”. Participants with correct answers were considered to have knowledge about the respective recommendation.

Cancer risk perception Cancer risk perception was assessed by two standardized questions. Participants were asked to express their perceived cancer risk in a percentage between 0 and 100. In addition they were asked to choose one out of 5 categories: ranging from a very low to a very high perceived cancer risk [32].

Symptoms of depression Symptoms of depression were measured by using the Dutch version of the Hospital Anxiety and Depression Scale (HADS) [33]. The HADS consists of 14 items assessing self-reported symptoms of anxiety (7 items) and depression (7 items) in the past week. Each item is scored on a 4-point Likert scale, ranging from 0 to 3, with higher scores indicating more symptoms. For the current study, only scores for symptoms of depression were used (because of the conceptual overlap with cancer worry). A total score can be calculated for symptoms of depression by adding up the scores on the 7 items. This total score ranges from 0 to 21, with higher scores indicating more symptoms [33].

Cancer worry Cancer worry was assessed using the Cancer Worry Scale (CWS), consisting of 8 items. The reliability and validity has shown to be good among breast and colorectal cancer survivors [34, 35]. The total score ranges between 8 and 32, with higher scores corresponding to more cancer worry.

Statistical analyses

The population for analysis consisted of participants with complete baseline data. Participants with missing data on one or more of the variables included in the analyses were excluded from the analyses.

Means with standard deviations (SD) and frequency tables were used to describe potential socio-demographic, health-related, and psychological determinants. Since the variables ‘age’ and ‘time since LS diagnosis’ were not normally distributed, these variables were incorporated in the statistical analyses as categorical variables. Age was categorized into the following categories based on the observed data distribution: 21–43 years; 44–54 years; and 55–73

years. Time since LS diagnosis was categorized into the following categories: 0–2 years; 2–4 years; and 4–20 years.

First, univariable logistic regression analyses were conducted with adherence to one of the WCRF recommendations on body weight, physical activity, or red and processed meat intake as dependent dichotomous variable, and a single potential determinant as independent variable. The following potential demographic determinants were included as independent variables: gender (male, female); age (21–43, 44–54, and 55–73 years), education level (low, medium, high), and marital status (partner, no partner). The following potential health-related determinants were included: years since LS diagnosis (0–2 years, 2–4 years, and 4–20 years), colon surgery (yes, no), personal cancer history (yes, no), and smoking status (current, ex-, never smoker). The following potential psychological determinants were included: awareness (yes, no) and knowledge of the recommendations (yes, no), symptoms of depression (continuous), cancer worry (continuous), and cancer risk perception (<50%, 50%, >50%).

Subsequently, multivariable logistic regression analyses were conducted with adherence to each recommendation as dependent variable, and as independent variables all socio-demographic, health-related, and psychological characteristics that were found to be statistically significantly associated with adherence in the univariable logistic regression analyses.

Statistical analyses were performed using IBM SPSS Statistics for Windows, version 24. P-values < 0.05 were considered to be statistically significant.

Results

Of the 218 LS patients who agreed to participate in the study, seven participants with missing data on one or more of the variables were excluded and 211 were included in the population for analysis. Participants were aged between 21 and 73 years (mean 48.2; SD 10.9), and 61.1% was female (N = 129) (Table 1). The number of years since LS diagnosis ranged between 0 and 20 years (mean 3.7; SD 2.7). 18% had had a type of colon surgery (colectomy n = 7, hemicolectomy n = 24, colon resection n = 7).

The majority of participants were aware of the influence of or had knowledge about the recommendation on body weight (73% and 64.5%, respectively) and physical activity (66.8% and 64.5%, respectively) in relation to cancer risk. Much less participants were aware of the influence of or had knowledge about the recommendation on red and processed meat intake in relation to cancer risk (37.4% and 14.2%, respectively).

Table 1 Demographic, health-related, behavior change and psychological characteristics in Lynch Syndrome patients (n=211)

	Total (n = 211) N (%)*	Cancer in personal history (n = 75) N (%)*	No cancer in personal history (n = 136) N (%)*
<i>Demographic characteristics</i>			
Gender	129(61.1)	45(60.0)	84(61.8)
Female	82(38.9)	30(40.0)	52(38.2)
Male			
Age at measurement	72(34.1)	10(13.3)	62(45.6)
21–43 years	74(35.1)	34(45.3)	40(29.4)
44–54 years	65(30.8)	31(41.3)	34(25.0)
55–73 years			
Educational level			
Low	20(9.5)	7(9.3)	13(9.6)
Medium	107(50.7)	43(57.3)	64(47.1)
High	84(39.8)	25(33.3)	59(43.4)
Partner	187(88.6)	9(12.0)	15(11.0)
Yes	24(11.4)	66(88.0)	121(89.0)
No			
<i>Health-related characteristics</i>			
Years since LS diagnosis	82(38.9)	27(36.0)	55(40.4)
0–2 years	63(29.9)	18(24.0)	45(33.1)
2–4 years	66(31.3)	30(40.0)	36(26.5)
4–20 years			
Colon surgery	173(82.0)	38(50.7)	135(99.3)
No colon surgery	38(18.0)	37(49.3)	1(0.7)
Colon surgery			
Smoking status	22(10.4)	8(10.7)	14(10.3)
Current smoker	92(43.6)	40(53.3)	52(38.2)
Ex-smoker	97(46.0)	27(36.0)	70(51.5)
Never smoker			
<i>Behavior change and psychological characteristics</i>			
Knowledge			
Weight recommendation			
Yes	111(52.6)	41(54.7)	70(51.5)
No	100(47.4)	34(45.3)	66(48.5)
Physical activity recommendation			
Yes	136(64.5)	50(66.7)	86(63.2)
No	75(35.5)	25(33.3)	50(36.8)
Meat intake recommendation			
Yes	30(14.2)	10(13.3)	20(14.7)
No	181(85.8)	65(86.7)	116(85.3)
Awareness			
Influence of overweight on cancer risk			
Yes	154(73.0)	56(74.7)	98(72.1)
No	57(27.0)	19(25.3)	38(27.9)
Influence of physical activity on cancer risk			
Yes	141(66.8)	52(69.3)	89(65.4)
No	70(33.2)	23(30.7)	47(34.6)
Influence of meat intake on cancer risk			
Yes	79(37.4)	35(46.7)	44(32.4)
No	132(62.6)	40(53.3)	92(67.6)
Symptoms of depression [Mean(SD)]			
Cancer worry [Mean(SD)]	2.78(3.13)	3.4(3.4)	2.4(3.0)
Cancer risk perception	13.8(4.22)	15.1(4.6)	13.1(3.8)
<50%	71(33.6)	24(32.0)	47(34.6)
50%	51(24.2)	16(21.3)	35(25.7)
>50%	89(42.2)	35(46.7)	54(39.7)

*Unless otherwise specified; M = mean; SD = standard deviation; BMI = Body Mass Index

Of the 211 participants, 35.5% had a cancer diagnosis in their personal medical history (n = 75), of which 37 had been

diagnosed with colorectal cancer, 17 with endometrium cancer, 4 with both colorectal and endometrium cancer, and 17

with other types of cancer. Compared with LS patients without a cancer diagnosis in their personal history, LS patients with a cancer diagnosis were older ($p < .000$), had more often had a type of colon surgery ($p < .000$), were more frequently aware of the influence of meat intake on cancer risk ($p = .04$), and had a higher mean score of depressive symptoms ($p = .037$) and cancer worry ($p = .001$). See Table 1.

Adherence to the recommendations

Out of the 211 LS patients, 50.2% adhered to the body weight recommendation, 78.7% adhered to the physical activity recommendation, and 33.6% adhered to the red and processed meat intake recommendation.

Determinants of adherence

Body weight recommendation

The univariable logistic regression analyses showed that age 44–54 vs. <44 years, medium and high vs. low educational level, and symptoms of depression were associated with adherence to the body weight recommendation (Table 2).

In the multivariable analyses, only age 44–54 vs. <44 years (OR 0.48, 95% CI: 0.24–0.94) and medium (OR 4.55, 95% CI: 1.34–15.5) and high (OR 6.41, 95% CI: 1.83–22.5) vs. low educational level remained statistically significantly associated with adherence to the body weight recommendation.

Physical activity recommendation

The univariable logistic regression analyses showed that age 55–73 vs. <44 years, ex-smoking vs. current smoking, and having vs. not having knowledge about the physical activity recommendation were associated with adherence to the physical activity recommendation (Table 3).

In the multivariable analyses, only having knowledge about the physical activity recommendation remained statistically significantly associated with adherence to this recommendation (OR 2.04, 95% CI: 1.04; 3.98).

Red and processed meat intake recommendation

The univariable logistic regression analyses showed that only having vs. not having knowledge about the red and processed meat intake recommendation was associated with adherence to the red and processed meat recommendation (Table 4; OR 2.62, 95% CI: 1.19; 5.74).

Discussion

This first quantitative explorative study on determinants of adherence to WCRF lifestyle and body weight recommendations for cancer prevention in LS patients showed that knowledge about the recommendations was a statistically significant determinant of adherence to the lifestyle recommendations on physical activity and red and processed meat intake. Being younger and having a higher level of education were associated with adherence to the recommendation on body weight.

Adherence to the body weight recommendation among LS patients in the current study was comparable to adherence in the general Dutch population in which 50% of those aged 18 and older adhered to the body weight recommendation [36]. As compared to an observational study in Dutch colorectal cancer survivors, adherence to the recommendations on body weight (50% vs. 34%), physical activity (78.7% vs. 73%), and red and processed meat (33.6% vs. 8%) was higher in the LS patients participating in the current study [37].

To our knowledge, no other studies have quantitatively investigated determinants of adherence to lifestyle and body weight recommendations in LS patients. The results of this first quantitative exploration of determinants of adherence are in accordance with our previous qualitative findings showing that having knowledge about the recommendations serves as a cue to action for adherence to lifestyle recommendations in LS patients [22]. Knowledge is incorporated as a determinant in multiple frequently used theories and models of health behavior change (e.g., the theory of planned behavior, the Health Belief Model, Social Cognitive Theory) [25]. In this study, knowledge was found to be a determinant of adherence to the recommendations on health behaviors (physical activity and red and processed meat intake), but not of adherence to the body weight recommendation. These findings may be explained by the theoretical proximity of the determinant knowledge to a certain health behavior (such as physical activity or red and processed meat intake) as opposed to an outcome of multiple lifestyle behaviors (body weight). Considering that adherence to the body weight recommendation is subject to adherence to recommendations on energy balancing behaviors (physical activity, sedentary behavior, and dietary intake), it seems plausible that knowledge is a more proximal determinant of health behaviors and a more distal determinant of adherence to the body weight recommendation (outcome of the health behaviors physical activity and diet quality). In other words, it makes sense that it's more difficult to influence (the result of) multiple lifestyle behaviors just by increasing knowledge than it is to influence a single lifestyle behavior. Hence, this could explain our finding that

Table 2 Demographic, health-related, behavior change and psychological characteristics of Lynch Syndrome patients (n=211) and associations with adherence to the WCRF recommendation on body weight [1]

	Non-adherent N = 105	Adherent N = 106	Univariable ^{[[2]]}	Multivariable ^{[[3]]}
	N(%)	N(%)	OR(95%CI)	OR(95%CI)
<i>Demographic characteristics</i>				
Gender	59(56.2)	70(66.0)	1.52(0.87–2.65)	
Female	46(43.8)	36(34.0)	1	
Male				
Age at measurement	28(26.7)	44(41.5)	1	1
21–43 years	43(41.0)	31(29.2)	0.46(0.24–	0.48(0.24–0.94)*
44–54 years	34(32.4)	31(29.2)	0.89)*	0.86(0.41–1.79)
55–73 years			0.58(0.29–1.14)	
Education level				
Low	16(15.2)	4(3.8)	1	1
Medium	56(53.3)	51(48.1)	3.64(1.14–	4.55(1.34–15.5)*
High	33(31.4)	51(48.1)	11.6)*	6.41(1.83–
			6.18(1.90–	22.5)**
			20.1)**	
Partner	97(92.4)	90(84.9)	0.46(0.19–1.14)	
Yes	8(7.6)	16(15.1)	1	
No				
<i>Health-related characteristics</i>				
Years since LS diagnosis	35(33.3)	47(44.3)	1	
0–2 years	33(31.4)	30(28.3)	0.68	
2–4 years	37(35.2)	29(27.4)	(0.35–1.31)	
4–20 years			0.58(0.30–1.12)	
Colon surgery	81(77.1)	92(86.8)	1	
No surgery	24(22.9)	14(13.2)	1.95(0.94–4.02)	
Surgery				
Cancer in personal history				
Yes	44(41.9)	31(29.2)	0.57(0.32–1.01)	
No	61(58.1)	75(70.8)	1	
Smoking status	12(11.4)	10(9.4)	1	
Current smoker	54(51.4)	38(35.8)	0.84(0.33–2.15)	
Ex-smoker	39(37.1)	58(54.7)	1.79(0.70–4.53)	
Never smoker				
<i>Behavior change and psychological characteristics</i>				
Knowledge				
Yes	55(52.4)	50(47.2)	1.02(0.59–1.75)	
No	50(47.6)	56(52.8)	1	
Awareness				
Yes	76(72.4)	78(73.6)	1.06(0.58–1.95)	
No	29(27.6)	28(26.4)	1	
Symptoms of depression[Mean(SD)] [4]	3.25(3.22)	2.32(2.98)	0.91(0.83–	0.94(0.85–1.03)
			0.99)*	
Cancer worry [Mean(SD)] [5]	14.4(4.42)	13.3(3.95)	0.94(0.88–1.00)	
Cancer risk perception	39(37.1)	32(30.2)	1	
<50%	27(25.7)	24(22.6)	1.08(0.53–2.23)	
50%	39(37.1)	50(47.2)	1.56(0.83–2.93)	
>50%				

*p < .05, **p < .01, ***p < .001

¹Body weight recommendation: Body Mass Index 18.5–24.9 kg/m²

²Odds ratios are derived from univariable logistic regression analyses with adherence to the weight recommendation (yes vs. no) as dependent variable and one sociodemographic, health-related or psychological characteristic as independent variable

³Odds ratios are derived from a multivariable logistic regression analysis with adherence to the weight recommendation (yes vs. no) as dependent variable and all statistically significant (p < .05) sociodemographic, cancer-related, and health-related characteristics in the univariable logistic regression analyses as independent variables

⁴Odds ratio per 1 unit increase in the depressive symptoms subscale of the Hospital Anxiety and Depression Scale

⁵Odds ratio per 1 unit increase in the Cancer Worry Scale

Table 3 Demographic, health-related, behavior change and psychological characteristics of Lynch Syndrome patients (n=211) and associations with adherence to the WCRF recommendation on physical activity [1]

	Non-adherent N = 45	Adherent N = 166	Univariable ^[2]	Multivariable ^[3]
	N(%)	N(%)	OR(95%CI)	OR(95%CI)
<i>Demographic characteristics</i>				
Gender	23(51.1)	106(63.9)	1.69(0.87–3.29)	
Female	22(48.9)	60(36.1)	1	
Male				
Age at measurement	16(35.6)	56(33.7)	1	1
21–43 years	23(51.1)	51(30.7)	0.63(0.30–1.33)	0.54(0.25–1.19)
44–54 years	6(13.3)	59(35.5)	2.81(1.03–	2.44(0.85–6.97)
55–73 years			7.69)*	
Education level				
Low	3(6.7)	17(10.2)	1	
Medium	22(48.9)	85(51.2)	0.68(0.18–2.54)	
High	20(44.4)	64(38.6)	0.57(0.15–2.13)	
Partner	42(93.3)	145(87.3)	0.49(0.14–1.73)	
Yes	3(6.7)	21(12.7)	1	
No				
<i>Health-related characteristics</i>				
Years since LS diagnosis	22(48.9)	60(36.1)	1	
0–2 years	12(26.7)	51(30.7)	1.56(0.70–3.46)	
2–4 years	11(24.4)	55(33.1)	1.83(0.82–4.13)	
4–20 years				
Colon surgery	33(73.3)	140(84.3)	1	
No surgery	12(26.7)	26(15.7)	1.96(0.90–4.28)	
Surgery				
Cancer in personal history				
Yes	20(44.4)	55(33.1)	0.62(0.32–1.21)	
No	25(55.6)	111(66.9)	1	
Smoking status	8(17.8)	14(8.4)	1	1
Current smoker	14(31.1)	78(47.0)	3.18(1.13–9.00)*	2.59(0.87–7.74)
Ex-smoker	23(51.1)	74(44.6)	1.84(0.69–4.93)	1.72(0.60–4.95)
Never smoker				
<i>Behavior change and psychological characteristics</i>				
Knowledge				
Yes	23(51.1)	113(68.1)	2.04(1.04–	2.22(1.09–4.52)*
No	22(48.9)	53(31.9)	3.98)*	1
			1	
Awareness				
Yes	27(60.0)	114(68.7)	1.46(0.74–2.89)	
No	18(40.0)	52(31.3)	1	
Symptoms of depression [Mean(SD)] [4]	3.42(2.86)	2.61(3.19)	0.93(0.84–1.02)	
Cancer worry [Mean(SD)] [5]	13.9 (4.68)	13.8(4.09)	0.99(0.92–1.07)	
Cancer risk perception	14(31.1)	57(34.3)	1	
<50%	13(28.9)	38(22.9)	0.72(0.30–1.70)	
50%	18(40.0)	71(42.8)	0.97(0.44–2.12)	
>50%				

* $p < .05$, ** $p < .01$, *** $p < .001$ ¹Physical activity recommendation: moderate to vigorous activities for at least 30 min a day, for a minimum of 5 days a week²Odds ratios are derived from univariable logistic regression analyses with adherence to the physical activity recommendation as dependent variable and one sociodemographic, health-related or psychological characteristic as independent variable³Odds ratios are derived from a multivariable logistic regression analysis with adherence to the physical activity recommendation (yes vs. no) as dependent variable and all statistically significant ($p < .05$) sociodemographic, health-related, and psychological characteristics in the univariable logistic regression analyses as independent variables⁴Odds ratio per 1 unit increase in the depressive symptoms subscale of the Hospital Anxiety and Depression Scale⁵Odds ratio per 1 unit increase in the Cancer Worry Scale

Table 4 Demographic, health-related, behavior change and psychological characteristics of Lynch Syndrome patients (n=211) and associations with adherence to the WCRF recommendation on red and processed meat intake [1]

	Non-adherent N = 140	Adherent N = 71	Univariable ^{[[2]]}	Multivariable ^{[[3]]}
	N(%)	N(%)	OR(95%CI)	OR(95%CI)
<i>Demographic characteristics</i>				
Gender	82(58.6)	47(66.2)	1.39(0.76–2.51)	
Female	58(41.4)	24(33.8)	1	
Male				
Age at measurement	48(34.3)	24(33.8)	1	
21–43 years	49(35.0)	25(35.2)	1.02(0.51–2.03)	
44–54 years	43(30.7)	22(31.0)	1.02(0.50–2.08)	
55–73 years				
Education level				
Low	14(10.0)	6(8.5)	1	
Medium	75(53.6)	32(45.1)	0.99(0.35–2.82)	
High	51(36.4)	33(46.5)	1.51(0.53–4.32)	
Partner	121(86.4)	66(93.0)	2.07(0.74–5.81)	
Yes	19(13.6)	5(7.0)	1	
No				
<i>Health-related characteristics</i>				
Years since LS diagnosis	58(41.4)	24(33.8)	1	
0–2 years	38(27.1)	25(35.2)	1.59(0.80–3.18)	
2–4 years	44(31.4)	22(31.0)	1.21(0.60–2.43)	
4–20 years				
Colon surgery	111(79.3)	62(87.3)	1	
No surgery	29(20.7)	9(12.7)	1.80(0.80–4.04)	
Surgery				
Cancer in personal history				
Yes	55(39.3)	20(28.2)	0.61(0.33–1.13)	
No	85(60.7)	51(71.8)	1	
Smoking status	19(13.6)	3(4.2)	1	
Current smoker	59(42.1)	33(46.5)	3.54(0.98–12.9)	
Ex-smoker	62(44.3)	35(49.3)	3.58(0.99–12.9)	
Never smoker				
<i>Behavior change and psychological characteristics</i>				
Knowledge				
Yes	14(10.0)	16(22.5)	2.62(1.19–	2.62(1.19–5.74)*
No	126(90.0)	55(77.5)	5.74)*	1
			1	
Awareness				
Yes	50(35.7)	29(40.8)	1.24(0.69–2.23)	
No	90(64.3)	42(59.2)	1	
Symptoms of depression [Mean(SD)] [4]	2.87(3.13)	2.61(3.16)	0.97(0.89–1.07)	
Cancer worry [Mean(SD)] [5]	14.1(4.52)	13.2(3.48)	0.94(0.88–1.01)	
Cancer risk perception	46(32.9)	25(35.5)	1	
<50%	32(22.9)	19(26.8)	1.09(0.52–2.31)	
50%	62(44.3)	27(38.0)	0.80(0.41–1.56)	
>50%				

* $p < .05$, ** $p < .01$, *** $p < .001$

¹Meat intake recommendation: <500 g/w red meat, <3 g/d processed meat

²Odds ratios are derived from univariable logistic regression analyses with adherence to the WCRF red and processed meat intake recommendation (yes vs. no) as dependent variable and one sociodemographic, health-related or psychological characteristic as independent variable

³The independent variable Knowledge is the only variable that was statistically significantly ($p < .05$) associated with adherence to the WCRF red and processed meat intake recommendation (yes vs. no) in the univariable logistic regression analyses

⁴Odds ratio per 1 unit increase in the depressive symptoms subscale of the Hospital Anxiety and Depression Scale

⁵Odds ratio per 1 unit increase in the Cancer Worry Scale

knowledge was found to be a statistically significant determinant of the health behaviors physical activity and red and processed meat intake, but not for the outcome of health behaviors (body weight).

The observed association between adherence to the body weight recommendation and educational level is in line with previous research. A large Canadian cross-sectional study examining determinants of adherence to WCRF recommendations in the general population, also found that higher education attainment was associated with higher odds of adhering to the recommendation for body weight [38].

It should be noted that most of the potential determinants of adherence included in this study did not show a statistically significant association with adherence to recommendations on body weight, physical activity, and red and processed meat intake. Contrary to our expectations, having a cancer diagnosis in one's personal medical history was not found to be statistically significantly associated with adherence. This seems to be in disagreement with the presumed window of opportunity for lifestyle change after a cancer diagnosis that has been described in the scientific literature on health behavior change after a cancer diagnosis [39]. In addition, time after LS diagnosis also was not found to be statistically significantly associated with adherence.

Strengths and limitations

A strength of this first quantitative study on determinants of adherence to WCRF lifestyle recommendations for cancer prevention in LS patients is the relatively large sample size ($n=211$) in relation to the number of LS patients (estimated 10-year prevalence of 3.316 in the Netherlands) [40, 41]. Other strengths include the extensive assessment of adherence to the recommendations and potential determinants and the use of widely-used validated questionnaires.

Several limitations should be considered when interpreting the results of this study. Our study sample consisted of LS patients who agreed to participate in a study about lifestyle and cancer risk (response rate 53%). LS patients who participated were more likely to be older, female, and to have had a previous diagnosis of cancer compared with those who did not participate. Therefore, our study sample may not be a representative sample of LS patients. In addition, our sample consisted of a relatively high proportion of highly educated individuals, which may limit the generalizability of our findings and may reflect an overestimation of the proportion of LS patients having knowledge about the recommendations. Furthermore, while interpreting our findings, it should be taken into account that adherence to lifestyle and body weight recommendations was assessed using self-report questionnaires, which may have led to over-reporting of healthy lifestyle behavior and under-reporting

of body weight, particularly among individuals with overweight or obesity [42, 43]. Additionally, the sample size ($n=211$) was too small to be able to enter all independent variables into one multivariable logistic regression analyses as the validated rule of thumb of a minimum of 10 participants per independent categorical variable in the smallest group would have been violated [44]. Therefore, only the independent variables that were statistically significantly associated with adherence were entered into the multivariable logistic regression analyses. It should also be noted that we did not distinguish between different MMR genes in our statistical analyses, while the cumulative cancer risk and the risk of different cancer types differs according to MMR gene mutation type [7]. Since we found that having been diagnosed with (any type of) cancer was not associated with adherence this is not expected to influence our results. Finally, it should be noted that there are many more possible determinants of health behavior change that we did not incorporate in this study that may have influenced adherence. Such possible determinants include for example social and environmental factors, which should be incorporated in future studies to provide a more comprehensive picture of the determinants of adherence to lifestyle recommendations in LS patients.

The results of this study confirm the importance of having knowledge about lifestyle recommendations and suggest that such knowledge should be promoted to achieve adherence. Our previous publication about the LiLy study has shown that knowledge about lifestyle recommendations can be increased by providing LS patients with WCRF-NL health promotion materials [21]. Health care providers involved in (follow-up) care for LS patients (such as genetic counsellors, clinical geneticists, gastro-enterologists, gynecologists) could easily incorporate providing WCRF-NL health promotion materials during counselling or surveillance visits with LS patients. Informing LS patients about lifestyle-related factors (including the preventive use of aspirin [45]) and cancer risk is in line with current guidelines for LS patients [16]. Increasing knowledge, by providing health promotion materials or referring to online health education material (e.g., via the international and national websites of the WCRF such as www.wcrf.org), is an important first step to achieve adherence. When health care professionals provide these materials, this is in itself an additional behavior change technique (credible source) [46]. However, as our previous study and many others have shown, health behavior change is not likely to be achieved by solely providing information [21, 47]. Although information provision is an important first step towards health behavior change, typically, a combination of multiple behavior change techniques and strategies targeting a multitude of health behavior determinants is needed to achieve and maintain health behavior

changes. Therefore, health care professionals treating LS patients could refer to other health care professionals specialized in health behavior change (such as a dietician, physical therapist, or a lifestyle coach). They could provide these additional behavior change techniques to achieve health behavior changes and to improve health outcomes in LS patients.

Conclusion

The results of this first quantitative study on determinants of adherence to WCRF lifestyle and body weight recommendations among LS patients confirm that knowledge about the recommendation is an important determinant for adherence to the recommendations on physical activity and red and processed meat intake. Results suggest that strategies to increase knowledge (e.g., providing health education materials) should be included in lifestyle promotion targeted at LS patients, along with behavior change techniques influencing other modifiable determinants of adherence.

Acknowledgements The World Cancer Research Fund (WCRF) financially supported this study (grant number: SG2013/1019): <http://www.wcrf.org>. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data Availability Data is available upon request from the corresponding author.

Declarations

Competing interests The authors have declared no conflicting interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Barrow E, Hill J, Evans DG (2013) Cancer risk in Lynch syndrome. *Fam Cancer* 12(2):229–240
- Jenkins MA, Baglietto L, Dowty JG et al (2006) Cancer risks for mismatch repair gene mutation carriers: a population-based early onset case-family study. *Clin Gastroenterol Hepatol* 4(4):489–498
- Steinke V, Engel C, Büttner R, Schackert HK, Schmiegel WH, Propping P (2013) Hereditary nonpolyposis colorectal cancer (HNPCC)/Lynch syndrome. *Deutsches Ärzteblatt International* 110(3):32
- Barrow E, Alduaij W, Robinson L et al (2008) Colorectal cancer in HNPCC: cumulative lifetime incidence, survival and tumour distribution. A report of 121 families with proven mutations. *Clin Genet* Sep 74(3):233–242. doi:<https://doi.org/10.1111/j.1399-0004.2008.01035.x>
- Jenkins MA, Baglietto L, Dowty JG et al (2006) Cancer risks for mismatch repair gene mutation carriers: a population-based early onset case-family study. *Clin Gastroenterol Hepatol* Apr 4(4):489–498. doi:<https://doi.org/10.1016/j.cgh.2006.01.002>
- Quehenberger F, Vasen HF, van Houwelingen HC (2005) Risk of colorectal and endometrial cancer for carriers of mutations of the hMLH1 and hMSH2 gene: correction for ascertainment. *J Med Genet* Jun 42(6):491–496. doi:<https://doi.org/10.1136/jmg.2004.024299>
- Dominguez-Valentin M, Sampson JR, Seppala TT et al (Jan 2020) Cancer risks by gene, age, and gender in 6350 carriers of pathogenic mismatch repair variants: findings from the Prospective Lynch Syndrome Database. *Genet Med* 22(1):15–25. doi:<https://doi.org/10.1038/s41436-019-0596-9>
- International Mismatch Repair C (2021) Variation in the risk of colorectal cancer in families with Lynch syndrome: a retrospective cohort study. *lancet Oncol* Jul 22(7):1014–1022. doi:[https://doi.org/10.1016/S1470-2045\(21\)00189-3](https://doi.org/10.1016/S1470-2045(21)00189-3)
- Wiseman M, The Second World Cancer Research Fund/American Institute for Cancer Research Expert Report (2008). Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective: Nutrition Society and BAPEN Medical Symposium on 'Nutrition support in cancer therapy'. *Proceedings of the Nutrition Society*. :67(3):253–256
- Park JG, Park YJ, Wijnen JT, Vasen HF (1999) Gene-environment interaction in hereditary nonpolyposis colorectal cancer with implications for diagnosis and genetic testing. *Int J Cancer* 82(4):516–519
- van Duijnhoven FJ, Botma A, Winkels R, Nagengast FM, Vasen HF, Kampman E (2013) Do lifestyle factors influence colorectal cancer risk in Lynch syndrome? *Fam Cancer* 12(2):285–293
- Donald N, Malik S, McGuire JL, Monahan KJ (2018) The association of low penetrance genetic risk modifiers with colorectal cancer in lynch syndrome patients: a systematic review and meta-analysis. *Fam Cancer* Jan 17(1):43–52. doi:<https://doi.org/10.1007/s10689-017-9995-8>
- Research WCRFAIfC. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Continues Update Project Expert Report 2018. Available at <http://www.dietandcancerreport.org/>
- Coletta AM, Peterson SK, Gatus LA et al (2019) Energy balance related lifestyle factors and risk of endometrial and colorectal cancer among individuals with lynch syndrome: a systematic review. *Fam Cancer* Oct 18(4):399–420. doi:<https://doi.org/10.1007/s10689-019-00135-7>
- Lazzeroni M, Bellerba F, Calvello M et al (2021) A Meta-Analysis of Obesity and Risk of Colorectal Cancer in Patients with Lynch Syndrome: The Impact of Sex and Genetics. *Nutrients* May 20(5). doi:<https://doi.org/10.3390/nu13051736>
- Seppala TT, Latchford A, Negroi I et al (2021) European guidelines from the EHTG and ESCP for Lynch syndrome: an updated third edition of the Mallorca guidelines based on gene and gender. *Br J Surg* May 27(5):484–498. doi:<https://doi.org/10.1002/bjs.11902>
- Meyerhardt JA, Heseltine D, Niedzwiecki D et al (2006) :24(22):3535-41. doi:<https://doi.org/10.1200/JCO.2006.06.0863>

18. Meyerhardt JA, Niedzwiecki D, Hollis D et al (2007) Association of dietary patterns with cancer recurrence and survival in patients with stage III colon cancer. *Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't. Jama* Aug 15(7):754–764. doi:<https://doi.org/10.1001/jama.298.7.754>
19. Meyerhardt JA, Niedzwiecki D, Hollis D et al (2008) Impact of body mass index and weight change after treatment on cancer recurrence and survival in patients with stage III colon cancer: findings from Cancer and Leukemia Group B 89803. *Randomized Controlled Trial Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't. J Clin Oncol* Sep 1(25):4109–4115. doi:<https://doi.org/10.1200/JCO.2007.15.6687>
20. Blanchard CM, Courneya KS, Stein K (2008) Cancer survivors' adherence to lifestyle behavior recommendations and associations with health-related quality of life: results from the American Cancer Society's SCS-II. *Research Support, Non-U.S. Gov't. J Clin Oncol* May 1(13):2198–2204. doi:<https://doi.org/10.1200/JCO.2007.14.6217>
21. Vrieling A, Visser A, Hoedjes M et al (2018) Increasing awareness and knowledge of lifestyle recommendations for cancer prevention in Lynch syndrome carriers: Randomized controlled trial. *Clin Genet* Jan 93(1):67–77. doi:<https://doi.org/10.1111/cge.13076>
22. Visser A, Vrieling A, Murugesu L, Hoogerbrugge N, Kampman E, Hoedjes M (2017) Determinants of adherence to recommendations for cancer prevention among Lynch Syndrome mutation carriers: A qualitative exploration. *PLoS ONE* 12(6):e0178205. doi:<https://doi.org/10.1371/journal.pone.0178205>
23. Wang X, Arafa A, Liu K, Eshak ES, Hu Y, Dong JY (2021) Combined healthy lifestyle and depressive symptoms: a meta-analysis of observational studies. *J Affect disorders* Jun 15:289:144–150. doi:<https://doi.org/10.1016/j.jad.2021.04.030>
24. Driessen KAJ, de Rooij BH, Vos MC et al (2021) Cancer-related psychosocial factors and self-reported changes in lifestyle among gynecological cancer survivors: cross-sectional analysis of PROFILES registry data. *Support Care Cancer* Aug 28. doi:<https://doi.org/10.1007/s00520-021-06433-0>
25. Michie S, West R, Campbell R, Brown J, Gainforth H (2014) *ABC of Behaviour Change Theories: an essential resource for researchers, policy makers, and practitioners*. Silverback Publishing
26. Jeruszka-Bielak M, Kollajtis-Dolowy A, Santoro A et al (2018) Are Nutrition-Related Knowledge and Attitudes Reflected in Lifestyle and Health Among Elderly People? A Study Across Five European Countries. *Front Physiol* 9:994. doi:<https://doi.org/10.3389/fphys.2018.00994>
27. Okosun IS, Davis-Smith M, Seale JP (2012) Awareness of diabetes risks is associated with healthy lifestyle behavior in diabetes free American adults: evidence from a nationally representative sample. *Prim Care Diabetes* Jul 6(2):87–94. doi:<https://doi.org/10.1016/j.pcd.2011.12.005>
28. Botma A, Nagengast FM, Braem M et al (2010) Body mass index increases risk of colorectal adenomas in men with Lynch syndrome: the GEOLynch cohort study. *J Clin Oncol* 28(28):4346–4353
29. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *Research Support US(1996) Gov't, Non-P.H.S. Journal of clinical epidemiology*. Dec ;49(12):1373-9
30. Wendel-Vos GW, Schuit AJ, Saris WH, Kromhout D (2003) Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 56(12):1163–1169
31. van Lee L, Geelen A, van Huysduynen EJH, de Vries JH, van't Veer P, Feskens EJ (2012) The Dutch Healthy Diet index (DHD-index): an instrument to measure adherence to the Dutch Guidelines for a Healthy Diet. *Nutr J* 11(1):49
32. Vrieling A, Visser A, Hoedjes M et al (2018) Increasing awareness and knowledge of lifestyle recommendations for cancer prevention in Lynch syndrome carriers: Randomized controlled trial. *Clin Genet* 93(1):67–77
33. Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta psychiatrica Scandinavica* Jun 67(6):361–370
34. Custers JAE, Gielissen MFM, Janssen SHV, de Wilt JHW, Prins JB (2016) Fear of cancer recurrence in colorectal cancer survivors. *Support Care Cancer* Feb 24(2):555–562. doi:<https://doi.org/10.1007/s00520-015-2808-4>
35. Custers JA, van den Berg SW, van Laarhoven HW, Bleiker EM, Gielissen MF, Prins JB (2014) The Cancer Worry Scale: detecting fear of recurrence in breast cancer survivors. *Cancer Nurs* Jan-Feb 37(1):E44–50. doi:<https://doi.org/10.1097/NCC.0b013e3182813a17>
36. Volksgezondheidszorg.info. RIVM. Accessed 21st of (2021) <https://www.volksgezondheidszorg.info/onderwerp/overgewicht/cijfers-context/samenvatting>
37. Winkels RM, van Lee L, Beijer S et al (2016) Adherence to the World Cancer Research Fund/American Institute for Cancer Research lifestyle recommendations in colorectal cancer survivors: results of the PROFILES registry. *Cancer Med* Sep 5(9):2587–2595. doi:<https://doi.org/10.1002/cam4.791>
38. Whelan HK, Xu JY, Vaseghi S, Lo Siou G, McGregor SE, Robson PJ (2017) Alberta's Tomorrow Project: adherence to cancer prevention recommendations pertaining to diet, physical activity and body size. *Public health nutrition* May 20(7):1143–1153. doi:<https://doi.org/10.1017/s1368980016003451>
39. Demark-Wahnefried W, Aziz NM, Rowland JH, Pinto BM (2005) Riding the crest of the teachable moment: promoting long-term health after the diagnosis of cancer. *J Clin Oncol* Aug 20(24):5814–5830. doi:<https://doi.org/10.1200/JCO.2005.01.230>
40. Stichting MLD(2021) Lynch Syndroom 28th of Accessed 28th of October, 2021. <https://www.mlds.nl/chronische-ziekten/lynch-syndroom/>
41. Volksgezondheidszorg.info. Het voorkomen van dikkedarmkanker. 28th of October(2021) Accessed 28th of October, 2021. <https://www.volksgezondheidszorg.info/onderwerp/dikkedarmkanker/cijfers-context/huidige-situatie#!node-het-v%C3%B3%C3%B3rkommen-van-dikkedarmkanker>
42. Gosse MA (2014) How accurate is self-reported BMI? *Nutr Bull* 39(1):105–114. doi:DOI: <https://doi.org/10.1111/mbu.12075>
43. Wehling H, Lusher J (2017) People with a body mass index ≥ 30 under-report their dietary intake: A systematic review. *J Health Psychol*. doi:<https://doi.org/10.1177/1359105317714318>
44. Wilson VanVoorhis CR, Morgan BL (2007) Understanding Power and Rules of Thumb for Determining Sample Sizes. *Tutorials in Quantitative Methods for Psychology* 3(2):43–50. <https://www.tqmp.org/RegularArticles/vol03-2/p043/>
45. Serrano D, Patrignani P, Stigliano V et al (2022) Aspirin Colorectal Cancer Prevention in Lynch Syndrome: Recommendations in the Era of Precision Medicine. *Genes (Basel)* Mar 3(3). doi:<https://doi.org/10.3390/genes13030460>
46. Michie S, Richardson M, Johnston M et al (2013) The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* Aug 46(1):81–95. doi:<https://doi.org/10.1007/s12160-013-9486-6>
47. Franz MJ, VanWormer JJ, Crain AL et al (Oct 2007) Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc* 107(10):1755–1767