

# Annual Review of Entomology

# The Biology and Ecology of Parasitoid Wasps of Predatory Arthropods

# Minghui Fei,<sup>1,\*</sup> Rieta Gols,<sup>2</sup> and Jeffrey A. Harvey<sup>3,4</sup>

- <sup>1</sup>Department of Entomology, College of Plant Protection, Nanjing Agricultural University, Nanjing, People's Republic of China; email: feimh@njau.edu.cn
- <sup>2</sup>Laboratory of Entomology, Wageningen University, Wageningen, The Netherlands; email: rieta.gols@wur.nl
- <sup>3</sup>Department of Terrestrial Ecology, Netherlands Institute of Ecology, Wageningen, The Netherlands; email: j.harvey@nioo.knaw.nl
- <sup>4</sup>Animal Ecology Section, Department of Ecological Sciences, Vrije Universiteit Amsterdam, Amsterdam. The Netherlands



### www.annualreviews.org

- · Download figures
- · Navigate cited references
- Keyword search
- · Explore related articles
- Share via email or social media

Annu. Rev. Entomol. 2023. 68:109-28

First published as a Review in Advance on October 5, 2022

The *Annual Review of Entomology* is online at ento.annualreviews.org

https://doi.org/10.1146/annurev-ento-120120-111607

Copyright © 2023 by the author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.

\*Corresponding author



# **Keywords**

biological control, ladybird, natural enemy, trophic cascade, spiders

### Abstract

Parasitoid wasps are important components of insect food chains and have played a central role in biological control programs for over a century. Although the vast majority of parasitoids exploit insect herbivores as hosts, others parasitize predatory insects and arthropods, such as ladybird beetles, hoverflies, lacewings, ground beetles, and spiders, or are hyperparasitoids. Much of the research on the biology and ecology of parasitoids of predators has focused on ladybird beetles, whose parasitoids may interfere with the control of insect pests like aphids by reducing ladybird abundance. Alternatively, parasitoids of the invasive ladybird *Harmonia axyridis* may reduce its harmful impact on native ladybird populations. Different life stages of predatory insects and spiders are susceptible to parasitism to different degrees. Many parasitoids of predators exhibit intricate physiological interrelationships with their hosts, adaptively manipulating host behavior, biology, and ecology in ways that increase parasitoid survival and fitness.

# Primary hyperparasitoid: a (koinobiont endo-) parasitoid that oviposits inside the larvae of another parasitoid that is in turn developing inside the body of a secondary herbivore host

# 1. INTRODUCTION

The study of multitrophic interactions is embedded in our understanding of community ecology and ecosystem functioning (58, 77, 122). Many of the best-studied multitrophic associations involve plant–insect herbivore–natural enemy interactions (129). Among the most important insect natural enemies are parasitoid wasps (Hymenoptera), which are often ubiquitous components of multitrophic interactions and ecological communities (53, 105). Parasitoids develop on or inside the bodies of other organisms (often other insects); the adults are free living and feed primarily on nectar or pollen, although some host-feed as well (52, 72). Because the development of parasitoid eggs and larvae is restricted by the ostensibly finite resources contained in an individual host, they are considered as model organisms for studying life history, development, and reproductive strategies (59, 71). Moreover, the importance of parasitoids in classical biological control programs in agroecosystems over the past century is well documented (65, 162).

Although studies of food chains in terrestrial food webs involving insects have focused on three levels, these food chains do not necessarily stop there and may extend to four or even more trophic levels (63). For example, the larvae and pupae of many parasitoids of insect herbivores are attacked by primary hyperparasitoids and secondary parasitoids, respectively (117). Hyperparasitoids are important components of plant–insect food chains, and their role in structuring insect communities has received considerable attention and has been reviewed several times in this journal (117, 144, 145). However, other natural enemies in the third trophic level may also harbor parasitoid assemblages. For instance, a diverse range of predatory insects and other arthropods such as ladybird beetles, hoverflies, lacewings, ground beetles, and spiders are also hosts to parasitoids. Because of the importance of some of these predators in mediating trophic cascades in natural communities and in biological control, increasing attention is being paid to their parasitoids to highlight both the applied and fundamental importance of these interactions.

In this review, we synthesize studies examining various aspects of the biology and ecology of parasitoids of predatory arthropods. The review begins with a detailed section examining various aspects of the biology and ecology of parasitoids of ladybird beetles, particularly those of the parasitoid wasp Dinocampus coccinellae, which has been the most comprehensively studied. We report on studies describing parasitoid guilds of different developmental stages of ladybird beetles. Lastly, we examine in this section how some parasitoids manipulate ladybird development and behavior in ways that optimize parasitoid fitness. In the subsequent sections, we address similar areas with parasitoids of other predatory insects such as hoverflies, lacewings, ants, and ground beetles. In the next-to-last section, we discuss parasitoids of spiders, including parasitoids that attack spider eggs, egg sacs, juveniles, and adults. We also describe host location strategies in parasitoids and how antagonistic encounters with spiders have led parasitoids to evolve strategies to overcome spider behavioral defenses. The adaptive manipulation of spider web-building behavior is also discussed. We conclude by discussing the role of parasitoids of predatory arthropods in biological control from both positive and negative perspectives and suggest avenues for future research. In this review, we only include non-stinging parasitoids within the Parasitica (68). Some stinging wasps in the Aculeata that also use spiders as hosts (e.g., Pompilidae) are also often described as parasitoids (i.e. 99, 141). However, for the sake of clarity, we exclude pompilids from this review.

# 2. PARASITOIDS OF PREDATORY LADYBIRD BEETLES

Most species in the large family Coccinellidae, which contains more than 6,000 species in approximately 360 genera, are predators of Sternorrhyncha (aphids, scale insects, whiteflies, and stinkbugs), but some species feed on other prey (e.g., caterpillars, ants, and mites), and some are

phytophagous or omnivorous (51). They are important biological control agents of aphids and coccids, in particular, which are more effectively controlled than aphids (32).

# 2.1. Parasitoids of Different Ladybird Beetle Life Stages

Parasitism of ladybird eggs is rare and appears to be restricted to phytophagous ladybird hosts in the Epilachninae by several eulophid parasitoid species (17). Because cannibalism among newly hatched carnivorous ladybird larvae is often high, the risk of predation of parasitized eggs is also high and has reduced selection for such interactions to evolve (17). Larval and pupal stages of predatory ladybird beetles are the most vulnerable stages to attack by parasitoids (19). The best-studied larval parasitoids are in the genus *Oomyzus* (Eulophidae: Tetrastichinae), particularly Oomyzus scaposus, which has a wide distribution and has been described as the most common parasitoid associated with ladybird beetles (19). Larvae of *Oomyzus* species are gregarious, and brood sizes tend to increase with increasing size or later instar of the host at the time of parasitoid oviposition (39, 139). Hymenopteran parasitoids are haplodiploid, with unfertilized eggs developing into males and fertilized eggs developing into females, and parasitoid mothers are able to control the sex of their offspring (27, 52). Sex ratios of both O. scaposus and Oomyzus spiraculus are highly female biased irrespective of host size or instar at parasitoid oviposition (39, 139). This suggests that the species are engaged in local mate competition in which just enough sons are produced to mate with the daughters (131). The genus Homalotylus (Encyrtidae) contains larval and pupal parasitoids that are specialized on parasitizing coccinellids (19). Species attacking small or early host stages are solitary, whereas those attacking large or late stages are gregarious (80). In a study of the gregarious Homalotylus eytelweinii, brood sizes increased with instar at parasitoid oviposition, but adult emergence was higher when early instar hosts were parasitized (57). This suggests that various life history traits are differentially affected by host instar at the time of parasitoid oviposition.

Other larval and pupal parasitoid species of ladybird beetles belong to the Chalcididae (e.g., *Ulga* species), Encyrtidae (genus *Cowperi*), Proctotrupidae (genus *Nothoserphus*), Pteromalidae, and Eulophidae (19). Detailed information on various aspects of the taxonomy and biology of these parasitoids has been provided in other, earlier reviews (17, 19).

# 2.2. Interaction Between Ladybird Beetles and Dinocampus coccinellae

Interactions between the cosmopolitan parasitoid D. coccinellae (Braconidae: Euphorinae) and its coccinellid hosts have been disproportionately well-studied. It is a solitary endoparasitoid of adult ladybird beetles mainly belonging to the subfamily Coccinellinae (19). Although it prefers to parasitize adult ladybird beetles, it may also parasitize pupae and larvae when adults are scarce (47). However, immature parasitoid larvae delay their development in immature ladybird stages and only egress from the imago (73). This host stage preference is considered to be adaptive because parasitism is more successful and development time is shorter when hosts are parasitized in the adult stage (109). Females can parasitize and successfully develop in a range of host species (54 reported species; 17, 18, 19) but have a preference for and are more successful in larger individuals, both across and within species (90, 123, 157). Moreover, D. coccinellae prefers to parasitize female over male hosts (12, 15, 96), even when correcting for size (25). This has been attributed to the fact that female hosts have a greater resource intake than male hosts and thus produce larger progeny (25). In addition, preference and performance are higher in young, newly egressed adults than in older adults emerging from diapause (96). The ability of D. coccinellae to discriminate between parasitized and unparasitized hosts is poorly developed and/or transient (111), as shown by the fact that rates of superparasitism in their hosts in the field can exceed 60% (16).

### **Endoparasitoid:**

a parasitoid that oviposits into the body of the host and whose larvae develop by feeding internally on host hemolymph and fat body

# Superparasitism:

parasitism of a host that has already been parasitized by the same (i.e., self) or another (i.e., conspecific) female parasitoid of the same species The larvae of *D. coccinellae* feed on teratocytes (73, 135), trophic cells that are released in the host hemocoel upon larval hatching and absorb nutrients from the host haemolymph (143) and fat body (135). Teratocytes are only produced by some parasitoid clades and also play a role in immunosuppression (143). Reduced production of these cells may compromise parasitism success in some host species such as *Harmonia axyridis* (42). Interestingly, adult ladybird hosts often survive parasitism and are able to reproduce successfully (103). Allowing the host to remain alive and active following parasitoid egression is adaptive when it increases parasitoid survival, for instance, when the parasitoid manipulates the behavior of the host to provide protection against parasitoids and predators (61, 102; see also Section 2.3).

To locate hosts over short distances, *D. coccinellae* relies on visual and olfactory cues (2, 112, 123). Movement of the host is the initial cue attracting female parasitoids, which explains why mobile hosts are often attacked more frequently than are inactive ones (11). Movement also facilitates parasitism by exposing the preferred site of oviposition on the host (123), where tissues are softer, allowing easy penetration by the ovipositor (123, 135); this site may vary with host species (112). Volatiles secreted by a gland under the ladybird coxae play a role in host recognition but do not seem to be important for oviposition to take place (123). Aposematically colored ladybird beetles have a distinct smell and, upon attack, excrete a fluid containing toxic alkaloids that function as a defense against predators (24). An extract of adult *Coccinella septempunctata*, as well as a pure alkaloid isolated from this extract that elicited a strong electrophysiological response in electroantennogram assays, was attractive to *D. coccinellae* in an olfactometer (2). These results suggest that host-derived chemicals play an important role in the steps leading up to host acceptance. However, the exact identity and source of the chemicals mediating each of the steps are not clear (whether volatile, contact, or those excreted upon attack) (96).

Dinocampus coccinellae has also been studied for its ability to control H. axyridis. This ladybird species has been widely introduced as a biological control agent, but due to its harmful effects on native coccinellids and adverse effects on human health and fruit production, it is now considered invasive in many parts of its introduced range (11, 14). Several studies have shown that H. axyridis is a poor host for D. coccinellae, and therefore, the effectiveness of this parasitoid to control this invasive species is limited (11, 22, 26, 31, 67, 127). Field observations of rates of parasitism of H. axyridis by D. coccinellae vary between 0% and 30%, whereas rates can reach up to almost 50% for other coccinellid hosts (18). Multivoltine parasitoid species, such as D. coccinellae, must synchronize their active period with that of their hosts (16), whose availability in turn depends on that of the host's prey. As a result of this and other factors, levels of parasitism may also vary over the course of a growing season (115, 138, 150). Host dissections showed that, in the field, parasitism rates of H. axyridis and Coleomegilla maculata, a native ladybird species in North America, were similar (67), but due to reduced host suitability, the rate of emergence of adult parasitoids from the former species is much lower (42). This result suggests that, in the field, D. coccinellae may not be able to discriminate between suitable and less suitable host species (67). Alternatively, in the field, D. coccinellae may also attack larval stages of H. axyridis, which are less aggressive (43) and more suitable for offspring development and survival (11, 43). Recent data suggest that populations of D. coccinellae in central Europe are overcoming host immunity: Parasitism rates of *H. axyridis* were found to be as high as 46% (83), which is remarkable given parasitism rates in its native range of 7.5% or less (18).

# 2.3. Host Behavioral Manipulation by Ladybird Parasitoids

Interactions between parasitoids and their hosts are profoundly intimate and frequently involve an array of changes in the biology, ecology, and behavior of hosts that are mediated by the parasitoid




Figure 1

(a) Adult female Dinocampus coccinellae ovipositing into adult ladybird beetle (Coccinella septempunctata).

(b) Mature larva of D. coccinellae egressing from adult ladybird beetle (Coleomegilla maculata). (c,d) Previously parasitized adult C. maculata beetle sitting atop cocoon of D. coccinellae. Photograph in panel a courtesy of Hans Smid (http://bugsinspace.nl). Photographs in panels b and c courtesy of Mathieu Bélanger Morin. Photograph in panel d courtesy of Anand Varma (https://www.varmaphoto.com/).

mother or her offspring (9). Endoparasitic koinobionts in a small number of braconid subfamilies exhibit adaptations whereby their hosts become phenotypic extensions of the parasitoids through some form of physiological manipulation (1). In this case, parasitized hosts are not killed during parasitism, but instead remain alive for up to several days after the mature parasitoid larvae egress, construct cocoons, and pupate (59). The zombified hosts thereafter act as bodyguards for the parasitoid cocoons, defending them against generalist predators or hyperparasitoids (56, 61). One of the best-studied host-parasitoid associations examining redirection of host behavior toward parasitoid cocoon guarding involves D. coccinellae and its ladybird host, C. maculata (101–103). The parasitoid female oviposits into adult ladybird beetles (Figure 1a), and the larvae primarily feed on host haemolymph and fat body before egressing from the abdomen of their host (Figure 1b). The parasitoid then spins its cocoon, after which the ladybird wraps its appendages around it, apparently shielding it from potential predators and hyperparasitoids (Figure 1c,d). Many ladybird beetles biosynthesize alkaloids, which act as repellents against predators (24), making ladybird beetles effective bodyguards for their parasitoids. As in many species that sequester or synthesize toxins in their body, ladybird beetles are aposematically colored and thus advertise their toxicity to natural enemies (146). Host behavioral manipulation appears to be mediated by an RNA virus,

Koinobiont:
a parasitoid that
attacks hosts that
continue to feed and
grow during the
course of parasitism,
only arresting host
development just prior
to parasitoid pupation

injected by the parasitoid female into the host at oviposition (29). This virus replicates in the host's nervous tissues, inducing neuropathological changes that affect behavior (29).

# 3. PARASITOIDS OF HOVERFLIES

Adult hoverflies (Syrphidae: Diptera) feed on nectar and pollen and are important pollinators of flowering plants in wild and agricultural ecosystems. The larvae are insectivorous, feeding on aphids and some other Homoptera (whiteflies and leafhoppers). Thus, both life stages provide important ecosystem services in agricultural systems (34).

The adult stages of many hoverfly species are mimics of stinging wasps and bees, reducing their risk of becoming prey themselves. However, the immature stages of syrphid flies are attacked by a range of hymenopteran parasitoids in the families Diplazontinae (Ichneumonidae), Figitidae (Cynipoidea), Encyrtidae and Pteromalidae (Chalcidoidea), and Megasplidae (Ceraphronoidea) (44, 74, 130). Species in the subfamily Diplazontinae are generally associated with Syrphidae (81), with the exception of the cosmopolitan *Diplazon laetatorius*, which has a broad host range (75, 164) and is also frequently recovered from parasitized hoverflies (44, 55, 70). Species in this subfamily are koinobiont parasitoids, attacking egg or early larval stages of their host and egressing as adults from the pupae (44).

The genus *Pachyneuron* (Chalcidoidea: Pteromalidae) contains species that are hyperparasitoids of parasitoids of sap-sucking insects, as well as primary and hyperparasitoids of their predators, including syrphid flies, coccinellid beetles, and chrysopids (lacewings) (49). It is not clear which host stage(s) these wasp species parasitize, but the pupae or cocoons are likely to be the most vulnerable (49).

Parasitism rates of syrphids in agricultural fields have been found to range from 5% to 45% (136), and parasitism by *D. laetatorius* alone can reach levels of more than 20% (70). Parasitoids of syrphids often attack more than one host species, i.e., they are considered generalists, although their host range may be overestimated (128).

### 4. PARASITOIDS OF LACEWINGS

Green (Neuroptera: Chrysopidae) and brown (Neuroptera: Hemerobiidae) lacewings are beneficial insects in agroecosystems, especially in orchards (142). The larval stages are carnivorous and feed on plant-sucking arthropods, such as aphids, soft scales (Hemiptera; Coccoidea), and mites, as well as other soft-bodied insect life stages (e.g., eggs and early larval stages) (142). Adults of brown lacewings and those of green lacewings in the genus *Chrysoperla* are omnivorous, whereas all other adult green lacewings are nonpredacious (28). Green lacewing species in the genera *Chrysoper* and *Chrysoperla* are most commonly used in integrated pest management programs, particularly species of the *Chrysoperla carnea* complex (common green lacewings), which are also frequently found in field crops (142). *Chrysoperla carnea* sensu latu is the best-studied lacewing species and is an important predator used in greenhouses. It is thus not surprising that most of the known lacewing parasitoids are those associated with this species.

Egg parasitoids of chrysopids predominantly belong to the genus *Telonomus* (Scelionidae) (3, 48, 76). In addition, *Trichogramma atopovirilia* (Trichogrammatidae) and *Myartsevaia chrysopae* (Encyrtidae) have been recovered from chrysopid eggs in Mexico (121), where the control of *Melanaphis sacchari* (sugarcane aphid) largely relies on the release of *C. carnea. Trichogramma* species are often used to control lepidopteran pests (137, 167), but parasitism of nontarget hosts, such as egg stages of lacewing predators, may reduce biological control efficacy when trichogrammatid host fidelity is limited or modified under field conditions (132).

Larval stages of chrysopids are predominantly attacked by chalcidoid parasitoids, e.g., *Catolaccus* sp. (Pteromalidae), *Tetrastychus* sp. and *Isodromus* sp. (Eulophidae), and *Perilampud* sp.

(Perilampidae) (3, 48, 76, 151). Perilampid species lay eggs in prey patches of their hosts. The first peripampid larval stage is free living (planidia) and externally attaches itself to the lacewing host larva, whereas the other parasitoids oviposit directly into the body of the host (48). The trophic status of perilampid species is not always clear, although they are often referred to as hyperparasitoids (3). Identified pupal parasitoids of *Chrysoperla* include *Baryscapus* sp. (Eulophidae) and *Gelis ilicicolator* (Ichneumonidae) (3, 76). Species in the genus *Chrysopophthorus* (Braconidae) are koinobiont endoparasitoids of adult green lacewings (3, 154). Interestingly, as in some adult parasitoids of ladybird beetles, the chrysopid host may survive, and females may continue to lay eggs after the *Chrysopophthorus* parasitoid larva has egressed (154). Parasitism rates of chrysopids in the field are highly variable but can reach levels of up to 80%, predominantly by *Tetrastycus* and *Telenomus* species, depending on biogeographical region (3, 48, 76).

Parasitoids of brown lacewings have been little studied. One exception is *Anacharis zealandica* (Figitidae), a solitary larval koinobiont endoparasitoid of the Tasmanian lacewing *Micromus tasmaniae*, which is an important predator of aphids in grasslands and on vegetable crops in Australia and New Zealand (69, 126). These studies demonstrated that the addition of floral resources can be used as an alternative resource for this omnivorous lacewing but can also provide nectar to *A. zealandica* parasitoids and increase their numbers. Interestingly, the addition of flower resources only affected biological control (negatively) at high aphid densities and not at low densities, showing that these multitrophic interactions are complex and context dependent.

### 5. PARASITOIDS OF PREDATORY GROUND BEETLES

Ground beetles (Carabidae) are a very well-studied family in the Coleoptera, with over 40,000 described species worldwide (94). They are abundant in most terrestrial habitats, including cold polar regions (94). The larvae and adults of most species are predators of other invertebrates, although some species are omnivorous as adults and may feed on plant materials like seeds and pollen (94). Many species are considered to be important as generalist predators in biological control programs against a range of pests (95).

Very little is known about egg parasitoids of carabids. Females of certain species in the genera *Dicaelus*, *Chlaenius*, and *Galerita* lay their eggs in mud or clay cells above the soil (79). Their eggs have been found to be parasitized by *Hoplogryon* (Scelionidae, Teleasinae). A common egg parasitoid of *Chlaenius* species is *Prosacantha caraborum* (Scelionidae) (79). Parasitoids in the Proctotrupidae are larval endoparasitoids that are often associated with Carabidae and other soil arthropods (149). In general, females in this family are wingless and dwell in the soil. Given that these parasitoids are efficient soil excavators, parasitism of carabid larvae most likely occurs in the host's burrow (23). The most common parasitoids of carabids in this family are the gregarious *Phaenoserphus viator* and the solitary *Phaenoserphus pallipes*, which have been recovered from *Pterostichus* and *Notiophilus* species (23, 149). Levels of parasitism by these species can reach 25% (23). Parasitism of adults is rare. For instance, in a study of parasitoids of adult riparian carabid beetles in the genus *Bembidion*, no hymenopteran parasitoids were found (4). The described parasitoids of adult carabid stages are braconid wasps in the genus *Microtomus*, which emerged from field-collected *Harpalus rufipes*, of which the larvae are herbivorous and the adults omnivorous, and carnivorous *Galeritus* sp. (23, 66, 125).

# 6. PARASITOIDS OF PREDATORY ANTS

Many species in the Formicidae are ground-dwelling predators or scavengers of arthropods. Some species may attack humans and livestock (e.g., *Solenopsis* sp.) (33). Ants often supplement their diet

### **Phoresy:**

an association whereby one organism disperses while attached to the body of another organism, which generally confers no fitness costs to the host

# **Ectoparasitoid:**

at parasitoid that attaches its egg(s) externally to the host at oviposition and whose larvae feed externally on host tissues

Idiobiont: a parasitoid that attacks non-growing host stages, such as eggs or pupae, or that permanently paralyzes the host, keeping host resources static throughout the interaction

with plant tissues and honeydew (163). Given their omnipresence and often broad dietary breadth, ants can be important biological control agents of agricultural pests (33, 163). However, some ant species require relatively high densities of honeydew-producing Hemiptera (Sternorrhyncha families) to supplement their energy needs and may feed on other beneficial arthropods.

Species in two hymenopteran parasitoid taxa have well-established relationships with ants: the family Eucharitidae (Chalcidoidea) and the tribe Neoneurinae (Braconidae) (38, 91). There are parasitoids of ants in other hymenopteran taxa, but the evidence demonstrating that they are true parasitoids of ants, and not parasitoids of myrmecophilous arthropods, is circumstantial (91). Eucharitid wasps parasitize larvae of ants in 53 genera distributed over three families (Oraseminae, Eucharitinae, and Gollumielinae) (91). Most eucharitids have a host range restricted to a single genus. Females deposit their tiny eggs singly (Oresema) or in clusters of up to 10,000 eggs (Stilbula tenuicornis) in or on specific plant tissues (leaves, buds), depending on the species (20). The first-instar larvae, or planidea, actively seek worker ants or prey of ants and use phoresy for transportation to the ant nest, where they attack mature larvae and pupae and develop as endo- or ectoparasitoids (91). They egress as adults from the host pupa and avoid detection by the ant by active or passive chemical mimicry of ant colony odors (91). Worker ants may even assist the adult to egress from the cocoon or remove partially opened cocoons from the nest without attacking them (7). Species in the tribe Neoneurinae are koinobiont endoparasitoids that attack adult worker ants and develop in the abdomen of their host (91). Full-grown Elasmosoma sp. larvae exit the host, often through the host anus, and pupate in the soil (118). Parasitism rates are highly variable but can reach levels up to 70% or higher for some parasitoid species (91).

# 7. PARASITOIDS OF SPIDERS

Spiders are among the most important and ubiquitous of terrestrial predatory arthropods (108). Currently, there are almost 50,000 described spider species from 129 families (30). With few exceptions, spiders are all predators of other invertebrates, and even small vertebrates (108, 124). They occur in a huge variety of habitats (153) and are considered to be important in biological control programs (124).

Although the vast majority of hymenopteran parasitoids attack other insects, a small number of parasitoid families contain species whose larvae are solitary or gregarious idiobionts and koinobionts that develop on or inside the eggs, juveniles, or adults of spiders. Most hymenopteran parasitoids of spiders are highly specialized and do not exploit other hosts (45). In this section, we discuss egg and egg sac parasitoids and parasitoids of juveniles or adults separately.

# 7.1. Parasitoids of Spider Eggs and Egg Sacs

The parasitoids of spider eggs and egg sacs that have been described come from only a few families. They exploit their host eggs in two quite distinct ways. In the first group, females are idiobionts that probe through the thin silk of spider egg sacs and oviposit into individual spider eggs that are clustered within the sacs. The parasitoid larvae develop solitarily within the eggs and emerge as adults by perforating the egg chorion. Species of solitary parasitoids in the family Platygastridae (subfamily Sceliioninae), such as *Idris flavicornis*, *Ceratobaeus* spp., and *Baeus* spp., and Eurytomidae (subfamily Eurytominae), such as *Philolema latrodecti*, are known to attack the eggs of spiders in several families, including the Lycosidae, Theridiidae, Segestriidae, and Desidae (6, 35, 78, 98, 114, 120). Little is known about the biology, development, or behavior of most spider egg parasitoids. An early study found that *Idris* sp. preferred younger over older egg sacs of the wolf spider, *Pardosa lapidicina* (35), and responded to silk odors of the host egg sac with vigorous tapping movements

of its antennae. This study also found that eggs close to the periphery of the egg sac were more susceptible to parasitism than eggs clustered in the center. By contrast, females of the platygastrid wasp *Ecthrodesis laorali* parasitized every egg in egg sacs of the intertidal spider, *Desis formidabilis*, when they were able to locate and exploit them but preferred microhabitats where its host was most abundant (114). As in many primary parasitoids of herbivorous insects, *E. laorali* and *Ceratobaeus* spp. are protandrous, and their offspring sex ratios are heavily biased in favor of females (5, 114, 156). This suggests that, when a female parasitizes multiple eggs within a single egg sac, the earlier-emerging males mate with female siblings under conditions of local mate competition (110).

The second group consists of parasitoids whose larvae attack multiple eggs simultaneously inside of fully cocooned egg sacs. The majority of these parasitoids are from the family Ichneumonidae (subfamily Cryptinae), although this behavior has also been observed in Pteromalus platyphilus (Pteromalidae) (45). Species in these clades are ectoparasitic idiobionts, and some species are also secondary hyperparasitoids of primary parasitoid cocoons (119). By far the most important parasitoids of spider egg sacs are in the genus Gelis; members of this genus have been recorded in egg sacs of spiders in several families, including Linyphiidae (155), Lycosidae (8, 21, 37), Agelendiae (8), Zodariidae (87), and Theridiosomatidae (89). Females perforate the thin silk of the egg sacs and lay their egg(s) directly onto the surface of the clustered eggs within them. After hatching, the parasitoid larvae feed externally on the eggs and may kill most but not necessarily all of them, although the number killed depends on the relative size of the parasitoid and the egg sac of the spider (13, 35). Most species of Gelis are wingless and mimic ants chemically, behaviorally, and morphologically (62). Despite lacking wings, they are apparently capable of parasitizing egg sacs of spiders that have been suspended via threads (89). For example, Gelis agilis females descend head-first down single threads of silk to reach the cocoons of the primary parasitoid Meteorus pulchricornis suspended at the end of them (60). Adult geline females, which attack primary parasitoid hosts, generally have very low fecundity (158) and produce large, yolky anhydropic eggs (92) that require proteins that are generally obtained by host-feeding by the adult female wasps (72). Although host-feeding from egg sacs has not yet been observed from gelines, it is likely that it also occurs. Other cryptines (i.e., Bathytrix brunnea, Trychosis cyperia) have occasionally also been reported as egg sac parasitoids of spiders (45, 107).

As with parasitoids of spider eggs, the biology and ecology of spiders that produce egg sacs have been little studied. It has been argued that spiders have evolved ecophysiological strategies related to the structure and placement of their egg sacs to resist attack from natural enemies (6). Some spiders even guard their egg sacs against predators and parasitoids (41). However, while these strategies are effective against many generalist predators, such as ants, strong reciprocal selection among specialist parasitoids has led them to become highly efficient at locating and exploiting unattended egg sacs of their spider hosts (6). For example, host searching behavior of Gelis fenestrans, a specialist parasitoid of egg sacs of the linypiid spider Erigone atra, was affected by both plant and host-related odors. In Y-tube experiments, parasitoids preferred odors of wheat over grass and were strongly attracted to silk from E. atra but not to silk of three other species of linyphiids, suggesting strong host specificity (155). Another solitary cryptine, T. ciperia, is a specialist egg sac parasitoid of the crab spider, Misumena vatia (Thomisidae), and parasitoid larvae invariably consume all of the host eggs before pupation. In this interaction, spider host egg mass was positively correlated with adult female, but not male, parasitoid size, although females were larger at eclosion and utilized host resources more efficiently than males (107). Offspring sex ratios of T. ciperia from field-collected egg sacs were also female biased. These admittedly limited data reveal that spider egg sac parasitoids exhibit traits (e.g., sexual size dimorphism, female-biased sex ratios) similar to those of hyperparasitoids and primary parasitoids of herbivorous hosts (52, 59).

### **Protandrous:**

species in which male parasitoids emerge earlier than sibling and nonsibling females from a host patch that facilitates mating when this is time limited

# Secondary parasitoid:

also known as a pseudohyperparasitoid; a (idiobiont ecto-) parasitoid that lays its egg on or inside the prepupae or pupae of a primary parasitoid within its preconstructed cocoon



Figure 2

(a) Adult female Sinarachna pallipes. (b) Mature larva of S. pallipes feeding on adult araneid spider, Eustala anastera. (c) Cocoon of S. pallipes suspended in the web of E. anastera. The hole in the top of the cocoon indicates that an adult parasitoid has recently emerged. (d) Prey capture orbicular web of the araneid Cyclosa conica. (e) Highly modified web of C. conica parasitized by Zatypota picticollis. Photographs in panels a-c courtesy of John Maxwell (https://bugguide.net/node/view/401720). Photographs in panels d-e courtesy of Stanislav Korenko.

This suggests that selection pressures affecting the expression of traits are similar across parasitoids in different trophic levels.

# 7.2. Parasitoids of Juvenile and Adult Spiders

Juvenile and adult spider stages are also attacked by a small but highly specialized group of parasitoids in the *Polysphicta* group (45) (**Figure 2a**). This is a monophyletic lineage of solitary koinobiont ectoparasitoids in the Pimplinae (Ichneumonidae), which contains approximately 190 parasitoid species from 21 genera (46). These parasitoids attack spiders in several families including the Agelenidae (100), Araneidae (36, 54, 82), Dictynidae (84), Tetragnathidae (85), and Theridiidae (40, 86, 148). Host specialization among polysphinctine species occurs at the spider family or even species level (88).

Recent studies have shed considerable light on the complex biology and behavior of these parasitoids. Attack behavior varies among different parasitoid species and involves both host subjugation and mitigation of spider defenses. Direct parasitoid attacks have been observed on spiders resting in exposed locations on the web (36), whereas protracted pursuits may occur when spiders are protected by silk threads in more structurally complex webs or when they escape inside retreats (54). In the latter case, the parasitoid female may adopt a sit-and-wait strategy until the spider leaves its retreat and the pursuit recommences (82). During the oviposition sequence, female parasitoids first inject venom into their spider host, paralyzing it for several moments (82, 165). Host envenomation sometimes requires precise and repeated insertions of the ovipositor into the spider's mouth, presumably reaching the subesophageal ganglion (82). The ovipositors of polysphinctine parasitoids are uniquely adapted for laying eggs onto the antero-lateral dorsal section of the spider's abdomen (147). In some species, the first-instar parasitoid larva remains inside the egg chorion, whereas later instars produce saddles on their seventh or eighth abdominal segments that anchor them to the host abdomen (Figure 2b). The larvae also possess hook-like appendages that are used to hold the silk lines of their host (165). Like egg-sac parasitoids in the Cryptinae and most other pimpline ichneumonids, polysphinctine wasps produce large, anhydropic eggs that presumably require exogenous proteins obtained via host-feeding behavior (72), although this has not yet been described in these parasitoids. During parasitism, the spider host continues foraging and web-building as usual (88). Once the parasitoid larva enters its final instar, it consumes most host tissues, avoiding only sclerotized portions of the spider body (e.g., appendages, cephalothorax). At this point, the mature parasitoid larva spins its cocoon on its host's web. In host spider species where there is pronounced sexual size dimorphism (i.e., large females

and dwarf males), female parasitoids tend to reject male spiders for oviposition because the limited amount of biomass on small hosts decreases offspring size, and thus fitness (86).

# 7.3. Host Behavioral Manipulation by Polysphinctine Parasitoids

As described above for ladybird beetles, some primary parasitoids manipulate the behavior of their hosts before killing them, turning them into bodyguards. Similarly, polysphinctine parasitoids manipulate the ecology and web-spinning behavior of their hosts to optimize their survival to eclosion. In some species, the parasitoid also induces overwintering behavior in its spider host and completes its development the following year (86). After parasitism, Zatypota sp. induces juveniles of the social theridiid spider, Anelosimus eximus, to disperse from communal webs and construct entirely new webs made up of much denser silk with limited openings, giving reduced access to predators or hyperparasitoids (40). Isolation from the communal web is presumably adaptive because it reduces the risk of the adult parasitoid being attacked by other A. eximus spiders at eclosion. Another parasitoid, Brachyzapus niookensis, induces its host, the funnel-web-building spider, Agelena limbata, to lay additional threads (or veils) of both fine and dense silk that cover the spider and protect the entrance of the funnel against generalist predators such as ants (100). The most intricate form of regulation among polysphinctines is documented among parasitoids of spiders that build orbicular webs. Species in several genera (e.g., Zatypota, Reclinervellus, Polysphincta) induce their araneid spider hosts to construct highly modified webs that differ dramatically from the capture webs of healthy spiders. The modified cocoon-webs spun by parasitized spiders use the pre-existing frame lines of the capture webs but lack its characteristic orb and contain a single, strong central thread on which the parasitoid either anchors or suspends its cocoon (Figure 2c-e). This apparently reduces the accessibility of the parasitoid to predators and perhaps hyperparasitoids. Changes in spider web-spinning behavior either are caused by chemicals injected into the host by the adult female parasitoid during oviposition or are introduced into the spider body by parasitoid larva early in the larva's development. When older parasitoid larvae are artificially removed from their spider hosts, the spiders will nevertheless construct altered cocoon webs (36), indicating that the behavioral change is persistent.

# 8. DISCUSSION AND FUTURE PERSPECTIVES

The ecology and biology of parasitoids of arthropod predators have received little attention, with the exception of a few systems such as the ladybird parasitoid D. coccinellae or Zatypota spp. parasitoids of spiders. Most papers on these parasitoids have focused on taxonomy and phylogeny. All predator groups reviewed in this article have parasitoids that are often highly specialized and thus are confined to specific families or genera of predatory hosts. This is particularly true for parasitoids of spiders and ants and can be explained by unique aspects of the biology of these predators. The factors that mediate foraging behavior of parasitoids of arthropod predators in natural environments have to date received little attention. Herbivore-induced plant volatiles are important foraging cues for parasitoids of insect herbivores (152). As the prey of ladybird beetles, hoverflies, and lacewings are herbivorous, prey-induced plant volatiles may also provide reliable cues to guide the parasitoids of these predators to their hosts. For instance, Orre et al. (113) showed that A. zealandica (Figitidae), a parasitoid of lacewing larvae, was attracted to traps baited with methyl salicylate, a ubiquitous compound in volatile blends emitted by plants in response to herbivory (116). Alternatively, host location may largely be mediated by direct cues related to the prey of the predator. There is some evidence that hoverfly larvae that have depleted a prey colony rely on aphid-related cues (e.g., honeydew), and not on plant-related cues, to find new prey patches on other plants (160). Females of ladybird beetles, lacewings, and hoverflies lay their eggs near aphid colonies and may use aphid-derived cues such as honeydew and/or aphid alarm pheromones when locating suitable oviposition sites (161). As prey abundance and colony size are determined by food plant quality, the host plant may only indirectly play a role in this case. How parasitoids of many other arthropod predators find their hosts is even less known.

Ladybird beetles, hoverflies, and lacewings are important biological control agents of aphids, which cause serious economic losses in agriculture not only via their feeding damage, but also by vectoring plant viruses. This may explain why these predators, particularly ladybird beetles, have been relatively well studied, although the importance of ants, ground beetles, and spiders in controlling agricultural pests is increasingly being recognized (10, 33, 166). Some aphid species are myrmecophiles and have evolved a mutualistic relationship with ants, in which ants protect aphids against their enemies in return for energy-rich resources in the form of honeydew (97, 140). Ants and aphid predators may engage in competitive interactions, especially when prey is scarce (97), and some species of aphid predators have evolved specific behavioral, morphological, or chemical




Figure 3

Aphid food web extended to the fifth trophic level. Different colors refer to the various trophic levels or natural enemy status [predators in blue and (hyper)parasitoids in purple and red], with the plant (first trophic level) in green and aphids (second trophic level) in yellow. Parasitoids of aphids are in the third trophic level, whereas those of aphid predators are in the fourth. All parasitoids can be attacked by their own parasitoids (hyperparasitoids). Direct interactions between food-web members are indicated by arrows, with predation in blue and parasitism in red. Other types of direct interactions are indicated by dark gray arrows. Interactions are not restricted to direct ones, but for simplicity, indirect interactions are not considered in this figure. Figure adapted with permission from Reference 50.

traits that facilitate their ability to live in the vicinity of aphid-attending ants (64, 97, 104, 106, 134, 159), which may even provide an enemy-free space for the aphid predators or render them less suitable hosts for their parasitoids (133, 134). For instance, the sticky wax filaments on the larvae of the coccinellid *Azya orbigera*, which feed on coccid prey, provided protection against attack by *Azteca instabilis* ants, which were aggressive towards a parasitoid of the coccinellid *Homalotylus shuvakhinae* (Chalcidoidea: Encyrtidae) (93).

The structure of the aphid-predator food web does not end there. As do parasitoids of herbivorous insects, parasitoids of predatory arthropods have their own parasitoid natural enemies. Because predatory arthropods are members of the third trophic level, their primary parasitoids expand the trophic chain to the fourth level, and their hyperparasitoids in turn expand it to the fifth level. Studies on these hyperparasitoids are scarce, but a few studies have included hyperparasitoid records when surveying parasitism of predatory arthropods [lacewings (3), ladybirds (19), and hoverflies (74)]. Interestingly, whereas hyperparasitoids of primary parasitoids of insect herbivores may compromise efficacy of biological control, hyperparasitoids of predator parasitoids may positively affect control by reducing the abundance of parasitoids attacking predacious biological control agents. To what extent this is the case under natural conditions is an open question. Ecological research should investigate species interactions under conditions of increasing complexity, considering all members of the system, to better understand and predict how changes in one compartment of the food web influences others and vice versa. In Figure 3, we expand the food web proposed by Gilbert (50) by adding parasitoids and hyperparasitoids of the various members in the food web. We realize that considering all of these components simultaneously may be difficult; however, when focusing only on bi- or tritrophic interactions, one should exercise caution when describing patterns or deriving conclusions if all members of the system have not been considered.

# **DISCLOSURE STATEMENT**

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

# **ACKNOWLEDGMENTS**

This review is dedicated to the memory of Edward O. Wilson and Simon Leather. This work was supported by the National Natural Science Foundation of China (NSFC; grant 31800355).

# LITERATURE CITED

- Adamo SA. 2019. Turning your victim into a collaborator: exploitation of insect behavioral control systems by parasitic manipulators. Curr. Opin. Insect Sci. 33:25–29
- Al Abassi S, Birkett MA, Pettersson J, Pickett JA, Wadhams LJ, Woodcock CM. 2001. Response of the ladybird parasitoid *Dinocampus coccinellae* to toxic alkaloids from the seven-spot ladybird, *Coccinella* septempunctata. J. Chem. Ecol. 27:33–43
- Al Rouechdi K, Panis A. 1981. Les parasites de Chrysoperla carnea Steph. (Neuroptera, Chrysopidae) sur Olivier en Provence. Agronomie 1:139–41
- Andersen J, Skorping A. 1991. Parasites of carabid beetles: prevalence depends on habitat selection of the host. Can. 7. Zool. 69:1216–20
- Austin A. 1984. The fecundity, development and host relationships of Ceratobaeus spp. (Hymenoptera: Scelionidae), parasites of spider eggs. Ecol. Entomol. 9:125–38
- Austin AD. 1985. The function of spider egg sacs in relation to parasitoids and predators, with special reference to the Australian fauna. 7. Nat. Hist. 19:359–76
- Ayre GL. 1962. Pseudometagea schwarzii (Ashm.) (Eucharitidae: Hymenoptera), a parasite of Lasius neoniger Emery (Formicidae: Hymenoptera). Can. 7. Zool. 40:157–64

- Barron JR. 1987. Adults and larvae of two new species of Gelis (Hymenoptera: Ichneumonidae) parasitizing eggs of spiders Agelenopsis potteri (Araneae: Agelenidae) and Lycosa rabida (Araneae: Lycosidae). Ann. Entomol. Soc. Am. 80:21–28
- Beckage NE, Gelman DB. 2004. Wasp parasitoid disruption of host development: implications for new biologically based strategies for insect control. Annu. Rev. Entomol. 49:299–330
- Benamú MA, Lacava M, García LF, Santana M, Viera C. 2017. Spiders associated with agroecosystems: roles and perspectives. In *Behaviour and Ecology of Spiders*, ed. C Viera, MO Gonzaga, pp. 275–302. Berlin: Springer
- Berkvens N, Moens J, Berkvens D, Samih MA, Tirry L, De Clercq P. 2010. Dinocampus coccinellae as a parasitoid of the invasive ladybird Harmonia axyridis in Europe. Biol. Control 53:92–99
- Bjørnson S. 2008. Natural enemies of the convergent lady beetle, Hippodamia convergens Guérin-Méneville: their inadvertent importation and potential significance for augmentative biological control. Biol. Control 44:305–11
- Bowden J, Buddle C. 2012. Egg sac parasitism of Arctic wolf spiders (Araneae: Lycosidae) from northwestern North America. 7. Arachnol. 40:348–50
- Brown PM, Thomas CE, Lombaert E, Jeffries DL, Estoup A, Handley L-JL. 2011. The global spread of Harmonia axyridis (Coleoptera: Coccinellidae): distribution, dispersal and routes of invasion. BioControl 56:623–41
- Cartwright B, Eikenbary RD, Angalet GW. 1982. Parasitism by Perilitus coccinellae [Hym.: Braconidae] of indigenous coccinellid hosts and the introduced Coccinella septempunctata [Col.: Coccinellidae], with notes on winter mortality. Entomophaga 27:237–43
- Ceryngier P. 2000. Overwintering of Coccinella septempunctata (Coleoptera: Coccinellidae) at different altitudes in the Karkonosze Mts, SW Poland. Eur. J. Entomol. 97:323–28
- Ceryngier P, Hodek I. 1996. Enemies of Coccinellidae. In Ecology of Coccinellidae, ed. I Hodek, A Honěk, pp. 319–50. Berlin: Springer
- Ceryngier P, Nedvěd O, Grez AA, Riddick EW, Roy HE, et al. 2018. Predators and parasitoids of the harlequin ladybird, *Harmonia axyridis*, in its native range and invaded areas. *Biol. Invasions* 20:1009–31
- Ceryngier P, Roy HE, Poland RL. 2012. Natural enemies of ladybird beetles. In *Ecology and Behaviour of the Ladybird Beetles (Coccinellidae)*, ed. I Hodek, HF van Emden, A Honěk, pp. 375–443. Oxford, UK: Blackwell
- Clausen CP. 1940. The oviposition habits of the Eucharidae (Hymenoptera). J. Wash. Acad. Sci. 30:504– 16
- Cobb LM, Cobb VA. 2004. Occurrence of parasitoid wasps, Baeus sp. and Gelis sp., in the egg sacs of the wolf spiders Pardosa moesta and Pardosa sternalis (Araneae, Lycosidae) in southeastern Idaho. Can. Field-Nat. 118:122–23
- Comont RF, Purse BV, Phillips W, Kunin WE, Hanson M, et al. 2014. Escape from parasitism by the invasive alien ladybird, *Harmonia axyridis. Insect Conserv. Divers.* 7:334

  –42
- 23. Critchley B. 1973. Parasitism of the larvae of some Carabidae (Coleoptera). 7. Entomol. 48:37-42
- Daloze D, Braekman J-C, Pasteels JM. 1994. Ladybird defence alkaloids: structural, chemotaxonomic and biosynthetic aspects (Col.: Coccinellidae). Chemoecology 5:173–83
- Davis DS, Stewart SL, Manica A, Majerus ME. 2006. Adaptive preferential selection of female coccinellid
  hosts by the parasitoid wasp *Dinocampus coccinellae* (Hymenoptera: Braconidae). *Eur. J. Entomol.* 103:41–
  45
- de Castro-Guedes C, de Almeida L. 2016. Laboratory investigations reveal that Harmonia axyridis (Coleoptera: Coccinellidae) is a poor host for Dinocampus coccinellae (Hymenoptera: Braconidae) in Brazil. 7. Insect Sci. 16:e55
- De La Filia AG, Bain SA, Ross L. 2015. Haplodiploidy and the reproductive ecology of Arthropods. Curr. Opin. Insect Sci. 9:36–43
- Devetak D, Klokocovnik V. 2016. The feeding biology of adult lacewings (Neuroptera): a review. Trends Entomol. 12:29–42
- Dheilly NM, Maure F, Ravallec M, Galinier R, Doyon J, et al. 2015. Who is the puppet master? Replication of a parasitic wasp-associated virus correlates with host behaviour manipulation. *Proc. R. Soc. Biol. Sci. B* 282:e20142773

- Dimitrov D, Hormiga G. 2021. Spider diversification through space and time. Annu. Rev. Entomol. 66:225–41
- 31. Dindo ML, Francati S, Lanzoni A, Di Vitantonio C, Marchetti E, et al. 2016. Interactions between the multicolored Asian lady beetle *Harmonia axyridis* and the parasitoid *Dinocampus coccinellae*. *Insects* 7:e67
- Dixon AFG, Hemptinne JL, Kindlmann P. 1997. Effectiveness of ladybirds as biological control agents: patterns and processes. *Entomophaga* 42:71–83
- Drummond F, Choate B. 2011. Ants as biological control agents in agricultural cropping systems. Terr. Arthropod Rev. 4:157–80
- Dunn L, Lequerica M, Reid CR, Latty T. 2020. Dual ecosystem services of syrphid flies (Diptera: Syrphidae): pollinators and biological control agents. Pest Manag. Sci. 76:1973–79
- Eason RR, Peck WB, Whitcomb W. 1967. Notes on spider parasites, including a reference list. J. Kans. Entomol. Soc. 40:422–34
- 36. Eberhard WG. 2000. Spider manipulation by a wasp larva. Nature 406:255-56
- Edgar WD. 1971. Aspects of the ecology and energetics of the egg sac parasites of the wolf spider Pardosa lugubris (Walckenaer). Oecologia 7:155–63
- 38. Feener DH. 2000. Is the assembly of ant communities mediated by parasitoids? Oikos 90:79-88
- Fei M, Hu H, Gols R, Liu S, Wan X, et al. 2021. Development and oviposition strategies in two congeneric gregarious larval-pupal endoparasitoids of the seven-spot ladybird, Coccinella septempunctata. Biol. Control 163:104756
- Fernandez-Fournier P, Straus S, Sharpe R, Avilés L. 2019. Behavioural modification of a social spider by a parasitoid wasp. *Ecol. Entomol.* 44:157–62
- Fink LS. 1987. Green lynx spider egg sacs: sources of mortality and the function of female guarding (Araneae, Oxyopidae). J. Arachnol. 15:231–39
- Firlej A, Lucas E, Coderre D, Boivin G. 2007. Teratocytes growth pattern reflects host suitability in a host-parasitoid assemblage. *Physiol. Entomol.* 32:181–87
- Firlej A, Lucas E, Coderre D, Boivin G. 2010. Impact of host behavioral defenses on parasitization efficacy of a larval and adult parasitoid. *BioControl* 55:339–48
- Fitton M, Rotheray G. 1982. A key to the European genera of diplazontine ichneumon-flies, with notes on the British fauna. Syst. Entomol. 7:311–20
- Fitton M, Shaw MR, Austin A. 1987. The Hymenoptera associated with spiders in Europe. Zool. J. Linn. Soc. 90:65–93
- Gauld ID, Dubois J. 2006. Phylogeny of the Polysphincta group of genera (Hymenoptera: Ichneumonidae; Pimplinae): a taxonomic revision of spider ectoparasitoids. Syst. Entomol. 31:529–64
- Geoghegan IE, Majerus T, Majerus ME. 1998. Differential parasitisation of adult and pre-imaginal Coccinella septempunctata (Coleoptera: Coccinellidae) by Dinocampus coccinellae (Hymenoptera: Braconidae). Eur. J. Entomol. 95:571–79
- 48. Gerling D, Bar D. 1985. Parasitization of *Chysoperla carnea* [Neuroptera, Chrysopidae] in cotton fields of Israel. *Entomophaga* 30:409–14
- 49. Gibson GAP. 2001. The Australian species of *Pachyneuron Walker* (Hymenoptera: Chalcidoidea: Pteromalidae). *J. Hymenoptera Res.* 10:29–54
- 50. Gilbert F. 2005. Syrphid aphidophagous predators in a food-web context. Eur. 7. Entomol. 102:325–33
- Giorgi JA, Vandenberg NJ, McHugh JV, Forrester JA, Ślipiński SA, et al. 2009. The evolution of food preferences in Coccinellidae. *Biol. Control* 51:215–31
- 52. Godfray HCJ. 1994. Parasitoids: Behavioral and Evolutionary Ecology. Princeton, NJ: Princeton Univ. Press
- 53. Godfray HCJ, Shimada M. 1999. Parasitoids as model organisms for ecologists. Popul. Ecol. 41:3-10
- Gonzaga MO, Sobczak JF. 2007. Parasitoid-induced mortality of *Araneus omnicolor* (Araneae, Araneidae) by *Hymenoepimecis* sp. (Hymenoptera, Ichneumonidae) in southeastern Brazil. *Naturwissenschaften* 94:223–27
- Greco C. 1997. Specificity and instar preference of *Diplazon laetatorius* (Hym.: Ichneumonidae) parasitizing aphidophagous syrphids (Dipt.: Syrphidae). *Entomophaga* 42:315–18
- 56. Grosman AH, Janssen A, de Brito EF, Cordeiro EG, Colares F, et al. 2008. Parasitoid increases survival of its pupae by inducing hosts to fight predators. *PLOS ONE* 3:e2276

- Guo Q, Meng L, Fei M, Li B. 2021. Oviposition and development of the gregarious parasitoid Homalotylus eytelweinii (Hymenoptera: Encyrtidae) in relation to host stage of predaceous ladybirds. Biocontrol Sci. Technol. 32:455–66
- Guzman LM, Germain RM, Forbes C, Straus S, O'Connor MI, et al. 2019. Towards a multi-trophic extension of metacommunity ecology. *Ecol. Lett.* 22:19–33
- Harvey JA. 2005. Factors affecting the evolution of development strategies in parasitoid wasps: the importance of functional constraints and incorporating complexity. *Entomol. Exp. Appl.* 117:1–13
- Harvey JA, Gols R, Tanaka T. 2011. Differing success of defense strategies in two parasitoid wasps in protecting their pupae against a secondary hyperparasitoid. *Ann. Entomol. Soc. Am.* 104:1005–11
- Harvey JA, Tanaka T, Kruidhof M, Vet LEM, Gols R. 2011. The "usurpation hypothesis" revisited: Dying caterpillar repels attack from a hyperparasitoid wasp. *Anim. Behav.* 81:1281–87
- 62. Harvey JA, Visser B, Lammers M, Marien J, Gershenzon J, et al. 2018. Ant-like traits in wingless parasitoids repel attack from wolf spiders. *7. Chem. Ecol.* 44:894–904
- Harvey JA, Wagenaar R, Bezemer TM. 2009. Interactions to the fifth trophic level: Secondary and tertiary parasitoid wasps show extraordinary efficiency in utilizing host resources. J. Anim. Ecol. 78:686–92
- Hayashi M, Nomura M. 2011. Larvae of the green lacewing Mallada desjardinsi (Neuroptera: Chrysopidae) protect themselves against aphid-tending ants by carrying dead aphids on their backs. App. Entomol. Zool. 46:407–13
- Heimpel GE, Mills NJ. 2017. Biological Control: Ecology and Applications. Cambridge, UK: Cambridge Univ. Press
- Holland JM. 2002. Carabid beetles: their ecology, survival and use in agroecosystems. In The Agroecology of Carabid Beetles, ed. JM Holland, pp. 1–40. Andover, UK: Intercept Ltd.
- Hoogendoorn M, Heimpel GE. 2002. Indirect interactions between an introduced and a native ladybird beetle species mediated by a shared parasitoid. *Biol. Control* 25:224–30
- Huber JT. 2017. Biodiversity of Hymenoptera. In *Insect Biodiversity: Science and Society*, ed. R Foottit, PH Adler, pp. 419–61. Hoboken, NJ: Wiley
- Jacometti M, Jørgensen N, Wratten S. 2010. Enhancing biological control by an omnivorous lacewing: Floral resources reduce aphid numbers at low aphid densities. *Biol. Control* 55:159–65
- Jankowska B. 2004. Parasitoids of aphidophagous Syrphidae occurring in cabbage aphid (Brevicoryne brassicae L.) colonies on cabbage vegetables. 7. Plant Prot. Res. 44:299–305
- Jervis MA, Ellers J, Harvey JA. 2008. Resource acquisition, allocation, and utilization in parasitoid reproductive strategies. Annu. Rev. Entomol. 53:361–85
- 72. Jervis MA, Kidd NAC. 1986. Host-feeding strategies in hymenopteran parasitoids. Biol. Rev. 61:395–434
- Kadono-Okuda K, Sakurai H, Takeda S, Okuda T. 1995. Synchronous growth of a parasitoid, Perilitus
  coccinellae, and teratocytes with the development of the host, Coccinella septempunctata. Entomol. Exp. Appl.
  75:145–49
- Kamal M. 1926. A study of some hymenopterous parasites of aphidophagous Syrphidae. J. Econ. Entomol. 19:721–30
- 75. Kareem AA, Abbas MA, Aljaafari RK, Al-Zurfi SM, Mouhsan ZM. 2019. New report of hoverfly parasite Diplazon laetatorius (Fabricius, 1781) (Ichneumonidae-Hymenoptera) from Iraq. In IOP Conference Series: Earth and Environmental Science, Vol. 388, The 4th International Conference on Agricultural Sciences (4thICAS), 17–18 November 2019, Agriculture College/University of Kerbala, Kerbala City, Iraq, art. 388. Bristol, UK: IOP Publ.
- Karut K, Kazak C, Arslan A, Şekeroğlu E. 2003. Natural parasitism of Chrysoperla carnea by hymenopterous parasitoids in cotton-growing areas of Çukurova, Turkey. Phytoparasitica 31:90–93
- Kawatsu K, Ushio M, van Veen FF, Kondoh M. 2021. Are networks of trophic interactions sufficient for understanding the dynamics of multi-trophic communities? Analysis of a tri-trophic insect food-web time-series. Ecol. Lett. 24:543–52
- Kessler A, Fokkinga A. 1973. Hymenopterous parasites in egg sacs of spiders of the genus Pardosa (Araneida, Lycosidae). Tijdschr. Entomol. 116:43–61
- King J. 1919. Notes on the biology of the carabid genera Brachynus, Galerita and Chlaenius. Ann. Entomol. Soc. Am. 12:382–88

- Klausnitzer B. 1969. Zur Kenntnis der Entomoparasiten mitteleuropaischer Coccinellidae. Abh. Ber. Naturkundemus. Görlitz 44:1–15
- Klopfstein S. 2014. Revision of the Western Palaearctic Diplazontinae (Hymenoptera, Ichneumonidae).
   Zootaxa 3801:1–143
- Kloss TG, Gonzaga MO, Roxinol JAM, Sperber CF. 2016. Host behavioural manipulation of two orbweaver spiders by parasitoid wasps. *Anim. Behav.* 111:289–96
- Knapp M, Řeřicha M, Maršíková S, Harabiš F, Kadlec T, et al. 2019. Invasive host caught up with a native parasitoid: field data reveal high parasitism of *Harmonia axyridis* by *Dinocampus coccinellae* in Central Europe. *Biol. Invasions* 21:2795–802
- 84. Korenko S. 2017. First record from Italy of *Zatypota anomala* (Ichneumonidae, Ephialtini), a parasitoid of the cribellate spider *Dictyna pusilla* (Araneae, Dictynidae). *Arachnol. Lett.* 54:1–4
- 85. Korenko S, Korenková B, Satrapová J, Hamouzová K, Belgers D. 2015. Modification of *Tetragnatha montana* (Araneae, Tetragnathidae) web architecture induced by larva of the parasitoid *Acrodactyla quadrisculpta* (Hymenoptera, Ichneumonidae, *Polysphincta* genus-group). *Zool. Stud.* 54:e40
- Korenko S, Pekár S. 2011. A parasitoid wasp induces overwintering behaviour in its spider host. PLOS ONE 6:e24628
- Korenko S, Schmidt S, Schwarz M, Gibson GA, Pekar S. 2013. Hymenopteran parasitoids of the anteating spider Zodarion styliferum (Simon) (Araneae, Zodariidae). ZooKeys 262:e3857
- 88. Korenko S, Spasojevic T, Pekár S, Walter GH, Korenková V, et al. 2018. One generalist or several specialist species? Wide host range and diverse manipulations of the hosts' web-building behaviour in the true spider parasitoid *Zatypota kauros* (Hymenoptera: Ichneumonidae). *Insect Conserv. Divers.* 11:587–99
- 89. Kostro-Ambroziak A, Kupryjanowicz J, Schwarz M. 2020. Who wins? Ray spiders (*Theridiosoma gemmosum*) (Araneae: Theridiosomatidae) versus egg sac parasitoids. *J. Arachnol.* 48:90–93
- Koyama S, Thomas CE, Takata M. 2013. Relationship between the size of the parasitoid wasp *Dinocam-pus coccinellae* (Hymenoptera: Braconidae) and host ladybird species (Coleoptera: Coccinellidae). *Trends Entomol*. 9:39–43
- Lachaud J-P, Pérez-Lachaud G. 2012. Diversity of species and behavior of Hymenopteran parasitoids of ants: a review. Psyche J. Entomol. 2012:e134746
- Le Ralec A. 1995. Egg contents in relation to host-feeding in some parasitic Hymenoptera. Entomophaga 40:87–93
- Liere H, Perfecto I. 2014. Cheating on a mutualism: indirect benefits of ant attendance to a coccidophagous coccinellid. Environ. Entomol. 37:143

  –49
- Lövei GL, Sunderland KD. 1996. Ecology and behavior of ground beetles (Coleoptera: Carabidae).
   Annu. Rev. Entomol. 41:231–56
- 95. Luff M. 1987. Biology of polyphagous ground beetles in agriculture. Agric. Zool. Rev. 2:237-78
- Majerus MEN, Geoghegan IE, Majerus TMO. 2000. Adaptive preferential selection of young coccinellid hosts by the parasitoid wasp *Dinocampus coccinellae* (Hymenoptera: Braconidae). *Eur. J. Entomol.* 97:161–64
- Majerus MEN, Sloggett JJ, Godeau J-F, Hemptinne J-L. 2007. Interactions between ants and aphidophagous and coccidophagous ladybirds. *Popul. Ecol.* 49:15–27
- 98. Marie J, Vetter RS. 2015. Establishment of the brown widow spider (Araneae: Theridiidae) and infestation of its egg sacs by a parasitoid, *Philolema latrodecti* (Hymenoptera: Eurytomidae), in French Polynesia and the Cook Islands. *7. Med. Entomol.* 52:1291–98
- Martins A, Gallão J, Bichuette M, Santos E. 2016. The first record of Notocyphus tyrannicus Smith, (Hymenoptera: Pompilidae) as parasitoid of Acanthoscurria ausserer, 1871 (Teraphosidae: Teraphosinae). Bruz. 7. Biol. 76:806–7
- Matsumoto R. 2009. "Veils" against predators: modified web structure of a host spider induced by an ichneumonid parasitoid, *Brachyzapus nikkoensis* (Uchida) (Hymenoptera). J. Insect Behav. 22:39–48
- Maure F, Brodeur J, Droit A, Doyon J, Thomas F. 2013. Bodyguard manipulation in a multipredator context: different processes, same effect. Behav. Process. 99:81–86
- 102. Maure F, Brodeur J, Ponlet N, Doyon J, Firlej A, et al. 2011. The cost of a bodyguard. Biol. Lett. 7:843–46
- Maure F, Doyon J, Thomas F, Brodeur J. 2014. Host behaviour manipulation as an evolutionary route towards attenuation of parasitoid virulence. *J. Evol. Biol.* 27:2871–75

- Milbrath LR, Tauber MJ, Tauber CA. 1993. Prey specificity in Chrysopa: an interspecific comparison of larval feeding and defensive behavior. Ecology 74:1384–93
- Mills N. 2009. Parasitoids. In Encyclopedia of Insects, ed. VH Resh, RT Cardé, pp. 748–51. San Diego: Academic. 2nd ed.
- 106. Mizuno M, Itioka T, Tatematsu Y, Itô Y. 1997. Food utilization of aphidophagous hoverfly larvae (Diptera: Syrphidae, Chamaemyiidae) on herbaceous plants in an urban habitat. *Ecol. Res.* 12:239–48
- Morse DH. 1994. The effect of host size on sex ratio in the ichneumonid wasp, Trychosis cyperia, a spider parasitoid. Am. Midl. Nat. 131:281–87
- Nyffeler M, Birkhofer K. 2017. An estimated 400–800 million tons of prey are annually killed by the global spider community. Sci. Nat. 104:e30
- Obrycki JJ, Tauber MJ, Tauber CA. 1985. Perilitus coccinellae (Hymenoptera: Braconidae): parasitization and development in relation to host-stage attacked. Ann. Entomol. Soc. Am. 78:852–54
- Ode PJ, Vyas DK, Harvey JA. 2022. Extrinsic inter- and intra-specific competition in parasitoid wasps. Annu. Rev. Entomol. 67:305–28
- Okuda T, Ceryngier P. 2000. Host discrimination in *Dinocampus coccinellae* (Hymenoptera: Braconidae), a solitary parasitoid of coccinellid beetles. *Appl. Entomol. Zool.* 35:535–39
- Orr CJ, Obrycki JJ, Flanders RV. 1992. Host-acceptance behavior of Dinocampus coccinellae (Hymenoptera: Braconidae). Ann. Entomol. Soc. Am. 85:722–30
- Orre GUS, Wratten SD, Jonsson M, Hale RJ. 2010. Effects of an herbivore-induced plant volatile on arthropods from three trophic levels in brassicas. *Biol. Control* 53:62–67
- 114. Owen CA, van Noort S, Compton SG, Coetzee JA. 2019. Nest site choice by the intertidal spider Desis formidabilis (Araneae: Desidae) and nest utilisation by its hymenopteran egg parasitoid. Ecol. Entomol. 44:62–70
- Paula DP, Togni PH, Costa VA, Souza LM, Sousa AA, et al. 2021. Scrutinizing the enemy release hypothesis: population effects of parasitoids on *Harmonia axyridis* and local host coccinellids in Brazil. *BioControl* 66:71–82
- Pichersky E, Gershenzon J. 2002. The formation and function of plant volatiles: perfumes for pollinator attraction and defense. Curr. Opin. Plant Biol. 5:237–43
- Poelman EH, Cusumano A, de Boer JG. 2022. The ecology of hyperparasitoids. Annu. Rev. Entomol. 67:143–61
- Poinar G. 2004. Behaviour and development of *Elasmosoma* sp. (Neoneurinae: Braconidae: Hymenoptera), an endoparasite of *Formica* ants (Formicidae: Hymenoptera). *Parasitology* 128:521–31
- Quicke DLJ. 2014. The Braconid and Ichneumonid Parasitoid Wasps: Biology, Systematics, Evolution and Ecology. Hoboken, NJ: Wiley
- Rádai Z, Popovici O, Vas Z, Fusu L. 2018. First record of the parasitoid *Idris flavicornis* (Hymenoptera: Scelionidae) from eggs of the wolf spider *Pardosa agrestis* (Araneae: Lycosidae). *Folia Entomol*. 79:101–6
- Ramírez-Ahuja MDL, Garza-González E, Talamas EJ, Gómez-Govea MA, Rodríguez-Pérez MA, et al. 2020. Parasitoids of Chrysopidae eggs in Sinaloa Mexico. *Insects* 11:e849
- Reiss J, Bridle JR, Montoya JM, Woodward G. 2009. Emerging horizons in biodiversity and ecosystem functioning research. Trends Ecol. Evol. 24:505–14
- Richerson JV, DeLoach CJ. 1972. Some aspects of host selection by *Perilitus coccinellae*. Ann. Entomol. Soc. Am. 65:834–39
- 124. Riechert SE, Lockley T. 1984. Spiders as biological control agents. Annu. Rev. Entomol. 29:299-320
- Rivard I. 1964. Notes on parasitism of ground beetles (Coleoptera: Carabidae) in Ontario. Can. J. Zool. 42:919–20
- 126. Robinson KA. 2009. *Use of floral resources by the lacewing* Micromus tasmaniae *and its parasitoid* Anacharis zealandica, *and the consequences for biological control by* M. tasmaniae. PhD Thesis, Lincoln Univ., PA
- Romero V, Zaviezo T, Grez Villarroel A. 2020. The invasive coccinellid Harmonia axyridis (Coleoptera: Coccinellidae) is a less suitable host for parasitism than resident species. Cienc. Investig. Agrar. 47:312–23
- Rotheray GE. 1984. Host relations, life cycles and multiparasitism in some parasitoids of aphidophagous Syrphidae (Diptera). Ecol. Entomol. 9:303–10
- Schoonhoven LM, van Loon JJA, Dicke M. 2005. Insect-Plant Biology. Oxford, UK: Oxford Univ. Press. 2nd ed.

- Scott EI. 1939. An account of the developmental stages of some aphidophagous Syrphidae (Dipt.) and their parasites (Hymenopt.). Ann. Appl. Biol. 26:509–32
- Shuker DM, Pen I, Duncan AB, Reece SE, West SA. 2005. Sex ratios under asymmetrical local mate competition: theory and a test with parasitoid wasps. Am. Nat. 166:301–16
- 132. Silva IMMS, Stouthamer R. 1999. Do sympatric *Trichogramma* species parasitize the pest insect *Heli-coverpa armigera* and the beneficial insect *Chrysoperla carnea* in different proportions? *Entomol. Exp. Appl.* 92:101–7
- Sloggett JJ, Webberley KM, Majerus ME. 2004. Low parasitoid success on a myrmecophilous host is maintained in the absence of ants. Ecol. Entomol. 29:123–27
- 134. Sloggett JJ, Wood RA, Majerus ME. 1998. Adaptations of Coccinella magnifica Redtenbacher, a myrme-cophilous coccinellid, to aggression by wood ants (Formica rufa group). I. Adult behavioral adaptation, its ecological context and evolution. 7. Insect Behav. 11:889–904
- Sluss R. 1968. Behavioral and anatomical responses of the convergent lady beetle to parasitism by *Perilitus coccinellae* (Schrank) (Hymenoptera: Braconidae). 7. Invertebr. Pathol. 10:9–27
- Smith HA, Chaney WE. 2007. A survey of syrphid predators of Nasonovia ribisnigri in organic lettuce on the central coast of California. 7. Econ. Entomol. 100:39

  –48
- Smith SM. 1996. Biological control with *Trichogramma*: advances, successes, and potential of their use. *Annu. Rev. Entomol.* 41:375–406
- 138. Soleimani S, Madadi H. 2015. Seasonal dynamics of: the pea aphid, Acyrthosiphon pisum (Harris), its natural enemies the seven spotted lady beetle Coccinella septempunctata Linnaeus and variegated lady beetle Hippodamia variegata Goeze, and their parasitoid Dinocampus coccinellae (Schrank). J Plant Prot. Res. 55:jppr-2015-0058
- Song H, Meng L, Li B. 2017. Fitness consequences of body-size-dependent parasitism in a gregarious parasitoid attacking the 7-spot ladybird, Coccinella septempunctata (Coleoptera: Coccinellidae). Biol. Control 113:73–79
- Stadler B, Dixon AFG. 2005. Ecology and evolution of aphid-ant interactions. Annu. Rev. Ecol. Evol. Sys. 36:345–72
- 141. Stanley E, Toscano-Gadea C, Aisenberg A. 2013. Spider hawk in sand dunes: Anoplius bicinctus (Hymenoptera: Pompilidae), a parasitoid wasp of the sex-role reversed spider Allocosa brasiliensis (Araneae: Lycosidae). 7. Insect Behav. 26:514–24
- 142. Stelzl M, Devetak D. 1999. Neuroptera in agricultural ecosystems. Agric. Ecosyst. Environ. 74:305-21
- 143. Strand MR. 2014. Teratocytes and their functions in parasitoids. Curr. Opin. Insect Sci. 6:68–73
- 144. Sullivan DJ. 1987. Insect hyperparasitism. Annu. Rev. Entomol. 32:49-70
- Sullivan DJ, Volkl W. 1999. Hyperparasitism: multitrophic ecology and behavior. Annu. Rev. Entomol. 44:291–315
- Summers K, Speed M, Blount J, Stuckert A. 2015. Are aposematic signals honest? A review. J. Evol. Biol. 28:1583–99
- 147. Takasuka K, Fritzén NR, Tanaka Y, Matsumoto R, Maeto K, Shaw MR. 2018. The changing use of the ovipositor in host shifts by ichneumonid ectoparasitoids of spiders (Hymenoptera, Ichneumonidae, Pimplinae). *Parasite* 25:17
- Takasuka K, Matsumoto R. 2011. Infanticide by a solitary koinobiont ichneumonid ectoparasitoid of spiders. Naturwissenschaften 98:529–36
- 149. Thiele H-U. 1977. Carabid Beetles in Their Environments: A Study on Habitat Selection by Adaptations in Physiology and Behaviour. Berlin: Springer
- Triltsch H. 1996. On the parasitization of the ladybird Coccinella septempunctata L. (Col., Coccinellidae).
   Appl. Entomol. 120:375–78
- 151. Trjapitzin VA, Shuvakhina EY. 2019. Contribution to the knowledge of the encyrtid-wasp genus Iso-dromus Howard, 1887 (Hymenoptera, Encyrtidae: Homalotylini), parasitoids of lacewings (Neuroptera, Chrysopidae) in the Palaearctic. Entomol. Rev. 99:1382–88
- Turlings TCJ, Erb M. 2018. Tritrophic interactions mediated by herbivore-induced plant volatiles: mechanisms, ecological relevance, and application potential. *Annu. Rev. Entomol.* 63:433–52
- 153. Turnbull A. 1973. Ecology of the true spiders (Araneomorphae). Annu. Rev. Entomol. 18:305–48

- 154. van Achterberg C. 1994. The Palaearctic species of the genus Chrysopophthorus Goidanich (Hymenoptera: Braconidae: Euphorinae). Zool. Meded. 68:301–7
- 155. van Baarlen P, Topping CJ, Sunderland KD. 1996. Host location by *Gelis festinans*, an eggsac parasitoid of the linyphiid spider *Erigone atra*. *Entomol. Exp. Appl.* 81:155–63
- 156. van Noort S, Masner L, Popovici O, Valerio AA, Taekul C, et al. 2014. Systematics and biology of the aberrant intertidal parasitoid wasp *Echtbrodesis lamorali* Masner (Hymenoptera: Platygastridae sl.): a parasitoid of spider eggs. *Invertebr. Syst.* 28:1–16
- Vansant H, Vasquez YM, Obrycki JJ, Sethuraman A. 2019. Coccinellid host morphology dictates morphological diversity of the parasitoid wasp *Dinocampus coccinellae*. Biol. Control 133:110–16
- Visser B, Le Lann C, Snaas H, Hardy IC, Harvey JA. 2014. Consequences of resource competition for sex allocation and discriminative behaviors in a hyperparasitoid wasp. *Behav. Ecol. Sociobiol.* 68:105–13
- Völkl W. 1995. Behavioral and morphological adaptations of the coccinellid, *Platynaspis luteorubra* for exploiting ant-attended resources (Coleoptera: Coccinellidae), 7. Insect Behav. 8:653–70
- Vosteen I, Gershenzon J, Kunert G. 2018. Dealing with food shortage: larval dispersal behaviour and survival on non-prey food of the hoverfly Episyrphus balteatus. Ecol. Entomol. 43:578–90
- 161. Vosteen I, Weisser WW, Kunert G. 2016. Is there any evidence that aphid alarm pheromones work as prey and host finding kairomones for natural enemies? *Ecol. Entomol.* 41:1–12
- Wang ZZ, Liu YQ, Shi M, Huang JH, Chen XX. 2019. Parasitoid wasps as effective biological control agents. J. Integr. Agric. 18:705–15
- 163. Way M, Khoo K. 1992. Role of ants in pest management. Annu. Rev. Entomol. 37:479-503
- 164. Weems HV. 1954. Natural enemies and insecticides that are detrimental to beneficial Syrphidae. Ohio J. Sci. 54:45–54
- Weng JL, Barrantes G. 2007. Natural history and larval behavior of the parasitoid Zatypota petronae (Hymenoptera: Ichneumonidae). 7. Hymenoptera Res. 16:326–35
- 166. Woodcock BA, Harrower C, Redhead J, Edwards M, Vanbergen AJ, et al. 2014. National patterns of functional diversity and redundancy in predatory ground beetles and bees associated with key UK arable crops. 7. Appl. Ecol. 51:142–51
- Zang LS, Wang S, Zhang F, Desneux N. 2021. Biological control with Trichogramma in China: history, present status, and perspectives. *Annu. Rev. Entomol.* 66:463–84



# Annual Review of Entomology

# Contents

Volume 68, 2023

Complex and Beautiful: Unraveling the Intricate Communication Systems Among Plants and Insects  *James H. Tumlinson**	1
Chemical Ecology of Floral Resources in Conservation Biological Control Stefano Colazza, Ezio Peri, and Antonino Cusumano	13
Management of Insect Pests with Bt Crops in the United States  Aaron J. Gassmann and Dominic D. Reisig	31
Iron Homeostasis in Insects  Maureen J. Gorman	51
Phoresy and Mites: More Than Just a Free Ride  Owen D. Seeman and David Evans Walter	69
Postcopulatory Behavior of Tephritid Flies  Diana Pérez-Staples and Solana Abraham	89
The Biology and Ecology of Parasitoid Wasps of Predatory Arthropods  Minghui Fei, Rieta Gols, and Jeffrey A. Harvey	109
Dehydration Dynamics in Terrestrial Arthropods: From Water Sensing to Trophic Interactions  Joshua B. Benoit, Kevin E. McCluney, Matthew J. DeGennaro,  and Julian A.T. Dow	129
Biology and Management of the Spotted Lanternfly, <i>Lycorma delicatula</i> (Hemiptera: Fulgoridae), in the United States <i>Julie M. Urban and Heather Leach</i>	151
Historical and Contemporary Control Options Against Bed Bugs,  Cimex spp.  Stephen L. Doggett and Chow-Yang Lee	169
Functional Diversity of Vibrational Signaling Systems in Insects  Meta Virant-Doberlet, Nataša Stritih-Peljhan, Alenka Žunič-Kosi,  and Jernej Polajnar	191
Forest Insect Biosecurity: Processes, Patterns, Predictions, Pitfalls  Helen F. Nahrung, Andrew M. Liebhold, Eckehard G. Brockerhoff,  and Davide Rassati	211

Stingless Bee (Apidae: Apinae: Meliponini) Ecology  David W. Roubik	31
Diapause in Univoltine and Semivoltine Life Cycles  Hideharu Numata and Yoshinori Shintani	:57
Early Monitoring of Forest Wood-Boring Pests with Remote Sensing  Youqing Luo, Huaguo Huang, and Alain Roques	77
Spodoptera frugiperda: Ecology, Evolution, and Management Options of an Invasive Species  Wee Tek Tay, Robert L. Meagher Jr., Cecilia Czepak, and Astrid T. Groot	:99
Molecular Mechanisms of Winter Survival  Nicholas M. Teets, Katie E. Marshall, and Julie A. Reynolds	19
Arthropod and Pathogen Damage on Fossil and Modern Plants:  Exploring the Origins and Evolution of Herbivory on Land  Conrad C. Labandeira and Torsten Wappler	41
The Resilience of Plant–Pollinator Networks  **Jordi Bascompte and Marten Scheffer**	63
The Mechanisms of Silkworm Resistance to the Baculovirus and Antiviral Breeding  Zhaoyang Hu, Feifei Zhu, and Keping Chen	81
Diversity, Form, and Postembryonic Development of Paleozoic Insects  *Jakub Prokop, André Nel, and Michael S. Engel	01
Molecular Mechanisms Underlying Host Plant Specificity in Aphids  Po-Yuan Shih, Akiko Sugio, and Jean-Christophe Simon	-31
Adaptive Plasticity of Insect Eggs in Response to Environmental Challenges Monika Hilker: Hassan Salem, and Nina E. Fatouros	÷51

# Errata

An online log of corrections to *Annual Review of Entomology* articles may be found at http://www.annualreviews.org/errata/ento

# Related Articles

# From the Annual Review of Animal Biosciences, Volume 10 (2022)

Translating Basic Research to Animal Agriculture George E. Seidel Jr.

Concepts and Consequences of a Core Gut Microbiota for Animal Growth and Development

Daphne Perlman, Marina Martínez-Álvaro, Sarah Moraïs, Ianina Altshuler, Live H. Hagen, Elie Jami, Rainer Roehe, Phillip B. Pope, and Itzhak Mizrahi

Host Genetic Determinants of the Microbiome Across Animals:

From Caenorhabditis elegans to Cattle Erica P. Ryu and Emily R. Davenport

Chagas Disease Ecology in the United States: Recent Advances in Understanding Trypanosoma cruzi Transmission Among Triatomines, Wildlife, and Domestic Animals and a Quantitative Synthesis of Vector–Host Interactions Rachel E. Busselman and Sarah A. Hamer

# From the *Annual Review of Genetics*, Volume 56 (2022)

The Genetics of Autophagy in Multicellular Organisms *Hong Zhang* 

# From the Annual Review of Microbiology, Volume 76 (2022)

Division and Transmission: Malaria Parasite Development in the Mosquito David S. Guttery, Mohammad Zeeshan, David J.P. Ferguson, Anthony A. Holder, and Rita Tewari

# From the Annual Review of Phytopathology, Volume 60 (2022)

Yellow Dwarf Viruses of Cereals: Taxonomy and Molecular Mechanisms W. Allen Miller and Zachary Lozier

Future of Bacterial Disease Management in Crop Production

Anuj Sharma, Peter Abrahamian, Renato Carvalho, Manoj Choudhary,

Mathews L. Paret, Gary E. Vallad, and Jeffrey B. Jones

Ecology of Yellow Dwarf Viruses in Crops and Grasslands: Interactions in the Context of Climate Change

Jasmine S. Peters, Beatriz A. Aguirre, Anna DiPaola, and Alison G. Power

# From the Annual Review of Virology, Volume 9 (2022)

Citrus Tristeza Virus: From Pathogen to Panacea Svetlana Y. Folimonova and Yong-Duo Sun

Advances in Understanding Neuropathogenesis of Rift Valley Fever Virus

Kaleigh A. Connors and Amy L. Hartman