

The effect of silicon on the antioxidant system of tomato seedlings exposed to individual and combined nitrogen and water deficit

Annals of Applied Biology

Machado, Joana; Fernandes, Ana Patricia G.; Bokor, Boris; Vaculík, Marek; Heuvelink, Ep et al

<https://doi.org/10.1111/aab.12849>

This publication is made publicly available in the institutional repository of Wageningen University and Research, under the terms of article 25fa of the Dutch Copyright Act, also known as the Amendment Taverne.


Article 25fa states that the author of a short scientific work funded either wholly or partially by Dutch public funds is entitled to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed using the principles as determined in the Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' project. According to these principles research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and / or copyright owner(s) of this work. Any use of the publication or parts of it other than authorised under article 25fa of the Dutch Copyright act is prohibited. Wageningen University & Research and the author(s) of this publication shall not be held responsible or liable for any damages resulting from your (re)use of this publication.

For questions regarding the public availability of this publication please contact openaccess.library@wur.nl

The effect of silicon on the antioxidant system of tomato seedlings exposed to individual and combined nitrogen and water deficit

Joana Machado^{1,2,3}  | Ana Patricia G. Fernandes¹ | Boris Bokor^{4,5} |
Marek Vaculík^{5,6} | Ep Heuvelink³ | Susana M. P. Carvalho¹ |
Marta W. Vasconcelos² 

¹GreenUPorto—Sustainable Agrifood Production Research Centre/Inov4Agro, DGAOT, Faculty of Sciences of University of Porto, Vairão, Portugal

²Universidade Católica Portuguesa, CBQF—Centro de Biotecnologia e Química Fina—Laboratório Associado, Escola Superior de Biotecnologia, Porto, Portugal

³Horticulture and Product Physiology, Wageningen University & Research, Wageningen, The Netherlands

⁴Comenius University Science Park, Bratislava, Slovakia

⁵Department of Plant Physiology, Faculty of Natural Sciences, Comenius University in Bratislava, Bratislava, Slovakia

⁶Institute of Botany, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Bratislava, Slovakia

Correspondence

Marta W. Vasconcelos, Universidade Católica Portuguesa, CBQF—Centro de Biotecnologia e Química Fina—Laboratório Associado, Escola Superior de Biotecnologia, Rua Diogo Botelho 1327, 4169-005 Porto, Portugal.
Email: mvasconcelos@ucp.pt

Funding information

European Cooperation in Science and Technology, Grant/Award Number: CA19116; Fundação para a Ciência e a Tecnologia, Grant/Award Numbers: SFRH/BD/116147/2016, SFRH/BD/05748/2020, UIDB/05748; Norte2020, Grant/Award Number: NORTE-01-0145-FEDER-000041

Abstract

Exploring sustainable strategies for improving crop water and nitrogen use efficiency is essential. Silicon (Si) has been reported as a beneficial metalloid for plants since it alleviates several abiotic stresses (including drought) by triggering the plants' antioxidant system. However, its role in mitigating the negative impact of nitrogen (N) deficit alone or when combined with water (W) deficit is not well studied. This study applied 0 or 2 mM of Na₂SiO₃ to 3-week-old tomato cv. Micro-Tom seedlings that were grown under the following conditions: control (CTR; 100%N + 100% Field Capacity), N deficit (N; 50% N + 100% Field Capacity), water deficit (W; 100% N + 50% Field Capacity) or combined stress (N + W; 50% N + 50% Field Capacity). The Si effect on tomato plant growth depended on the type of stress. Si could only alleviate stress caused by N + W deficit resulting in a higher root dry weight (by 28%), total dry weight (by 23%) and root length (by 37%). Alongside this, there was an increase in the antioxidant (AOX) system activity with the root activity of ascorbate peroxidase and catalase enzymes being enhanced by 48% and by 263%, respectively. Si application also enhanced AOX enzyme activity when tomato plants were subjected to individual deficits but to a lesser extent. In conclusion, Si-treated tomato plants could efficiently modulate their AOX networks in a situation of combined N and water limitation, thus mitigating some of the adverse effects of this combined stress.

KEYWORDS

abiotic stress, antioxidant system, cv Micro-Tom, phenolics, plant growth, silicon

1 | INTRODUCTION

Considering the increasing demand for food and the need for more environmentally friendly agriculture (Jhariya et al., 2019), finding alternative and sustainable solutions to ensure efficient use of water and

nitrogen (N) is a great priority, as these two resources have a huge impact on crop yield (Fernandes et al., 2022). Besides the direct consequences of climate change on water availability, this also brings an additional threat to food security as several studies have demonstrated that the continuous increase of the atmospheric

CO₂ concentration negatively impacts the N status of most C3 plants, leading to lower yield and nutritional crop value (Gojon, 2017).

Numerous morphological, physiological and metabolic changes are common in plants subjected to N or drought stress (reviewed by Machado et al., 2022). Commercial tomato cultivars (*Solanum lycopersicum* L.) are quite sensitive to these deficits at all stages of plant development (Du et al., 2018; Heuvelink et al., 2020) meaning that large amounts of fertilisers and irrigation must be used during their production cycle (Sandhu et al., 2021). Indeed, recent studies that analysed a broad range of tomato genotypes (including wild relatives, landraces, heirlooms and hybrids) concluded that although some variability exists among them, in general this crop is highly sensitive to combined N and water deficits (Machado et al., 2023a; Ruggiero et al., 2022). Moreover, considering the increased consumption of fresh tomato and tomato-derived products and the economic importance of this crop worldwide (Cicco, 2019), it is of high importance to study its responses to these abiotic stresses.

A sustainable method that is receiving great attention in recent years, due to its role on alleviating the negative effects of several abiotic stresses, is the application of silicon (Si) (Bokor et al., 2021; Nunes da Silva et al., 2022). For instance, supplementation with Si has been adopted as an effective strategy for relieving drought stress and improving the drought resistance of several crops (recently reviewed by Bokor et al., 2021; Malik et al., 2021; Thorne et al., 2020; Verma et al., 2022; Wang et al., 2021). In contrast, knowledge about the role of Si in alleviating nutrient deficiencies is far scarcer, as pointed out by Ali et al. (2020) and Pavlovic et al. (2021). In recent studies, it was shown that the application of Si had a positive impact on N uptake, assimilation, utilisation and remobilisation, which in turn can contribute to improving N utilisation within plant tissues (Minden et al., 2020; Pavlovic et al., 2021). Nevertheless, most of this research was conducted on Si-accumulating species such as rice and wheat. Concerning Si non-accumulating species (Si excluders), such as tomato, which exhibit passive absorption and much less accumulation of Si (Hoffmann et al., 2020; Liang et al., 2007), the literature is limited.

One of the key mechanisms proposed by Liang et al. (2007) for the alleviation of abiotic stress in higher plants using Si has to do with the stimulation of the antioxidant system (AOX). Previous reports demonstrated that, exogenously applied, Si can enhance the AOX system under drought stress, including in tomato plants, where a Si-mediated significant decrease of reactive oxygen species (ROS) was observed (Shi et al., 2016). In the same study, the authors reported an increase in the activity of the AOX enzymes superoxide dismutase (SOD) and catalase (CAT) as well as in the concentration of the non-enzymatic antioxidant molecules ascorbic acid and glutathione. For wheat and maize, under drought stress, it has also been shown that Si decreases oxidative damage, ROS, phenolics concentration and ascorbate peroxidase (APX) activity, but increases SOD and CAT (Gong et al., 2005; Parveen et al., 2019; Sattar et al., 2020; Xu et al., 2017). Peroxidase (POD) showed some variability in responses: while some authors found no significant differences (Gong et al., 2005), others noticed a decrease (Xu et al., 2017) or reported an increase in the activity of this enzyme (Parveen et al., 2019; Sattar et al., 2020). Concerning the effects of Si on crops grown at N deficit, most studies focus on the role of Si on N uptake and transport directly and less on the AOX system, which needs further investigation (Ali et al., 2020). Additionally, to the best of our

knowledge, so far there are no studies on the effects of Si on combined N and water deficit, which is surprising given that often plants do not face only one individual abiotic stress, but rather a combination of them. In fact, recently it has been shown that there is a differential behavior between tomato plants grown under combined N and water deficit and single deficits, as well as between plant organs (roots and shoots), which highlights that plants grown under combined stresses employ specific response mechanisms that cannot be elucidated from the responses obtained when these stressors are applied individually (Machado et al., 2023b; Machado et al., 2023c). For instance, AOX system was shown to be highly relevant for tomato acclimation to these combined stresses, with the plants presenting a higher AOX response than the ones subjected to the individual deficits (Machado et al., 2023c). This requires that more research explores this complex acclimation mechanism (García-Martí et al., 2019; Hussain et al., 2018; Pandey et al., 2015). Thus, we hypothesise that tomato responses to Si supplementation under combined N and W deficit cannot be directly predicted from the responses obtained for the individual stresses and that the AOX system plays an important role in the defence mechanism employed by tomato.

This study aimed to: (i) evaluate if Si supplementation can be used as a strategy to help tomato plants to cope with N deficit alone and/or when combined with water deficit; (ii) understand the impact of Si on the AOX responses (antioxidant activity, secondary metabolism responses and enzymatic activity) in plants subjected to single or combined N and W deficit.

2 | MATERIALS AND METHODS

2.1 | Plant material and growth conditions

The experiment was carried out in a growth chamber under controlled environmental conditions (photoperiod: 16 h light; temperature: 25°C day/23°C night; photosynthetic photon flux density [PPFD]: 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$; relative humidity: 70%) according to a complete randomised design and using tomato plants cv. 'Micro-Tom'.

After surface sterilisation with 70% (v/v) ethanol (10 min) and 20% (v/v) sodium hypochlorite (5% v/v active chloride; 6 min), and following multiple washes with deionised water, seeds were sown in potting soil and allowed to germinate. At the stage of third leaf appearance (approximately 3 weeks after sowing), seedlings were selected for uniformity and transplanted to single-plant pots (10 cm high, 10 cm diameter) filled with 0.1–1.5 mm-grade vermiculite (50 g). At this stage plants were divided into eight experimental groups, resulting from the combination of four levels of deficit conditions (control [CTR; 10.5 mM N; 100% field capacity], N deficit [N; 5.3 mM N], water deficit [W; 50% field capacity] and combined stress [N + W; 5.3 mM N; 50% field capacity]) and two levels of Si supply (0 mM or 2 mM of sodium silicate [Na_2SiO_3] solution (27% SiO_2 (w/w) dissolved in 14% NaOH (w/v); Sigma). Immediately before transplanting, all pots were irrigated to 100% field capacity (FC; determined using the soil gravimetric water content method; Joshi et al., 2021), by adding approximately 230 mL of the nutrient solution supplied with 0 or 2 mM of Na_2SiO_3 , depending on the treatment. Two types of

nutrient solutions were used. A full nutrient solution was applied to the CTR and W deficit plants (10 mM NO_3^- ; 0.5 mM NH_4^+ ; 1.9 mM H_2PO_4^- ; 6.1 mM K^+ ; 3.6 mM Ca^{2+} ; 1.6 mM SO_4^{2-} ; 2.5 mM Mg^{2+} ; 2.6 mM Cl^- ; 0.5 mM HCO_3^- [pH 5.8 and electrical conductivity—EC—of 2.0 dS m^{-1}]). Plants subjected to single N deficit or combined N + W deficit were irrigated with a nutrient solution containing half the amount of N (5 mM NO_3^- ; 0.3 mM NH_4^+ ; 1.9 mM H_2PO_4^- ; 6.1 mM K^+ ; 3.6 mM Ca^{2+} ; 4.8 mM SO_4^{2-} ; 2.8 mM Mg^{2+} ; 5.5 mM Cl^- ; 0.5 mM HCO_3^- [pH 5.8 and EC of 2.1 dS m^{-1}]). Both nutrient solutions had the same composition in terms of micronutrients (35 μM Fe-EDDHA; 10 μM Mn^{2+} ; 20.1 μM B^{3+} ; 0.9 μM Cu^{2+} ; 5.0 μM Zn^{2+} ; 0.5 μM MoO_4^{2-}). Following seedling transplanting, all the pots' surface was covered with black plastic to prevent evaporation.

During the experimental period each pot from the CTR and N treatment (with or without Si supplementation), was weighed and re-watered (with distilled water) on a daily basis, to maintain water content at 100% FC. In contrast, for plants under single or combined water deficit (W and N + W) with 0 or 2 mM of Na_2SiO_3 , no additional irrigation was supplied, resulting in a progressive decrease of water content to 50% of FC (day 10) which was then kept at this point (with distilled water for an additional 17 days). At day 27 post-transplantation, all the plants were harvested and used for the analysis of morphological and physiological parameters ($n = 6$). For the biochemical analysis, roots were carefully washed, and plants were quickly separated into roots and leaves being immediately frozen in liquid nitrogen and stored at -80°C until analyses. For the analyses ($n = 3$), plant organs were ground and homogenised in liquid nitrogen using a mortar and pestle. Each biological replicate resulted from a pool of three plants that were ground together.

2.2 | Biometric traits

At the end of the experimental period, plant height, as well as the maximum root length, were recorded. Total leaf number was obtained by counting leaves larger than 1 cm and total leaf area was measured using a LI-3100C area meter (LI-COR, Lincoln, Nebraska, USA). The dry weight (48 h at 105°C in a ventilated oven) of leaves, stems and roots were determined. Shoot-root ratio (S/R, ratio of the shoot to root dry weight; g g^{-1} DW) was calculated.

2.3 | Total antioxidant capacity and non-enzymatic components

2.3.1 | Preparation of the extracts

For the analysis of the total antioxidant activity (ABTS [2,2-azinobis-(3-ethylbenzothiazoline-6-sulphonic acid)] free radical scavenging assay) and non-enzymatic components (total soluble polyphenols and flavonoids), 300 or 250 mg of frozen samples of roots and leaves, respectively, were extracted with 1.2 or 1.5 mL of 80% methanol, respectively, under cold conditions. Samples were centrifuged at 15000 \times g for 15 min and the resulting supernatant was used for all the analyses.

2.3.2 | ABTS free radical scavenging assay

ABTS assay was performed following the protocol described by Gonçalves et al. (2009), with slight modifications for application using a 96-well microplate. Briefly, ABTS $^{\cdot+}$ solution was generated through a chemical oxidation reaction with potassium persulphate, and its concentration adjusted with methanol to an initial absorbance of 0.700 (± 0.020) at 734 nm (Multiskan GO Microplate Spectrophotometer, Thermo Fisher Scientific Inc., Waltham, MA, USA). To 20 μL of the sample or standard solution, 180 μL of the adjusted ABTS $^{\cdot+}$ solution was added. The mixture was then incubated for 5 min at 30°C , the absorbance was measured at 734 nm and the results were obtained by calculating the reduction of percentage in sample absorbance (ASPL) with respect to the control (ACTL) using Equation 1. Trolox was used as a standard to prepare a calibration curve and the results were expressed as mmol Trolox equivalents/g FW. All reactions were performed in triplicate, and the radical stock solution was freshly prepared and filtered with a 0.45 μm syringe filter.

$$\text{ABTS scavenging (\%)} = (A_{\text{CTL}} - A_{\text{SPL}}) * \frac{100}{A_{\text{CTL}}} \quad (1)$$

2.3.3 | Total phenolics and flavonoids

Determination of total phenolic contents was performed using the Folin-Ciocalteu reagent following Ramos et al. (2019), with slight modifications. Briefly, 20 μL of extract or standard was mixed with 80 μL of Folin-Ciocalteu reagent previously diluted 1:10 and 100 μL of 7.5% (w/v) sodium carbonate was added to the mixture in a 96-well plate. After 60 min at room temperature and dark conditions, absorbance was measured at 750 nm. Results were expressed as mg gallic acid equivalents/g of fresh weight (mg GAE/g FW). Each reaction was performed in triplicate.

Total flavonoids were quantified through the aluminium chloride method adapted from Zafar et al. (2016). Briefly, an aliquot of each extract (40 μL) was mixed with 20 μL of 10% (w/v) aluminium chloride and 20 μL of 1 M potassium acetate solutions. Thereafter, distilled water was added to obtain a final volume of 200 μL in a 96-well plate. After 30 min of incubation in dark conditions and room temperature, the absorbance at 415 nm was measured by using a microplate reader. The calibration curve was drawn by using quercetin as standard and the flavonoid concentration was established in μg quercetin equivalents/g FW (μg QE/g FW).

2.4 | Enzymatic assay

2.4.1 | Enzyme extraction and activity quantification: SOD, CAT and APX

The extraction of the AOX enzymes was performed under cold conditions according to Fidalgo et al. (2011), using frozen samples of around 250 mg. After centrifugation, the supernatants were collected and used

for both protein and enzyme quantification following the protocols described by Soares et al. (2019). Briefly, protein quantification was performed as described by Bradford (1976), using bovine serum albumin as standard. Superoxide dismutase (SOD, EC.1.15.1.1) activity was quantified with a spectrophotometric assay based on the inhibition of photochemical reduction of nitroblue tetrazolium (NBT) (Donahue et al., 1997). Absorbance was read at 560 nm and results expressed as units of SOD/mg of protein, in which one unit of SOD corresponds to the amount of enzyme needed to inhibit the photochemical reduction of NBT by 50% (Spormann et al., 2023). Catalase (CAT, EC.1.11.1.6) activity was determined spectrophotometrically by measuring CAT-mediated H_2O_2 degradation ($\epsilon_{240\text{nm}} = 39.4 \text{ mM}^{-1} \text{ cm}^{-1}$) (Aebi, 1984). Absorbance measurements were recorded every 5 s for 60 s, and results were expressed as $\text{mmol H}_2\text{O}_2 \text{ min}^{-1}/\text{mg}$ of protein. Ascorbate peroxidase (APX, EC.1.11.1.11) activity was determined by monitoring the oxidation of ascorbic acid (AsA) for 1 min, with readings every 5 s ($\epsilon_{290\text{nm}} = 0.49 \text{ M}^{-1} \text{ cm}^{-1}$) (Nakano & Asada, 1981); results were expressed as $\text{mmol dehydroascorbate (DHA) min}^{-1}/\text{mg}$ of protein. The protocols for quantifying CAT and APX activity were adapted to UV-microplates following the adaptation from Soares, Pereira, et al. (2019).

2.5 | Statistical analysis

Results were expressed as mean \pm standard error of the mean (SEM). The effects of deficit level and Si supply on plant growth parameters and AOX system were analysed with a two-way ANOVA, using deficit level, silicon supply and their interaction as fixed factors. Mean separation was based on Fisher's protected least significant difference (LSD) test at $p = .05$ using IBM SPSS Statistics 26 and GraphPad Prism 8.

3 | RESULTS

3.1 | Plant growth

The deficit level significantly ($p < .001$) affected all plant growth parameters (Table S1; Figures 1 and 2), but Si supplementation significantly ($p < .001$) increased root length at all deficit levels (21% CTR, 18% N, 30%W, 37% N + W; Table S1; Figure 2b). Deficit level and Si supplementation showed a significant interaction for RDW ($p = .002$) and TDW ($p = .021$). When Si was added to plants grown under W deficit a significant reduction of 15% occurred in both RDW and TDW. On the other hand, plants grown under N + W deficit and supplemented with Si significantly increased their RDW by 29% and their TDW by 23%.

3.2 | Total antioxidant activity (ABTS assay), non-enzymatic components: Total phenolics and flavonoids concentration

Neither the deficit level, the Si supplementation, nor their interaction significantly affected the roots or leaves total AOX activity (ABTS

assay; $p > .05$; Table S2; Figure 3a,b). However, phenolics concentration in roots was significantly affected either by the Si supplementation ($p = .024$) or the interaction between the deficit level and the Si supply ($p = .003$; Table S2). When Si was added to CTR plants a significant increase of 83% was found (Figure 3c). A significant ($p = .030$) interaction between the deficit level and the Si supplementation was also found for the flavonoid's concentration in the leaves, which was decreased in plants grown in CTR conditions and supplemented with Si by 31% (Table S2 and Figure 3f).

3.3 | Enzymatic components: SOD, CAT and APX

All enzymes' activity, both in roots and leaves, was significantly ($p < .001$) affected by the deficit level. Si supplementation alone only significantly affected CAT roots activity ($p < .001$) and APX and SOD activity in leaves ($p = .003$ and $p < .001$, respectively; Table S2).

Deficit level and Si supplementation showed a significant interaction for APX, both in roots and leaves ($p < .001$ and $p = .003$, respectively; Table S2). While in CTR plants Si addition led to a 52% significantly lower activity, in plants grown under N or N + W deficit and supplemented with Si, APX activity was actually increased by 98% or by 48%, respectively (Figure 4a). An interaction between both factors also occurred for the APX activity in the leaves ($p = .003$; Table S2), being significantly higher (by 103%) in CTR plants grown with Si supplementation (Figure 4b). The activity of CAT in roots was also significantly increased by the interaction between deficit level and Si supplementation ($p < .0001$; Table S2). The N + W plants showed the highest increase (by 263%), followed by N (by 176%) and W (by 24%) deficits (Figure 4c). Concerning SOD activity, only in leaves an interaction between both factors was found ($p < .0001$; Table S2). An increase in CTR (by 18%) and in W deficit (by 14%) plants occurred in SOD activity when Si was supplied in the nutrient solution (Figure 4f).

4 | DISCUSSION

To the best of our knowledge, this is the first study that compares the benefits of Si application on the AOX response of young tomato plants subjected to individual N or W deficit or combined N + W deficit. By studying a set of plant growth traits and AOX biochemical parameters, we showed that individual and combined N + W deficit hampered plant growth, particularly when no Si was added to the nutrient solution (Table S1; Figures 1 and 2), and that tomato plants grown under N + W deficit were able to modulate their AOX network in the presence of Si (Table S1; Figure 4).

It has been reported that Si plays an important role in plant functioning under stressful conditions, yet under optimal conditions its role is often reported as being minimal or non-existent (reviewed by Coskun et al., 2019). Accordingly, concerning plant growth, in our study, the addition of Si did not cause significant effects in CTR plants, except for root length, which was increased (Table S1 and Figures 1 and 2). The positive effects of Si on root morphological traits, including root length, were already observed for several crops and were




FIGURE 1 Root dry weight (a), stem dry weight (b), leaf dry weight (c), total dry weight (d), shoot-to-root ratio (e) and leaf area (f) of tomato plants cv. 'Micro-Tom' grown for 27 days under four levels of deficit (control [CTR; 10.5 mM N + 100% W], nitrogen deficit [N; 5.3 mM N + 100%W], water deficit [W; 10.5 mM N + 50% W] and combined nitrogen and water deficit [N + W; 5.3 mM N + 50% W]) without (0 mM Na_2SiO_3) or with (2 mM Na_2SiO_3) silicon supplementation. Data presented are mean \pm SEM ($n = 6$). Bars represent Fisher's protected LSD test ($p = .05$) for the significant interaction deficit level \times silicon supplementation and for the independent effects when the interaction was not statistically significant. D, deficit level; LSD, least significant difference; N, nitrogen; Si, silicon supplementation; W, water.

recently reviewed by Tripathi et al. (2021). Oppositely, Si addition positively affected growth and biomass production in several plant species under environmental stress conditions (Bokor et al., 2021;

Coskun et al., 2019). However, the literature is very scarce on the effects of Si supplementation in Si-excluder plants, such as tomato, subjected to N deficiency. In our study, Si addition to N-stressed




FIGURE 2 Plant height (a) and root length (b) of tomato plants cv. 'Micro-Tom' grown for 27 days under four levels of deficit (control [CTR; 10.5 mM N + 100% W], nitrogen deficit [N; 5.3 mM N + 100%W], water deficit [W; 10.5 mM N + 50% W] and combined nitrogen and water deficit [N + W; 5.3 mM N + 50% W]) without (0 mM Na₂SiO₃) or with (2 mM Na₂SiO₃) silicon supplementation. Data presented are mean \pm SEM ($n = 6$). Bars represent Fisher's protected LSD test ($p = .05$) for the significant interaction deficit levels \times silicon supplementation and for the independent effects when the interaction was not statistically significant. D, deficit level; LSD, least significant difference.

plants did not significantly benefit any of the analysed growth traits (Table S1; Figures 1 and 2). The same was previously reported by Haddad et al. (2018), who also found that for the Si-excluder *Brassica napus* under N starvation, the Si treatment did not affect plant biomass. Furthermore, Leal et al. (2021) came to the same conclusion for cotton plants subjected to N deficiency and supplemented with Si. On the other hand, the positive effects of Si in mitigating the W deficit are widely reported (recently reviewed by Thorne et al., 2020; Bokor et al., 2021; Malik et al., 2021; Wang et al., 2021; Verma et al., 2022). However, in our work, Si addition to plants grown under single W deficit did not show noticeable effects, except for RDW and TDW where a significant reduction was found, and for root length which was increased (Table S1; Figures 1 and 2). This difference in the results could be due to different experimental design conditions, such as the form and concentration of Si given to the plants, the type of substrate used, or even using different tomato genotypes. For example, Zhang et al. (2018) and Shi et al. (2016) used potassium silicate instead of sodium silicate to conduct their experiment, while Shi et al. (2014) utilised sodium silicate but at a lower concentration of 0.5 mM. Furthermore, a study by Ali et al. (2018) demonstrated that tomato genotypes subjected to drought stress responded differently to the application of sodium silicate at a concentration of 1.5 mM, with the tolerant genotype having no beneficial effects on growth upon Si application. Finally, in most of the studies (Ali et al., 2018; Shi et al., 2014, 2016; Zhang et al., 2018), drought stress was imposed on tomato plants using polyethylene glycol and not through the decrease of watering per se which may also influence the results. Interestingly, although under individual deficit (N or W deficit) Si addition to the nutrient solution did not result in significant benefits, under combined

N + W deficit it led to an improvement in growth traits such as increased RDW, TDW and root length (Table S1; Figures 1 and 2). This, in turn, may be linked to the stimulation of the AOX system (Table S2; Figure 4) as already observed for other crops under abiotic stress (Bu et al., 2016; Geng et al., 2018; Kim et al., 2017; Shi et al., 2014).

It is known that under abiotic stress, plants have a complex AOX system for scavenging toxic ROS, which can be classified as non-enzymatic and enzymatic AOX systems, that work *in tandem* to counteract the overproduction of these by-products (Laxa et al., 2019; Soares, Carvalho, et al., 2019). Si application has been reported as a sustainable way of increasing the tolerance of plants under several abiotic stresses by boosting plant defence responses, such as those of the AOX system (reviewed by Kim et al., 2017). In our work, the total AOX capacity (ABTS) and the non-enzymatic system (phenolics and flavonoids) did not significantly change after Si addition, except for phenolics content in the roots, which was increased, and for the flavonoid content in the leaves, which was decreased, in CTR plants (Figure 3). Increasing evidence shows that Si influences the concentration of phenolic compounds (Hajiboland et al., 2017) which is supported by our findings in CTR plants. Such increase has also been reported for unstressed strawberry plants (Hajiboland et al., 2018).

However, the enzymatic system responded both to the imposed deficit as well as to the Si supplementation, demonstrating a higher importance of the enzymatic part of the AOX system in the deficits studied (Table S2; Figure 4). In CTR plants supplemented with Si, APX was decreased in roots but increased in leaves, while CAT did not significantly differ in any plant organ, and SOD only increased in leaves (Figure 4). This suggests that Si impacted the tomato AOX response




FIGURE 3 Antioxidant capacity measured by ABTS in roots (a) and leaves (b), total phenolics in roots (c) and leaves (d) and total flavonoids in roots (e) and leaves (f) of tomato plants cv. 'Micro-Tom' grown for 27 days under four levels of deficit (control [CTR; 10.5 mM N + 100% W], nitrogen deficit [N; 5.3 mM N + 100%W], water deficit [W; 10.5 mM N + 50% W] and combined nitrogen and water deficit [N + W; 5.3 mM N + 50% W]) without (0 mM Na₂SiO₃) or with (2 mM Na₂SiO₃) silicon supplementation. Data presented are mean ± SEM ($n = 3$). Bars represent Fisher's protected LSD test ($p = .05$) for the significant interaction deficit levels × silicon supplementation and for the independent effects when the interaction was not statistically significant. ABTS, 2,2-azinobis-(3-ethylbenzothiazoline-6-sulphonic acid); D, deficit level; LSD, least significant difference.

even in CTR conditions, as previously reported by Vega et al. (2019) after observing that Si application stimulated the AOX system of two healthy barley cultivars. Interestingly, although after Si addition, APX

and CAT were increased in the roots of plants subjected to all the imposed deficits, it was in the N + W deficit where the Si application showed a higher impact (Figure 4a,c). Shi et al. (2016) observed that Si




FIGURE 4 APX activity in roots (a) and leaves (b), CAT activity in roots (c) and leaves (d) and SOD activity in roots (e) and leaves (f) of tomato plants cv. 'Micro-Tom' grown for 27 days under four levels of deficit (control [CTR; 10.5 mM N + 100% W], nitrogen deficit [N; 5.3 mM N + 100%W], water deficit [W; 10.5 mM N + 50% W] and combined nitrogen and water deficit [N + W; 5.3 mM N + 50% W]) without (0 mM Na₂SiO₃) or with (2 mM Na₂SiO₃) silicon supplementation. Data presented are mean ± SEM ($n = 3$). Bars represent Fisher's protected LSD test ($p = .05$) for the significant interaction deficit levels × silicon supplementation and for the independent effects when the interaction was not statistically significant. APX, ascorbate peroxidase; CAT, catalase; D, deficit level; LSD, least significant difference; SOD, superoxide dismutase.

improved the growth and development of tomato plants under drought stress by improving root hydraulic conductance due to an increase in the AOX system response. In this study, the general improvement of the AOX system on the roots of combined stress

tomato plants may therefore be the reason why this group of plants responded better to the imposed stress. On the other hand, Si addition increased the activity of SOD in the leaves of plants subjected to W deficit, but not in the other stress treatments, where no significant

changes were observed (Figure 4c,f). Although Si was already shown to increase the activities of SOD, CAT and APX in crops such as wheat, tomato, chickpea, rapeseed, sugarcane and sunflower under drought conditions (Gunes et al., 2007, 2008; Shi et al., 2016; Xu et al., 2017; Verma et al., 2020), we could not find literature reports about the effect of Si on the AOX system of plants subjected to individual N or combined N + W deficit. Our results suggest that Si differentially affects the response of the enzymatic system in plants grown under individual stress or in a combination of stresses, the latter occurring most often. While for individual W deficit, Si seems to have stimulated the enzymatic machinery partly in roots and partly in the leaves, for combined N + W deficit, and to a certain extent for N deficit, it seems to have acted mostly at the root level, where a high increase was seen in the activity of APX and CAT enzymes. This greater impact of Si on the roots of plants grown under combined N + W deficit may be due to the fact that both the N deficit and the W deficit are primarily perceived by roots (Bengough et al., 2011; Sun et al., 2020). On the other hand, since Si was also applied via the root, it may have led to a greater activation here, as a first line of defence, which could only later be extended to the leaf. Indeed, the significant role of Si on root development during stress has been previously noted (Hattori et al., 2005) and was also observed in our study (Figure 2b). The root growth stimulation by Si application may be the result of root elongation by increasing the extensibility of cell walls (Hattori et al., 2003).

5 | CONCLUSIONS

Root Si application on tomato seedlings can partly alleviate the stress caused by combined N and water deficit as demonstrated by the increase in growth traits such as RDW, and root length, which ultimately contributed to their higher TDW. The better performance of plants grown under combined N + W deficit and supplemented with Si seems to be related, at least in part, to the AOX system of the roots (where the stresses were first experienced), as shown by the increased activity of CAT and APX enzyme. Plants' response to combined stresses cannot be extrapolated from the responses observed for the individual deficits, because for plants subjected to either N or W deficit the impact of Si was not so evident. These results are highly relevant if we consider that: (i) plants often do not face only one individual abiotic stress, but rather a combination of them and (ii) combined stress was the treatment where the highest decreases in growth traits were found, and therefore the treatment where the application of Si would bring more benefits. Comprehension of the full mechanisms involved in the complex interactions between these two abiotic stresses and Si application is still in its infancy. It is also not clear yet to what extent the present findings can be extrapolated to other crops and stress levels.

ACKNOWLEDGEMENTS

The authors would like to thank Fundação para a Ciência e a Tecnologia (FCT) for Joana Machado and Ana Patricia G. Fernandes PhD scholarship

(SFRH/BD/116147/2016 and SFRH/BD/05748/2020, respectively). This research was also supported by national funds through FCT within the scope of UIDB/05748/2020, UIDP/05748/2020, UIDB/50016/2020, Norte2020 (NORTE-01-0145-FEDER-000041) and Acordo de Cooperação Científica FCT/Eslováquia–2019/2020, DRI 5160 (SK-PT-18-0020). The work is also a part of cooperation supported by European Cooperation in Science and Technology, grant Nr. COST CA19116.

ORCID

Joana Machado  <https://orcid.org/0000-0002-8222-7211>

Marta W. Vasconcelos  <https://orcid.org/0000-0002-5110-7006>

REFERENCES

- Aebi, H. (1984). Catalase in vitro. *Methods in Enzymology*, 105, 121–126.
- Ali, N., Réthoré, E., Yvin, J. C., & Hosseini, S. A. (2020). The regulatory role of silicon in mitigating plant nutritional stresses. *Plants (Basel)*, 9(12), 1779. <https://doi.org/10.3390/plants9121779>
- Ali, N., Schwarzenberg, A., Yvin, J.-C., & Hosseini, S. A. (2018). Regulatory role of silicon in mediating differential stress tolerance responses in two contrasting tomato genotypes under osmotic stress. *Frontiers in Plant Science*, 9, 1475. <https://doi.org/10.3389/fpls.2018.01475>
- Bengough, A. G., McKenzie, B. M., Hallett, P. D., & Valentine, T. A. (2011). Root elongation, water stress, and mechanical impedance: A review of limiting stresses and beneficial root tip traits. *Journal of Experimental Botany*, 62(1), 59–68. <https://doi.org/10.1093/jxb/erq350>
- Bokor, B., Santos, C. S., Kostoláni, D., Machado, J., da Silva, M. N., Carvalho, S. M. P., Vaculík, M., & Vasconcelos, M. W. (2021). Mitigation of climate change and environmental hazards in plants: Potential role of the beneficial metalloid silicon. *Journal of Hazardous Materials*, 416, 126193. <https://doi.org/10.1016/j.jhazmat.2021.126193>
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248–254.
- Bu, R., Xie, J., Yu, J., Liao, W., Xiao, X., Lv, J., Wang, C., Ye, J., & Calderón-Urrea, A. (2016). Autotoxicity in cucumber (*Cucumis sativus* L.) seedlings is alleviated by silicon through an increase in the activity of antioxidant enzymes and by mitigating lipid peroxidation. *Journal of Plant Biology*, 59, 247–259. <https://doi.org/10.1007/s12374-016-0526-1>
- Cicco, A. D. (2019). The fruit and vegetable sector in the EU—A statistical overview Eurostat 2019. https://ec.europa.eu/eurostat/statistics-explained/index.php/The_fruit_and_vegetable_sector_in_the_EU_-_a_statistical_overview
- Coskun, D., Deshmukh, R., Sonah, H., Menzies, J. G., Reynolds, O., Ma, J. F., Kronzucker, H. J., & Bélanger, R. R. (2019). The controversies of silicon's role in plant biology. *New Phytologist*, 221, 67–85. <https://doi.org/10.1111/nph.15343>
- Donahue, J. L., Okpodu, C. M., Cramer, C. L., Grabau, E. A., & Alscher, R. G. (1997). Responses of antioxidants to paraquat in pea leaves (relationships to resistance). *Plant Physiology*, 113, 249–257. <https://doi.org/10.1104/pp.113.1.249>
- Du, Y. D., Niu, W. Q., Gu, X. B., Zhang, Q., & Cui, B. J. (2018). Water- and nitrogen-saving potentials in tomato production: A meta-analysis. *Agricultural Water Management*, 210, 296–303. <https://doi.org/10.1016/j.agwat.2018.08.035>
- Fernandes, A. P. G., Machado, J., Fernandes, T. R., Vasconcelos, M. W., & Carvalho, S. M. P. (2022). Water and nitrogen fertilization management in the light of climate change: Impacts on food security and product quality. In V. Kumar, A. K. Srivastava, & P. Suprasanna (Eds.), *Plant nutrition and food security in the era of climate change* (pp. 147–178). Elsevier. <https://doi.org/10.1016/B978-0-12-822916-3.00013-5>
- Fidalgo, F., Freitas, R., Ferreira, R., Pessoa, A. M., & Teixeira, J. (2011). *Solanum nigrum* L. antioxidant defence system isozymes are regulated

- transcriptionally and post-translationally in Cd-induced stress. *Environmental and Experimental Botany*, 72, 312–319. <https://doi.org/10.1016/j.envexpbot.2011.04.007>
- García-Martí, M., Piñero, M. C., García-Sánchez, F., Mestre, T. C., López-Delacalle, M., Martínez, V., & Rivero, R. M. (2019). Amelioration of the oxidative stress generated by simple or combined abiotic stress through the K and Ca²⁺ supplementation in tomato plants. *Antioxidants (Basel)*, 8(4), 81. <https://doi.org/10.3390/antiox8040081>
- Geng, A., Wang, X., Wu, L., Wang, F., Wu, Z., Yang, H., Chen, Y., Wen, D., & Liu, X. (2018). Silicon improves growth and alleviates oxidative stress in rice seedlings (*Oryza sativa* L.) by strengthening antioxidant defense and enhancing protein metabolism under arsenic acid exposure. *Ecotoxicology and Environmental Safety*, 158, 266–273. <https://doi.org/10.1016/j.ecoenv.2018.03.050>
- Gojon, A. (2017). Nitrogen nutrition in plants: Rapid progress and new challenges. *Journal of Experimental Botany*, 68(10), 2457–2462. <https://doi.org/10.1093/jxb/erx171>
- Gonçalves, B., Falco, V., Moutinho-Pereira, J., Bacelar, E., Peixoto, F., & Correia, C. (2009). Effects of elevated CO₂ on grapevine (*Vitis vinifera* L.): Volatile composition, phenolic content, and in vitro antioxidant activity of red wine. *Journal of Agricultural and Food Chemistry*, 57(1), 265–273. <https://doi.org/10.1021/jf8020199>
- Gong, H., Zhu, X., Chen, K., Wang, S., & Zhang, C. L. (2005). Silicon alleviates oxidative damage of wheat plants in pots under drought. *Plant Science*, 169(2), 313–321. <https://doi.org/10.1016/j.plantsci.2005.02.023>
- Gunes, A., Pilbeam, D. J., Inal, A., Bagci, E. G., & Coban, S. (2007). Influence of silicon on antioxidant mechanisms and lipid peroxidation in chickpea (*Cicer arietinum* L.) cultivars under drought stress. *Journal of Plant Interactions*, 2(2), 105–113.
- Gunes, A., Pilbeam, D. J., Inal, A., & Coban, S. (2008). Influence of silicon on sunflower cultivars under drought stress, I: Growth, antioxidant mechanisms, and lipid peroxidation. *Communications in Soil Science and Plant Analysis*, 39(13–14), 1885–1903.
- Haddad, C., Arkoun, M., Jamois, F., Schwarzenberg, A., Yvin, J.-C., Etienne, P., & Lainé, P. (2018). Silicon promotes growth of *Brassica napus* L. and delays leaf senescence induced by nitrogen starvation. *Frontiers in Plant Science*, 9, 516. <https://doi.org/10.3389/fpls.2018.00516>
- Hajiboland, R., Bahrami-Rad, S., & Poschenrieder, C. (2017). Silicon modifies both a local response and a systemic response to mechanical stress in tobacco leaves. *Biologia Plantarum*, 61, 187–191. <https://doi.org/10.1007/s10535-016-0633-3>
- Hajiboland, R., Moradtalab, N., Eshaghi, Z., & Feizy, J. (2018). Effect of silicon supplementation on growth and metabolism of strawberry plants at three developmental stages. *New Zealand Journal of Crop and Horticultural Science*, 46(2), 144–161. <https://doi.org/10.1080/01140671.2017.1373680>
- Hattori, T., Inanaga, S., Arak, I. H., An, P., Morita, S., Luxová, M., & Lux, A. (2005). Application of silicon enhanced drought tolerance in *Sorghum bicolor*. *Physiologia Plantarum*, 123, 459–466. <https://doi.org/10.1111/j.1399-3054.2005.00481.x>
- Hattori, T., Inanaga, S., Tanimoto, E., Lux, A., Luxová, M., & Sugimoto, Y. (2003). Silicon-induced changes in viscoelastic properties of sorghum root cell walls. *Plant and Cell Physiology*, 44, 743–749. <https://doi.org/10.1093/pcp/pcg090>
- Heuvelink, E., Okello, R. C. O., Peet, M., Giovannoni, J. J., & Dorais, M. (2020). Chapter 7: Tomato. In H. C. Wien & H. Stützel (Eds.), *Physiology of vegetable crops* (2nd ed., pp. 138–178). CAB International.
- Hoffmann, J., Berni, R., Hausman, J. F., & Guerriero, G. (2020). A review on the beneficial role of silicon against salinity in non-accumulator crops: Tomato as a model. *Biomolecules*, 10(9), 1284. <https://doi.org/10.3390/biom10091284>
- Hussain, H. A., Hussain, S., Khaliq, A., Ashraf, U., Anjum, S. A., Men, S., & Wang, L. (2018). Chilling and drought stresses in crop plants: Implications, cross talk, and potential management opportunities. *Frontiers in Plant Science*, 9, 393. <https://doi.org/10.3389/fpls.2018.00393>
- Jhariya, M. K., Banerjee, A., Meena, R. S., & Yadav, D. K. (2019). Agriculture, forestry and environmental sustainability: A way forward. In M. Jhariya, A. Banerjee, R. Meena, & D. Yadav (Eds.), *Sustainable agriculture, forest and environmental management*. Springer. https://doi.org/10.1007/978-981-13-6830-1_1
- Joshi, S., Thoday-Kennedy, E., Daetwyler, H. D., Hayden, M., Spangenberg, G., & Kant, S. (2021). High-throughput phenotyping to dissect genotypic differences in safflower for drought tolerance. *PLoS One*, 16(7), e0254908. <https://doi.org/10.1371/journal.pone.0254908>
- Kim, Y.-H., Khan, A. L., Waqas, M., & Lee, I.-J. (2017). Silicon regulates antioxidant activities of crop plants under abiotic-induced oxidative stress: A review. *Frontiers in Plant Science*, 8, 510. <https://doi.org/10.3389/fpls.2017.00510>
- Laxa, M., Liebthal, M., Telman, W., Chibani, K., & Dietz, K.-J. (2019). The role of the plant antioxidant system in drought tolerance. *Antioxidants*, 8(4), 94. <https://doi.org/10.3390/antiox8040094>
- Leal, A. J. F., Barreto, R. F., Mello Prado, R., Silva Junior, G. B., Barreto, C. F., dos Santos, L. C. N., & Campos, C. N. S. (2021). Silicon attenuates abiotic stress caused by ammonium toxicity but not nitrogen deficiency in cotton plants. *Journal of Agronomy and Crop Science*, 207(4), 774–781. <https://doi.org/10.1111/jac.12499>
- Liang, Y. C., Sun, W. C., Zhu, Y. G., & Christie, P. (2007). Mechanisms of silicon-mediated alleviation of abiotic stresses in higher plants: A review. *Environmental Pollution*, 147, 422–428. <https://doi.org/10.1016/j.envpol.2006.06.008>
- Machado, J., Fernandes, A. P. G., Fernandes, T. R., Heuvelink, E., Vasconcelos, M. W., & Carvalho, S. M. P. (2022). Drought and nitrogen stress effects and tolerance mechanisms in tomato: A review. In V. Kumar, A. K. Srivastava, & P. Suprasanna (Eds.), *Plant nutrition and food security in the era of climate change* (pp. 315–359). Elsevier. <https://doi.org/10.1016/B978-0-12-822916-3.00014-7>
- Machado, J., Vasconcelos, M. W., Soares, C., Fidalgo, F., Heuvelink, E., & Carvalho, S. M. P. (2023b). Young tomato plants respond differently under single or combined mild nitrogen and water deficit: an insight into morphophysiological responses and primary metabolism. *Plants*, 12, 1181.
- Machado, J., Vasconcelos, M. W., Soares, C., Fidalgo, F., Heuvelink, E., & Carvalho, S. M. P. (2023c). Enzymatic non-enzymatic antioxidant responses of young tomato plants (cv. Micro-Tom) to single and combined mild nitrogen and water deficit: not the sum of the parts. *Antioxidants*, 12, 375.
- Machado, J., Heuvelink, E., Vasconcelos, M. W., Cunha, L. M., Finkers, R., & Carvalho, S. M. P. (2023a). Exploring tomato phenotypic variability under combined nitrogen and water deficit. *Plant and Soil*, 1–16.
- Malik, M. A., Wani, A. H., Mir, S. H., Rehman, I. U., Tahir, I., Ahmad, P., & Rashid, I. (2021). Elucidating the role of silicon in drought stress tolerance in plants. *Plant Physiology and Biochemistry*, 165, 187–195. <https://doi.org/10.1016/j.plaphy.2021.04.021>
- Minden, V., Schaller, J., & Olde Venterink, H. (2020). Plants increase silicon content as a response to nitrogen or phosphorus limitation: A case study with *Holcus lanatus*. *Plant and Soil*, 462, 95–108. <https://doi.org/10.1007/s11104-020-04667-1>
- Nakano, Y., & Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant & Cell Physiology*, 22, 867–880. <https://doi.org/10.1093/oxfordjournals.pcp.a076232>
- Nunes da Silva, M., Machado, J., Osorio, J., Duarte, R., & Santos, C. S. (2022). Non-essential elements and their role in sustainable agriculture. *Agronomy*, 12, 888. <https://doi.org/10.3390/agronomy12040888>
- Pandey, P., Ramegowda, V., & Senthil-Kumar, M. (2015). Shared and unique responses of plants to multiple individual stresses and stress combinations: Physiological and molecular mechanisms. *Frontiers in Plant Science*, 6, 723. <https://doi.org/10.3389/fpls.2015.00723>

- Parveen, A., Liu, W., Hussain, S., Asghar, J., Perveen, S., & Xiong, Y. (2019). Silicon priming regulates morpho-physiological growth and oxidative metabolism in maize under drought stress. *Plants (Basel)*, 8(10), 431. <https://doi.org/10.3390/plants8100431>
- Pavlovic, J., Kostic, L., Bosnic, P., Kirkby, E. A., & Nikolic, M. (2021). Interactions of silicon with essential and beneficial elements in plants. *Frontiers in Plant Science*, 12, 697592. <https://doi.org/10.3389/fpls.2021.697592>
- Ramos, P. A. B., Moreirinha, C., Silva, S., Costa, E. M., Veiga, M., Coscueta, E., Santos, S. A. O., Almeida, A., Pintado, M. M., Freire, C. S. R., Silva, A. M. S., & Silvestre, A. J. D. (2019). The health-promoting potential of *Salix* spp. bark polar extracts: Key insights on phenolic composition and in vitro bioactivity and biocompatibility. *Antioxidants*, 8, 609. <https://doi.org/10.3390/antiox8120609>
- Ruggiero, A., Punzo, P., Van Oosten, M. J., Cirillo, V., Esposito, S., Costa, A., Maggio, A., Grillo, S. & Batelli, G. (2022). Transcriptomic and splicing changes underlying tomato responses to combined water and nutrient stress. *Frontiers in Plant Science*, 13, 974048. <https://doi.org/10.3389/fpls.2022.974048>
- Sandhu, R. K., Boyd, N. S., Zotarelli, L., Agehara, S., & Peres, N. (2021). Effect of planting density on the yield and growth of intercropped tomatoes and peppers in Florida. *HortScience*, 56, 286–290. <https://doi.org/10.21273/HORTSCI15567-20>
- Sattar, A., Sher, A., Ijaz, M., Ul-Allah, S., Butt, M., Irfan, M., Rizwan, S. M., Ali, H., & Cheema, M. A. (2020). Interactive effect of biochar and silicon on improving morpho-physiological and biochemical attributes of maize by reducing drought hazards. *Journal of Soil Science and Plant Nutrition*, 20, 1819–1826. <https://doi.org/10.1007/s42729-020-00253-7>
- Shi, Y., Zhang, Y., Han, W., Feng, R., Hu, Y., Guo, J., & Gong, H. (2016). Silicon enhances water stress tolerance by improving root hydraulic conductance in *Solanum lycopersicum* L. *Frontiers in Plant Science*, 7, 196. <https://doi.org/10.3389/fpls.2016.00196>
- Shi, Y., Zhang, Y., Yao, H., Wu, J., Sun, H., & Gong, H. (2014). Silicon improves seed germination and alleviates oxidative stress of bud seedlings in tomato under water deficit stress. *Plant Physiology and Biochemistry*, 78, 27–36. <https://doi.org/10.1016/j.plaphy.2014.02.009>
- Soares, C., Carvalho, M. E. A., Azevedo, R. A., & Fidalgo, F. (2019). Plants facing oxidative challenges—A little help from the antioxidant networks. *Environmental and Experimental Botany*, 161, 4–25. <https://doi.org/10.1016/j.envexpbot.2018.12.009>
- Soares, C., Pereira, R., Spormann, S., & Fidalgo, F. (2019). Is soil contamination by a glyphosate commercial formulation truly harmless to non-target plants?—Evaluation of oxidative damage and antioxidant responses in tomato. *Environmental Pollution*, 247, 256–265. <https://doi.org/10.1016/j.envpol.2019.01.063>
- Spormann, S., Soares, C., Martins, V., Azenha, M., Gerós, H., & Fidalgo, F. Early activation of antioxidant responses in ni-stressed tomato cultivars determines their resilience under co-exposure to drought. *Journal of Plant Growth Regulation*, 42, 877–891 (2023).
- Sun, X., Chen, F., Yuan, L., & Mi, G. (2020). The physiological mechanism underlying root elongation in response to nitrogen deficiency in crop plants. *Planta*, 251, 84. <https://doi.org/10.1007/s00425-020-03376-4>
- Thorne, S. J., Hartley, S. E., & Maathuis, F. J. M. (2020). Is silicon a panacea for alleviating drought and salt stress in crops? *Frontiers in Plant Science*, 11, 1221. <https://doi.org/10.3389/fpls.2020.01221>
- Tripathi, P., Subedi, S., Khan, A. L., Chung, Y.-S., & Kim, Y. (2021). Silicon effects on the root system of diverse crop species using root phenotyping technology. *Plants*, 10, 885. <https://doi.org/10.3390/plants10050885>
- Vega, I., Nikolic, M., Pontigo, S., Godoy, K., Mora, M. D. L. L., & Cartes, P. (2019). Silicon improves the production of high antioxidant or structural phenolic compounds in barley cultivars under aluminum stress. *Agronomy*, 9, 388. <https://doi.org/10.3390/agronomy9070388>
- Verma, K. K., Liu, X.-H., Wu, K.-C., Singh, R. K., Song, Q.-Q., Malviya, M. K., Song, X.-P., Singh, P., Verma, C. L., & Li, Y.-R. (2020). The impact of silicon on photosynthetic and biochemical responses of sugarcane under different soil moisture levels. *Silicon*, 12(6), 1355–1367.
- Verma, K. K., Song, X. P., Lin, B., Guo, D. J., Singh, M., Rajput, V. D., Singh, R. K., Singh, P., Sharma, A., Malviya, M. K., Chen, G. L., & Li, Y. R. (2022). Silicon induced drought tolerance in crop plants: Physiological adaptation strategies. *Silicon*, 14(6), 2473–2487. <https://doi.org/10.1007/s12633-021-01071-x>
- Wang, M., Wang, R., Mur, L. A. J., Ruan, J., Shen, Q., & Guo, S. (2021). Functions of silicon in plant drought stress responses. *Horticulture Research*, 8, 254. <https://doi.org/10.1038/s41438-021-00681-1>
- Xu, L., Islam, F., Ali, B., Pei, Z., Li, J., Ghani, M. A., & Zhou, W. (2017). Silicon and water-deficit stress differentially modulate physiology and ultrastructure in wheat (*Triticum aestivum* L.). *3 Biotech*, 7(4), 273. <https://doi.org/10.1007/s13205-017-0904-5>
- Zafar, H., Ali, A., Ali, J. S., Haq, I. U., & Zia, M. (2016). Effect of ZnO nanoparticles on *Brassica nigra* seedlings and stem explants: Growth dynamics and antioxidative response. *Frontiers in Plant Science*, 7, 535. <https://doi.org/10.3389/fpls.2016.00535>
- Zhang, Y., Shi, Y., Gong, H., Zhao, H., Li, H., Hu, Y., & Wang, Y. (2018). Beneficial effects of silicon on photosynthesis of tomato seedlings under water stress. *Journal of Integrative Agriculture*, 17(10), 2151–2159. [https://doi.org/10.1016/s2095-3119\(18\)62038-6](https://doi.org/10.1016/s2095-3119(18)62038-6)

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Machado, J., Fernandes, A. P. G., Bokor, B., Vaculík, M., Heuvelink, E., Carvalho, S. M. P., & Vasconcelos, M. W. (2023). The effect of silicon on the antioxidant system of tomato seedlings exposed to individual and combined nitrogen and water deficit. *Annals of Applied Biology*, 1–11. <https://doi.org/10.1111/aab.12849>