

Wind dispersed tree species have greater maximum height

Global Ecology and Biogeography

Slik, Ferry; Pinho, Bruno X.; Griffith, Daniel M.; Webb, Edward; Raghubanshi, Akhilesh Singh et al

<https://doi.org/10.1111/geb.13878>

This publication is made publicly available in the institutional repository of Wageningen University and Research, under the terms of article 25fa of the Dutch Copyright Act, also known as the Amendment Taverne.

Article 25fa states that the author of a short scientific work funded either wholly or partially by Dutch public funds is entitled to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed using the principles as determined in the Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' project. According to these principles research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and / or copyright owner(s) of this work. Any use of the publication or parts of it other than authorised under article 25fa of the Dutch Copyright act is prohibited. Wageningen University & Research and the author(s) of this publication shall not be held responsible or liable for any damages resulting from your (re)use of this publication.

For questions regarding the public availability of this publication please contact openaccess.library@wur.nl

Wind dispersed tree species have greater maximum height

Ferry Slik¹  | Bruno X. Pinho^{2,3}  | Daniel M. Griffith⁴ | Edward Webb^{5,6} | Akhilesh Singh Raghubanshi⁷ | Adriano C. Quaresma^{8,9} | Aida Cuni Sanchez^{10,11} | Aisha Sultana¹² | Alexandre F. Souza¹³ | Andreas Ensslin¹⁴ | Andreas Hemp¹⁵  | Andrew Lowe¹⁶ | Andrew R. Marshall^{17,18,19} | Kamalakumari Anitha²⁰ | Anne Mette Lykke²¹ | Armadyanto²² | Asyraf Mansor^{23,24} | Atsri K. Honam²⁵ | Axel D. Poulsen²⁶ | Ben Sparrow²⁷ | Benjamin J. W. Buckley²⁸ | Bernat Ripoll Capilla²⁹ | Bianca Weiss Albuquerque⁹ | Christine B. Schmitt³⁰ | Dharmalingam Mohandass³¹ | Diogo S. B. Rocha³² | Douglas Sheil^{33,34} | Eduardo A. Pérez-García³⁵ | Eduardo Catharino³⁶ | Eduardo van den Berg³⁷ | Ervan Rutishauser³⁸ | Fabian Brambach³⁹  | Felipe Zamborlini Saiter^{40,41} | Feyera Senbeta⁴² | Florian Wittmann⁴³ | Francesco Rovero⁴⁴ | Francisco Mora Ardila⁴⁵ | Frans Bongers³³ | Gabriella M. Fredriksson⁴⁶ | Gemma Rutten² | Gerard Imani⁴⁷ | Gerardo A. Aymard Corredor^{48,49} | Giselda Durigan⁵⁰ | Gopal Shukla⁵¹ | Guadalupe Williams-Linera⁵² | Heike Culmsee⁵³ | Hendrik Segah⁵⁴ | Iñigo Granzow-de la Cerda⁵⁵ | Jamuna S. Singh⁵⁶ | James Grogan⁵⁷ | Jan Reitsma⁵⁸ | Jean-François Bastin⁵⁹ | Jeremy Lindsell^{60,61} | Jerome Millet⁶² | Joao Roberto dos Santos⁶³ | Jochen Schoengart⁹ | John H. Vandermeer⁶⁴ | John Herbohn⁶⁵ | Jon Lovett^{66,67} | Jorge A. Meave³⁵ | José Roberto Rodrigues Pinto⁶⁸ | Juan Carlos Montero⁶⁹ | Kalle Ruokolainen^{70,71} | Khairil Bin Mahmud^{72,73} | Layon O. Demarchi⁹ | Lourens Poorter⁷⁴  | Luis Bernacci⁷⁵ | Manichanh Satdichanh⁷⁶ | Marcio Seiji Suganuma⁷⁷ | Maria T. F. Piedade⁹ | Mariarty A. Niun⁷⁸ | Mark E. Harrison^{79,80} | Mark Schulze⁸¹ | Markus Fischer² | Michael Kessler⁸² | Miguel Castillo⁸³ | Mohammad Shah Hussain¹² | Moses B. Libalah^{84,85} | Muhammad Ali Imron⁸⁶ | Narayanaswamy Parthasarathy⁸⁷ | Naret Seuaturien⁸⁸ | Natalia Targhetta⁹ | Ni Putu Diana Mahayani⁸⁶ | Nigel C. A. Pitman⁸⁹ | Orlando Rangel⁹⁰ | Pantaleo Munishi⁹¹ | Patricia Balvanera⁴⁵ | Peter Ashton⁹² | Pia Parolin⁹³ | Polyanna da Conceição Bispo⁹⁴ | Priya Davidar⁴⁵ | Rahayu Sukri⁹⁶ | Rahmad Zakaria^{23,97} | Rama Chandra Prasad⁹⁸ | Ravi K. Chaturvedi⁹⁹ | Robert Steinmetz¹⁰⁰  | Rodrigo Muñoz¹⁰¹ | Rozainah Mohamad Zakaria¹⁰² | Saara J. DeWalt¹⁰³ | Hoang Van Sam¹⁰⁴ | Samir Rolim¹⁰⁵ | Sharif Ahmed Mukul^{106,107} | Siti Maimunah¹⁰⁸ | Swapan Kumar Sarker¹⁰⁹ | Terry Sunderland^{110,111} | Thomas Gillespie¹¹² | Tinde van Andel¹¹³ | Tran Van Do¹¹⁴  | Wanlop Chutipong¹¹⁵ |

For affiliations refer to page 8.

Runguo Zang¹¹⁶ | Xiaobo Yang¹¹⁷ | Xinghui Lu¹¹⁸ | Yves Laumonier¹¹⁹ |
Zhila Hemati¹²⁰

Correspondence

Ferry Slik, Environmental and Life Sciences, Faculty of Science, Universiti Brunei Darussalam, Jalan Tungku Link, Gadong BE1410, Brunei Darussalam. Email: ferryslik@hotmail.com; johan.slik@ubd.edu.bn

Funding information

Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq, Grant/Award Number: 520053/1998-2; ARC Future Fellowship, Grant/Award Number: FT170100279; US Fish and Wildlife Service Great Apes Conservation Fund; Leverhulme Trust, Valuing the Arc project; The Orangutan Project; Arcus Foundation

Handling Editor: Arndt Hampe

Abstract

Aim: We test the hypothesis that wind dispersal is more common among emergent tree species given that being tall increases the likelihood of effective seed dispersal.

Location: Americas, Africa and the Asia-Pacific.

Time period: 1970–2020.

Major taxa studied: Gymnosperms and Angiosperms.

Methods: We used a dataset consisting of tree inventories from 2821 plots across three biogeographic regions (Americas, Africa and Asia-Pacific), including dry and wet forests, to determine the maximum height and dispersal strategy of 5314 tree species. A web search was used to determine whether species were wind-dispersed. We compared differences in tree species maximum height between biogeographic regions and examined the relationship between species maximum height and wind dispersal using logistic regression. We also tested whether emergent tree species, that is species with at least one individual taller than the 95% height percentile in one or more plots, were disproportionately wind-dispersed in dry and wet forests within each biogeographic region.

Results: Our dataset provides maximum height values for 5314 tree species, of which more than half (2914) had no record of this trait in existing global databases. We found that, on average, tree species in the Americas have lower maximum heights compared to those in Africa and the Asia Pacific. The probability of wind dispersal increased significantly with tree species maximum height and was significantly higher among emergent than non-emergent tree species in both dry and wet forests in all three biogeographic regions.

Main conclusion: Wind dispersal is more prevalent in tall, emergent tree species than in non-emergent species and may thus be an important factor in the evolution of tree species maximum height. By providing the most comprehensive dataset so far of tree species maximum height and wind dispersal strategies, this study paves the way for advancing our understanding of the eco-evolutionary drivers of tree size.

KEYWORDS

emergent trees, evolution, functional traits, seed dispersal, tree species maximum height, wind dispersal

1 | INTRODUCTION

Tree species maximum height is closely linked to competition for light, a fundamental resource for tree growth, survival and reproduction. Yet, if light exerted the predominant selection pressure on trees in closed-canopy forests, one would expect to find little variation in interspecific maximum heights as smaller species would be shaded out and eventually disappear from the community (D'Andrea et al., 2019, 2020; Scheffer & van Nes, 2006). On the other hand, species emerging above the canopy are subject to other selective pressures like exposure

to wind, lightning and elevated levels of evaporation, which also limit their abundance (Jackson et al., 2021; Larjavaara, 2014; Margrove et al., 2015; Olson et al., 2018). However, maximum heights of co-existing tree species vary considerably, reflecting a range of life history strategies (Kohyama, 1993; Kohyama et al., 2003; Rieger et al., 2020; Souza, 2021). Some species complete their entire life cycle in the forest understorey, while others tower above their neighbours as emergents (Banin et al., 2012; Iida et al., 2014). Selective pressures in addition to light and to the risks associated with emergence must therefore account for the variation found in tree maximum heights.

In this study, we explore the hypothesis that emergent tree species—those capable of growing taller than the surrounding canopy—are more likely to be wind-dispersed than non-emergent species. For dispersal to be effective, seeds must generally be carried away from the parent tree to avoid negative distance- and density-dependence, and to colonize favourable sites for regeneration (Comita et al., 2014). Thus, it should be advantageous for wind-dispersed species to be tall because wind speeds increase logarithmically with height above the canopy (de Santana et al., 2017; Smith et al., 2015, 2016). Release of wind-dispersed fruits and seeds at elevated heights increases the probability of long-distance dispersal, which can facilitate successful establishment (Heydel et al., 2014; Nathan et al., 2002; Wu et al., 2023). Therefore, wind-dispersed tree species likely have an evolutionary advantage by being emergent, as this condition may offset the increased mortality risks that result from their size (Jackson et al., 2021; Larjavaara, 2014; Margrove et al., 2015; Olson et al., 2018). However, the relationship between emergence and wind dispersal has not been examined for a large group of species at a global scale.

Here we test this relationship using a dataset of tree surveys with height measurements that covers most of the (sub-)tropics and some temperate forests in the Americas, Africa and Asia-Pacific, thus encompassing a wide variety of tree species in three more or less independent biogeographic regions (Cazzolla Gatti et al., 2022; Slik et al., 2015). We also provide an appendix containing maximum heights for more than 5000 tree species to promote further study on the ecology and evolution of tree species maximum height.

2 | MATERIALS AND METHODS

2.1 | Tree height data

We used tree inventory data (diameter at breast height ≥ 10 cm, excluding monocots and Cactaceae) from 2821 individual old-growth

forest plots varying in size from 0.003 to 18 ha, which covered 246 one-degree latitude/longitude grid cells across the Americas, Africa and Asia Pacific (Figure 1; Appendix 1). Species' taxonomy was updated using the Taxonomic Name Resolution Service (<https://tnrs.biendata.org/>). We used the Plants of the World Online database (<https://powo.science.kew.org/>) to resolve any remaining taxonomic issues.

The dataset contained heights for a total of 7913 tree species. To ensure an adequate sample size for each species, we only considered those with a minimum of 10 reported height values, which reduced the number of species for analysis to 5314 (Appendix 2). To reduce measurement errors and outliers, we used the upper 10-percentile height value as a proxy for the maximum height of each species (Appendix 2). We compared these values with tree height data available in the two largest global plant trait databases: TRY (Kattge et al., 2020) and BIEN (Maitner et al., 2018). Together these databases contained 2400 of the species included in our dataset, of which 926 had lower and 1474 had higher maximum height than the 90-percentile height value calculated from our plots (Appendix 2). Our dataset thus provided maximum height values for an additional 2914 tree species with no record of this trait in either TRY or BIEN. For consistency, all analyses were done using our 90-percentile height values as a proxy for maximum height. Maximum height values reported in TRY and BIEN are shown in Appendix 2.

2.2 | Wind dispersal syndrome data

We considered species that have fruits or seeds with wings, hairs or other structures that aid flight, as well as minute (<1 mm) seeds released from dehiscent dry fruits, as being wind-dispersed. To determine the fruit or seed types for each species included in our dataset, we used the search string [species name + fruit] in Google images. If no images were found, we scanned through the first 20 hits that




FIGURE 1 Location of 246 one-degree latitude/longitude locations comprising 2821 forest plots with tree height data used in this study. Sites were classified as wet (annual rainfall ≥ 1800 mm, blue dots) and dry (<1800 mm, red dots) forests.

provided textual information on the species to see if there were species descriptions that included fruit or dispersal type. If that did not resolve the issue, we used the search string [genus name + fruit] to determine the dispersal type at the genus level, and assigned that to the species given that dispersal type is a phylogenetically conserved trait that is generally shared among congeners (Chazdon et al., 2003). In cases where the fruit type was a dehiscent dry fruit (e.g. pods, follicles, capsules), we repeated the same search method using the search string [species name + seed], and so on. This procedure allowed us to classify all species in a binary format as either wind-dispersed or not (Appendix 2).

2.3 | Tree species maximum height between biogeographic regions

We used a Kruskal–Wallis test followed by Dunn's post-hoc test to determine whether species maximum tree heights differed between the three main biogeographic regions (Africa, Americas and Asia-Pacific). To avoid potential bias resulting from the higher prevalence of dry forests among the biogeographic regions (Miles et al., 2006), with dry forests tending to have smaller trees than wet forests, we repeated this analysis including only the 10% tallest tree species in each biogeographic region. To map patterns in species maximum height across sample locations, we used the species list of each location in combination with the species' 90-percentile height values to determine the median maximum height for each location in our dataset.

2.4 | Tree species maximum height and wind dispersal

To compare the portion of species overall with that of wind-dispersed species across height classes, we partitioned the tree species of each biogeographic region into 5-m maximum height intervals. To examine the relationship between species maximum height and wind dispersal, we applied generalized linear models with binomial (Bernoulli) distribution and the logit function to conduct logistic regressions of wind dispersal (binary variable) as a function of maximum height (continuous variable) of tree species with a maximum height ≥ 15 m. We accounted for the potential bias due to physiognomic structural differences between dry and wet forests by analysing these two forest types separately in each biogeographic region. Using climatic information from WorldClim version 2.0 (Fick & Hijmans, 2017), we defined dry and wet sites as those with mean annual precipitation lower or higher than 1800 mm, respectively (Figure 1).

2.5 | Emergent tree species and wind dispersal

We defined emergent species as those with at least one individual taller than the 95% height percentile in one or more sample plots for each region-forest type combination (Appendix 3). Pearson's Chi-squared

tests with Yates' continuity correction was used to test whether emergent species had higher proportions of wind dispersal than non-emergent species in wet and dry forests for each biogeographic region.

All statistical analyses were performed with PAST (paleontological statistics software package for education and data analysis) v4.06b (Hammer et al., 2001) and R 4.3.1 (R Core Team, 2023).

3 | RESULTS

3.1 | Maximum tree height differences between biogeographic regions

Maximum height of all tree species was largest in the Asia-Pacific region (median=23.9 m), followed by Africa (median=22.0 m) and the Americas (median=18.0 m) (Kruskal–Wallis test $H=377.8$, $n=5290$, $p<0.0001$) (Figure 2). According to Dunn's post-hoc test, the three biogeographic regions differed significantly from one another in median maximum tree height with Bonferroni corrected $p<0.0018$ for the least significant comparison. Maximum height of the 10% tallest species was largest in Africa (median=40.6 m), followed by the Asia-Pacific (median=38.7 m) and the Americas (median=30.0 m) (Kruskal–Wallis test $H=272.0$, $n=529$, $p<0.0001$) (Figure 2). The three biogeographic regions were again significantly different from one another in median maximum height of the 10% tallest species with Bonferroni corrected $p=0.0165$ for the least significant comparison.

When mapped across locations, the equatorial regions of Africa and the Asia Pacific as well as sites in southern Australia showed higher median tree species maximum heights than forests in the Americas and non equatorial regions (Figure 3).

3.2 | Tree species maximum height and wind dispersal

The percentage of wind-dispersed tree species increased towards higher maximum height classes in all biogeographic regions, despite the declining portion of species in those classes. The percentage of wind-dispersed species also increased slightly towards the smallest maximum height classes, especially in the Americas (Figure 4).

Analysing tree species with a maximum height ≥ 15 m, we found a strong and significant increase in the probability of being wind-dispersed with increasing species maximum height in all three biogeographic regions and in dry and wet forests (Figure 5). This relationship was strongest in the Asia-Pacific region, followed by Africa and the Americas.

3.3 | Emergent tree species and wind dispersal in dry versus wet forests

When tree species were classified as emergent versus non-emergent based on heights measured within each region-forest type




FIGURE 2 Distribution of species maximum heights across the three biogeographical regions for all tree species (left) and the 10% tallest tree species (right). The middle line represents the median and the lower and upper boxes represent the first and third quartiles, respectively. The lighter-coloured areas represent the distribution of maximum tree heights around the median. Number of tree species included in the analysis: Africa, 769; Americas, 1879; and Asia-Pacific, 2642.




FIGURE 3 Median of tree species maximum height (m) based on species lists per location.

combination, emergent species were significantly more likely to be wind-dispersed than non-emergent species in both dry and wet forests across all three biogeographic regions (Figure 6).

4 | DISCUSSION

4.1 | Tree species maximum height among biogeographic regions

Tree species in the Americas had lower maximum heights, on average, than those in Africa and the Asia-Pacific. This difference was especially pronounced when only the 10% tallest tree species in each biogeographic region were compared, with the median height approximately 10m taller in Africa and the Asia-Pacific than in the Americas. This does not appear to be related to the proportion of

wind-dispersed tree species on each continent, which, based on our data, was 10.4% in Africa, 14.0% in the Americas and 17.5% in the Asia-Pacific. It does, however, agree with earlier findings that tropical American forests are generally shorter, show less variation in tree height and contain less tree biomass than their African and Asian-Pacific counterparts (Banin et al., 2012; Dial et al., 2004; Dudley & DeVries, 1990; Slik et al., 2013). Even after accounting for environmental variation between these three regions, Banin et al. (2012) found that the tree height differences persisted, with trees typically shorter in the Americas.

Ascertaining why maximum tree heights are generally lower in the Americas remains an intriguing question. Studies show that canopy level and tree heights are highest in areas with stable climates that have sufficient rainfall to offset transpiration (Banin et al., 2012; Givnish et al., 2014; Gorgens et al., 2020; Larjavaara, 2014; Mao et al., 2020; Marks et al., 2016; Venter et al., 2017), that is areas




FIGURE 4 Comparison of the percentage of tree species in each biogeographic region in 5 m maximum height intervals (orange bars) and the percentage of those species that are wind-dispersed (blue bars). Number of tree species included in the analysis: Africa, 769; Americas, 1879; and Asia-Pacific, 2642.




FIGURE 5 Logistic regressions of the probability of being wind-dispersed as a function of tree species maximum height for the three biogeographic regions in dry and wet forests, for species with a maximum height ≥ 15 m. Asia-Pacific dry: Estimate 0.141, Std. Error 0.017, z-value 8.525, $p < 2e-16$, R^2 0.177; Asia-Pacific wet: Estimate 0.140, Std. Error 0.010, z-value 14.650, $p < 2e-16$, R^2 0.137; Americas dry: Estimate 0.074, Std. Error 0.022, z-value 3.391, $p < 0.0007$, R^2 0.022; Americas wet: Estimate 0.092, Std. Error 0.015, z-value 6.157, $p < 7.42e-10$, R^2 0.044; Africa dry: Estimate 0.112, Std. Error 0.020, z-value 5.698, $p < 1.21e-08$, R^2 0.128; Africa wet: Estimate 0.122, Std. Error 0.020, z-value 6.179, $p < 6.46e-10$, R^2 0.140.

where hydraulic vulnerability is lowest (Olson et al., 2018), as well as fertile, stable soils that are not located on alluvial sites (Gorgens et al., 2020; Jackson et al., 2021; Margrove et al., 2015). Tall forests

also tend to occur in areas where wind speeds and the number of lightning strikes are low and that enjoy at least 130 sunny days per year (de Lima et al., 2023; Gorgens et al., 2020). Differences in




FIGURE 6 Comparison of the number of emergent species among trees that are non-wind and wind-dispersed in dry and wet forests across the Asia-Pacific, Africa and Americas. Emergent species were significantly more likely to be wind-dispersed in both forest types within all three regions (Pearson's Chi-squared test with Yates' continuity correction).

maximum tree heights among biogeographic regions is thus likely related to environmental conditions conducive for growing tall, which suggests that American forests generally lack such favourable conditions relative to their African and Asian-Pacific counterparts. Further research is needed to determine how the relationship between tree species maximum heights and particular environmental conditions has played out at both the continental scale and over evolutionary time scales.

4.2 | Probability of wind dispersal increases with tree species maximum height

While a positive association between tree species' maximum height and wind dispersal has long been observed in closed canopy forests at local scales (Chazdon et al., 2003; Hughes et al., 1994; Slik et al., 2013; Suzuki & Ashton, 1996; Yamada & Suzuki, 1999), we found this association to be widespread in both dry and wet forests throughout three large biogeographic regions that have limited species overlap (Cazzolla Gatti et al., 2022; Slik et al., 2015). This relationship makes sense ecologically as tree height is physically linked to dispersal distance in wind-dispersed trees, even in the absence of wind (Augspurger et al., 2017; Hughes et al., 1994; Smith et al., 2015, 2016). Being emergent aids wind dispersal because these trees are more exposed to wind above the canopy, especially during storms (Heydel et al., 2014). Wind-dispersed seeds can be lifted to great heights during storms and dispersed over distances of several hundred meters or even farther (Heydel et al., 2014). Although such events are infrequent, most trees live for hundreds of years and thus a few such storm events over the reproductive lifespan of a tree might be enough for successful long-distance dispersal and establishment.

Interestingly, we also detected an increase in the proportion of wind-dispersed tree species towards the smallest maximum height classes, especially in the Americas. This may reflect species occurring

in open woodlands and savannah-like habitats where maximum tree height is often limited to 5–15 m and trees are spaced far apart. Wind dispersal is likely to provide a successful strategy under such conditions given that individual trees, despite their relatively short stature, would still be exposed to wind.

4.3 | Emergent tree species are disproportionately wind-dispersed in dry and wet forests

Emergent tree species were significantly more likely to be wind-dispersed than non-emergent species in both dry and wet forests across the three biogeographic regions. Based on the *p*-values from our chi-squared tests, the association between emergent species and wind dispersal was stronger in wet forests than dry forests in the Asia-Pacific and Americas. This is not surprising given that being emergent is probably not as important in dry (especially deciduous) forests where wind is not obstructed by a dense canopy compared to wet forests. Other factors like water stress in tall trees, strong winds, and the relative scarcity of animal frugivores are also likely to affect the strength of this relationship in dry forests (Correa et al., 2015). Nevertheless, the consistency of the association across forest types and regions is probably a result of the fact that height is positively related to dispersal distance in wind-dispersed tree species (Augspurger et al., 2017; Hughes et al., 1994; Smith et al., 2015, 2016; Thomson et al., 2011).

4.4 | Areas of future inquiry

Our results raise several questions for further research. First, they suggest that wind dispersal may be an important factor in the evolution of maximum tree height. The three biogeographic regions have negligible species overlap and also differ considerably in their generic compositions (Cazzolla Gatti et al., 2022; Slik et al., 2015, 2018),

implying largely independent evolutionary histories. Borrowing from work on traits impinging on tree height, such as hydraulic efficiency (Liu et al., 2019) and those related to seed dispersal, like terminal velocity and seed release timing (Wright et al., 2008), phylogenetic comparative analysis could provide a more integrated picture of the evolution of different life-history traits and reproductive associations in trees. A related question is how this association evolved, for example, whether wind dispersal facilitated the evolution of emergence or vice versa. Phylogenetic methods using tests of contingent evolution could trace the evolutionary pathways that led to this association in different wind-dispersed lineages (Larson-Johnson, 2016). Another area of inquiry is how maximum height and wind dispersal covary across finer biogeographic and environmental gradients than those explored here, for example in habitats with different canopy structures. We hope that this study and the large list of tree species maximum heights provided in the supplementary material will stimulate research to address these and other interesting questions on the relationship between wind dispersal and tree maximum height.

AFFILIATIONS

¹Environmental and Life Sciences, Faculty of Science, Universiti Brunei Darussalam, Gadong, Brunei Darussalam

²Institute of Plant Sciences, University of Bern, Bern, Switzerland

³AMAP, Université Montpellier, INRAe, CIRAD, CNRS, IRD, Montpellier, France

⁴Laboratorio de Ecología Tropical y Servicios Ecosistémicos (EcoSs Lab), Departamento de Ciencias Biológicas y Agropecuarias, Universidad Técnica Particular de Loja, Loja, Ecuador

⁵Department of Forest Sciences, Viikki Tropical Resources Institute, University of Helsinki, Helsinki, Finland

⁶Institute for Sustainability, University of Helsinki, Helsinki, Finland

⁷Institute of Environment and Sustainable Development, Banaras Hindu University, Varanasi, India

⁸Department of Wetland Ecology, Institute of Geography and Geoecology, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

⁹Ecologia, Monitoramento e Uso Sustentável de Áreas Úmidas (MAUA), Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil

¹⁰Department of Geography and Environment, Wentworth Way, University of York, Heslington, UK

¹¹Department of International Environmental and Development Studies (NORAGRIC), Norwegian University of Life Sciences, Ås, Norway

¹²Biodiversity Parks Programme, CEMDE, University of Delhi, Delhi, India

¹³Departamento de Ecologia, CB, Universidade Federal do Rio Grande do Norte (UFRN), Natal, Brazil

¹⁴Geneva Conservatory and Botanical Garden, Chemin de l'Impératrice 1, Chambésy, Switzerland

¹⁵Department of Plant Systematics, University of Bayreuth, Bayreuth, Germany

¹⁶School of Biological Sciences, University of Adelaide, Adelaide, South Australia, Australia

¹⁷Forest Research Institute, University of the Sunshine Coast, Sippy Downs, Queensland, Australia

¹⁸Department of Environment and Geography, University of York, York, UK

¹⁹Flamingo Land Ltd., Kirby Misperton, UK

²⁰Rainforest Traditions, Kochi, India

²¹Department of Ecoscience, Aarhus University, Aarhus, Denmark

²²Yayasan Borneo Nature Indonesia, Palangka Raya, Indonesia

²³School of Biological Sciences, Universiti Sains Malaysia, Penang, Malaysia

²⁴Centre for Marine and Coastal Studies, Universiti Sains Malaysia, Penang, Malaysia

²⁵Laboratoire de Recherche Forestière, Faculté Des Sciences, Université de Lomé-Togo, Lomé, Togo

²⁶Royal Botanical Garden Edinburgh, Edinburgh, UK

²⁷TERN, The University of Adelaide, Adelaide, South Australia, Australia

²⁸Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, UK

²⁹Borneo Nature Foundation International, Penryn, UK

³⁰Physical Geography with a Focus on Human-Environment Research, University of Passau, Passau, Germany

³¹Novel Research Academy, Puducherry, India

³²International Institute for Sustainability / IIS, Rio de Janeiro, Brazil

³³Department of Environmental Sciences, Wageningen University and Research, Wageningen, The Netherlands

³⁴Center for International Forestry Research (CIFOR), Kota Bogor, Indonesia

³⁵Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad de México, Mexico

³⁶Instituto de Botânica de São Paulo, São Paulo, Brazil

³⁷Departamento de Ecologia e Conservação, Universidade Federal de Lavras, Lavras, Brazil

³⁸InfoFlora, Geneva Botanical Garden, Geneva, Switzerland

³⁹Biodiversity, Macroecology and Biogeography, Georg-August-University Göttingen, Göttingen, Germany

⁴⁰Instituto Federal Do Espírito Santo, Espírito Santo, Brazil

⁴¹Instituto Nacional da Mata Atlântica, Espírito Santo, Brazil

⁴²Department of Biological Sciences, Faculty of Sciences, Botswana University of Agriculture and Natural Resources, Sebele, Botswana

⁴³Department of Floodplain Ecology, Institute of Geography and Geoecology, Karlsruhe Institute of Technology, Rastatt, Germany

⁴⁴Department of Biology, University of Florence, Trento, Italy

⁴⁵Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Morelia, Mexico

⁴⁶Pro Natura Foundation, Balikpapan, Indonesia

⁴⁷Biology Department, Université Officielle de Bukavu, Bukavu, Democratic Republic of the Congo

⁴⁸UNELLEZ-Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT), Mesa de Cavacas, Venezuela

⁴⁹Compensation International Progress S.A.-Ciprogress Greenlife, Bogotá D.C., Colombia

⁵⁰Laboratório de Ecologia e Hidrologia Florestal, Floresta Estadual de Assis, Instituto de Pesquisas Ambientais, Assis, Brazil

⁵¹Department of Forestry, North Eastern Hill University, Tura, India

⁵²Functional Ecology, Instituto de Ecología, Xalapa, Mexico

⁵³DBU Natural Heritage, German Federal Foundation for the Environment, Osnabrück, Germany

⁵⁴Forestry Department, Faculty of Agriculture, University of Palangka Raya, Palangka Raya, Indonesia

⁵⁵Real Jardín Botánico - CSIC, Madrid, Spain

⁵⁶Ecosystems Analysis Laboratory, Department of Botany, Banaras Hindu University, Varanasi, India

⁵⁷Smith College Botanic Garden, Northampton, Massachusetts, USA

⁵⁸Bureau Waardenburg, Culemborg, The Netherlands

⁵⁹Gembloux Agro Bio-Tech, University of Liege, Gembloux, Belgium

⁶⁰The RSPB, Sandy, Bedfordshire, UK

⁶¹A Rocha International, Cambridge, UK

⁶²Agence française pour la biodiversité, Vincennes, France

⁶³Remote Sensing Department (DSR), National Institute for Space Research (INPE), São Paulo, Brazil

⁶⁴Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA

⁶⁵Tropical Forests and People Research Centre, University of the Sunshine Coast, Maroochydore, Queensland, Australia

⁶⁶School of Geography, University of Leeds, Leeds, UK

⁶⁷Royal Botanic Gardens, Kew, Surrey, UK

⁶⁸Departamento de Engenharia Florestal, Universidade de Brasília, Brasília, Brazil

⁶⁹Instituto Boliviano de Investigación Forestal (IBIF), Santa Cruz de la Sierra, Bolivia

⁷⁰Department of Biology, University of Turku, Turku, Finland

⁷¹Department of Biology, Aarhus University, Aarhus, Denmark

⁷²Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia (UPM), Seri Kembangan, Malaysia

- ⁷³Faculty of Bioresources and Food Sciences, Universiti Sultan Zainal Abidin, Besut, Malaysia
- ⁷⁴Forest Ecology and Forest Management Group, Wageningen University & Research, Wageningen, The Netherlands
- ⁷⁵São Paulo Agribusiness Technology Agency, APTA Regional de Pindorama, São Paulo, Brazil
- ⁷⁶School of Life Sciences/RCUH, Department of Natural Resources and Environmental Management, University of Hawai'i at Mānoa, Mānoa, USA
- ⁷⁷Colégio de Aplicação (CA), Centro de Ciências da Educação (CED), Universidade Federal de Santa Catarina (UFSC), Florianópolis, Brazil
- ⁷⁸Technical Management Unit of Mungku Baru Forest, Universitas Muhammadiyah Palangkaraya, Palangkaraya, Indonesia
- ⁷⁹Department of Earth and Environmental Science, Faculty of Environment, Science and Economy, University of Exeter, Penryn, UK
- ⁸⁰School of Geography, Geology and the Environment, University of Leicester, Leicester, UK
- ⁸¹H.J. Andrews Experimental Forest, Blue River, Oregon, USA
- ⁸²Systematic and Evolutionary Botany, University of Zurich, Zurich, Switzerland
- ⁸³El Colegio de la Frontera Sur, San Cristóbal de las Casas, Mexico
- ⁸⁴Department of Plant Biology, Faculty of Science, University of Yaoundé I, Yaoundé, Cameroon
- ⁸⁵Laboratory of Plant Systematics and Ecology, Higher Teachers' Training College, University of Yaoundé I, Yaoundé, Cameroon
- ⁸⁶Faculty of Forestry, Universitas Gadjah Mada, Yogyakarta, Indonesia
- ⁸⁷Department of Ecology and Environmental Sciences, Pondicherry University, Puducherry, India
- ⁸⁸WWF Thailand, Bangkok, Thailand
- ⁸⁹Science and Education, Field Museum of Natural History, Chicago, Illinois, USA
- ⁹⁰Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia
- ⁹¹Department of Ecosystems and Conservation, Sokoine University of Agriculture, Morogoro, Tanzania
- ⁹²Bullard Professor of Forestry Emeritus, Harvard University, Cambridge, USA
- ⁹³Max Planck Institute of Animal Behavior, Radolfzell, Germany
- ⁹⁴Department of Geography, School of Environment Education and Development, University of Manchester, Manchester, UK
- ⁹⁵Sigur Nature Trust, Nilgiris, India
- ⁹⁶Institute for Biodiversity and Environmental Research, Universiti Brunei Darussalam, Gadong, Brunei Darussalam
- ⁹⁷Centre for Global Sustainability Studies, Universiti Sains Malaysia, Penang, Malaysia
- ⁹⁸International Institute of Information Technology, Hyderabad, India
- ⁹⁹Center for Integrative Conservation Xishuangbanna Tropical Botanical Garden Chinese Academy of Sciences, Mengla, People's Republic of China
- ¹⁰⁰World Wildlife Fund Thailand, Bangkok, Thailand
- ¹⁰¹Wageningen Environmental Research, Wageningen University & Research, Wageningen, The Netherlands
- ¹⁰²Institute of Biological Sciences, University of Malaya, Kuala Lumpur, Malaysia
- ¹⁰³Department of Plant and Microbial Biology, College of Biological Sciences, University of Minnesota – Twin Cities, St. Paul, Minnesota, USA
- ¹⁰⁴Centre of Biodiversity, Vietnam National University of Forestry, Xuan Mai, Vietnam
- ¹⁰⁵Laboratory of Dendrology and Tropical Silviculture, Agroforestry Science Training Center, Federal University of Southern Bahia, Ilhéus, Brazil
- ¹⁰⁶Tropical Forests and People Research Centre, University of the Sunshine Coast, Maroochydore DC, Queensland, Australia
- ¹⁰⁷Department of Environment and Development Studies, United International University, Dhaka, Bangladesh
- ¹⁰⁸Faculty of Forestry, Instiper Yogyakarta, Yogyakarta, Indonesia
- ¹⁰⁹Department of Forestry & Environmental Science, Shahjalal University of Science & Technology, Sylhet, Bangladesh
- ¹¹⁰Faculty of Forestry, University of British Columbia, Vancouver, British Columbia, Canada
- ¹¹¹Centre for International Forestry Research (CIFOR), Bogor, Indonesia
- ¹¹²Department of Geography, University of California, Los Angeles,

- California, USA
- ¹¹³Naturalis, Leiden, The Netherlands
- ¹¹⁴Silviculture Research Institute, Vietnamese Academy of Forest Sciences, Hanoi, Vietnam
- ¹¹⁵Conservation Ecology Program, Pilot Plant Development and Training Institute, King Mongkut's University of Technology Thonburi, Bangkok, Thailand
- ¹¹⁶Institute of Ecology and Nature Conservation, Chinese Academy of Forestry, Beijing, China
- ¹¹⁷College of Ecology and Environment, Hainan University, Haikou City, China
- ¹¹⁸College of Agronomy and Agricultural Engineering, Liaocheng University, Liaocheng, China
- ¹¹⁹Center for International Forestry Research (CIFOR), Bogor, Indonesia
- ¹²⁰Department of Natural Resources and Watershed Management, Shabestar, East Azerbaijan, Iran

ACKNOWLEDGEMENTS

We like to thank the following collaborators for providing tree plot data for this study: Kazuki Miyamoto, Andrea Permana, Andres Avella, Eddie Lenza de Oliveira, Emanuel Martin, Faustin Boyemba, Girirai, Hansjoerg Woll, Indiana Coronado, Jianwei Tang, John Katembo, Kipiro Damas, Miriam van Heist, Navendu Page, Nicolas Barbier, Timothy J. Killeen, Timothy Whitfeld and Victor Adekunle. We also thank Johannes Jamaludin for helping with the preparation of some of the figures.

FUNDING INFORMATION

Vegetation plots in Tanzania were established under funding from the Leverhulme Trust under the Valuing the Arc project. Marshall was funded by ARC Future Fellowship FT170100279. Plots in southern Brazil (Rio Grande do Sul) were established under funding from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, grant 520053/1998-2), and AFS thanks Solon J. Longhi for kindly agreeing to share this data. BJWB, HS, M, MAI and SM thank The Orangutan Project, Arcus Foundation, and US Fish and Wildlife Service Great Apes Conservation Fund for financial support and BRIN (formerly RISTEK) and KPH for permissions and support. Moses Libalah was funded by the IFS grant number I-1-D-5621-2. JRRP thank CNPq for productivity grant support (Process 312571/2021-6).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

All data used in this study are provided in [Appendices 1–3](#).

ORCID

- Ferry Slik  <https://orcid.org/0000-0003-3988-7019>
- Bruno X. Pinho  <https://orcid.org/0000-0002-6588-3575>
- Andreas Hemp  <https://orcid.org/0000-0002-5369-2122>
- Fabian Brambach  <https://orcid.org/0000-0002-3203-2248>
- Lourens Poorter  <https://orcid.org/0000-0003-1391-4875>
- Robert Steinmetz  <https://orcid.org/0000-0001-6860-8715>
- Tran Van Do  <https://orcid.org/0000-0001-9059-5842>
- Runguo Zang  <https://orcid.org/0000-0001-5392-8193>

REFERENCES

- Augsburger, C. K., Franson, S. E., & Cushman, K. C. (2017). Wind dispersal is predicted by tree, not diaspore, traits in comparisons of Neotropical species. *Functional Ecology*, *31*, 808–820.
- Banin, L., Feldpausch, T. R., Phillips, O. L., Baker, T. R., Lloyd, J., Affum-Baffoe, K., Arets, E. J. M. M., Berry, N. J., Bradford, M., Brienen, R. J. W., Davies, S., Drescher, M., Higuchi, N., Hilbert, D. W., Hladik, A., Iida, Y., Salim, K. A., Kassim, A. R., King, D. A., ... Lewis, S. L. (2012). What controls tropical forest architecture? Testing environmental, structural and floristic drivers. *Global Ecology and Biogeography*, *21*, 1179–1190.
- Cazzolla Gatti, R., Reich, P. B., Gamarra, J. G. P., Crowther, T., Hui, C., Morera, A., Bastin, J. F., de-Miguel, S., Nabuurs, G. J., Svenning, J. C., Serra-Diaz, J. M., Merow, C., Enquist, B., Kamenetsky, M., Lee, J., Zhu, J., Fang, J., Jacobs, D. F., Pijanowski, B., ... Liang, J. (2022). The number of tree species on Earth. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(6), e2115329119.
- Chazdon, R. L., Careaga, S., Webb, C., & Vargas, O. (2003). Community and phylogenetic structure of reproductive traits of woody species in wet tropical forests. *Ecological Monographs*, *73*(3), 331–348.
- Comita, L. S., Queenborough, S. A., Murphy, S. J., Eck, J. L., Xu, K., Krishnadas, M., Beckman, N., Zhu, Y., & Gómez-Aparicio, L. (2014). Testing predictions of the Janzen–Connell hypothesis: a meta-analysis of experimental evidence for distance- and density-dependent seed and seedling survival. *Journal of Ecology*, *102*(4), 845–856.
- Correa, D. F., Alvarez, E., & Stevenson, P. R. (2015). Plant dispersal systems in Neotropical forests: Availability of dispersal agents or availability of resources for constructing zoochorous fruits? *Global Ecology and Biogeography*, *24*(2), 203–214.
- D'Andrea, R., Guittar, J., O'Dwyer, J. P., Figueroa, H., Wright, S. J., Condit, R., & Ostling, A. (2020). Counting niches: Abundance by trait patterns reveal niche partitioning in a Neotropical forest. *Ecology*, *101*, e03019.
- D'Andrea, R., Riolo, M., & Ostling, A. M. (2019). Generalizing clusters of similar species as a signature of coexistence under competition. *PLoS Computational Biology*, *15*, 1–19.
- de Lima, R. B., Görgens, E. B., da Silva, D. A. S., de Oliveira, C. P., Batista, A. P. B., Ferreira, R. L. C., Costa, F. R., Ferreira de Lima, R. A., da Silva Aparício, P., de Abreu, J. C., da Silva, J. A. A., & Phillips, O. L. (2023). Giants of the Amazon: How does environmental variation drive the diversity patterns of large trees? *Global Change Biology*, *17*, 4861–4879.
- de Santana, R. A. S., Dias-Junior, C. Q., do Vale, R. S., Tota, J., & Fitzjarrald, D. R. (2017). Observing and modeling the vertical wind profile at multiple sites in and above the Amazon rain forest canopy. *Advances in Meteorology*, *2017*, 5436157.
- Dial, R., Bloodworth, B., Lee, A., Boyne, P., & Heys, J. (2004). The distribution of free space and its relation to canopy composition at six forest sites. *Forest Science*, *50*(3), 312–325.
- Dudley, R., & DeVries, P. (1990). Tropical rain forest structure and the geographical distribution of gliding vertebrates. *Biotropica*, *22*, 432–434.
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, *37*, 4302–4315.
- Givnish, T. J., Wong, S. C., Stuart-Williams, H., Holloway-Phillips, M., & Farquhar, G. D. (2014). Determinants of maximum tree height in Eucalyptus species along a rainfall gradient in Victoria, Australia. *Ecology*, *95*(11), 2991–3007.
- Gorgens, E., Nunes, M. H., Jackson, T., Coomes, D., Keller, M., Reis, C. R., Valbuena, R., Rosette, J., de Almeida, D. R. A., Gimenez, B., Cantinho, R., Motta, A. Z., Assis, M., de Souza Pereira, F. R., Spanner, G., Higuchi, N., & Ometto, J. P. (2020). Resource availability and disturbance shape maximum tree height across the Amazon. *Global Change Biology*, *27*, 177–189.
- Hammer, O., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for educational and data analysis. *Palaeontologia Electronica*, *4*, 1–8.
- Heydel, F., Cunze, S., Bernhardt-Romermann, M., & Tackenberg, O. (2014). Long-distance seed dispersal by wind: Disentangling the effects of species traits, vegetation types, vertical turbulence and wind speed. *Ecological Research*, *29*, 641–651.
- Hughes, L., Dunlop, M., French, K., Leishman, M. R., Rice, B., Rodgeron, L., & Westoby, M. (1994). Predicting dispersal spectra: A minimal set of hypotheses based on plant attributes. *Journal of Ecology*, *82*, 933–950.
- Iida, Y., Poorter, L., Sterck, F., Kassim, A. R., Potts, M. D., Kubo, T., & Kohyama, T. S. (2014). Linking size dependent growth and mortality with architectural traits across 145 co-occurring tropical tree species. *Ecology*, *95*(2), 353–363.
- Jackson, T. D., Shenkin, A. F., Majalap, N., Bin Jami, J., Bin Sailim, A., Reynolds, G., Coomes, D. A., Chandler, C. J., Boyd, D. S., Burt, A., Wilkes, P., Disney, M., & Malhi, Y. (2021). The mechanical stability of the world's tallest broadleaf trees. *Biotropica*, *53*, 110–120.
- Kattge, J., Bönsch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner, G. D. A., Aakala, T., Abedi, M., Acosta, A. T. R., Adamidis, G. C., Adamson, K., Aiba, M., Albert, C. H., Alcántara, J. M., Alcázar, C. C., Aleixo, I., Ali, H., ... Wirth, C. (2020). TRY plant trait database—Enhanced coverage and open access. *Global Change Biology*, *26*, 119–188.
- Kohyama, T. (1993). Size-structured tree populations in gap-dynamic forest—The forest architecture hypothesis for the stable coexistence of species. *Journal of Ecology*, *81*, 131–143.
- Kohyama, T., Suzuki, E., Partomihardjo, T., Yamada, T., & Kubo, T. (2003). Tree species differentiation in growth, recruitment and allometry in relation to maximum height in a Bornean mixed dipterocarp forest. *Journal of Ecology*, *91*, 797–806.
- Larjavaara, M. (2014). The world's tallest trees grow in thermally similar climates. *New Phytologist*, *202*, 344–349.
- Larson-Johnson, K. (2016). Phylogenetic investigation of the complex evolutionary history of dispersal mode and diversification rates across living and fossil Fagales. *New Phytologist*, *209*, 418–435.
- Liu, H., Gleason, S. M., Hao, G., Hua, L., He, P., Goldstein, G., & Ye, Q. (2019). Hydraulic traits are coordinated with maximum plant height at the global scale. *Science Advances*, *5*, eaav1332.
- Maitner, B. S., Boyle, B., Casler, N., Condit, R., Donoghue, J., II, Durán, S. M., Guaderrama, D., Hinchliff, C. E., Jørgensen, P. M., Kraft, N. J. B., McGill, B., Merow, C., Morueta-Holme, N., Peet, R. K., Sandel, B., Schildhauer, M., Smith, S. A., Svenning, J. C., Thiers, B., ... Enquist, B. J. (2018). The BIEN R package: A tool to access the Botanical Information and Ecology Network (BIEN) database. *Methods in Ecology and Evolution*, *9*, 373–379.
- Mao, L., Swenson, N. G., Sui, X., Zhang, J., Chen, S., Li, J., Peng, P., Zhou, G., & Zhang, X. (2020). The geographic and climatic distribution of plant height diversity for 19,000 angiosperms in China. *Biodiversity and Conservation*, *29*, 487–502.
- Margrove, J. A., Burslem, D. F. R. P., Ghazoul, J., Khoo, E., Kettle, C. J., & Maycock, C. R. (2015). Impacts of an extreme precipitation event on dipterocarp mortality and habitat filtering in a Bornean tropical rain forest. *Biotropica*, *47*(1), 66–76.
- Marks, C. O., Muller-Landau, H. C., & Tilman, D. (2016). Tree diversity, tree height and environmental harshness in eastern and western North America. *Ecology Letters*, *19*(7), 743–751.
- Miles, L., Newton, A. C., DeFries, R. S., Ravilious, C., May, I., Blyth, S., Kapos, V., & Gordon, J. E. (2006). A global overview of the conservation status of tropical dry forests. *Journal of Biogeography*, *33*(3), 491–505.
- Nathan, R., Katul, G. G., Horn, H. S., Thomas, S. M., Oren, R., Avissar, R., Pacala, S. W., & Levin, S. A. (2002). Mechanisms of long-distance dispersal of seeds by wind. *Nature*, *418*, 409–413.

- Olson, M. E., Soriano, D., Rosell, J. A., Anfodillo, T., Donoghue, M. J., Edwards, E. J., León-Gómez, C., Dawson, T., Camarero Martínez, J. J., Castorena, M., Echeverría, A., Espinosa, C. I., Fajardo, A., Gazol, A., Isnard, S., Lima, R. S., Marcati, C. R., & Méndez-Alonzo, R. (2018). Plant height and hydraulic vulnerability to drought and cold. *Proceedings of the National Academy of Sciences of the United States of America*, 115(29), 7551–7556.
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Rüger, N., Condit, R., Dent, D. H., DeWalt, S. J., Hubbell, S. P., Lichstein, J. W., Lopez, O. R., Wirth, C., & Fariori, C. E. (2020). Demographic tradeoffs predict tropical forest dynamics. *Science*, 368(6487), 165–168.
- Scheffer, M., & van Nes, E. H. (2006). Self-organized similarity, the evolutionary emergence of groups of similar species. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 6230–6235.
- Slik, J. W. F., Arroyo-Rodríguez, V., Aiba, S., Alvarez-Loayza, P., Alves, L. F., Ashton, P., Balvanera, P., Bastian, M. L., Bellingham, P. J., van den Berg, E., Bernacci, L., da Conceição Bispo, P., Blanc, L., Böhning-Gaese, K., Boeckx, P., Bongers, F., Boyle, B., Bradford, M., Brearley, F. Q., ... Venticinque, E. M. (2015). An estimate of the number of tropical tree species. *Proceedings of the National Academy of Sciences of the United States of America*, 112, 7472–7477.
- Slik, J. W. F., Franklin, J., Arroyo-Rodríguez, V., Field, R., Aguilar, S., Aguirre, N., Ahumada, J., Aiba, S. I., Alves, L. F., Anitha, K., Avella, A., Mora, F., Aymard C., G. A., Báez, S., Balvanera, P., Bastian, M. L., Bastin, J. F., Bellingham, P. J., van den Berg, E., ... Zang, R. (2018). Phylogenetic classification of the world's tropical forests. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 1837–1842.
- Slik, J. W. F., Paoli, G., McGuire, K., Amaral, I., Barroso, J., Bastian, M., Blanc, L., Bongers, F., Boundja, P., Clark, C., Collins, M., Dauby, G., Ding, Y., Doucet, J. L., Eler, E., Ferreira, L., Forshed, O., Fredriksson, G., Gillet, J. F., ... Zweifel, N. (2013). Large trees drive forest aboveground biomass variation in moist lowland forests across the tropics. *Global Ecology and Biogeography*, 22, 1261–1271.
- Smith, J. R., Bagchi, R., Ellens, J., Kettle, C. J., Burslem, D. F. R. P., Maycock, C. R., Khoo, E., & Ghazoul, J. (2015). Predicting dispersal of auto-gyrating fruit in tropical trees: A case study from the Dipterocarpaceae. *Ecology and Evolution*, 5, 1794–1801.
- Smith, J. R., Bagchi, R., Kettle, C. J., Maycock, C., Khoo, E., & Ghazoul, J. (2016). Predicting the terminal velocity of dipterocarp fruit. *Biotropica*, 48(2), 154–158.
- Souza, A. F. (2021). A review of the structure and dynamics of araucaria mixed forests in southern Brazil and northern Argentina. *New Zealand Journal of Botany*, 59(1), 2–54.
- Suzuki, E., & Ashton, P. S. (1996). Sepal and nut size ratio of fruits of Asian Dipterocarpaceae and its implications for dispersal. *Journal of Tropical Ecology*, 12, 853–870.
- Thomson, F. J., Moles, A. T., Auld, T. D., & Kingsford, R. T. (2011). Seed dispersal distance is more strongly correlated with plant height than with seed mass. *Journal of Ecology*, 99, 1299–1307.
- Venter, M., Dwyer, J., Dieleman, W., Ramachandra, A., Gillieson, D., Laurance, S., Cernusak, L. A., Beehler, B., Jensen, R., & Bird, M. I. (2017). Optimal climate for large trees at high elevations drives patterns of biomass in remote forests of Papua New Guinea. *Global Change Biology*, 23, 4873–4883.
- Wright, S. J., Trakhtenbrot, A., Bohrer, G., Detto, M., Katul, G. G., Horvitz, N., Muller-Landau, H. C., Jones, F. A., & Nathan, R. (2008). Understanding strategies for seed dispersal by wind under contrasting atmospheric conditions. *Proc Natl Acad Sci USA*, 105(49), 19084–19089.
- Wu, Z.-Y., Milne, R. I., Liu, J., Nathan, R., Corlett, R. T., & Li, D. Z. (2023). The establishment of plants following long-distance dispersal. *Trends in Ecology & Evolution*, 38(3), 289–300.
- Yamada, T., & Suzuki, E. (1999). Comparative morphology and allometry of winged diaspores among the Asian Sterculiaceae. *Journal of Tropical Ecology*, 15, 619–635.

BIOSKETCH

Ferry Slik is a professor at the Environmental and Life Sciences Department at Universiti Brunei Darussalam. His research focusses on tropical tree ecology and conservation. He is also the curator of the university herbarium (UBDH) and works on the tree checklist and flora of Brunei Darussalam.

Bruno X. Pinho is currently a postdoc at the Institute of Plant Sciences in the University of Bern. He is interested in elucidating the causes and consequences of biodiversity changes across scales in human-modified landscapes.

Daniel M. Griffith is an associate professor in the Department of Biological and Agricultural Sciences at the Universidad Técnica Particular de Loja, Ecuador. His research focuses on conservation biology, tropical forest restoration, and the sustainable management of forest and agricultural landscapes in the Neotropics.

Edward Webb is a tropical ecologist specializing in vegetation ecology, tropical tree community dynamics, conservation and forest management. His geographic areas of expertise are Pacific tropical islands, and South and Southeast Asian forests. He is the Director of the Viikki Tropical Resources Institute in the Department of Forest Sciences at the University of Helsinki.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Slik, F., Pinho, B. X., Griffith, D. M., Webb, E., Raghubanshi, A. S., Quaresma, A. C., Cuni Sanchez, A., Sultana, A., Souza, A. F., Ensslin, A., Hemp, A., Lowe, A., Marshall, A. R., Anitha, K., Lykke, A. M., Armadyanto, Mansor, A., Honam, A. K., Poulsen, A. D., ... Hemati, Z. (2024). Wind dispersed tree species have greater maximum height. *Global Ecology and Biogeography*, 33, e13878. <https://doi.org/10.1111/geb.13878>