


CORRECTION

Open Access



Correction: H₂-driven xylitol production in *Cupriavidus necator* H16

Tytti Jämsä^{1*} , Nico J. Claassens² , Laura Salusjärvi¹  and Antti Nyssölä¹ 

Correction: *Microbial Cell Factories* (2024) 23:345

<https://doi.org/10.1186/s12934-024-02615-7>

In the methods section, the phrase “flushing with 100% H₂ at 0.5 mL/min for 2 min.” should read “flushing with 100% H₂ at 0.5 L/min for 2 min.”

Published online: 19 January 2025

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1186/s12934-024-02615-7>.

*Correspondence:

Tytti Jämsä

tytti.jamsa@vtt.fi

¹VTT Technical Research Centre of Finland Ltd., Tekniikantie 21,
02150 Espoo, Finland

²Laboratory of Microbiology, Wageningen University, Stippeneng 4,
6708WE Wageningen, The Netherlands



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.